

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-Benzyl-N-cyclohexylbenzene-sulfonamideZeeshan Haider,^a Islam Ullah Khan,^{a*} Muhammad Zia-ur-Rehman^b and Muhammad Nadeem Arshad^a^aDepartment of Chemistry, Government College University, Lahore 54000, Pakistan, and ^bApplied Chemistry Research Centre, PCSIR Laboratories Complex, Ferozpur Road, Lahore 54600, Pakistan

Correspondence e-mail: iukhan.gcu@gmail.com


Received 4 November 2009; accepted 6 November 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.037; wR factor = 0.094; data-to-parameter ratio = 20.7.

In the title compound, $\text{C}_{19}\text{H}_{23}\text{NO}_2\text{S}$, the cyclohexyl ring exists in a chair form. The dihedral angle between the two terminal phenyl rings is $86.70(6)^\circ$. No significant interactions are observed except for a weak intramolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bond.

Related literature

For the biological activity of sulfonamides, see: Innocenti *et al.* (2004); Ozbek *et al.* (2007); Parari *et al.* (2008); Ratish *et al.* (2009); Selnam *et al.* (2001); For related structures, see: Khan *et al.* (2009); Zia-ur-Rehman *et al.* (2009) Gowda *et al.* (2007*a,b,c*). For bond length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{19}\text{H}_{23}\text{NO}_2\text{S}$
 $M_r = 329.44$
 Orthorhombic, $P2_12_12_1$
 $a = 9.1996(5)$ Å
 $b = 11.0406(5)$ Å
 $c = 17.1897(9)$ Å
 $V = 1745.94(15)$ Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.20$ mm⁻¹
 $T = 296$ K
 $0.39 \times 0.34 \times 0.28$ mm

Data collection

Bruker APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.928$, $T_{\max} = 0.948$
 10897 measured reflections
 4306 independent reflections
 3507 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.094$
 $S = 1.03$
 4306 reflections
 208 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.24$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.24$ e Å⁻³
 Absolute structure: Flack (1983),
 1853 Friedel pairs
 Flack parameter: 0.03 (7)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C7}-\text{H7}\cdots\text{O2}$	0.98	2.34	2.874 (2)	113

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: PLATON (Spek, 2009) and Mercury (Macrae *et al.*, 2006); software used to prepare material for publication: SHELXTL (Sheldrick, 2008) and local programs.

The authors are grateful to the Higher Education Commission of Pakistan for the financial support to purchase the diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2484).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Gowda, B. T., Foro, S. & Fuess, H. (2007*a*). *Acta Cryst.* **E63**, o2339.
- Gowda, B. T., Foro, S. & Fuess, H. (2007*b*). *Acta Cryst.* **E63**, o2570.
- Gowda, B. T., Foro, S. & Fuess, H. (2007*c*). *Acta Cryst.* **E63**, o2597.
- Innocenti, A., Antel, J., Wurl, M., Scozzafava, A. & Supuran, C. T. (2004). *Bioorg. Med. Chem. Lett.* **14**, 5703–5707.
- Khan, I. U., Haider, Z., Zia-ur-Rehman, M., Arshad, M. N. & Shafiq, M. (2009). *Acta Cryst.* **E65**, o2867.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Ozbek, N., Katircioğ lu, H., Karacan, N. & Baykal, T. (2007). *Bioorg. Med. Chem.* **15**, 5105–5109.
- Parari, M. K., Panda, G., Srivastava, K. & Puri, S. K. (2008). *Bioorg. Med. Chem. Lett.* **18**, 776–781.
- Ratish, G. I., Javed, K., Ahmad, S., Bano, S., Alam, M. S., Pillai, K. K., Singh, S. & Bagchi, V. (2009). *Bioorg. Med. Chem. Lett.* **19**, 255–258.
- Selnam, P., Chandramohan, M., Clercq, E. D., Witvrouw, M. & Pannecouque, C. (2001). *Eur. J. Pharm. Sci.* **14**, 313–316.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Zia-ur-Rehman, M., Choudary, J. A., Elsegood, M. R. J., Siddiqui, H. L. & Khan, K. M. (2009). *Eur. J. Med. Chem.* **44**, 1311–1316.

supplementary materials

Acta Cryst. (2009). E65, o3053 [doi:10.1107/S1600536809046960]

N-Benzyl-*N*-cyclohexylbenzenesulfonamide

Z. Haider, I. U. Khan, M. Zia-ur-Rehman and M. N. Arshad

Comment

Sulfonamides are familiar as biologically active compounds and possess anti-microbial (Ozbek *et al.*, 2007; Parari *et al.*, 2008), anti-inflammatory (Ratish *et al.*, 2009), anti HIV (Selnam *et al.*, 2001) and carbonic anhydrase inhibition activity (Innocenti *et al.*, 2004). In the present paper, the structure of *N*-cyclohexyl-*N*-propyl benzene sulfonamide has been determined as part of a research program involving the synthesis of various sulfur containing heterocycles (Zia-ur-Rehman *et al.*, 2009; Khan *et al.*, 2009).

In the molecule of (**I**) (Fig. 1), bond lengths and bond angles are almost similar to those in the related molecules (Gowda *et al.*, 2007*a,b,c*) and are within normal ranges (Allen *et al.*, 1987). The benzene rings are essentially planar while cyclohexane ring is in the chair form. The dihedral angles between the phenyl (C1–C6) & benzyl ring (C14–C19), the phenyl (C1–C6) ring & the mean plane of cyclohexyl ring (C7–C12), and the benzyl (C14–C19) ring & the mean plane cyclohexyl ring (C7–C12) are 86.70 (6), 42.43 (8) and 55.47 (8)°, respectively, while the r.m.s. deviation for the phenyl (C1–C6), benzyl (C14–C19) and cyclohexyl (C7–C12) rings are 0.0021, 0.0036 and 0.2365 Å, respectively. An intramolecular C—H···O hydrogen bond gives rise to a five-membered hydrogen bonded ring (Table 1).

Experimental

A mixture of *N*-cyclohexylbenzene sulfonamide (1 g, 0.43 mmol), sodium hydride (0.21 g, 0.88 mmol) and *N,N*-dimethylformamide (10 ml) was stirred at room temperature for half an hour followed by addition of benzyl chloride (0.114 g, 0.90 mmol). Stirring was continued further for a period of three hours and the contents were poured over crushed ice. Precipitated product was isolated, washed and crystallized from methanol-water mixture (50:50).

Refinement

All H atoms were identified in a difference map and then were treated as riding (C—H = 0.93 or 0.97 Å), with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of (**I**), with displacement ellipsoids at the 50% probability level.

N-Benzyl-*N*-cyclohexylbenzenesulfonamide

Crystal data

$C_{19}H_{23}NO_2S$	$F_{000} = 704$
$M_r = 329.44$	$D_x = 1.253 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: P 2ac 2ab	Cell parameters from 3979 reflections
$a = 9.1996 (5) \text{ \AA}$	$\theta = 2.3\text{--}21.8^\circ$
$b = 11.0406 (5) \text{ \AA}$	$\mu = 0.20 \text{ mm}^{-1}$
$c = 17.1897 (9) \text{ \AA}$	$T = 296 \text{ K}$
$V = 1745.94 (15) \text{ \AA}^3$	Blocks, colourless
$Z = 4$	$0.39 \times 0.34 \times 0.28 \text{ mm}$

Data collection

Bruker APEXII CCD area-detector diffractometer	4306 independent reflections
Radiation source: fine-focus sealed tube	3507 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.023$
$T = 296 \text{ K}$	$\theta_{\text{max}} = 28.3^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.2^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -12 \rightarrow 11$
$T_{\text{min}} = 0.928$, $T_{\text{max}} = 0.948$	$k = -11 \rightarrow 14$
10897 measured reflections	$l = -22 \rightarrow 17$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.037$	$w = 1/[\sigma^2(F_o^2) + (0.049P)^2 + 0.1094P]$
$wR(F^2) = 0.094$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.03$	$(\Delta/\sigma)_{\text{max}} < 0.001$
4306 reflections	$\Delta\rho_{\text{max}} = 0.24 \text{ e \AA}^{-3}$
208 parameters	$\Delta\rho_{\text{min}} = -0.24 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none
Secondary atom site location: difference Fourier map	Absolute structure: Flack (1983), 1853 Friedel pairs
	Flack parameter: 0.03 (7)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.3444 (2)	0.27298 (15)	0.91369 (9)	0.0423 (4)
C2	0.4528 (2)	0.2866 (2)	0.96899 (11)	0.0596 (5)
H2	0.5302	0.2325	0.9709	0.072*
C3	0.4447 (3)	0.3813 (2)	1.02106 (13)	0.0737 (7)
H3	0.5170	0.3910	1.0583	0.088*
C4	0.3305 (3)	0.4615 (2)	1.01838 (13)	0.0720 (7)
H4	0.3252	0.5250	1.0539	0.086*
C5	0.2244 (3)	0.4475 (2)	0.96314 (14)	0.0676 (6)
H5	0.1478	0.5023	0.9611	0.081*
C6	0.2295 (2)	0.3532 (2)	0.91061 (11)	0.0542 (5)
H6	0.1566	0.3438	0.8736	0.065*
C7	0.61067 (18)	0.20568 (15)	0.77713 (10)	0.0396 (4)
H7	0.6361	0.1620	0.8249	0.048*
C8	0.6733 (2)	0.33204 (18)	0.78485 (14)	0.0587 (5)
H8A	0.6285	0.3732	0.8286	0.070*
H8B	0.6528	0.3782	0.7381	0.070*
C9	0.8375 (2)	0.3243 (2)	0.79731 (14)	0.0647 (6)
H9A	0.8777	0.4054	0.8002	0.078*
H9B	0.8572	0.2838	0.8463	0.078*
C10	0.9099 (2)	0.2560 (2)	0.73209 (13)	0.0653 (6)
H10A	0.8985	0.3008	0.6839	0.078*
H10B	1.0131	0.2489	0.7428	0.078*
C11	0.8458 (2)	0.1321 (2)	0.72278 (14)	0.0717 (6)
H11A	0.8666	0.0845	0.7689	0.086*
H11B	0.8908	0.0922	0.6787	0.086*
C12	0.6807 (2)	0.13740 (19)	0.71029 (12)	0.0574 (5)
H12A	0.6593	0.1780	0.6616	0.069*
H12B	0.6416	0.0559	0.7077	0.069*
C13	0.3750 (2)	0.28365 (15)	0.71472 (9)	0.0407 (4)
H13A	0.4319	0.3571	0.7088	0.049*
H13B	0.2811	0.3064	0.7358	0.049*
C14	0.35265 (19)	0.22767 (13)	0.63549 (9)	0.0385 (3)
C15	0.2610 (2)	0.12964 (17)	0.62532 (12)	0.0556 (5)
H15	0.2141	0.0962	0.6681	0.067*
C16	0.2385 (3)	0.0808 (2)	0.55228 (14)	0.0647 (6)
H16	0.1760	0.0153	0.5463	0.078*
C17	0.3074 (3)	0.1283 (2)	0.48873 (12)	0.0646 (6)
H17	0.2917	0.0955	0.4396	0.077*

supplementary materials

C18	0.4001 (3)	0.2249 (2)	0.49795 (12)	0.0668 (6)
H18	0.4484	0.2568	0.4551	0.080*
C19	0.4216 (2)	0.27463 (17)	0.57071 (11)	0.0535 (5)
H19	0.4833	0.3407	0.5762	0.064*
N1	0.44909 (16)	0.20318 (12)	0.77103 (8)	0.0402 (3)
O1	0.21214 (15)	0.12969 (13)	0.81853 (8)	0.0590 (4)
O2	0.43915 (17)	0.05891 (11)	0.88027 (8)	0.0592 (4)
S1	0.35656 (5)	0.15399 (4)	0.84502 (2)	0.04355 (12)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0453 (9)	0.0462 (9)	0.0352 (8)	-0.0024 (8)	0.0063 (8)	0.0056 (7)
C2	0.0564 (12)	0.0720 (13)	0.0504 (11)	0.0075 (11)	-0.0061 (10)	0.0019 (10)
C3	0.0780 (16)	0.0892 (17)	0.0539 (12)	-0.0098 (14)	-0.0089 (12)	-0.0100 (12)
C4	0.0928 (19)	0.0669 (13)	0.0564 (12)	-0.0073 (14)	0.0093 (14)	-0.0126 (11)
C5	0.0761 (15)	0.0617 (13)	0.0651 (14)	0.0124 (12)	0.0139 (12)	-0.0008 (11)
C6	0.0508 (11)	0.0621 (11)	0.0495 (10)	0.0073 (10)	0.0009 (8)	0.0008 (10)
C7	0.0378 (9)	0.0447 (9)	0.0364 (8)	0.0043 (7)	-0.0017 (7)	0.0039 (7)
C8	0.0437 (11)	0.0513 (10)	0.0812 (14)	0.0000 (9)	0.0033 (10)	-0.0147 (10)
C9	0.0432 (11)	0.0695 (13)	0.0815 (14)	-0.0033 (10)	-0.0042 (10)	-0.0146 (12)
C10	0.0351 (10)	0.0957 (17)	0.0651 (13)	0.0074 (11)	-0.0003 (9)	-0.0016 (13)
C11	0.0513 (12)	0.0869 (16)	0.0768 (14)	0.0229 (13)	0.0006 (11)	-0.0247 (13)
C12	0.0501 (11)	0.0611 (11)	0.0610 (12)	0.0101 (9)	-0.0006 (9)	-0.0166 (10)
C13	0.0418 (9)	0.0388 (8)	0.0416 (9)	0.0036 (7)	-0.0034 (7)	0.0009 (7)
C14	0.0357 (8)	0.0376 (7)	0.0423 (8)	0.0045 (7)	-0.0080 (7)	0.0032 (6)
C15	0.0549 (11)	0.0551 (11)	0.0570 (11)	-0.0136 (10)	-0.0014 (9)	0.0008 (9)
C16	0.0671 (14)	0.0558 (11)	0.0712 (14)	-0.0153 (11)	-0.0111 (11)	-0.0097 (11)
C17	0.0855 (16)	0.0613 (12)	0.0469 (11)	0.0000 (11)	-0.0204 (11)	-0.0056 (10)
C18	0.0957 (18)	0.0657 (14)	0.0390 (10)	-0.0094 (13)	-0.0041 (10)	0.0078 (10)
C19	0.0662 (13)	0.0471 (10)	0.0472 (11)	-0.0115 (10)	-0.0069 (9)	0.0067 (9)
N1	0.0387 (8)	0.0455 (8)	0.0365 (7)	0.0043 (6)	-0.0002 (6)	0.0043 (6)
O1	0.0469 (7)	0.0690 (9)	0.0610 (8)	-0.0151 (7)	0.0052 (6)	-0.0039 (7)
O2	0.0711 (9)	0.0436 (7)	0.0629 (8)	0.0059 (7)	0.0071 (7)	0.0168 (6)
S1	0.0449 (2)	0.0415 (2)	0.0442 (2)	-0.0025 (2)	0.0035 (2)	0.00529 (19)

Geometric parameters (\AA , $^\circ$)

C1—C6	1.380 (3)	C10—H10B	0.9700
C1—C2	1.386 (3)	C11—C12	1.535 (3)
C1—S1	1.7696 (18)	C11—H11A	0.9700
C2—C3	1.378 (3)	C11—H11B	0.9700
C2—H2	0.9300	C12—H12A	0.9700
C3—C4	1.375 (3)	C12—H12B	0.9700
C3—H3	0.9300	C13—N1	1.480 (2)
C4—C5	1.370 (3)	C13—C14	1.510 (2)
C4—H4	0.9300	C13—H13A	0.9700
C5—C6	1.379 (3)	C13—H13B	0.9700
C5—H5	0.9300	C14—C19	1.382 (2)

C6—H6	0.9300	C14—C15	1.383 (2)
C7—N1	1.490 (2)	C15—C16	1.382 (3)
C7—C8	1.515 (3)	C15—H15	0.9300
C7—C12	1.518 (2)	C16—C17	1.368 (3)
C7—H7	0.9800	C16—H16	0.9300
C8—C9	1.528 (3)	C17—C18	1.374 (3)
C8—H8A	0.9700	C17—H17	0.9300
C8—H8B	0.9700	C18—C19	1.380 (3)
C9—C10	1.506 (3)	C18—H18	0.9300
C9—H9A	0.9700	C19—H19	0.9300
C9—H9B	0.9700	N1—S1	1.6240 (14)
C10—C11	1.498 (3)	O1—S1	1.4299 (14)
C10—H10A	0.9700	O2—S1	1.4305 (14)
C6—C1—C2	120.51 (18)	C12—C11—H11A	109.3
C6—C1—S1	119.94 (14)	C10—C11—H11B	109.3
C2—C1—S1	119.54 (15)	C12—C11—H11B	109.3
C3—C2—C1	119.3 (2)	H11A—C11—H11B	107.9
C3—C2—H2	120.4	C7—C12—C11	109.44 (16)
C1—C2—H2	120.4	C7—C12—H12A	109.8
C4—C3—C2	120.6 (2)	C11—C12—H12A	109.8
C4—C3—H3	119.7	C7—C12—H12B	109.8
C2—C3—H3	119.7	C11—C12—H12B	109.8
C5—C4—C3	119.6 (2)	H12A—C12—H12B	108.2
C5—C4—H4	120.2	N1—C13—C14	114.01 (13)
C3—C4—H4	120.2	N1—C13—H13A	108.7
C4—C5—C6	121.0 (2)	C14—C13—H13A	108.7
C4—C5—H5	119.5	N1—C13—H13B	108.7
C6—C5—H5	119.5	C14—C13—H13B	108.7
C5—C6—C1	119.05 (19)	H13A—C13—H13B	107.6
C5—C6—H6	120.5	C19—C14—C15	118.11 (16)
C1—C6—H6	120.5	C19—C14—C13	120.72 (15)
N1—C7—C8	113.74 (14)	C15—C14—C13	121.17 (16)
N1—C7—C12	111.15 (15)	C16—C15—C14	120.77 (19)
C8—C7—C12	111.24 (15)	C16—C15—H15	119.6
N1—C7—H7	106.8	C14—C15—H15	119.6
C8—C7—H7	106.8	C17—C16—C15	120.4 (2)
C12—C7—H7	106.8	C17—C16—H16	119.8
C7—C8—C9	109.69 (17)	C15—C16—H16	119.8
C7—C8—H8A	109.7	C16—C17—C18	119.5 (2)
C9—C8—H8A	109.7	C16—C17—H17	120.2
C7—C8—H8B	109.7	C18—C17—H17	120.2
C9—C8—H8B	109.7	C17—C18—C19	120.1 (2)
H8A—C8—H8B	108.2	C17—C18—H18	119.9
C10—C9—C8	111.14 (18)	C19—C18—H18	119.9
C10—C9—H9A	109.4	C18—C19—C14	121.02 (18)
C8—C9—H9A	109.4	C18—C19—H19	119.5
C10—C9—H9B	109.4	C14—C19—H19	119.5
C8—C9—H9B	109.4	C13—N1—C7	119.61 (14)
H9A—C9—H9B	108.0	C13—N1—S1	118.13 (12)

supplementary materials

C11—C10—C9	111.26 (19)	C7—N1—S1	118.29 (11)
C11—C10—H10A	109.4	O1—S1—O2	119.39 (9)
C9—C10—H10A	109.4	O1—S1—N1	107.48 (8)
C11—C10—H10B	109.4	O2—S1—N1	107.38 (8)
C9—C10—H10B	109.4	O1—S1—C1	107.04 (9)
H10A—C10—H10B	108.0	O2—S1—C1	107.20 (8)
C10—C11—C12	111.71 (18)	N1—S1—C1	107.90 (7)
C10—C11—H11A	109.3		
C6—C1—C2—C3	0.1 (3)	C16—C17—C18—C19	-1.0 (4)
S1—C1—C2—C3	178.36 (17)	C17—C18—C19—C14	1.0 (3)
C1—C2—C3—C4	0.0 (3)	C15—C14—C19—C18	-0.2 (3)
C2—C3—C4—C5	-0.4 (4)	C13—C14—C19—C18	-179.12 (19)
C3—C4—C5—C6	0.7 (4)	C14—C13—N1—C7	90.42 (18)
C4—C5—C6—C1	-0.6 (3)	C14—C13—N1—S1	-112.35 (15)
C2—C1—C6—C5	0.3 (3)	C8—C7—N1—C13	49.3 (2)
S1—C1—C6—C5	-178.03 (16)	C12—C7—N1—C13	-77.18 (18)
N1—C7—C8—C9	175.43 (16)	C8—C7—N1—S1	-107.91 (16)
C12—C7—C8—C9	-58.1 (2)	C12—C7—N1—S1	125.63 (14)
C7—C8—C9—C10	56.9 (2)	C13—N1—S1—O1	37.74 (14)
C8—C9—C10—C11	-56.2 (3)	C7—N1—S1—O1	-164.73 (12)
C9—C10—C11—C12	55.8 (3)	C13—N1—S1—O2	167.36 (12)
N1—C7—C12—C11	-174.82 (17)	C7—N1—S1—O2	-35.11 (15)
C8—C7—C12—C11	57.3 (2)	C13—N1—S1—C1	-77.38 (14)
C10—C11—C12—C7	-56.0 (3)	C7—N1—S1—C1	80.15 (14)
N1—C13—C14—C19	-115.08 (18)	C6—C1—S1—O1	-25.17 (17)
N1—C13—C14—C15	66.1 (2)	C2—C1—S1—O1	156.51 (15)
C19—C14—C15—C16	-0.5 (3)	C6—C1—S1—O2	-154.38 (14)
C13—C14—C15—C16	178.40 (19)	C2—C1—S1—O2	27.30 (17)
C14—C15—C16—C17	0.5 (3)	C6—C1—S1—N1	90.25 (16)
C15—C16—C17—C18	0.3 (4)	C2—C1—S1—N1	-88.07 (16)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C7—H7 \cdots O2	0.98	2.34	2.874 (2)	113

Fig. 1

