



## Early View

Research letter

# Negative association between fatigue and signs of sleep apnoea in patients after COVID-19

M. C. Hayden, G. Schwarzl, M. Limbach, S. Mitrea, M. Schuler, D. Nowak, K. Schultz

Please cite this article as: Hayden MC, Schwarzl G, Limbach M, *et al.* Negative association between fatigue and signs of sleep apnoea in patients after COVID-19. *ERJ Open Res* 2022; in press (<https://doi.org/10.1183/23120541.00289-2022>).

This manuscript has recently been accepted for publication in the *ERJ Open Research*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJOR online.

Copyright ©The authors 2022. This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact [permissions@ersnet.org](mailto:permissions@ersnet.org)

# **Negative association between fatigue and signs of sleep apnoea in patients after COVID-19**

Hayden, M.C.<sup>1,+</sup>, Schwarzl, G.<sup>1,+,\*</sup>, Limbach, M.<sup>1</sup>, Mitrea, S.<sup>1</sup>, Schuler, M.<sup>2</sup>, Nowak, D.<sup>3</sup>. & Schultz, K.<sup>1</sup>

<sup>1</sup> Klinik Bad Reichenhall der Deutschen Rentenversicherung Bayern-Süd, Bad Reichenhall, Germany

<sup>2</sup> Department für Angewandte Gesundheitswissenschaften, Hochschule für Gesundheit, Bochum, Germany

<sup>3</sup> Institut und Poliklinik für Arbeits-, Sozial- und Umweltmedizin, Klinikum der Ludwig-Maximilians-Universität München, München, Germany

<sup>+</sup> These authors contributed equally to the manuscript.

<sup>\*</sup> Corresponding author:

**Dr. med.univ. Gabriele Schwarzl**

Klinik Bad Reichenhall der Deutschen Rentenversicherung Bayern Süd

Salzburger Str. 8-11

D-83435 Bad Reichenhall

Germany, EU

Tel: 0049 8651 / 709 – 531

Mail: [gabriele.schwarzl@klinik-bad-reichenhall.de](mailto:gabriele.schwarzl@klinik-bad-reichenhall.de)

Fax: 0049 8651 / 709 – 565

## **Take home message:**

Compared to the general population, there are increased scores of the apnoea-hypopnoea-index in patients recovering from COVID-19. Yet, there is a negative correlation to symptoms of fatigue and no significant correlation to daytime sleepiness.

## **Introduction**

Both during and after the acute phase, COVID-19 is associated with a variety of clinical symptoms, many of which may persist even several months after the infection [1, 2]. One of the most common symptoms that is likely to persist is fatigue [1, 3]. Despite intensive research, little is known about the factors that contribute to the development and persistence of fatigue during and after COVID-19 [3]. Since sleep apnoea and particularly more severe forms are commonly associated with tiredness and exertion, an increased rate of sleep apnoea may explain at least in part the commonly mentioned fatigue symptoms.

Various studies have shown associations between COVID-19 and sleep apnoea [4]. Besides similar risk factors, there are indications that patients affected by sleep apnoea are at a greater risk of developing more severe forms of COVID-19, of being hospitalized, and of dying of the disease [5]. On the other hand, COVID-19 may also increase the risk of developing sleep apnoea due to thrombotic effects and fibrotic changes [4]. There are few studies that examined the prevalence of sleep apnoea in COVID-19 patients, suggesting higher rates than in the general population [6]. However, the link to persistent symptoms of fatigue has so far not been studied yet.

Therefore, the current investigation was conducted with two goals:

- a) Determining the risk for sleep apnoea in form of an increased apnoea-hypopnoea-index (AHI) in patients recovering from COVID-19
- b) Investigating the associations between the AHI and symptoms of fatigue

## **Methods**

The patients were investigated at the Klinik Bad Reichenhall, Germany, in the framework of a research project on effects of a 3–6-week inpatient pulmonary rehabilitation on Long-COVID symptoms. The study protocol was approved by the ethics committee of the medical faculty of

the Ludwig-Maximilians-Universität München, Germany (No. 20-326) and is registered in the German register of clinical studies (DRKS00023180). Participation in the study was voluntary and all participants gave their written informed consent. The recruiting time lasted from April 2020 until April 2021 [7]. All participants were asked to undergo polygraphy screening (Miniscreen, Löwenstein Medical Diagnostics or SOMNOtouch RESP, SOMNOmedics). If the screening detected an AHI of  $\geq 15$ , a more detailed examination using polysomnography (Alice, Löwenstein Medical Diagnostics or SOMNO HD eco, SOMNOmedics) was conducted. Furthermore, we assessed the daytime sleepiness with the Epworth Sleepiness Scale (ESS) [8] and symptoms of fatigue with the Brief Fatigue Inventory (BFI) [9]. These data were intercorrelated along with the body mass index (BMI), sociodemographic variables, and symptom rating scales for dyspnoea, both at rest and under exertion. For an advanced examination, we used a linear regression analysis with fatigue as criterion and all variables that were significantly correlated as predictors to investigate whether bivariate correlations between AHI- and BFI-scores depend on associations between BFI-scores and other variables.

## **Results**

143 patients (39.9% ♀; mean age =  $54.1 \pm 10.4$  years; mean BMI  $30.2 \pm 5.9$  kg/m<sup>2</sup>) underwent polygraphy screening. Comorbidities were common, with only 11 patients (7.7%) having no comorbidity at all. The most common comorbidities were cardiovascular disease (58.0%) and metabolic disorders (53.8%). Only 25.2% of the sample revealed an AHI of  $< 5$ , indicating no clinically relevant sleep apnoea. The percentages for mild ( $5 \leq \text{AHI} < 15$ ), moderate ( $15 \leq \text{AHI} < 30$ ), and severe sleep apnoea ( $\text{AHI} \geq 30$ ) were 39.9%, 21.7%, and 13.3%, respectively.

The results of the correlation and regression analyses are presented in Table 1. There was no significant correlation between the AHI and the ESS. Furthermore, symptoms of fatigue were

negatively correlated with the AHI. Subgroup analyses did not reveal any divergences for age groups, sex, or clusters of the most common comorbidities (cardiovascular disease, metabolic disorders, obesity, orthopaedic disorders, pulmonary diseases, and psychiatric disorders). An additional screening of the scatter plot did not indicate non-linear associations. The regression model was significant with about 34% of variance explained ( $R^2=0.371$ , corrected  $R^2=0.339$ ,  $F(6, 126)=11.793$ ,  $p<0.001$ ).

## **Discussion**

As expected, data revealed increased AHI-scores in the sample, with three quarters of the patients having an AHI of  $\geq 5$  and a third of the patients having an AHI of  $\geq 15$ . Interestingly, the association between the AHI and fatigue turned out to be negative and the AHI scores were not intercorrelated with self-perceived daytime sleepiness.

Putting the results into context, the prevalence of increased AHI-scores that we detected, exceed the prevalence in the German general population of 46% for an AHI  $\geq 5$  and of 21% for an AHI  $\geq 15$  [10]. Regarding daytime sleepiness, it is important to consider that previous research has suggested that some patients suffering from sleep apnoea do not suffer from daytime sleepiness [11]. For example, in the Sleep Heart Health Study, a majority of participants with moderate to severe sleep disordered breathing did not report sleepiness [12]. Despite these results, an non-significant correlation of  $r<0.1$  diverges from the norm, since the severity of sleep apnoea is one of the most important predictors for excessive daytime sleepiness and the AHI is known to correlate with both subjective and objective daytime sleepiness [11]. Similar to the aforementioned analysis, the negative association between the AHI and symptoms of fatigue that we detected in our sample, is hard to interpret. Both included analyses indicate that higher scores of the AHI go along with fewer self-perceived symptoms of fatigue. Even though several studies have highlighted various links between COVID-19 and sleep apnoea, such as similar risk factors or the risk of reciprocal

intensification [4, 5], to the best of our knowledge, a negative correlation between the two constructs has not been reported before. It is important to consider possible publication biases, yet the results appear counter-intuitive. Influences of selection bias and non-linear associations are rather unlikely, since further investigations of the data did not reveal any indications of these two common factors of influence. Nevertheless, sampling effects in the referral to pulmonary rehabilitation may be a possible confounding variable. Furthermore, another explanation may be that commonly used screening instruments, such as the BFI, are not suitable for the detection of COVID-related symptoms of fatigue. Therefore, future analyses should evaluate our results, using broader assessment strategies than only patient reported outcomes. In addition, it is advisable for future studies to focus on possible confounding variables.

In summary, our results confirm previously reported increased rates of the AHI in patients after COVID-19, yet they suggest negative associations to symptoms of fatigue and no associations to daytime sleepiness. Considering the heterogeneous burden of disease that is associated with COVID-19, a closer investigation of the interaction with sleep apnoea and symptoms of fatigue, including an analysis of potential moderators and mediators, is highly advisable.

## **Disclosure Statement**

Financial Disclosure: none. (The study was carried out with the Bad Reichenhall Clinic's regular financial resources without any additional external funding.)

Non-financial Disclosure: none.

The study has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki). Participation in the study was voluntary and all

participants gave their written informed consent. The study protocol was approved by the ethics committee of the medical faculty of the Ludwig-Maximilians-Universität München, Germany (No. 20-326) and is registered in the German register of clinical studies (DRKS00023180).

**Table 1: Correlation matrix and regression coefficients**

Correlation							
	<u>BFI</u>	<u>ESS</u>	<u>BMI</u>	<u>Dyspnoea at rest</u>	<u>Dyspnoea on exertion</u>	<u>Age</u>	<u>Gender</u>
<u>AHI</u>	$r_s = -0.188$ $p = 0.033$	$r_s = 0.022$ $p = 0.797$	$r_s = 0.367$ $p < 0.001$	$r_s = -0.070$ $p = 0.431$	$r_s = -0.063$ $p = 0.473$	$r_s = 0.165$ $p = 0.048$	$r_s = -0.170$ $p = 0.043$
<u>BFI</u>		$r_s = 0.183$ $p = 0.039$	$r_s = 0.183$ $p = 0.039$	$r_s = 0.495$ $p < 0.001$	$r_s = 0.521$ $p < 0.001$	$r_s = -0.148$ $p = 0.096$	$r_s = 0.205$ $p = 0.020$
<u>ESS</u>			$r_s = 0.054$ $p = 0.524$	$r_s = 0.371$ $p < 0.001$	$r_s = 0.097$ $p = 0.272$	$r_s = -0.201$ $p = 0.017$	$r_s = 0.017$ $p = 0.844$
<u>BMI</u>				$r_s = 0.064$ $p = 0.469$	$r_s = 0.217$ $p = 0.013$	$r_s = -0.060$ $p = 0.478$	$r_s = -0.022$ $p = 0.790$
<u>Dyspnoea at rest</u>					$r_s = 0.507$ $p < 0.001$	$r_s = -0.101$ $p = 0.252$	$r_s = 0.154$ $p = 0.081$
<u>Dyspnoea on exertion</u>						$r_s = -0.029$ $p = 0.743$	$r_s = 0.042$ $p = 0.632$
<u>Age</u>							$r_s = -0.097$ $p = 0.251$

  

Regression				
<u>Predictor</u>	<u>b</u>	<u>95% CI</u>	<u>β</u>	<u>p</u>
Dyspnoea on exertion	0.250	0.103 – 0.396	0.291	0.001
Dyspnoea at rest	0.290	0.116 – 0.463	0.290	0.001
AHI	-0.031	-0.056 – 0.006	-0.193	0.016
BMI	0.051	-0.005 – 0.107	0.147	0.074
Gender	0.568	-0.043 – 1.118	0.136	0.068
ESS	0.008	-0.063 – 0.080	0.017	0.821

95% CI = 95% confidence interval; b = regression coefficient; β = standardized regression coefficient.  
 AHI = apnoea-hypopnea-index; BFI = Brief Fatigue Inventory; BMI = body mass index; ESS = Epworth Sleepiness Scale.  
 Values in bold indicate statistically significant results of  $p < 0.05$  (two-sided).



## Literature

1. Förster C, Colombo MG, Wetzel AJ, Martus P, Joos S. Persisting Symptoms after COVID-19: Prevalence and Risk Factors in a Population-Based Cohort. *Dtsch. Arztebl. Int.* 2022; 119: 167–174.
2. Zarei M, Bose D, Nouri-Vaskeh M, Tajiknia V, Zand R, Ghasemi M. Long-term side effects and lingering symptoms post COVID-19 recovery. *Rev. Med. Virol.* 2022; 32.
3. Rao S, Benzouak T, Gunpat S, Burns RJ, Tahir TA, Jolles S, Kisely S. Fatigue Symptoms Associated With COVID-19 in Convalescent or Recovered COVID-19 Patients; a Systematic Review and Meta-Analysis. *Ann. Behav. Med.* [Internet] 2022; 56: 219–234 Available from: <https://academic.oup.com/abm/article/56/3/219/6403353>.
4. Miller MA, Cappuccio FP. A systematic review of COVID-19 and obstructive sleep apnoea. *Sleep Med. Rev.* [Internet] Elsevier Ltd; 2021; 55: 101382 Available from: <https://doi.org/10.1016/j.smr.2020.101382>.
5. Bhat S, Chokroverty S. Sleep disorders and COVID-19. *Sleep Med.* [Internet] Elsevier Ltd; 2021; : 1–9 Available from: <https://doi.org/10.1016/j.sleep.2021.07.021>.
6. Goldstein CA, Rizvydeen M, Conroy DA, O'Brien LM, Gupta G, Somers EC, Sharma P, Golob JL, Troost JP, Burgess HJ. The prevalence and impact of pre-existing sleep disorder diagnoses and objective sleep parameters in patients hospitalized for COVID-19. *J. Clin. Sleep Med.* [Internet] 2021; 17: 1039–1050 Available from: <http://jcsm.aasm.org/doi/10.5664/jcsm.9132>.
7. Hayden MC, Limbach M, Schuler M, Merkl S, Schwarzl G, Jakab K, Nowak D, Schultz K. Effectiveness of a Three-Week Inpatient Pulmonary Rehabilitation Program for Patients after COVID-19: A Prospective Observational Study. *Int. J. Environ. Res. Public Health* [Internet] 2021; 18: 9001 Available from: <https://www.mdpi.com/1660->

4601/18/17/9001.

8. Bloch KE, Schoch OD, Zhang JN, Russi EW. German Version of the Epworth Sleepiness Scale. *Respiration* 1999; 66: 440–447.
9. Radbruch L, Sabatowski R, Elsner F, Everts J, Mendoza T, Cleeland C. Validation of the German Version of the Brief Fatigue Inventory. *J. Pain Symptom Manage.* [Internet] 2003; 25: 449–458 Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0885392403000733>.
10. Fietze I, Laharnar N, Obst A, Ewert R, Felix SB, Garcia C, Gläser S, Glos M, Schmidt CO, Stubbe B, Völzke H, Zimmermann S, Penzel T. Prevalence and association analysis of obstructive sleep apnea with gender and age differences - Results of SHIP-Trend. *J. Sleep Res.* [Internet] 2019; 28: e12770 Available from: <https://onlinelibrary.wiley.com/doi/10.1111/jsr.12770>.
11. Zhou J, Camacho M, Tang X, Kushida CA. A review of neurocognitive function and obstructive sleep apnea with or without daytime sleepiness. *Sleep Med.* [Internet] Elsevier B.V.; 2016; 23: 99–108 Available from: <http://dx.doi.org/10.1016/j.sleep.2016.02.008>.
12. Kapur VK, Baldwin CM, Resnick HE, Gottlieb DJ, Nieto FJ. Sleepiness in patients with moderate to severe sleep-disordered breathing. *Sleep* 2005; 28: 472–477.
13. Walsleben JA, Kapur VK, Newman AB, Shahar E, Bootzin RR, Rosenberg CE, O'Connor G, Nieto FJ. Sleep and Reported Daytime Sleepiness in Normal Subjects: the Sleep Heart Health Study. *Sleep* 2004; 27: 293–298.