

## Research Article

# An Anatomical Basis for the Myofascial Trigger Points of the Abductor Hallucis Muscle

**Juliano T. Wada** , **Flavia Akamatsu** , **Flavio Hojaij**, **Ana Itezerote**, **José Carlos Scarpa**, **Mauro Andrade**, and **Alfredo Jacomo**

*Department of Surgery-Human Structural Topography-School of Medicine, University of São Paulo, São Paulo 01246-903, Brazil*

Correspondence should be addressed to Juliano T. Wada; [julianowada@usp.br](mailto:julianowada@usp.br)

Received 24 August 2019; Revised 7 December 2019; Accepted 27 December 2019; Published 23 January 2020

Academic Editor: Nihal Apaydin

Copyright © 2020 Juliano T. Wada et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Myofascial pain syndrome is characterized by pain and a limited range of joint motion caused by muscle contracture related to motor-end-plate dysfunction and the presence of myofascial trigger points (MTrPs). It is the most frequent cause of musculoskeletal pain, with a worldwide prevalence varying between 13.7% and 47%. Of the patients with myofascial pain syndrome, approximately 17% have pain in the medial hindfoot area. The abductor hallucis muscle is located in the medial, posterior region of the foot and is related to painful plantar syndromes. The objective of this study was to describe the distribution of the medial plantar nerve and their anatomical relationship with MTrPs found in the literature. Thirty abductor hallucis muscles were dissected from 15 human cadavers (8 males and 7 females). The muscles were measured, and the distribution data of the medial plantar nerve branches in each quadrant were recorded. For statistical analysis, we used generalized estimation equations with a Poisson distribution and a log logarithm function followed by Bonferroni multiple comparisons of the means. The data are expressed as the mean  $\pm$  standard deviation. The level of significance was adjusted to 5% ( $p < 0.05$ ). A high concentration of nerve branches was observed in the first quadrant (Q1) of the abductor hallucis muscle, which is the same area in which the MTrPs are described. The topography of the entry points of the branches of the medial plantar nerve to the abductor hallucis muscle correlates with the topography of the muscular trigger points. The anatomical structure of the MTrPs may be useful for a better understanding of the pathophysiology of myofascial disorders and provide a basis for surgical and clinical treatments.

## 1. Introduction

Chronic myofascial pain is the most frequent cause of skeletal muscle pain with prevalence in the world population varying between 13.7% and 47% [1, 2]. Pain is a global public health problem because it directly impacts the individual, the family, and the workplace as it results in a decrease in functional activity [3, 4]. Myofascial pain syndrome may be associated with myofascial trigger points (MTrPs) [5, 6]. MTrPs are foci of muscles that have intense sensitivity and irritability, are located predominantly near the motor end plates, and have palpable tensile band characteristics mediated by the local response of reflex muscle contraction upon palpation of muscle fibers. Increased local irritability gives rise to pain and sensory changes that may be local or

referred [7, 8]. The integrated MTrPs hypothesis postulates that, in myofascial pain, the motor endplates release excessive acetylcholine which is evidenced histopathologically by the presence of sarcomere shortening [9].

The abductor hallucis muscle has been gaining prominence in the most recent publications on plantar pain [10, 11]. It is a superficial, triangular, flat, bipenniform muscle that fills the foot medial arch and is responsible for sustaining the arch and hallux abduction and flexion [12, 13]. The muscle originates from flexor retinaculum, medial process of the calcaneus tuberosity, and intermuscular septum between itself and flexor hallucis brevis; the insertion is into the medial aspect of the proximal hallux phalanx, the collateral ligament of the proximal metatarsophalangeal joint, and the medial sesamoid bone

[14, 15]. The abductor hallucis muscle is innervated by a medial plantar nerve, which is a branch of the tibial nerve, and the vascular supply is made by a medial malleolar network, medial calcaneus branches of the lateral plantar artery, medial plantar artery, first plantar metatarsal artery, and perforations of the arterial plantar arch [16, 17].

According to Simons et al. [18], there are three areas in the abductor hallucis muscle that have MTrPs. The areas are concentrated in the region of the medial hindfoot and referred pain spreads throughout the medial aspect of the foot (Figure 1). The main epidemiological studies found that 17.4% had foot pains [19], most of them in the region of the medial hindfoot [20, 21].

The authors have sought anatomical approaches to support the theory of myofascial pain syndromes (MPSs) citing the presence of muscle trigger points and providing a pathophysiological explanation [6, 18]. However, some recent researchers refute this approach, concluding that there is no concrete and well-founded pathophysiological explanation for MPS, since, although the muscles have trigger points, patients often do not present with symptoms or some type of electromyographic muscle alteration [22–24].

Another important finding described in a recent study demonstrated that there may be a relationship between the entry points of the nerves to the muscle belly and the myofascial trigger points [25, 26]. A detailed description of the anatomy remains one of the best ways to better understand the pathophysiology and clinical applicability of myofascial trigger points. Nevertheless, the lack of detailed anatomical information still constitutes a major setback for a complete understanding of the physiopathology and the larger clinical applicability of MTrPs.

To better understand MTrPs pathophysiology and to provide more anatomical data for the treatment application based on the hypothesis that trigger points can be related to the muscle innervation, the objective of this study was to observe the anatomical correlation between the clinically described MTrPs and the entry point of the branches of the medial plantar nerve into the abductor hallucis.

## 2. Methods

This study was approved by the Ethics Committee of the Medical School for the Analysis of Research Projects Protocol No. 027/16. To study the interrelationship between the number of entry points of medial plantar nerve branches to the abductor hallucis muscle and the MTrPs of the muscle, human adult cadavers (8 males and 7 females) prepared with 4% phenolic acid and 0.5% formaldehyde were dissected. The cadavers were obtained from a body donation program managed by the Discipline of Human Structural Topography of the Department of Surgery of the University, Medical School. The criteria of the 15 cadavers and 30 feet dissected are as follows: preserved anatomical structures; absence of dissection or previous surgical incision in the feet; age between 40 and 80 years; BMI between 20 kg/m<sup>2</sup> and 25 kg/m<sup>2</sup>; absence of signs of previous surgeries in the regions of interest; and absence of severe pathologies, anatomical deformities, and processes of necroses,




FIGURE 1: Medial view of the feet, trigger points, and referred pain of the abductor hallucis muscle from Simons et al. [18].

hemorrhages, and fibroses that make adequate study of the tissues of the lower limbs impossible (Table 1).

The feet selected met all the study criteria; dissection was performed stratigraphically. Flaps of skin and subcutaneous tissue were folded to expose the abductor muscle of the hallux. The tendons of the long flexors of the digitorum and hallucis longus as well as the tibial artery and vein were separated, and in some cases removed, to identify and isolate the tibial, medial, and lateral plantar nerve. The abductor hallucis muscle was dissected and folded, and the deep surface was carefully dissected to preserve the nerve branches. All entry points of the branches of the medial plantar nerve on the lateral side of the muscle and the medial face of the muscle were demarcated with pins. The boundaries of the quadrants [26, 27] were traced with surgical wires, and the anatomical references were as follows: the X axis (anteroposterior) began at the origin (medial tuberosity of the calcaneus) and ended at the insertion (base of the first proximal phalanx), and the Y axis (transversal) originated from the midpoint of the X axis, close to the navicular bone (Figure 2). The number of entry points of each of the branches of the medial plantar nerve into the muscle in each quadrant was demarcated, and the statistical analysis was performed by a blind evaluator to avoid the observer bias. All the anatomical dissection steps were documented photographically by means of a Nikon D52 camera (Nikon Corporation; Tokyo Japan).

Statistical analysis was performed using SPSS (version 22, IBM). The sample size calculation was based on the results of the first 10 feet evaluated by the pilot project; with this sample, the difference between quadrants 1 and 2 (Q1 and Q2, respectively) was 1.26 points on average with a variability of 1.5 points. The sample used results in a power of 85% and an alpha value of 5%, suggesting a sample size of 26 feet total.

For the quadrant analysis, generalized estimation equations with the Poisson distribution and log logarithm function followed by Bonferroni multiple comparisons were used to evaluate the number of entry points of the medial plantar nerve into the hallucis abductor muscle in the

TABLE 1: Baseline sample characteristics.

Anthropometric data	Anatomic dissection group
Gender (M/F)	8/7
Age (years)	60.8 ± 10.7
Height (m)	1.68 ± 0.7
Weight (kg)	69.1 ± 11.6
BMI (kg/m <sup>2</sup> )	24.2 ± 4.3

Note. Data are presented as mean ± standard deviation. M/F: male, female; BMI: body mass index; m: meters; kg: kilograms.




FIGURE 2: The axes and quadrants of the plantar surface of the feet. 1° quadrant (1Q): posterolateral; 2° quadrant (2Q): posteromedial; 3° quadrant (3Q): anteromedial; 4° quadrant (4Q): anterolateral.

muscle’s respective quadrants. Data are presented as the mean ± standard deviation, and the level of significance was adjusted to 5% ( $p < 0.05$ ) for all tests.

### 3. Results

Only one foot ( $n = 1$ ) was excluded from the study because of an orthopedic surgery with intense tissue fibrosis in the ankle that prevented adequate anatomical evaluation; therefore, 29 feet were used for the analysis. The anatomical analysis showed that all of the entries of the branches of the medial plantar nerve were concentrated in the posterior foot area near the calcaneus bone and that both the third and fourth quadrants (3Q and 4Q, respectively) did not contain any entry points of the branches of the medial plantar nerve into the muscle belly. In contrast, the first quadrant (1Q, posterolateral) presented a statistically significant higher number of entry points of the branches of the medial plantar nerve relative to the second quadrant (2Q, posteromedial) ( $2.37 \pm 1.84$  points versus  $1.11 \pm 1.16$  points) ( $p < 0.05$ ) (Figure 3). After dissection, it was possible to observe that the medial plantar nerve penetrates the lateral side of the abductor hallucis muscle and crosses the entire second quadrant (medial, 2Q) before innervating the posterolateral (first quadrant, 1Q) region of the muscle. The nerve divides on average into 3 branches to penetrate the muscle in the middle of the muscle belly; briefly, the muscle receives all its innervation in only 1/4 of its total muscle area (Figure 4).




FIGURE 3: Number of entry points of the medial plantar nerve into the abductor hallucis muscle belly. 1° quadrant (1Q): posterolateral; 2° quadrant (2Q): posteromedial; 3° quadrant (3Q): anteromedial; 4° quadrant (4Q): anterolateral.

### 4. Discussion

The relationship between the entry points of nerves into the muscle belly and the myofascial trigger points has already been reported by Akamatsu et al. [25, 26]. Nerves are responsible for muscle contraction through the release of acetylcholine from the motor endplate. All the mechanisms involved in the nodular tense band in the muscular belly (i.e., MTrPs) are not completely elucidated, and motor plate disorders caused by an increase of acetylcholine may be one of the causes [28]. Electromyography (EMG) studies have demonstrated that acute and chronic muscle pain experience alters the distribution of activity of the muscle and may cause a shift of the peak muscle activity [29, 30]. An active MTrP causes a clinical pain complaint [18], and the International Association for the Study of Pain [31] refers MTrPs related to many pain conditions and the plantar pain is mainly caused by muscular dysfunction [32]. In this study, it was demonstrated through anatomical dissections that the branches of the medial plantar nerve penetrate the abductor hallucis muscle mainly in the first quadrant (Q1, posterolateral). The first quadrant (1Q, posterolateral) in both groups presented a significant difference relative to the other plantar areas.

The anatomical entry points of the medial plantar nerve into the abductor hallucis muscle corresponded to the described areas of the myofascial trigger points (MTrPs).

In addition, this region of the medial hindfoot is the same as described by Simon et al. [18] and Saggini et al. [33] for the trigger point of the hallucis muscle. The nerve divides on average into 3 branches before penetrating the muscle in the middle of the muscle belly, a similar result as Saitou et al. [34]. The main epidemiological studies found that 17.4% had foot pains [19], most of them in the region of the medial hindfoot [20, 21].

Several studies have examined the pathway of the medial plantar nerve but only up to the margin of the abductor hallucis muscle [35, 36]. The studies that more clearly




FIGURE 4: Medial view of the left foot (A and B) and medial plantar nerve branch penetration in the posterolateral region muscle belly. The abductor hallucis muscle was retracted to expose its inner side to view the muscle innervation.

describe the plantar nerves are from Delfaut et al. [37] and Alshami et al. [35], but they used magnetic resonance imaging only, which fails to demonstrate in detail the innervation of the muscle. Saitou et al. [34] and Minetto et al. [38] demonstrated that the motor endplate of the abductor muscle of the hallux is located mainly in the middle of the muscle and in the posterior region near the muscle origin. We found an anatomical match between the abductor hallucis muscle trigger points area and the anatomical entries of the medial plantar nerve branches into the muscle belly. According to the data described in the present study, it is believed that the muscle region with high nerve concentration (first quadrant, 1Q) may be related to the pathophysiology of myofascial syndromes.

The present study was the first to relate the entry points of the medial plantar nerve into the belly of the abductor hallucis muscle with the myofascial trigger points and to evaluate the standard location of the nerves using the quadrant approach. The location of the trigger points in relation to the entry points of the medial plantar nerve branches into the abductor muscle of the hallux is a logical explanation for understanding the dysfunction of the neurological activity of the MTrPs.

## 5. Conclusion

The branches of the medial plantar nerve penetrate mostly into the first muscle quadrant (1Q, posterolateral) of the abductor muscle of the hallux and represent a strong relationship between nerve anatomy and the location of the myofascial trigger points.

## Abbreviations

TN: Tibial nerve  
 MPN: Medial plantar nerve  
 MM: Medial malleolus  
 CB: Calcaneus bone  
 NB: Navicular bone  
 AH: Abductor hallucis  
 1Q: First quadrant

2Q: Second quadrant  
 3Q: Third quadrant  
 4Q: Fourth quadrant

## Data Availability

The data used to support the findings of this study are included within the article, but if it is necessary to consult all the data of the table with each of the measurements of the entrance of nerves per quadrant of the corpses feet, they are available from the corresponding author upon request.

## Conflicts of Interest

The authors have no financial or other potential conflicts of interest in the authorship or publication of this paper.

## Authors' Contributions

All authors have contributed at all stages of this study: Juliano T. Wada is a significant manuscript writer and contributed to the study concept and design, anatomy dissections, and data acquisition. Flavia Akamatsu is a significant manuscript reviser and performed the study concept and design, data acquisition, and data analysis and interpretation. Flávio Hojaij is a significant manuscript reviser and contributed to the data acquisition and data analysis and interpretation. Ana Itezerote performed anatomy dissections, data acquisition, and data interpretation. José Carlos Scarpa did anatomy dissections, study concept, and data interpretation. Mauro Andrade is a significant manuscript reviser and contributed to the study concept and design and data acquisition. Alfredo Jacomo is a significant manuscript writer and performed the study concept and design and data analysis and interpretation.

## References

- [1] A. M. Elliott, B. H. Smith, K. I. Penny, W. Cairns Smith, and W. Alastair Chambers, "The epidemiology of chronic pain in

- the community," *The Lancet*, vol. 354, no. 9186, pp. 1248–1252, 1999.
- [2] M. A. Cimmino, C. Ferrone, and M. Cutolo, "Epidemiology of chronic musculoskeletal pain," *Best Practice & Research Clinical Rheumatology*, vol. 25, no. 2, pp. 173–183, 2011.
  - [3] O. Gureje, M. Von Korff, G. E. Simon, and R. Gater, "Persistent pain and well-being," *JAMA*, vol. 280, no. 2, p. 147, 1998.
  - [4] A. C. Williams, C. Eccleston, and S. Morley, "Psychological therapies for the management of chronic pain (excluding headache) in adults," *The Cochrane Database of Systematic Reviews*, vol. 11, no. 11, Article ID CD007407, 2012.
  - [5] J. Z. Srbely, "New trends in the treatment and management of myofascial pain syndrome," *Current Pain and Headache Reports*, vol. 14, no. 5, pp. 346–352, 2010.
  - [6] S. Money, "Pathophysiology of trigger points in myofascial pain syndrome," *Journal of Pain & Palliative Care Pharmacotherapy*, vol. 31, no. 2, pp. 158–159, 2017.
  - [7] K. Takamoto, S. Sakai, E. Hori et al., "Compression on trigger points in the leg muscle increases parasympathetic nervous activity based on heart rate variability," *The Journal of Physiological Sciences*, vol. 59, no. 3, pp. 191–197, 2009.
  - [8] M. A. Giamberardino, G. Affaitati, A. Fabrizio, and R. Costantini, "Myofascial pain syndromes and their evaluation," *Best Practice & Research Clinical Rheumatology*, vol. 25, no. 2, pp. 185–198, 2011.
  - [9] D. G. Simons, "Review of enigmatic MTrPs as a common cause of enigmatic musculoskeletal pain and dysfunction," *Journal of Electromyography and Kinesiology*, vol. 14, no. 1, pp. 95–107, 2004.
  - [10] E. Brenner, "Insertion of the abductor hallucis muscle in feet with and without hallux valgus," *The Anatomical Record*, vol. 254, no. 3, pp. 429–434, 1999.
  - [11] R. K. Chittoria, H. Pratap, and S. H. Yekappa, "Abductor hallucis: anatomical variation and its clinical implications in the reconstruction of chronic nonhealing ulcers and defects of foot," *Advances in Wound Care*, vol. 4, no. 12, pp. 719–723, 2015.
  - [12] K. S. Nam, J. W. Kwon, and O.-Y. Kwon, "The relationship between activity of abductor hallucis and navicular drop in the one-leg standing position," *Journal of Physical Therapy Science*, vol. 24, no. 11, pp. 1103–1106, 2012.
  - [13] M.-H. Kim, C.-H. Yi, J.-H. Weon, H.-S. Cynn, D.-Y. Jung, and O.-Y. Kwon, "Effect of toe-spread-out exercise on hallux valgus angle and cross-sectional area of abductor hallucis muscle in subjects with hallux valgus," *Journal of Physical Therapy Science*, vol. 27, no. 4, pp. 1019–1022, 2015.
  - [14] S. Stranding, *Gray's Anatomy*, Churchill Livingstone, London, UK, 40th edition, 2008.
  - [15] Y. S. Wong, "Influence of the abductor hallucis muscle on the medial arch of the foot: a kinematic and anatomical cadaver study," *Foot & Ankle International*, vol. 28, no. 5, pp. 617–620, 2007.
  - [16] T. H. Lui, "Endoscopic decompression of the first branch of the lateral plantar nerve and release of the plantar aponeurosis for chronic heel pain," *Arthroscopy Techniques*, vol. 5, no. 3, pp. e589–e594, 2016.
  - [17] S. Stranding, *Gray's Anatomy: the Anatomical Basis of Clinical Practice*, Churchill Livingstone, London, UK, 40th edition, 2008.
  - [18] D. G. Simons, J. G. Travell, and L. S. Simons, *Myofascial Pain and Dysfunction: The Trigger Point Manual: Lower Half of Body*, Lippincott Williams & Wilkins, Philadelphia, PA, USA, 1999.
  - [19] C. L. Hill, T. K. Gill, H. B. Menz, and A. W. Taylor, "Prevalence and correlates of foot pain in a population-based study: the North West Adelaide health study," *Journal of Foot and Ankle Research*, vol. 1, no. 1, p. 2, 2008.
  - [20] H. B. Menz, A. B. Dufour, V. A. Casey et al., "Foot pain and mobility limitations in older adults: the framingham foot study," *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, vol. 68, no. 10, pp. 1281–1285, 2013.
  - [21] B. D. Chatterton, S. Muller, and E. Roddy, "Epidemiology of posterior heel pain in the general population: cross-sectional findings from the clinical assessment study of the foot," *Arthritis Care & Research*, vol. 67, no. 7, pp. 996–1003, 2015.
  - [22] R. M. Bennett and D. L. Goldenberg, "Fibromyalgia, myofascial pain, tender points and trigger points: splitting or lumping?" *Arthritis Research & Therapy*, vol. 13, no. 3, pp. 117–127, 2011.
  - [23] F. E. Akamatsu, S. Saleh, H. T. Pinesi et al., "Anatomical basis of the myofascial trigger points of the trapezius muscle," *International Journal of Morphology*, vol. 31, no. 3, pp. 915–920, 2013.
  - [24] J. L. Quintner, G. M. Bove, and M. L. Cohen, "A critical evaluation of the trigger point phenomenon," *Rheumatology (Oxford)*, vol. 54, no. 2, pp. 392–399, 2015.
  - [25] F. E. Akamatsu, B. R. Ayres, S. O. Saleh et al., "Trigger points: an anatomical substratum," *BioMed Research International*, vol. 2015, Article ID 623287, 29 pages, 2015.
  - [26] F. E. Akamatsu, T. M. Yendo, C. Rhode et al., "Anatomical basis of the myofascial trigger points of the gluteous maximus muscle," *BioMed Research International*, vol. 2017, Article ID 4821968, 5 pages, 2017.
  - [27] B. Saban and Y. Masharawi, "Pain threshold tests in patients with heel pain syndrome," *Foot & Ankle International*, vol. 37, no. 7, pp. 730–736, 2016.
  - [28] R. D. Gerwin, J. Dommerholt, and J. P. Shah, "An expansion of Simons' integrated hypothesis of trigger point formation," *Current Pain and Headache Reports*, vol. 8, no. 6, pp. 468–475, 2004.
  - [29] D. Falla, H. Andersen, B. Danneskiold-Samsøe, L. Arendt-Nielsen, and D. Farina, "Adaptations of upper trapezius muscle activity during sustained contractions in women with fibromyalgia," *Journal of Electromyography and Kinesiology*, vol. 20, no. 3, pp. 457–464, 2010.
  - [30] D. Falla, L. Arendt-Nielsen, and D. Farina, "The pain-induced change in relative activation of upper trapezius muscle regions is independent of the site of noxious stimulation," *Clinical Neurophysiology*, vol. 120, no. 1, pp. 150–157, 2009.
  - [31] IASP, "IASP international association for the study of pain: global year against musculoskeletal pain," *Musculoskeletal Pain Fact Sheets*, vol. 8, no. 6, pp. 468–475, 2017.
  - [32] H. Riel, M. Cotchett, E. Delahunt et al., "Is 'plantar heel pain' a more appropriate term than 'plantar fasciitis?' Time to move on," *British Journal of Sports Medicine*, vol. 51, no. 22, pp. 1576–1577, 2017.
  - [33] R. Saggini, R. G. Bellomo, G. Affaitati, D. Lapenna, and M. A. Giamberardino, "Sensory and biomechanical characterization of two painful syndromes in the heel," *The Journal of Pain*, vol. 8, no. 3, pp. 215–222, 2007.
  - [34] K. Saitou, T. Masuda, D. Michikami, R. Kojima, and M. Okada, "Innervation zone of the upper and lower limb muscles estimated by using multichannel surface EMG," *Journal of Human Ergology (Tokyo)*, vol. 29, no. 1-2, pp. 35–52, 2000.
  - [35] A. M. Alshami, T. Souvlis, and M. W. Coppieters, "A review of plantar heel pain of neural origin: differential diagnosis and

- management,” *Manual Therapy*, vol. 13, no. 2, pp. 103–111, 2008.
- [36] S. K. Ghosh, S. Raheja, and A. Tuli, “Potential sites of compression of tibial nerve branches in foot: a cadaveric and imaging study,” *Clinical Anatomy*, vol. 26, no. 6, pp. 768–779, 2013.
- [37] E. M. Delfaut, X. Demondion, A. Bieganski, M.-C. Thiron, H. Mestdagh, and A. Cotten, “Imaging of foot and ankle nerve entrapment syndromes: from well-demonstrated to unfamiliar sites,” *RadioGraphics*, vol. 23, no. 3, pp. 613–623, 2003.
- [38] M. A. Minetto, A. Botter, D. De Grandis, and R. Merletti, “Time and frequency domain analysis of surface myoelectric signals during electrically-elicited cramps,” *Neurophysiologie Clinique/Clinical Neurophysiology*, vol. 39, no. 1, pp. 15–25, 2009.