



Case report

Sudden emergence and spread of cutaneous larva migrans in Sudan: A case series calls for urgent actions

Ayman Ahmed^{a,b,c}, Maisa Aldai Hemaida^d, Amel Ahmed Hagelnur^e, Hala Fathi Eltigani^f, Emmanuel Edwar Siddig^{d,f,g,*}

^a Institute of endemic diseases, University of Khartoum, Khartoum, Sudan

^b Swiss Tropical and Public Health Institute (Swiss TPH), Allschwil, Switzerland

^c University of Basel, Petersplatz 1, Basel, Switzerland

^d Unit of Applied Medical Sciences, Faculty of Medical Laboratory Sciences, University of Khartoum, Khartoum, Sudan

^e General Director, Tropical Medicine Hospital, Omdurman, Khartoum State, Sudan

^f The Mycetoma Research Center, University of Khartoum, Khartoum, Sudan

^g ErasmusMC, University Medical Center Rotterdam, Department of Medical Microbiology and Infectious Diseases, Rotterdam, the Netherlands



ARTICLE INFO

Keywords:

Cutaneous larva migrans
Skin diseases Zoonoses
Integrated surveillance and Response system
One Health
Sudan

ABSTRACT

Cutaneous larva migrans (CLM) is a self-limiting dermatosis parasitic disease that is caused by zoonotic hookworm mainly infects cats, dogs, and accidentally humans. The disease affects hosts through the invasion and the migration of the hookworm larva into top layers of the skin. The disease occurs in tropical and subtropical regions; where people commonly are infected when sitting or barefoot walking on places contaminated with feces of infected cats or dogs. Due to self-limiting nature of the disease, the exact burden and prevalence are commonly underestimated. In this communication, we investigated all cases presented to the skin-diseases outpatient clinic at the reference hospital of Tropical Diseases Medicine in Khartoum state between January 2019 and January 2021. This is the first-ever a case series report of cutaneous larva migrans in Sudan. We identified 15 cases of CLM presented with rash (100 %), redness of skin (67 %), and only adults were presented with larva crawling under their skins (27 %). Infection sites were (53 %) leg, (40 %) foot, only 7 % showed abdominal infection. The majority of patients were children or young adults, 47 % of them are ≤ 5 year-old, male to female ratio is 2.75:1. Duration of infection was one to three weeks and all patients fully recovered after treatment with albendazole. This urges for One Health interventions including deworming cats and dogs, improving water, sanitation, and hygiene, community-engagement and raising awareness in area with risk of infection.

Introduction

Cutaneous larva migrans (CLM) is a zoonotic neglected tropical disease of public health importance that is caused by nematodes helminths [1]. Multiple types of hookworms are involved in the development of CLM [2]. The disease is commonly transmitted through the contamination of soil with animal feces (mainly dogs and cats particularly in beaches) that contain eggs, once it hatch it break through humans skin and migrate through the body [2,3]. CLM mainly affected body organs that are directly in contact with the contaminated soil including feet, legs, and hands; however, other body site can be involved [4]. CLM is a major public health problem in developing countries of tropical and sub-tropical weather. Nevertheless, few studies have

demonstrated the growing burden of the disease and spreading out the tropical and tropical regions, yet it highlighted the severe neglect and underestimation [5]. The disease is characterized by the appearance of erythematous, serpiginous, pruritic, and skin eruption. In addition to the health and economic burdens, this has added a social burden that is indicated by social stigma and isolation among patients [6].

In general, CLM is not fatal disease and most time it is self-limited infection [2]. This might have largely contributed in the underestimation of the disease burden in addition to the common confusion among healthcare providers of CLM with other dermal diseases of non-larval cutaneous migrations such as loiasis and scabies as well as the Guinea worm disease [2].

CLM has never been reported from Sudan previously. Therefore, here

* Corresponding author at: Unit of Applied Medical Sciences, Faculty of Medical Laboratory Sciences, University of Khartoum, Khartoum, Sudan.

E-mail address: emanwell-eds3@hotmail.com (E.E. Siddig).

<https://doi.org/10.1016/j.idcr.2023.e01789>

Received 20 February 2023; Received in revised form 27 April 2023; Accepted 2 May 2023

Available online 5 May 2023

2214-2509/© 2023 Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

In this short communication, we provide the first report CLM from Sudan describing 15 cases of CLM among Sudanese patients that were presented at the Sudan referral hospital for tropical disease, in Omdurman locality, Khartoum state, Sudan between January 2019 and January 2022. We have reviewed all diagnostic reports from the hospital about CLM and extracted cases information to be reported in an attempt to identify risk factors associated with and hotspots of CLM infection transmission.

Case presentation

We have identified 15 cases of CLM that were presented at the dermatology clinic at tropical medicine hospital, Omdurman city, Khartoum state, Sudan between January 2019 and January 2021. All patients are residents in Omdurman city, the western side of Khartoum state. Patients were clinically presented with itching (100 %), skin redness (67 %), and larva crawling inside the skin (27 %) (Fig. 1).

Our demographic analysis shows that patient age ranged very widely between three months and 53 years old, with average age 15 years, and the majority (60 %) were children younger than eight years old. The male to female sex ratio was 2.75:1 with 73 % male and 27 % female (Table 1).

The duration of the disease at the presentation varied from one to three weeks, with 53 % of CLM infections persisted for three weeks and 40 % for two weeks, while only a single patient (7 %) was presented to the clinic on the first week of disease onset (Table 1).

Leg was the most affected site by CLM infection (53 %), followed by the foot (40 %) and only a single case was presented with abdominal infection (7 %) (Table 1).

For the case management, albendazole was prescribed for all patients to take once a day for three weeks duration. All patients respond well for medical treatment and the complaint had regressed and recovered.

Discussion

In this communication, we report for first time cases of CLM from Sudan. This report highlights a sudden emergence and spread of CLM in Khartoum state. The geographical clustering and rapid emergence of cases suggest the development of a disease outbreak, however, considering the limitations in our surveillance, case tracking, and information management systems as well as the tendency of local people to not disclose their travel history, an epidemiological link between the patients wasn't established [7,8]. Particularly that the majority of the patients (60 %) are children under eight years old suggest that the transmission of CLM might be associated with playgrounds where children and their attending guardian have visited for outdoor activities.



Fig. 1. Clinical presentation of cutaneous larva migrans.

Table 1

Demographic and clinical characteristics of the cutaneous larva migrans cases that presented at Tropical Diseases Hospital in Sudan between January 2019 and January 2021.

Case No.	Age	Gender	Site	Duration (weeks)	Symptoms
Case 1	3 months	Male	Foot	3 weeks	Itching with redness of skin
Case 2	1 year	Male	Leg	2 weeks	Itchy rash
Case 3	2 years	Female	Foot	3 weeks	Itching with redness of skin
Case 4	2 years	Male	Foot	3 weeks	Itching with redness of skin
Case 5	3 years	Female	Leg	3 weeks	Itching with redness of skin
Case 6	4 years	Female	Leg	2 weeks	Itching with redness of skin
Case 7	5 years	Male	Abdomen	2 weeks	Itching with redness of skin
Case 8	6 years	Male	Leg	3 weeks	Itching with redness of skin
Case 9	7 years	Female	Foot	2 weeks	Itching with redness of skin
Case 10	18 years	Male	Leg	1 week	Itching with redness of skin
Case 11	18 years	Male	Leg	3 weeks	Itching with larva crawling inside the skin
Case 12	25 years	Male	Leg	2 weeks	Itching with redness of skin
Case 13	35 years	Male	Leg	3 weeks	Itching with larva crawling inside the skin
Case 14	48 years	Male	Foot	3 weeks	Itching with larva crawling inside the skin
Case 15	53 years	Male	Foot	2 weeks	Itching with larva crawling inside the skin

However, the fact that all adult (age of 18 years and over) patients are male might be attributed to social and healthcare disparities and inequalities that discourage female from seeking healthcare services. This skewed sex ratio is commonly observed across different infectious diseases in the country [9–12]. Additionally, the relatively high involvement of male in outdoor activities in places that usually contaminated with cats and dogs feces such as beaches and playgrounds is exposing them to higher risk of infection [6].

These cases were differentially diagnosed and confirmed mainly by highly trained dermatologists, however clinicians with such specialized training are not available in most of hospitals in the country including those in the capital city forget about remote areas. Therefore, this lack of expertise might be one of the reasons behind the lack of reporting CLM from Sudan before. Therefore, further research is needed to identify high risk areas, risk factors associated with CLM transmission, and improving the surveillance and reporting systems in the country. Also, healthcare providers in areas at risk need to be trained on the diagnosis, case management, and reporting. They particularly need to be trained on distinguishing CLM from other clinical mimics such as larva currens, dermatitis due to cercarial, contact dermatitis, scabies and migratory myiasis [2].

CLM infection is commonly a self-limiting condition with complete healing occurs at the 5th or 6th week after the onset of the disease due to the death of the larvae [12]. The ideal management is treating the symptoms such as pruritic rashes can be treated by the administration of topical or orally antihistamines, and the administration of albendazole for the larvae [12–16].

There is a growing reports about detection of CLM among return travelers, this is very alarming because with climate change and global warming, introduction of CLM-causing hookworms into novel areas might lead to the establishment of the disease locally and expanding it is

geographical distribution [14–17]. Similarly in Sudan, the emergence and spread of CLM in the country that we report here for the first time is associated with substantial socioeconomic, geopolitical, and climate changes that are highlighted by the escalation of political instability that led to growing violence and devastating economic crisis, and extreme weather events effected the country since late 2018 [18–20]. Several outbreaks of infectious diseases were driven by these changes including chikungunya, Cholera, Rift Valley fever, dengue, Hepatitis E, and malaria [18–20]. Considering the zoonotic nature of CLM, a One Health approach with integrated surveillance and response systems among humans, animals, and the environment is needed for effective prevention and control strategy [20]. Nowadays more than ever, we need to invest in developing and implementing effective preventive and control measures of infectious diseases particularly zoonoses including CLM as evidence has shown the devastating health and economic impacts of these infections among populations in endemic countries. These preventive and control measure should be accompanied by social engagement and health education to raise awareness about the disease and the associated risk factors and improve personal protection as well as reduce the associated stigma.

Ethics approval and consent to participate

Written, informed consent to publish history, findings, and images for educational purposes were obtained from the patients.

Consent for publication

Written informed consent was obtained from the patients for publication of this case report and any accompanying images.

Funding

Not Applicable.

CRedit authorship contribution statement

Ayman Ahmed: Conceived, design, analyzed, wrote the first draft, revised and approved the final draft. **Maisa Aldai Hemaïda:** wrote the first draft, revised and approved the final draft. **Amel Ahmed Hagelnur:** wrote the first draft, revised and approved the final draft. **Hala Fathi Eltigani:** Conceived, design, analyzed, wrote the first draft, revised and approved the final draft. **Emmanuel Edwar Siddig:** Conceived, design, analyzed, wrote the first draft, revised and approved the final draft.

Conflict of Interest

The authors declare that they have no competing interests.

Data availability

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgements

Not applicable.

References

- [1] Heukelbach J, Feldmeier H. Epidemiological and clinical characteristics of hookworm-related cutaneous larva migrans. *Lancet Infect Dis* 2008;8:302–9.
- [2] Maxfield L, Crane J.S. Cutaneous Larva Migrans. [Updated 2022 Oct 12]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK507706/>.
- [3] Reichert F, Pilger D, Schuster A, Lesschaff H, Guedes de Oliveira S, Ignatius R, Feldmeier H. Prevalence and risk factors of hookworm-related cutaneous larva migrans (HrCLM) in a resource-poor community in Manaus, Brazil. *PLoS Negl Trop Dis* 2016;10(3):e0004514.
- [4] Tammimga N, Bierman WFW, de Vries PJ. Cutaneous larva migrans acquired in Brittany, France. *Emerg Infect Dis* 2009;15:1856–8.
- [5] Hla Aye M.T., Kyaw A.Y., Rubel A.R., Han M.B., Mani B.I., Chong V.H. Cutaneous Larva Migrans. *QJM*. 2022 Aug 12;hcac193.
- [6] Ju T, Vander Does A, Ingrassi G, Norton SA, Yosipovitch G. Tropical parasitic itch in returned travellers and immigrants from endemic areas. *J Eur Acad Dermatol Venereol* 2022;36(12):2279–90.
- [7] Ahmed A. Urgent call for a global enforcement of the public sharing of health emergencies data: lesson learned from serious arboviral disease epidemics in Sudan. *Int Health* 2020;12(4):238–40.
- [8] Aljak ER, Eldigail M, Mahmoud I, Elhassan RM, Elduma A, Ibrahim AA, Ali Y, Weaver SC, Ahmed A. The first laboratory-confirmed imported infections of SARS-CoV-2 in Sudan. *Trans R Soc Trop Med Hyg* 2021;115(1):103–9.
- [9] Ahmed A, Elbashir A, Mohamed AA, et al. Socioeconomic impacts of elimination of onchocerciasis in Abu-Hamed focus, northern Sudan: lessons after elimination. *BMC Res Notes* 2020;13:256. <https://doi.org/10.1186/s13104-020-05101-6>.
- [10] Ahmed A, Eldigail M, Elduma A, Breima T, Dietrich I, Ali Y, Weaver SC. First report of epidemic dengue fever and malaria co-infections among internally displaced persons in humanitarian camps of North Darfur, Sudan. *Int J Infect Dis* 2021;108:513–6.
- [11] Ahmed A, Mahmoud I, Eldigail M, Elhassan RM, Weaver SC. The emergence of Rift Valley Fever in gedaref state urges the need for a cross-border one health strategy and enforcement of the international health regulations. *Pathogens* 2021;10(7):885.
- [12] Stufano A, Foti C, Lovreglio P, Romita P, De Marco A, Lia RP, Otranto D, Iatta R. Occupational risk of cutaneous larva migrans: A case report and a systematic literature review. *PLoS Negl Trop Dis* 2022;16(5):e0010330.
- [13] Sow D, Soro F, Javelle E, Simon F, Parola P, Gautret P. Epidemiological profile of cutaneous larva migrans in travelers returning to France between 2003 and 2015. *Travel Med Infect Dis* 2017;20:61–4.
- [14] Heukelbach J, Gomide M, Araújo FJR, Pinto NS, Santana RD, Brito JR, Feldmeier H. Cutaneous larva migrans and tungiasis in international travelers exiting Brazil: an airport survey. *J Travel Med* 2007;14(6):374–80.
- [15] Gill N, Somayaji R, Vaughan S. Exploring tropical infections: a focus on cutaneous larva migrans. *Adv Ski Wound Care* 2020;33(7):356–9.
- [16] Coello RD, Pazmiño BJ, Reyes EO, Rodríguez EX, Rodas EI, Dávila AX, Rodas JP, Cedeño PP. A case of cutaneous larva migrans in a child from Vinces, Ecuador. *Am J Case Rep* 2019;20:1402–6.
- [17] González FCG, Galilea ONM, Pizarro CK. Larva migrans cutánea autóctona en Chile. A propósito de un caso [Autochthonous cutaneous larva migrans in Chile. A case report]. *Rev Chil Pedia* 2015;86(6):426–9.
- [18] Ahmed A, Mohamed NS, Siddig EE, Algaily T, Sulaiman S, Ali Y. The impacts of climate change on displaced populations: a call for action. *J Clim Change Health* 2021;3:100057.
- [19] Ahmed A, Ali Y, Elduma A, Eldigail MH, Mhmoud RA, Mohamed NS, Ksiazek TG, Dietrich I, Weaver SC. Unique outbreak of Rift Valley fever in Sudan, 2019. *Emerg Infect Dis* 2020;12:3030.
- [20] Zinsstag J, Hediger K, Osman YM, Abukhattab S, Crump L, Kaiser-Grolimund A, Mauti S, Ahmed A, Hattendorf J, Bonfoh B, Heitz-Tokpa K. The promotion and development of one health at Swiss TPH and its greater potential. *Diseases* 2022;10(3):65.