



Draft Genome Sequence of *Xanthobacter tagetidis* ATCC 700314^{T}

Syeda A. Fatima,^a (DAbigail E. Goen,^b (DKyle S. MacLea^{a,b,c}

Biotechnology Program, University of New Hampshire, Manchester, New Hampshire, USA
Biology Program, University of New Hampshire, Manchester, New Hampshire, USA
CDepartment of Life Sciences, University of New Hampshire, Manchester, New Hampshire, USA

ABSTRACT Xanthobacter tagetidis is a thiophene-degrading bacterium associated with root balls of the plant genus *Tagetes*, which includes marigolds. It is a Gramnegative facultatively autotrophic bacterium with pleomorphic morphology exhibiting bent and branching rods. From strain TagT2C^T (= ATCC 700314^T), we report a genome assembly of 4,945,221 bp and a 69.5% G+C content.

Xanthobacter tagetidis TagT2C^T (= ATCC 700314^T) was isolated from the root balls of the French marigold *Tagetes patula* and was later found in the root balls and soil compost of other members of the genus *Tagetes*, including the African marigold *T. erecta* and the South American marigold *T. minuta*, known in Quechua as wakatay. *X. tagetidis* is a Gram-negative yellow-pigmented facultatively autotrophic bacterium that grows optimally at 28 to 30°C (1). Like other members of the genus *Xanthobacter*, *X. tagetidis* exhibits pleomorphism in its morphology, with its typical rod shape exhibiting substantial irregularity, including bends and branching (1–3). Because the host *Tagetes* species produce copious and diverse thiophene compounds, *X. tagetidis* is metabolically versatile and can grow as a chemolithotrophic autotroph on thiosulfate and other inorganic sulfur compounds, as a heterotroph on thiophene-2-carboxylic acid, acetic acid, and α -ketoglutaric acid, and as a mixotroph on thiosulfate in combination with thiophene-2-carboxylic acid and/or acetic acid (2). This ability has been noted for its potential and demonstrated utility in the biodegradation of sulfur compounds, including anthropogenic pollutants, in the soil (1, 2, 4, 5).

Lyophilized Xanthobacter tagetidis ATCC 700314^T was obtained from ATCC (Manassas, VA, USA). Bacteria from a pure culture were grown in tryptic soy broth at 30°C for 48 hours. Genomic DNA was extracted using a DNA minikit (Qiagen, Valencia, CA, USA), and DNA quality and quantity were tested using the DeNovix double-stranded DNA (dsDNA) broad-range assay kit (DeNovix, Wilmington, DE, USA). Purified DNA was fragmented and tagged using the Nextera DNA library prep kit (Illumina, San Diego, CA, USA). The library DNA was then sequenced on an Illumina HiSeq 2500 instrument at the Hubbard Center for Genome Studies at the University of New Hampshire (Durham, NH, USA). The 250-bp reads were trimmed computationally using Trimmomatic (6) in paired-end mode with a window size of 4, quality requirement of 15, and minimum read length of 36. SPAdes version 3.11.0 (7) with default settings was used to assemble the trimmed reads into contigs. After the removal of small contigs of <500 bp, the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (8) was used to predict genes and annotate them from the 87 contigs in this draft genome of X. tagetidis. The genome had a total sequence length of 4,945,221 bp with a 69.5% G+C content, somewhat smaller than its closest sequenced relative, Xanthobacter autotrophicus Py2 (5,308,894 bp and a 67.5% G+C content on the main chromosome). The total number of genes was 4,556, out of which 4,447 were protein-coding genes. It also contained 53 RNA genes (3 rRNAs,

Citation Fatima SA, Goen AE, MacLea KS. 2019. Draft genome sequence of *Xanthobacter tagetidis* ATCC 700314^T. Microbiol Resour Announc 8:e00242-19. https://doi.org/10.1128/ MRA.00242-19.

Editor Julia A. Maresca, University of Delaware Copyright © 2019 Fatima et al. This is an openaccess article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Kyle S. MacLea, Kyle.MacLea@UNH.edu.

Received 6 March 2019 Accepted 19 June 2019 Published 11 July 2019 46 tRNAs, and 4 noncoding RNAs [ncRNAs]) and 56 pseudogenes. The largest contig was found to be 613,008 bp long, and the N_{50} value was 229,695 bp.

Data availability. The *Xanthobacter tagetidis* ATCC 700314^T whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number RCTF00000000. The version described in this paper is the first version, RCTF01000000. The raw Illumina data from BioProject PRJNA495830 were submitted to the NCBI Sequence Read Archive (SRA) under the project accession number SRP187584 and the SRA run accession number SRR8670595.

ACKNOWLEDGMENTS

Sequencing and bioinformatics analysis were undertaken at the Hubbard Center for Genome Studies at UNH, supported by NH-INBRE, with the kind assistance of Kelley Thomas, Devin Thomas, Jordan Ramsdell, Toni Westbrook, and Stephen Simpson. K.S.M. thanks Kate Emma Silvia for her hard work and helpful discussions. This work was a project of the Microbiology Education through Genome Annotation-New Hampshire (MEGA-NH) program.

UNH Manchester provided funds for purchasing the bacterial strain through a Manchester undergraduate project support grant to S.A.F. Sequencing costs were supported by New Hampshire-INBRE through an Institutional Development Award (IDeA), P20GM103506, from the National Institute of General Medical Sciences of the NIH. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

- Padden AN, Rainey FA, Kelly DP, Wood AP. 1997. Xanthobacter tagetidis sp. nov., an organism associated with Tagetes species and able to grow on substituted thiophenes. Int J Syst Bacteriol 47:394–401. https://doi .org/10.1099/00207713-47-2-394.
- Padden AN, Kelly DP, Wood AP. 1998. Chemolithoautotrophy and mixotrophy in the thiophene-2-carboxylic acid-utilizing Xanthobacter tagetidis. Arch Microbiol 169:249–256. https://doi.org/10.1007/s002030050568.
- Doronina NV, Trotsenko YA. 2003. Reclassification of "Blastobacter viscosus" 7d and "Blastobacter aminooxidans" 14a as Xanthobacter viscosus sp. nov. and Xanthobacter aminoxidans sp. nov. Int J Syst Evol Microbiol 53:179–182. https://doi.org/10.1099/ijs.0.02231-0.
- Stubner S, Wind T, Conrad R. 1998. Sulfur oxidation in rice field soil: activity, enumeration, isolation and characterization of thiosulfate-oxidizing bacteria. Syst Appl Microbiol 21:569–578. https://doi.org/10.1016/S0723-2020(98) 80069-6.
- 5. Ali N, Dashti N, Salamah S, Sorkhoh N, Al-Awadhi H, Radwan S. 2016.

Dynamics of bacterial populations during bench-scale bioremediation of oily seawater and desert soil bioaugmented with coastal microbial mats. Microb Biotechnol 9:157–171. https://doi.org/10.1111/1751-7915.12326.

- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30:2114–2120. https://doi.org/10 .1093/bioinformatics/btu170.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol 19:455–477. https://doi.org/10.1089/cmb.2012.0021.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Ciufo S, Li W. 2013. Prokaryotic Genome Annotation Pipeline, p 173–186. *In* Beck J, Benson D, Coleman J, Hoeppner M, Johnson M, Maglott M, Mizrachi I, Morris R, Ostell J, Pruitt K, Rubinstein W, Sayers E, Sirotkin K, Tatusova T (ed), The NCBI handbook, 2nd ed. NCBI, Bethesda, MD.