

Current Understanding of the Role of Nesfatin-1

Martha A. Schalla¹ and Andreas Stengel^{1,2}

¹Charité Center for Internal Medicine and Dermatology, Department for Psychosomatic Medicine, Charité-Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, 10117 Berlin; and ²Department of Psychosomatic Medicine and Psychotherapy, Medical University Hospital Tübingen, 72076 Tübingen, Germany

ORCID numbers: 0000-0003-3294-4340 (A. Stengel).

Nesfatin-1 was discovered in 2006 and implicated in the regulation of food intake. Subsequently, its widespread central and peripheral distribution gave rise to additional effects. Indeed, a multitude of actions were described, including modulation of gastrointestinal functions, glucose and lipid metabolism, thermogenesis, mediation of anxiety and depression, as well as cardiovascular and reproductive functions. Recent years have witnessed a great increase in our knowledge of these effects and their underlying mechanisms, which will be discussed in the present review. Lastly, gaps in knowledge will be highlighted to foster further studies.

Copyright © 2018 Endocrine Society

This article has been published under the terms of the Creative Commons Attribution Non-Commercial, No-Derivatives License (CC BY-NC-ND; <https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Freeform/Key Words: food intake, hypothalamus, nucleobindin 2, stomach

Nesfatin-1 is an 82–amino acid polypeptide derived from the precursor protein nucleobindin 2 (NUCB2), whose processing also yields nesfatin-2 and -3, two peptides with so far unknown functions [1]. The 29-amino acid mid-fragment of nesfatin-1, nesfatin-1₃₀₋₅₉, has been identified as the active core of nesfatin-1, also exerting an anorexigenic effect [2]. Although full-length nesfatin-1 contains cleavage sites at the respective amino acids, cleavage has not been shown *in vivo*.

NUCB2/nesfatin-1 (most antibodies do not distinguish between NUCB2 and processed nesfatin-1, and therefore we refer to the analyte as NUCB2/nesfatin-1) was first detected in food intake regulatory brain nuclei such as the paraventricular nucleus, arcuate nucleus, and nucleus of the solitary tract [1], and subsequent studies extended this distribution to numerous other brain nuclei in the rat [3] and mouse [4]. It is to note that NUCB2/nesfatin-1 has been detected primarily in the soma and primary dendrites of neurons, whereas less immunoreactivity was observed in nerve fibers [5], possibly pointing to an autocrine or paracrine rather than endocrine mode of action. Nonetheless, nesfatin-1 was shown to cross the blood-brain barrier in both directions [6, 7], supporting a humoral route of signaling. Further corroborating this assumption, NUCB2/nesfatin-1 has also been detected in the periphery, namely in the gastric mucosa [8], adipose tissue [9], pancreatic beta cells [10], testes [11], ovaries [12], uterus, epididymis [13] and cardiomyocytes [14]. The stomach was identified as main source of peripheral NUCB2/nesfatin-1, with NUCB2 mRNA levels higher

Abbreviations: AKT, protein kinase B; AMPK, AMP-activated protein kinase; BMI, body mass index; CREB, cAMP response element-binding protein; CRF, corticotropin-releasing factor; CRF₂, corticotropin-releasing factor receptor 2; DIO, diet-induced obesity; E, embryonic day; FGF21, fibroblast growth factor 21; GSK, glycogen synthase kinase; ICV, intracerebroventricular; JAK2, Janus kinase 2; m-TOR, mammalian target of rapamycin; NPY, neuropeptide Y; NUCB2, nucleobindin 2; p-ACC, phosphate/acetyl-CoA carboxylase; p-AMPK, phosphorylated 5' AMP-activated protein kinase; PI3K, phosphoinositide-3-kinase; pm-TOR, phosphorylated mammalian target of rapamycin; STAT3, signal transducer and activator of transcription 3; UCP1, uncoupling protein 1.

than in other peripheral organs or the brain [8]. Additional analyses indicated the expression of NUCB2/nesfatin-1 in gastric endocrine X/A-like cells coexpressed with the food intake-stimulating hormone ghrelin [8] in rats, a finding later confirmed in humans [15].

Despite the increasing knowledge of the expression and regulation of the ligand, the receptor mediating nesfatin-1's actions has not been identified. A recent study using autoradiography showed nesfatin-1 binding in the brain, including the cortex, paraventricular nucleus of the hypothalamus, area postrema, dorsal motor nucleus of the vagus nerve, and cerebellum as well as in peripheral endocrine organs, namely the pituitary, stomach, small intestine, pancreas, adrenal gland, testes, and visceral adipose tissue and also in the heart, skeletal muscle, lungs, liver and kidneys [16]. This widespread binding of ^{125}I -nesfatin-1 as a surrogate for the putative expression of the receptor further supports the assumption of a pleiotropic action of the peptide. On a molecular level evidence points toward the mediation of nesfatin-1's effects *via* a G_i protein-coupled receptor with nesfatin-1 inducing a Ca^{2+} inflow through L-type, N-type, or P/Q-type Ca^{2+} channels [17, 18] and promoting the phosphorylation of cAMP responding element-binding protein (CREB) in neuroblastoma cells *in vitro*, additionally stimulating mitogen-activated protein kinase signaling [19]. In contrast, in cardiomyocytes G_q is the supposed receptor type because nesfatin-1 suppressed L-type Ca^{2+} channel functioning by means of protein kinase C [20].

In the present review we describe the pleiotropic effects of nesfatin-1 and highlight underlying mechanisms with a focus on recent developments. We also discuss gaps in knowledge to demonstrate the need for further studies. A PubMed search has been performed and all studies with the terms “nesfatin-1,” “NUCB2,” and “nucleobindin 2” screened.

1. Implications of Nesfatin-1 in the Regulation of Food Intake

In the landmark study of Oh-I *et al.* [1], nesfatin-1 was described as anorexigenic peptide exerting a robust reduction of dark phase food intake after third ventricular injection, with a reduction of body weight gain after repeated injection. This effect is likely to be of physiological importance because blockade of endogenous nesfatin-1 with an antisense oligonucleotide [1] or knockdown of NUCB2 specifically in the paraventricular nucleus of the hypothalamus [21] resulted in a stimulation of food intake and a subsequent increase in body weight in rats [1, 21]. The anorexigenic effect after brain ventricular injection was confirmed in subsequent studies by independent groups in rats [22–24], mice [25, 26], pigs, [27], and goldfish [28], pointing toward a robust effect. Interestingly, a recent study using mice with NUCB2 knockout in several brain areas and peripheral tissues failed to demonstrate an effect on food intake and body weight compared with control mice, leading to the conclusion by the authors that nesfatin-1 does not affect food intake [29]. However, the authors did not check the expression of NUCB2/nesfatin-1 in the stomach, a major source of the peptide as described earlier. This expression might well explain the circulating levels of NUCB2/nesfatin-1 still observed in these mice. Lastly, the possibility of a compensatory action by other food intake regulatory peptides under conditions of long-term loss of a hormone also must be kept in mind.

In contrast to the brain effects, conflicting data exist on the peripheral effects of nesfatin-1, with some studies showing no effect [24], whereas others report a decrease in food intake after intraperitoneal (IP) injection of high doses in mice [2]. Similarly, chronic subcutaneous infusion of nesfatin-1 at high doses *via* osmotic minipumps was able to inhibit food intake in rats [30]. Overall, the anorexigenic effect is more readily observed after central injection of low doses of the peptide, giving rise to a predominantly central mode of anorexigenic action.

In addition to the food intake-reducing effect, intracerebroventricularly (ICV) injected nesfatin-1 was also shown to reduce water intake, an effect likely to present a physiological action of the peptide because pretreatment with an antinesfatin-1 antisense oligonucleotide led to an increased drinking response to angiotensin II [31].

Subsequent studies investigated the microstructure underlying the reduction of food intake by using an automated system that allows the continuous monitoring of food intake in

undisturbed rodents. In mice, ICV injected nesfatin-1 reduced meal size, indicating increased satiation and meal frequency associated with a prolongation of intermeal intervals, reflecting higher satiation [25]. Interestingly, ICV injected nesfatin-1_{30–59}, described as the active core of nesfatin-1 [2], increased satiety as well whereas satiation was not altered in mice [32], giving rise to differential receptor binding or activation. Lastly, in rats nesfatin-1_{30–59} induced satiation but not satiety [33], pointing toward species differences, a finding confirmed after microinjection of nesfatin-1 into the lateral parabrachial nucleus in rats [34]. Although the effect of nesfatin-1_{30–59} was retained during diet-induced obesity (DIO) in rats [33], giving rise to a leptin-independent signaling as suggested before [1, 2], the underlying microstructure differed, with increased satiety [33] suggesting different downstream signaling under conditions of obesity.

When nesfatin-1 was injected ICV in rats, its anorexigenic effect was abolished by pretreatment with the selective corticotropin-releasing factor (CRF) 2 antagonist astressin₂-B [24] indicating downstream mediation *via* corticotropin-releasing factor receptor 2 (CRF₂) signaling. This finding was also confirmed in chicks with astressin-B, a CRF_{1/2} antagonist [35]. Because the melanocortin receptor 3/4 antagonist SHU9119, with anorexigenic melanocortin 4 signaling being well established upstream of CRF [36, 37], attenuated [34] or blocked [1] nesfatin-1's anorexigenic effect, nesfatin-1 might act *via* melanocortin → CRF signaling to inhibit food intake.

Moreover, nesfatin-1 has been shown to inhibit neuropeptide Y (NPY)-containing cells *in vitro* [38]. Because NUCB2/nesfatin-1 partly colocalizes with NPY in the arcuate nucleus [39], this effect might involve an autocrine mode of signaling. Also, mammalian target of rapamycin (*m*-TOR) is greatly colocalized with NUCB2/nesfatin-1 in the arcuate nucleus [39]. Based on the finding that *m*-TOR has been shown to reduce NPY mRNA expression [40], nesfatin-1 might also signal *via* this pathway in an endocrine manner to reduce food intake. Peripherally, a decrease of *m*-TOR signaling has been shown to reduce the expression of NUCB2/nesfatin-1 *in vitro* and *in vivo* [41], whereas an increase of gastric phosphorylated mammalian target of rapamycin (pm-TOR)/*m*-TOR expression was associated with an increased gastric secretion of NUCB2/nesfatin-1, resulting in elevated levels of circulating NUCB2/nesfatin-1 [42] probably involved in the observed suppression of food intake. Interestingly, in the brain a decrease of pm-TOR levels in the dorsal motor nucleus of the vagus nerve induced by fourth ventricular injection of nesfatin-1 was associated with a reduction of food intake [43], pointing toward a differential central or peripheral regulation of *m*-TOR.

Also, fibroblast growth factor 21 (FGF21) is likely to be involved in the mediation of nesfatin-1's anorexigenic effect because FGF21 increased the expression of NUCB2 mRNA in the paraventricular nucleus, increased [Ca²⁺]_i in NUCB2/nesfatin-1-containing neurons, and the suppression of food intake induced by ICV injection of FGF21 was absent in NUCB2 knockout mice [44].

Nesfatin-1 has been recently implicated in the endogenous inhibition of food intake induced by IP application of cisplatin because cisplatin-activated NUCB2/nesfatin-1 neurons in the hypothalamus and brainstem and the cisplatin-induced reduction of food intake were attenuated by ICV injection of nesfatin-1/NUCB2-antisense [45], giving rise to a pathophysiological role of nesfatin-1 in chemotherapy-induced hypophagia. Similarly, nesfatin-1 has also been implicated in the oxytocin-induced suppression of food intake because IP injected oxytocin increased the number of activated NUCB2/nesfatin-1 neurons in the paraventricular nucleus, arcuate nucleus, and nucleus of the solitary tract, and the oxytocin-induced anorexigenic effect was attenuated by ICV injection of antisense nesfatin-1 [46]. Conversely, ICV injection of nesfatin-1 activated oxytocin-positive neurons in the paraventricular nucleus of the hypothalamus, as assessed with *c*-Fos and stimulated the release of oxytocin *in vitro* [47]. Lastly, an oxytocin receptor antagonist blocked the anorexigenic effect of nesfatin-1 [22, 47] indicating a strong interaction between nesfatin-1 and oxytocin in the inhibition of food intake.

Repeated injections of nesfatin-1 into the third brain ventricle [1] or the lateral parabrachial nucleus [34] decreased body weight gain, probably *via* a stimulation of uncoupling protein 1 (UCP1) expression in brown adipose tissue [34].

The regulation of NUCB2/nesfatin-1 supports a role for this peptide as a physiological modulator of food intake. Whereas fasting led to a decrease in hypothalamic NUCB2 mRNA levels [1, 48], refeeding activated NUCB2/nesfatin-1-positive neurons in the supraoptic nucleus, associated with an increase in NUCB2 mRNA levels [48]. In line with these findings, plasma NUCB2/nesfatin-1 levels decreased during 24-hour fasting and were restored after refeeding in rats [8]. *In vitro*, glucose elevated NUCB2 mRNA expression and NUCB2/nesfatin-1 secretion from cultured stomach ghrelinoma cells [49], probably contributing to the observed increase in plasma levels. Interestingly, in goldfish intestinal cells NUCB2 mRNA was decreased by glucose *in vitro* [50], pointing toward species differences. Also, sex differences seem to play a role because male mice receiving a high-fat diet displayed a decrease in serum NUCB2/nesfatin-1 levels [49], whereas this diet did not alter gastric NUCB2 mRNA expression in female mice [51]. Lastly, in zebrafish receiving a high-fat diet, supplementation with *Lactobacillus rhamnosus* reduced gut NUCB2/nesfatin-1 levels [52], pointing toward a microbiota-associated regulation of nesfatin-1 to be further investigated.

In obese children, no difference in NUCB2/nesfatin-1 plasma levels was detected between the fasting and postprandial state [53]. Whether this also holds true under normal weight conditions remains to be investigated. Moreover, an investigation of dynamic meal-related changes in circulating NUCB2/nesfatin-1 in humans is still lacking. Chronic changes in body weight also affect the levels of circulating NUCB2/nesfatin-1, with decreased levels in underweight and anorexic subjects [54, 55]. Because rats with activity-based anorexia, an animal model for anorexia nervosa, showed an activation of nesfatin-1 immunoreactive neurons in several brain nuclei involved in the mediation of food intake, gastrointestinal functions, and the response to stress [56], nesfatin-1 might be implicated in the development or maintenance of this condition, a hypothesis to be further investigated.

Conversely, circulating NUCB2/nesfatin-1 levels were shown to be elevated in obese adults [9, 57], probably because of increased expression of NUCB2/nesfatin-1 in the stomach [15]. These circulating levels were observed to be decreased after bariatric surgery-associated weight loss [58]. However, other studies described a negative association of circulating NUCB2/nesfatin-1 and body mass index (BMI) [55, 59, 60] and an increase after bariatric surgery in humans [61] and also mice [62], giving rise to additional, confounding factors such as sex or comorbidities like diabetes. Also, chronic malnutrition in children was reported to upregulate serum NUCB2/nesfatin-1 concentrations [63]. Whether this effect reflects metabolism- or stress-associated changes remains to be investigated.

2. Implications of Nesfatin-1 in Gastrointestinal Functions

As observed earlier for several peptides regulating food intake, nesfatin-1 has been shown to alter gastrointestinal functions. Early on, an inhibitory action of nesfatin-1 on antral and duodenal motility [26] and a reduction of gastric emptying was shown after ICV injection [24], a finding later also observed after microinjection of the peptide into the central nucleus of the amygdala [64], basomedial amygdala [65], arcuate nucleus [66], paraventricular nucleus of the hypothalamus [67], lateral hypothalamus [68], and ventromedial hypothalamus in rodents [69], giving rise to the sites of action. Because gastric distension induced c-Fos expression in NUCB2/nesfatin-1-positive neurons not only in the nucleus of the solitary tract [70] but also in the ventromedial hypothalamus [71], and the injection of an antinesfatin-1 antibody prevented a nucleus accumbens-induced ventromedial hypothalamic neuronal firing [69], along with the finding that rats with lesioned ventromedial hypothalamic nuclei showed increased gastric emptying [72], ventromedial nucleus nesfatin-1 signaling might well play a role in the endogenous regulation of gastric motility.

Because intravenous (IV) application of nesfatin-1 was able to reduce gastric contractions in dogs [73], the effect might also be mediated in the periphery, as recently hypothesized, by the distribution of the receptor as assessed by autoradiography [16]. A recent study reported that chronic gastric electrical stimulation used as a novel attempt to treat intractable vomiting and nausea reduced circulating NUCB2/nesfatin-1 levels [74], possibly indicating

nesfatin-1 as pathophysiological contributor to this disease. This hypothesis should be followed up in future studies.

Besides an effect on motility, nesfatin-1 affects secretory functions: ICV injected nesfatin-1 reduced 2-deoxy-D-glucose-stimulated gastric acid production [75] associated with a lower expression of H^+/K^+ -ATPase in obese rats [76]. Moreover, nesfatin-1 seems to be implicated in inflammatory processes, because IP [77] or IV [78] injected nesfatin-1 improved healing in rat models of gastric ulcer, associated with a decrease in $TNF-\alpha$ and $IL-1\beta$ and mediated by downstream cyclooxygenase 2 signaling [79, 80].

3. Implications of Nesfatin-1 in Blood Glucose Homeostasis

Early on, an implication of nesfatin-1 in glucose homeostasis was suggested. Besides its expression in the stomach, NUCB2/nesfatin-1 has also been detected in endocrine islets of the pancreas [8, 81, 82] colocalized with insulin in rodents [10, 83], dogs [84], pigs [84], and humans [85].

In rats, glucose challenge led to a release of NUCB2/nesfatin-1 from isolated pancreatic islets *in vitro* [86], probably by activating L-type Ca^{2+} channels [87] and inhibiting $K_v2.1$ channels [88]; however, the release of insulin was more pronounced [83]. Nesfatin-1 further increased the glucose-stimulated insulin release *in vitro* [86], giving rise to a direct effect supported by the assumed nesfatin-1 receptor in the pancreas as suggested by autoradiography [16], an effect leading to a reduction in glucose levels in mice [85, 89]. The glucose-lowering effect after oral glucose intake [90] probably also involves a nesfatin-1-induced release of glucagon-like peptide 1, because preproglucagon mRNA was upregulated by nesfatin-1 in the enteroendocrine STC-1 cell line *in vitro* [91]. Moreover, subcutaneously infused nesfatin-1 increased insulin-stimulated phosphorylation of protein kinase B (AKT) in skeletal muscle, adipose tissue, and liver and glucose transporter type 4 membrane translocation in skeletal muscle and adipose tissue of mice [92]. IV infusion of nesfatin-1 in streptozotocin-induced type 2 diabetic mice decreased blood glucose and insulin resistance by increasing the expression of phosphorylated 5' AMP-activated protein kinase (*p*-AMPK) and acetyl-CoA carboxylase (*p*-ACC) of skeletal muscles [93]. Conversely, knockdown of hypothalamic NUCB2/nesfatin-1 increased hepatic glucose flux and reduced glucose uptake of peripheral tissues under conditions of normal weight and in obese rats, effects associated with decreased hepatic insulin receptors, insulin receptor substrate 1, and AKT kinase phosphorylation and phosphorylation of *m*-TOR, signal transducer and activator of transcription 3 (STAT3), and the suppressor of cytokine signaling 3 [94]. Simultaneously, hepatic glucose-6-phosphatase and phosphoenolpyruvate carboxykinase levels were increased [94]. Lastly, ICV injection of nesfatin-1 led to increased muscle glucose absorption, insulin receptor signaling through the AKT/AMP-activated protein kinase (AMPK)/transducer of regulated CREB protein 2 pathway, decreased gluconeogenesis and hepatic mRNA and protein expression and activity of phosphoenolpyruvate carboxykinase [95].

In immunohistochemical studies using c-Fos as marker for neuronal activity, hypoglycemia after peripheral insulin administration induced an activation of nesfatin-1-expressing neurons in the arcuate nucleus, paraventricular nucleus, lateral hypothalamic area, dorsal motor nucleus of the vagus, and nucleus of the solitary tract [96]. Retrograde tracing with fluorogold showed that those activated neurons in the dorsal motor nucleus of the vagus project to the pancreas and stomach, suggesting that nesfatin-1 is implicated in central glucose sensing and initiates the hormonal and physiological response to hypoglycemia [96]. Accordingly, nesfatin-1 inhibited almost 90% of the gastric distension inhibitory and activated >75% of gastric distension excitatory neurons in the dorsal vagal complex [97]. Similarly, nesfatin-1 injected into the lateral parabrachial nucleus activated the majority of glucose-inhibited glucosensing neurons [34]. Moreover, in the paraventricular nucleus glucose and insulin increased calcium concentrations in nesfatin-1 immunoreactive neurons, leading to activation of those initiating satiety [98]. In contrast, in the nucleus of solitary tract nesfatin-1 did not trigger the response of glucosensing neurons when the glycemic state changed [99].

Nesfatin-1's expression was altered under conditions of type 2 diabetes mellitus with reduced pancreatic NUCB2/nesfatin-1 peptide expression in rodents with genetically determined diabetes [83], whereas in DIO mice the NUCB2/nesfatin-1 peptide concentration was increased [86]. In humans, NUCB2 mRNA expression was reduced in islets of subjects with type 2 diabetes [85]. However, circulating NUCB2/nesfatin-1 levels were not altered under conditions of type 2 diabetes in rats [83], whereas in streptozotocin-induced type 2 diabetic mice [93] and in humans levels were higher in patients with type 2 diabetes or impaired glucose tolerance than in healthy controls [100], giving rise to species differences. However, in patients with metabolic syndrome, including insulin resistance, NUCB2/nesfatin-1 concentrations were lower than in controls [101]. A recent meta-analysis failed to show differences in circulating NUCB2/nesfatin-1 concentrations between controls and patients with type 2 diabetes [102]. Subgroup analyses showed significantly higher levels in patients newly diagnosed with type 2 diabetes without any treatment and lower levels in patients receiving antidiabetic medication [102], indicating an influence of medication on NUCB2/nesfatin-1 signaling. In line with this assumption, in a study of 100 patients with type 2 diabetes mellitus the oral glucose-lowering agent saxagliptin increased circulating NUCB2/nesfatin-1 along with C-peptide and improved insulin resistance as well as metabolic profile, BMI and blood pressure [103]. The observed negative correlation between nesfatin-1 and homeostatic model assessment for insulin resistance and hemoglobin A1c [103] emphasizes the beneficial effect of nesfatin-1 on glucose homeostasis.

The c.1012C>G polymorphism of the NUCB2 gene has been associated with a reduced risk of type 2 diabetes in the Chinese Han population [104]. In patients with type 2 diabetes mellitus the GG phenotype was related to lower BMI and fasting plasma glucose levels [104]. In accordance, in patients with metabolic syndrome the CG and GG genotypes were found less frequently, giving rise to a reduced risk of metabolic syndrome associated with the GG genotype and G allele [105].

In patients with gestational diabetes mellitus serum NUCB2/nesfatin-1 concentrations, NUCB2/nesfatin-1 levels in cord blood and protein expression of NUCB2/nesfatin-1 in subcutaneous fat were higher than in healthy pregnant subjects [106]. Inconsistently, a previous [107] and a subsequent [108] study showed the opposite regulation, with decreased serum NUCB2/nesfatin-1 levels and a positive association between NUCB2/nesfatin-1 and gestational week [108]. These discrepant results might be due to, although subtle, differences in age (29.6 ± 5.3 vs. 32.1 ± 6.2 years), BMI (31.0 ± 5.5 vs. 33.8 ± 6.5 kg/m²), and advancement of pregnancy (26.2 ± 1.8 vs. 23.0 ± 7.8 weeks), to be considered in future studies.

Laparoscopic Roux-en-Y gastric bypass or sleeve gastrectomy was shown to reduce BMI, waist circumference, hip circumference, waist-to-hip ratio, fasting blood glucose, and hemoglobin A1c levels, homeostatic model assessment for insulin resistance, and plasma insulin C-peptide after 1 year in patients with type 2 diabetes [109]. Glucagon stimulation reduced plasma levels of NUCB2/nesfatin-1, with higher levels observed in nonremitters compared with remitters [109]. Whether nesfatin-1 plays a pathogenetic role under these conditions or is compensatorily increased to improve glucose homeostasis will have to be further investigated in longitudinal studies.

4. Implications of Nesfatin-1 in Energy Metabolism

It was shown early on that nesfatin-1 is able to elevate core body temperature in rats after ICV injection [23], also modulating energy expenditure, dry heat, brown adipose tissue growth, and tail temperature, probably mediated *via* downstream melanocortin 3/4 signaling [110]. Moreover, cold ambient temperature activated nesfatin-1 immunoreactive neurons in the paraventricular nucleus [23]. Because knockout of the transcription factor Yin Yang 1 increased energy expenditure and oxygen consumption in beige and white fat depots along with an increased expression of NUCB2 mRNA in brown adipose tissue [111], NUCB2/nesfatin-1 might be physiologically involved in the regulation of energy expenditure. Lastly, nesfatin-1 was shown to stimulate brown adipocyte differentiation *via* m-TOR signaling and

upregulation of UCP1 mRNA expression [112]; therefore, nesfatin-1 might well be involved in long-term changes of energy expenditure. Because nesfatin-1 reduces food intake and increases energy expenditure, it induces a negative energy balance, which might be relevant in states of overnutrition but also might reflect conditions of stress, ultimately leading to wasting or cachexia, a hypothesis to be further investigated.

With regard to lipid metabolism, chronic subcutaneous infusion of nesfatin-1 reduced cholesterol and triglyceride levels in mouse plasma, an effect independent of food intake and body weight [113]. In diabetic mice, IV injected nesfatin-1 was able to normalize free fatty acids and body weight after 6 days, accompanied by an elevation of *p*-AMPK and *p*-ACC expression levels in skeletal muscle, indicating improved free fatty acid utilization [93]. Moreover, lipid accumulation was reduced by nesfatin-1 in cultured hepatocytes *in vitro*, probably because of a decrease of lipogenesis-relevant genes such as peroxisome proliferator-activated receptor- γ and sterol-regulatory element-binding protein 1 and enzymes including fatty acid synthase and glycerol-3-phosphatase acyltransferase, whereas β -oxidation-related genes were increased [113]. Because circulating levels of NUCB2/nesfatin-1 were reduced in patients with nonalcoholic fatty liver disease [114], the peptide might play a pathogenetic role under these conditions. This role should be further investigated along with the therapeutic potential in this disease.

5. Implications of Nesfatin-1 in the Development of Anxiety and Depression

Nesfatin-1 was shown to exert an anxiogenic effect after ICV injection in rats [115], a finding subsequently confirmed after IP injection [116]. A rat model inducing anxiety by using maternal separation and acute gastric irritation in early life and sequential stress in adulthood displayed increased NUCB2/nesfatin-1 peptide levels in the hippocampus, gastric fundus, and plasma, possibly contributing to the increased anxiety under these conditions [117]. Interestingly, in humans a correlation between circulating NUCB2/nesfatin-1 levels and self-reported anxiety was observed, with a positive association in women [118, 119], whereas in men this association was negative [120], pointing toward a sex-specific regulation of the peptide. However, the reduction of anxiety under inpatient treatment conditions did not significantly alter circulating NUCB2/nesfatin-1 levels [121]. Whether a longer (>4 weeks) observation time is necessary or whether patients with more severe anxiety disorders should be included to detect changes in NUCB2/nesfatin-1 warrants further investigation. Because anxiety and food intake are intimately linked, it might also be speculated that nesfatin-1 primarily affects anxiety and subsequently food intake. This speculation should be followed up, for example, by assessment of food intake after coapplication of nesfatin-1 and anxiolytic drugs.

Besides the correlation with self-reported anxiety, circulating NUCB2/nesfatin-1 was also associated with reported depression in a mixed-sex population [122], with NUCB2/nesfatin-1 levels rising with increasing severity of depression [123, 124], and correlated with inflammatory markers such as IL-6 and C-reactive protein as well as corticosterone under these conditions [125]. Higher NUCB2/nesfatin-1 levels were subsequently shown in patients reporting depression associated with subclinical hypothyroidism [126], leading to the hypothesis of nesfatin-1 being involved in these symptoms. Also, an animal model of gastric cancer comorbid with depression displayed higher NUCB2/nesfatin-1 peptide levels in the hippocampus, midbrain and plasma compared with undisturbed controls [127]. Nesfatin-1, at least based on animal data, is likely to play a role in the development of depressive symptoms, because ICV injection of the peptide was shown to reduce the consumption of a palatable snack in a novel environment as a surrogate for anhedonic behavior in rats [115]. This finding was subsequently extended to a peripheral effect as rats showed increased immobility in the forced swim test after IP injection of the peptide [116]. Whether nesfatin-1 plays a causal role in the development of depressive symptoms in humans as well remains to be investigated.

Interestingly, a positive association between circulating NUCB2/nesfatin-1 and depression has been described in women [118, 119], whereas in men this correlation was absent [120]. The

sex-specific alteration of the peptide was also described in a study on depression-associated suicide victims: while NUCB2 mRNA expression in the Edinger-Westphal nucleus was elevated in men, it was lower in women than in control subjects who died without any diagnosed neurodegenerative or psychiatric disorder [128]. This sex-specific regulation, possibly associated with a sex steroid dependency, as described below, should be followed up in future studies.

6. Implications of Nesfatin-1 in Cardiovascular Functions

ICV injection of nesfatin-1 was shown to increase blood pressure in rats [129, 130], an effect also observed after microinjection into the paraventricular nucleus of the hypothalamus [131] and probably mediated by increased central parasympathetic tone [132], assumed to mediate the observed bradycardia [133], and by sympathetic nerve outflow [134], and blocked by pretreatment with the melanocortin 3/4 antagonist SHU9119 [129, 130] or the α -adrenergic antagonist phentolamine [130]. Moreover, pretreatment with an oxytocin antagonist, ornithine vasotocin [22], or a CRF₂ antagonist, astressin₂-B [135], prevented nesfatin-1's hypertensive effect. Because the hypertensive action of CRF, but not α -melanocyte-stimulating hormone, was prevented by pretreatment with ornithine vasotocin [135], nesfatin-1 might act *via* downstream CRF \rightarrow oxytocin \rightarrow melanocortin 3/4 signaling to increase blood pressure. Also, microinjection of nesfatin-1 into the nucleus of the solitary tract increased blood pressure in rats [136]. Further corroborating the role of nesfatin-1 in the regulation of blood pressure, mice overexpressing NUCB2 displayed an increased systolic, diastolic, and mean blood pressure [137]. Conversely, knockdown of NUCB2 specifically in the paraventricular nucleus blunted the high-salt diet-stimulated increase in systolic blood pressure [21]. Likewise, silencing of paraventricular NUCB2/nesfatin-1 signaling counteracted the rise in systolic blood pressure observed in agouti-related peptide/3-phosphoinositide-dependent protein kinase-1 knockout mice [138].

Peripheral nesfatin-1 exerts pronounced effects on the cardiovascular system, namely an increase in blood pressure after IP administration in mice [139] and rats [140] or IV injection in rats [141], with inhibited relaxation of peripheral blood vessels probably contributing to this effect [141]. This effect is likely associated with stimulated signaling of the phosphoinositide-3-kinase (PI3K)/AKT/*m*-TOR pathway and phosphorylation of Janus kinase 2 (JAK2)/STAT3, resulting in proliferation, migration, and phenotype switch of vascular smooth muscle cells from a contractile to a synthetic state by elevating the mRNA and protein expression of matrix metalloproteinase 2 and 9 while reducing peroxisome proliferator-activated receptor- γ [142, 143]. Because NUCB2 mRNA expression was increased in the media of the aorta of spontaneously hypertensive rats [142], nesfatin-1 might play a pathogenetic role under these conditions. In line with this assumption, circulating NUCB2/nesfatin-1 levels were higher in patients with essential hypertension than in normotensive controls and correlated with systolic blood pressure [144]; therefore, nesfatin-1 has been suggested as a risk factor for obesity-associated hypertension (OR 1.5) [145]. Also, patients with polycystic ovary syndrome displayed increased circulating NUCB2/nesfatin-1 concentrations, which correlated with systolic and diastolic blood pressure levels [146].

In the heart nesfatin-1 decreases contractility and relaxation as assessed in isolated rat heart preparations [147]. In zebrafish, IP injected nesfatin-1 reduced end diastolic volume and cardiac output associated with decreased heart rate [148], whereas in rat [149] and goldfish [150] a positive inotropic effect was observed *ex vivo* [150], possibly reflecting the difference between whole body and organ or tissue conditions. These effects might well be locally mediated, because NUCB2 mRNA expression has been detected in the heart of mouse [14], rat [14], zebrafish [148], and human [14], a finding corroborated by NUCB2/nesfatin-1 protein expression in rat and human cardiomyocytes [14]. Because the receptor is assumed to be expressed in the heart, as recently suggested by autoradiography in rats [16], nesfatin-1 might act in the heart in an autocrine or paracrine manner.

The cardioprotective effects were first suggested after *ex vivo* experiments in Langendorff-perfused rat heart preparations, as indicated by reduced infarct size, lactate dehydrogenase

release and posts ischemic contracture [147]. These effects are probably mediated by myocardial upregulation of *p*-AKT/AKT and *p*-glycogen synthase kinase (GSK)-3 β /GSK-3 β , resulting in a reduction of apoptotic and necrotic cells under conditions of isoproterenol-induced myocardial infarction [151]. Consequently, cardiac troponin and proinflammatory cytokines were lower in rats receiving IP nesfatin-1 than in those treated with vehicle [151]. However, another study reported a nesfatin-1-induced apoptosis of cardiomyocytes in isolated neonatal rat hearts mediated *via* reduced AKT inactivation and increased expression of apoptogenic protein 1 and caspase-3 [152]. Whether this discrepancy is caused by dosage differences (1 μ M *vs.* 100 pM/L) warrants further investigation. Lastly, in a human study investigating subjects with rheumatoid arthritis, nesfatin-1 correlated positively with rheumatoid factor, matrix metalloproteinase-2, and plaque stability mediator and negatively with carotid intima-media thickness [153], giving rise to a protective effect of nesfatin-1 under these conditions.

7. Implications of Nesfatin-1 in Reproductive Functions

Expression of NUCB2 mRNA has been detected in rat [11], mouse [11], dog [154], and human [11] testes, and NUCB2/nesfatin-1 protein has been identified in the interstitium next to the seminiferous tubules [8], probably representing interstitial Leydig cells, with increased levels in the transition from puberty to adulthood stimulated by pituitary LH [11]. Similarly, in female mice NUCB2 mRNA expression was detected in the ovary, with NUCB2/nesfatin-1 peptide expression in the theca and interstitial cells of the ovary [13], the peptide expression has been confirmed in birds [155]. In female rats hypothalamic NUCB2 mRNA levels and NUCB2/nesfatin-1 protein content increased during pubertal transition [156]. Also, in humans circulating NUCB2/nesfatin-1 levels increased in girls with premature thelarche compared with the prepubertal state [157], leading to the hypothesis of a role for nesfatin-1 in gonadal development.

Indeed, ICV injection of nesfatin-1 resulted in an elevation of circulating gonadotropins, whereas ICV infusion of an anti-NUCB2 antisense oligonucleotide delayed vaginal opening and decreased ovarian weights as a sign of hampered puberty in these animals [156].

In male rats, nesfatin-1 increased human chorionic gonadotropin-stimulated testosterone secretion from testicular explants *ex vivo* [11]. ICV injection of nesfatin-1 resulted in a reduced expression of hypothalamic GnRH, kisspeptin, pituitary FSH β , LH β , and testicular steroidogenic acute regulatory protein, whereas it markedly elevated the expression of 3 β -hydroxysteroid dehydrogenase, 17 β -hydroxysteroid dehydrogenase, and cytochrome P450 mRNA in the testes of pubertal rats [158]. Conversely, testosterone stimulated mouse NUCB2 mRNA and NUCB2/nesfatin-1 protein expression in hypothalamic and pituitary cells *in vitro* [159]. Further corroborating the testosterone-dependent regulation of nesfatin-1, castrated mice displayed reduced NUCB2 mRNA expression in the pituitary, a finding reversed by testosterone treatment [160].

Immunohistochemical expression analysis of hemochorial mouse and human placenta demonstrated a wide distribution of NUCB2/nesfatin-1, namely in the ectoplacental zone, parietal trophoblast giant cells and early spongiotrophoblast from embryonic day (E) 7.5 to E9.5 and additionally at E10.5 to E12.5 in the developing labyrinth [161]. Glycogen trophoblast cells, syncytiotrophoblast, sinusoidal trophoblast giant cells and fetal capillary endothelial cells of the labyrinth expressed NUCB2/nesfatin-1 at high levels from E12.5 [161]. In human pregnancy, nesfatin-1 was highly expressed in syncytiotrophoblast throughout all three trimesters [161], giving rise to a modulating role of nesfatin-1 in glucose homeostasis also during pregnancy. Nesfatin-1 was also recently implicated in the maintenance of pregnancy because uterine expression levels of NUCB2 mRNA and NUCB2/nesfatin-1 protein levels were increased in a mouse spontaneous abortion model [162], a hypothesis to be further investigated.

8. Conclusions

The present review highlights nesfatin-1 as a pleotropic polypeptide (Table 1) with a well-established anorexigenic action and an inhibitory effect on gastrointestinal motility acting on

Table 1. Central and Peripheral Effects of Nesfatin-1

	Central Effects	Mediation	Peripheral Effects	Mediation
Feeding behavior	Food intake ↓ (acute) [1, 21, 24], water intake ↑ [31]	CRF ₂ - [24] and histamine- [35], melanocortin- [1] dependent [35], pm-TOR-related [67]	Food intake ↓/= (acute and chronic) [2, 30]	Leptin-independent [2]
Gastrointestinal functions	Gastric emptying ↓ [24, 67] Gastric motility ↓ [69] Acid output ↓ [69]	Melanocortin receptor-dependent ↓ [67] H+/K+-ATPase expression ↓ [76]	Gastric contractions ↓ [73] Gastric ulcer healing ↑ [78–80]	Cyclooxygenase 2-dependent [79]
Glucose homeostasis and metabolism	Muscle glucose absorption, insulin receptor signaling ↑ [95], gluconeogenesis ↓ [95], hypothalamic glucosensing neurons ↑/↓ [34, 97], core body temperature ↑ [23], energy expenditure ↑ [110], dry heat ↓ [110], brown adipose tissue ↑ [110], tail temperature ↑ [110], body weight gain (chronic) ↓ [1, 34]	AKT/AMPK/transducer of regulated CREB protein 2 [95], melanocortin 3/4 signaling [110], UCP-1 expression and <i>m</i> -TOR signaling [112], melanocortin system, UCP1 expression [34]	Glucose-stimulated insulin secretion ↑ [90], glucagon secretion ↑ [85], insulin sensitivity ↑ [90], blood glucose ↓ [92], blood glucose and insulin resistance ↓ [93], brown adipocyte differentiation ↑ [112], UCP1 expression ↑ [112], plasma cholesterol + triacylglyceride ↓ [113], lipid accumulation ↓ [113], free fatty utilization ↑ [92]	K _v 2.1 channel-dependent [88], AMPK and <i>p</i> -ACC phosphorylation [93], <i>m</i> -TOR signaling [112], AMPK phosphorylation [113]
Anxiety and depression	Anxiety and depressive-like behavior (acute) ↑ [115]		Anxiety-like behavior (chronic) ↑ [116]	
Cardiovascular functions	Blood pressure ↑ [130–132, 136], heart rate ↑/↓ [132], [133]	CRF- [22], sympathetic- [134], parasympathetic- [132, 133], melanocortin- [22] dependent	Blood pressure ↑ [139, 140], contractility of heart and vessels ↑ [141, 147], end diastolic volume, cardiac output, heart rate ↓ [148], aorta relaxation ↑ [149, 150], cardioprotection ↑ [147, 151], apoptosis of cardiomyocytes ↑/↓ [151, 152], proliferation, migration, phenotype switch of vascular smooth muscle cells ↑ [143]	PI3K/AKT/ <i>m</i> -TOR, JAK2/STAT3 signaling [142], endothelium, NO- and guanylatecyclase-dependent [149], <i>p</i> -AKT/AKT and <i>p</i> -GSK-3β/GSK-3β [143], AKT, apoptogenic protein 1 and caspase-3 [152], PI3K/AKT/ <i>m</i> -TOR pathway and phosphorylation of JAK2/STAT3 [142]
Reproductive functions	Pubertal transition ↑ [156]	Circulating gonadotropins ↑ [156, 158], expression of genes for GnRH, kisspeptin, FSH, LH, testicular steroidogenic acute regulatory protein ↓ [158], expression of genes for 3β-HSD, 17β-HSD, cytochrome P450 ↑ [158]	Human chorionic gonadotropin-dependent testosterone secretion ↑ [158]	

Abbreviations: ↑, increase/stimulation; ↓, decrease/inhibition; =, no effect; NO, nitric oxide.




Figure 1. Central effects of nesfatin-1. Mediation of nesfatin-1's effects in different brain nuclei. The blue areas depict central nesfatin-1 binding sites as assessed by autoradiography, giving rise to the expression of the nesfatin-1 receptor [16]. ↑, increase/stimulation; ↓, decrease/inhibition.

different brain nuclei (Fig. 1). Moreover, nesfatin-1 plays a role in glucose homeostasis as a negative regulator of glucose levels and is also involved in energy expenditure by increasing thermogenesis. In addition, nesfatin-1 might be involved in the development of anxiety and depression as well, with a differential regulation in male and female patients highlighting the need for the study of sex differences. Its most striking peripheral effects are related to the cardiovascular system, thereby improving cardiac contractility, aorta relaxation, and cardioprotection. Lastly, peripheral nesfatin-1 is involved in gonadal development and during pregnancy. The identification of the yet unknown nesfatin-1 receptor will represent a great leap forward in the understanding of nesfatin-1's physiology and will allow us to better investigate the effects underlying the different actions summarized here.

Acknowledgments

We acknowledge support by Deutsche Forschungsgemeinschaft and Open Access Publishing Fund of Charité University Berlin.

Financial Support: This work was supported by funding of the German Research Foundation (STE 1765/3-2) and Charité University Funding (UFF 89/441-176, to A.S.).

Correspondence: Andreas Stengel, MD, PhD, Psychosomatic Medicine, Osianderstr. 5, 72076 Tübingen, Germany. E-mail: andreas.stengel@med.uni-tuebingen.de.

Disclosure Summary: The authors have nothing to disclose.

References and Notes

1. Oh-I S, Shimizu H, Satoh T, Okada S, Adachi S, Inoue K, Eguchi H, Yamamoto M, Imaki T, Hashimoto K, Tsuchiya T, Monden T, Horiguchi K, Yamada M, Mori M. Identification of nesfatin-1 as a satiety molecule in the hypothalamus. *Nature*. 2006;443(7112):709–712.

2. Shimizu H, Oh IS, Hashimoto K, Nakata M, Yamamoto S, Yoshida N, Eguchi H, Kato I, Inoue K, Satoh T, Okada S, Yamada M, Yada T, Mori M. Peripheral administration of nesfatin-1 reduces food intake in mice: the leptin-independent mechanism. *Endocrinology*. 2009;**150**(2):662–671.
3. Goebel M, Stengel A, Wang L, Lambrecht NW, Taché Y. Nesfatin-1 immunoreactivity in rat brain and spinal cord autonomic nuclei. *Neurosci Lett*. 2009;**452**(3):241–246.
4. Goebel-Stengel M, Wang L, Stengel A, Taché Y. Localization of nesfatin-1 neurons in the mouse brain and functional implication. *Brain Res*. 2011;**1396**:20–34.
5. Goebel-Stengel M, Wang L. Central and peripheral expression and distribution of NUCB2/nesfatin-1. *Curr Pharm Des*. 2013;**19**(39):6935–6940.
6. Pan W, Hsueh H, Kastin AJ. Nesfatin-1 crosses the blood-brain barrier without saturation. *Peptides*. 2007;**28**(11):2223–2228.
7. Price CJ, Hoyda TD, Samson WK, Ferguson AV. Nesfatin-1 influences the excitability of paraventricular nucleus neurons. *J Neuroendocrinol*. 2008;**20**(2):245–250.
8. Stengel A, Goebel M, Yakubov I, Wang L, Witcher D, Coskun T, Taché Y, Sachs G, Lambrecht NW. Identification and characterization of nesfatin-1 immunoreactivity in endocrine cell types of the rat gastric oxyntic mucosa. *Endocrinology*. 2009;**150**(1):232–238.
9. Ramanjaneya M, Chen J, Brown JE, Tripathi G, Hallschmid M, Patel S, Kern W, Hillhouse EW, Lehnert H, Tan BK, Rande HS. Identification of nesfatin-1 in human and murine adipose tissue: a novel depot-specific adipokine with increased levels in obesity. *Endocrinology*. 2010;**151**(7):3169–3180.
10. Gonzalez R, Tiwari A, Unniappan S. Pancreatic beta cells colocalize insulin and nesfatin immunoreactivity in rodents. *Biochem Biophys Res Commun*. 2009;**381**(4):643–648.
11. García-Galiano D, Pineda R, Ilhan T, Castellano JM, Ruiz-Pino F, Sánchez-Garrido MA, Vazquez MJ, Sangiao-Alvarellos S, Romero-Ruiz A, Pinilla L, Diéguez C, Gaytán F, Tena-Sempere M. Cellular distribution, regulated expression, and functional role of the anorexigenic peptide, NUCB2/nesfatin-1, in the testis. *Endocrinology*. 2012;**153**(4):1959–1971.
12. Gonzalez R, Shepperd E, Thiruppugazh V, Lohan S, Grey CL, Chang JP, Unniappan S. Nesfatin-1 regulates the hypothalamo-pituitary-ovarian axis of fish. *Biol Reprod*. 2012;**87**(4):84.
13. Kim J, Chung Y, Kim H, Im E, Lee H, Yang H. The tissue distribution of nesfatin-1/NUCB2 in mouse. *Dev Reprod*. 2014;**18**(4):301–309.
14. Feijóo-Bandín S, Rodríguez-Penas D, García-Rúa V, Mosquera-Leal A, Otero MF, Pereira E, Rubio J, Martínez I, Seoane LM, Gualillo O, Calaza M, García-Caballero T, Portolés M, Roselló-Lletí E, Diéguez C, Rivera M, González-Juanatey JR, Lago F. Nesfatin-1 in human and murine cardiomyocytes: synthesis, secretion, and mobilization of GLUT-4. *Endocrinology*. 2013;**154**(12):4757–4767.
15. Stengel A, Hofmann T, Goebel-Stengel M, Lembke V, Ahnis A, Elbelt U, Lambrecht NW, Ordemann J, Klapp BF, Kobelt P. Ghrelin and NUCB2/nesfatin-1 are expressed in the same gastric cell and differentially correlated with body mass index in obese subjects. *Histochem Cell Biol*. 2013;**139**(6):909–918.
16. Prinz P, Goebel-Stengel M, Teuffel P, Rose M, Klapp BF, Stengel A. Peripheral and central localization of the nesfatin-1 receptor using autoradiography in rats. *Biochem Biophys Res Commun*. 2016;**470**(3):521–527.
17. Brailoiu GC, Dun SL, Brailoiu E, Inan S, Yang J, Chang JK, Dun NJ. Nesfatin-1: distribution and interaction with a G protein-coupled receptor in the rat brain. *Endocrinology*. 2007;**148**(10):5088–5094.
18. Iwasaki Y, Nakabayashi H, Kakei M, Shimizu H, Mori M, Yada T. Nesfatin-1 evokes Ca²⁺ signaling in isolated vagal afferent neurons via Ca²⁺ influx through N-type channels. *Biochem Biophys Res Commun*. 2009;**390**(3):958–962.
19. Ishida E, Hashimoto K, Shimizu H, Okada S, Satoh T, Kato I, Yamada M, Mori M. Nesfatin-1 induces the phosphorylation levels of cAMP response element-binding protein for intracellular signaling in a neural cell line. *PLoS One*. 2012;**7**(12):e50918.
20. Ying J, Zhang Y, Gong S, Chang Z, Zhou X, Li H, Tao J, Zhang G. Nesfatin-1 suppresses cardiac L-type Ca²⁺ channels through melanocortin type 4 receptor and the novel protein kinase C theta isoform pathway. *Cell Physiol Biochem*. 2015;**36**(2):555–568.
21. Nakata M, Gantulga D, Santoso P, Zhang B, Masuda C, Mori M, Okada T, Yada T. Paraventricular NUCB2/nesfatin-1 supports oxytocin and vasopressin neurons to control feeding behavior and fluid balance in male mice. *Endocrinology*. 2016;**157**(6):2322–2332.
22. Yosten GL, Samson WK. The anorexigenic and hypertensive effects of nesfatin-1 are reversed by pretreatment with an oxytocin receptor antagonist. *Am J Physiol Regul Integr Comp Physiol*. 2010;**298**(6):R1642–R1647.

23. Könczöl K, Pintér O, Ferenczi S, Varga J, Kovács K, Palkovits M, Zelena D, Tóth ZE. Nesfatin-1 exerts long-term effect on food intake and body temperature. *Int J Obes (Lond)*. 2012;**36**(12):1514–1521.
24. Stengel A, Goebel M, Wang L, Rivier J, Kobelt P, Mönnikes H, Lambrecht NW, Taché Y. Central nesfatin-1 reduces dark-phase food intake and gastric emptying in rats: differential role of corticotropin-releasing factor2 receptor. *Endocrinology*. 2009;**150**(11):4911–4919.
25. Goebel M, Stengel A, Wang L, Taché Y. Central nesfatin-1 reduces the nocturnal food intake in mice by reducing meal size and increasing inter-meal intervals. *Peptides*. 2011;**32**(1):36–43.
26. Atsuchi K, Asakawa A, Ushikai M, Ataka K, Tsai M, Koyama K, Sato Y, Kato I, Fujimiya M, Inui A. Centrally administered nesfatin-1 inhibits feeding behaviour and gastroduodenal motility in mice. *Neuroreport*. 2010;**21**(15):1008–1011.
27. Lents CA, Barb CR, Hausman GJ, Nonneman D, Heidorn NL, Cisse RS, Azain MJ. Effects of nesfatin-1 on food intake and LH secretion in prepubertal gilts and genomic association of the porcine NUCB2 gene with growth traits. *Domest Anim Endocrinol*. 2013;**45**(2):89–97.
28. Gonzalez R, Kerbel B, Chun A, Unniappan S. Molecular, cellular and physiological evidences for the anorexigenic actions of nesfatin-1 in goldfish. *PLoS One*. 2010;**5**(12):e15201.
29. Ravussin A, Youm YH, Sander J, Ryu S, Nguyen K, Varela L, Shulman GI, Sidorov S, Horvath TL, Schultze JL, Dixit VD. Loss of nucleobindin-2 causes insulin resistance in obesity without impacting satiety or adiposity. *Cell Reports*. 2018;**24**(5):1085–1092.e6.
30. Mortazavi S, Gonzalez R, Ceddia R, Unniappan S. Long-term infusion of nesfatin-1 causes a sustained regulation of whole-body energy homeostasis of male Fischer 344 rats. *Front Cell Dev Biol*. 2015;**3**:22.
31. Yosten GL, Redlinger L, Samson WK. Evidence for a role of endogenous nesfatin-1 in the control of water drinking. *J Neuroendocrinol*. 2012;**24**(7):1078–1084.
32. Stengel A, Goebel-Stengel M, Wang L, Kato I, Mori M, Taché Y. Nesfatin-1(30-59) but not the N- and C-terminal fragments, nesfatin-1(1-29) and nesfatin-1(60-82) injected intracerebroventricularly decreases dark phase food intake by increasing inter-meal intervals in mice. *Peptides*. 2012;**35**(2):143–148.
33. Prinz P, Teuffel P, Lembke V, Kobelt P, Goebel-Stengel M, Hofmann T, Rose M, Klapp BF, Stengel A. Nesfatin-130-59 injected intracerebroventricularly differentially affects food intake microstructure in rats under normal weight and diet-induced obese conditions. *Front Neurosci*. 2015;**9**:422.
34. Yuan JH, Chen X, Dong J, Zhang D, Song K, Zhang Y, Wu GB, Hu XH, Jiang ZY, Chen P. Nesfatin-1 in the lateral parabrachial nucleus inhibits food intake, modulates excitability of glucosensing neurons, and enhances UCP1 expression in brown adipose tissue. *Front Physiol*. 2017;**8**:235.
35. Heidarzadeh H, Zendejdel M, Babapour V, Gilanpour H. The effect of nesfatin-1 on food intake in neonatal chicks: role of CRF1/CRF2 and H1/ H3 receptors. *Vet Res Commun*. 2018;**42**(1):39–47.
36. Lu L, Liu Z, Huang M, Zhang Z. Dopamine-dependent responses to cocaine depend on corticotropin-releasing factor receptor subtypes. *J Neurochem*. 2003;**84**(6):1378–1386.
37. Kawashima S, Sakihara S, Kageyama K, Nigawara T, Suda T. Corticotropin-releasing factor (CRF) is involved in the acute anorexic effect of alpha-melanocyte-stimulating hormone: a study using CRF-deficient mice. *Peptides*. 2008;**29**(12):2169–2174.
38. Price CJ, Samson WK, Ferguson AV. Nesfatin-1 inhibits NPY neurons in the arcuate nucleus. *Brain Res*. 2008;**1230**:99–106.
39. Inhoff T, Stengel A, Peter L, Goebel M, Taché Y, Bannert N, Wiedenmann B, Klapp BF, Mönnikes H, Kobelt P. Novel insight in distribution of nesfatin-1 and phospho-mTOR in the arcuate nucleus of the hypothalamus of rats. *Peptides*. 2010;**31**(2):257–262.
40. Shimizu H, Arima H, Ozawa Y, Watanabe M, Banno R, Sugimura Y, Ozaki N, Nagasaki H, Oiso Y. Glucocorticoids increase NPY gene expression in the arcuate nucleus by inhibiting mTOR signaling in rat hypothalamic organotypic cultures. *Peptides*. 2010;**31**(1):145–149.
41. Li Z, Xu G, Li Y, Zhao J, Mulholland MW, Zhang W. mTOR-dependent modulation of gastric nesfatin-1/NUCB2. *Cell Physiol Biochem*. 2012;**29**(3-4):493–500.
42. Folgueira C, Barja-Fernandez S, Prado L, Al-Massadi O, Castela C, Pena-Leon V, Gonzalez-Saenz P, Baltar J, Baamonde I, Leis R, Dieguez C, Pagotto U, Casanueva FF, Tovar SA, Nogueiras R, Seoane LM. Pharmacological inhibition of cannabinoid receptor 1 stimulates gastric release of nesfatin-1 via the mTOR pathway. *World J Gastroenterol*. 2017;**23**(35):6403–6411.
43. Zhang W, Zhang C, Fritze D, Chai B, Li J, Mulholland MW. Modulation of food intake by mTOR signalling in the dorsal motor nucleus of the vagus in male rats: focus on ghrelin and nesfatin-1. *Exp Physiol*. 2013;**98**(12):1696–1704.
44. Santoso P, Nakata M, Shiizaki K, Boyang Z, Parmila K, Otgon-Uul Z, Hashimoto K, Satoh T, Mori M, Kuro OM, Yada T. Fibroblast growth factor 21, assisted by elevated glucose, activates paraventricular nucleus NUCB2/nesfatin-1 neurons to produce satiety under fed states. *Sci Rep*. 2017;**7**(1):45819.

45. Akiyama Y, Yoshimura M, Nishimura K, Nishimura H, Sonoda S, Ueno H, Mitojima Y, Saito R, Maruyama T, Nonaka Y, Hashimoto H, Uezono Y, Hirata K, Ueta Y. Activation of central nesfatin-1/NucB2 after intraperitoneally administered cisplatin in rats. *Biochem Biophys Res Commun*. 2017; **490**(3):794–799.
46. Saito R, Sonoda S, Ueno H, Motojima Y, Yoshimura M, Maruyama T, Hashimoto H, Tanaka K, Yamamoto Y, Kusuhara K, Ueta Y. Involvement of central nesfatin-1 neurons on oxytocin-induced feeding suppression in rats. *Neurosci Lett*. 2017; **655**:54–60.
47. Maejima Y, Sedbazar U, Suyama S, Kohno D, Onaka T, Takano E, Yoshida N, Koike M, Uchiyama Y, Fujiwara K, Yashiro T, Horvath TL, Dietrich MO, Tanaka S, Dezaki K, Oh IS, Hashimoto K, Shimizu H, Nakata M, Mori M, Yada T. Nesfatin-1–regulated oxytocinergic signaling in the paraventricular nucleus causes anorexia through a leptin-independent melanocortin pathway. *Cell Metab*. 2009; **10**(5):355–365.
48. Kohno D, Nakata M, Maejima Y, Shimizu H, Sedbazar U, Yoshida N, Dezaki K, Onaka T, Mori M, Yada T. Nesfatin-1 neurons in paraventricular and supraoptic nuclei of the rat hypothalamus coexpress oxytocin and vasopressin and are activated by refeeding. *Endocrinology*. 2008; **149**(3):1295–1301.
49. Mohan H, Ramesh N, Mortazavi S, Le A, Iwakura H, Unniappan S. Nutrients differentially regulate nucleobindin-2/nesfatin-1 in vitro in cultured stomach ghrelinoma (MGN3-1) cells and in vivo in male mice. *PLoS One*. 2014; **9**(12):e115102.
50. Bertucci JI, Blanco AM, Canosa LF, Unniappan S. Glucose, amino acids and fatty acids directly regulate ghrelin and NUCB2/nesfatin-1 in the intestine and hepatopancreas of goldfish (*Carassius auratus*) in vitro. *Comp Biochem Physiol A Mol Integr Physiol*. 2017; **206**:24–35.
51. Kentish SJ, Li H, Frisby CL, Page AJ. Nesfatin-1 modulates murine gastric vagal afferent mechanosensitivity in a nutritional state dependent manner. *Peptides*. 2017; **89**:35–41.
52. Falcinelli S, Rodiles A, Hatéf A, Picchiatti S, Cossignani L, Merrifield DL, Unniappan S, Carnevali O. Dietary lipid content reorganizes gut microbiota and probiotic *L. rhamnosus* attenuates obesity and enhances catabolic hormonal milieu in zebrafish. *Sci Rep*. 2017; **7**(1):5512.
53. Anık A, Çatlı G, Abacı A, Küme T, Bober E. Fasting and postprandial levels of a novel anorexigenic peptide nesfatin in childhood obesity. *J Pediatr Endocrinol Metab*. 2014; **27**(7-8):623–628.
54. Ogiso K, Asakawa A, Amitani H, Nakahara T, Ushikai M, Haruta I, Koyama K, Amitani M, Harada T, Yasuhara D, Inui A. Plasma nesfatin-1 concentrations in restricting-type anorexia nervosa. *Peptides*. 2011; **32**(1):150–153.
55. Ustabas Kahraman F, Vehapoğlu A, Özgen İT, Terzioğlu Ş, Cesur Y, Dündaröz R. Correlation of brain neuropeptide (nesfatin-1 and orexin-A) concentrations with anthropometric and biochemical parameters in malnourished children. *J Clin Res Pediatr Endocrinol*. 2015; **7**(3):197–202.
56. Scharner S, Prinz P, Goebel-Stengel M, Lommel R, Kobelt P, Hofmann T, Rose M, Stengel A. Activity-based anorexia activates nesfatin-1 immunoreactive neurons in distinct brain nuclei of female rats. *Brain Res*. 2017; **1677**:33–46.
57. Tan BK, Hallschmid M, Kern W, Lehnert H, Randeve HS. Decreased cerebrospinal fluid/plasma ratio of the novel satiety molecule, nesfatin-1/NUCB-2, in obese humans: evidence of nesfatin-1/NUCB-2 resistance and implications for obesity treatment. *J Clin Endocrinol Metab*. 2011; **96**(4):E669–E673.
58. Lee WJ, Chen CY, Ser KH, Chong K, Chen SC, Lee PC, Liao YD, Lee SD. Differential influences of gastric bypass and sleeve gastrectomy on plasma nesfatin-1 and obestatin levels in patients with type 2 diabetes mellitus. *Curr Pharm Des*. 2013; **19**(32):5830–5835.
59. Anwar GM, Yamamah G, Ibrahim A, El-Lebedy D, Farid TM, Mahmoud R. Nesfatin-1 in childhood and adolescent obesity and its association with food intake, body composition and insulin resistance. *Regul Pept*. 2014; **188**:21–24.
60. Tsuchiya T, Shimizu H, Yamada M, Osaki A, Oh IS, Ariyama Y, Takahashi H, Okada S, Hashimoto K, Satoh T, Kojima M, Mori M. Fasting concentrations of nesfatin-1 are negatively correlated with body mass index in non-obese males. *Clin Endocrinol (Oxf)*. 2010; **73**(4):484–490.
61. Dogan U, Bulbuler N, Cakır T, Habibi M, Mayir B, Koc U, Aslaner A, Ellidag HY, Gomceli I. Nesfatin-1 hormone levels in morbidly obese patients after laparoscopic sleeve gastrectomy. *Eur Rev Med Pharmacol Sci*. 2016; **20**(6):1023–1031.
62. He R, Yin Y, Li Y, Li Z, Zhao J, Zhang W. Esophagus-duodenum gastric bypass surgery improves glucose and lipid metabolism in mice. *EBioMedicine*. 2018; **28**:241–250.
63. Acar S, Çatlı G, Küme T, Tuhan H, Gürsoy Çalan Ö, Demir K, Böber E, Abacı A. Increased concentrations of serum nesfatin-1 levels in childhood with idiopathic chronic malnutrition. *Turk J Med Sci*. 2018; **48**(2):378–385.
64. Wang Q, Guo F, Sun X, Gao S, Li Z, Gong Y, Xu L. Effects of exogenous nesfatin-1 on gastric distention-sensitive neurons in the central nucleus of the amygdala and gastric motility in rats. *Neurosci Lett*. 2014; **582**:65–70.

65. Xu L, Wang Q, Guo F, Pang M, Sun X, Gao S, Gong Y. Nesfatin-1 signaling in the basomedial amygdala modulates the gastric distension-sensitive neurons discharge and decreases gastric motility via melanocortin 3/4 receptors and modified by the arcuate nucleus. *Eur J Pharmacol.* 2015;**764**:164–172.
66. Li ZL, Xu L, Sun XR, Guo FF, Gong YL, Gao SL. Central nesfatin-1 influences the excitability of ghrelin-responsive gastric distension neurons in the arcuate nucleus and reduces gastric motility in rats. *Eur J Neurosci.* 2013;**38**(11):3636–3643.
67. Guo FF, Xu L, Gao SL, Sun XR, Li ZL, Gong YL. The effects of nesfatin-1 in the paraventricular nucleus on gastric motility and its potential regulation by the lateral hypothalamic area in rats. *J Neurochem.* 2015;**132**(3):266–275.
68. Xu L, Wang H, Gong Y, Pang M, Sun X, Guo F, Gao S. Nesfatin-1 regulates the lateral hypothalamic area melanin-concentrating hormone-responsive gastric distension-sensitive neurons and gastric function via arcuate nucleus innervation. *Metabolism.* 2017;**67**:14–25.
69. Gao S, Guo F, Sun X, Zhang N, Gong Y, Xu L. The inhibitory effects of nesfatin-1 in ventromedial hypothalamus on gastric function and its regulation by nucleus accumbens. *Front Physiol.* 2017;**7**:634.
70. Bonnet MS, Ouelaa W, Tillement V, Trouslard J, Jean A, Gonzalez BJ, Gourcerol G, Dallaporta M, Troadec JD, Mounien L. Gastric distension activates NUCB2/nesfatin-1-expressing neurons in the nucleus of the solitary tract. *Regul Pept.* 2013;**187**:17–23.
71. Feng H, Wang Q, Guo F, Han X, Pang M, Sun X, Gong Y, Xu L. Nesfatin-1 influences the excitability of gastric distension-responsive neurons in the ventromedial hypothalamic nucleus of rats. *Physiol Res.* 2017;**66**(2):335–344.
72. Tian ZB, Deng RJ, Sun GR, Wei LZ, Kong XJ, Ding XL, Jing X, Zhang CP, Ge YL. Expression of gastrointestinal nesfatin-1 and gastric emptying in ventromedial hypothalamic nucleus- and ventrolateral hypothalamic nucleus-lesioned rats. *World J Gastroenterol.* 2014;**20**(22):6897–6905.
73. Watanabe A, Mochiki E, Kimura A, Kogure N, Yanai M, Ogawa A, Toyomasu Y, Ogata K, Ohno T, Suzuki H, Kuwano H. Nesfatin-1 suppresses gastric contractions and inhibits interdigestive migrating contractions in conscious dogs. *Dig Dis Sci.* 2015;**60**(6):1595–1602.
74. Meleine M, Melchior C, Prinz P, Penformis A, Coffin B, Stengel A, Ducrotté P, Gourcerol G. Gastrointestinal peptides during chronic gastric electrical stimulation in patients with intractable vomiting. *Neuromodulation.* 2017;**20**(8):774–782.
75. Xia ZF, Fritze DM, Li JY, Chai B, Zhang C, Zhang W, Mulholland MW. Nesfatin-1 inhibits gastric acid secretion via a central vagal mechanism in rats. *Am J Physiol Gastrointest Liver Physiol.* 2012;**303**(5):G570–G577.
76. Yang GT, Zhao HY, Kong Y, Sun NN, Dong AQ. Study of the effects of nesfatin-1 on gastric function in obese rats. *World J Gastroenterol.* 2017;**23**(16):2940–2947.
77. Kolgazi M, Cantali-Ozturk C, Deniz R, Ozdemir-Kumral ZN, Yuksel M, Sirvanci S, Yeğen BC. Nesfatin-1 alleviates gastric damage via direct antioxidant mechanisms. *J Surg Res.* 2015;**193**(1):111–118.
78. Kalayci M, Kocdor MA, Kuloglu T, Sahin I, Sarac M, Aksoy A, Yardim M, Dalkilic S, Gursu O, Aydin S, Akkoc RF, Ugras M, Artas G, Ozercan IH, Ugur K, Aydin S. Comparison of the therapeutic effects of sildenafil citrate, heparin and neuropeptides in a rat model of acetic acid-induced gastric ulcer. *Life Sci.* 2017;**186**:102–110.
79. Kolgazi M, Ozdemir-Kumral ZN, Cantali-Ozturk C, Demirci EK, Yuksel M, Sirvanci S, Yeğen BC. Anti-inflammatory effects of nesfatin-1 on acetic acid-induced gastric ulcer in rats: involvement of cyclo-oxygenase pathway. *J Physiol Pharmacol.* 2017;**68**(5):765–777.
80. Szlachcic A, Majka J, Strzalka M, Szmyd J, Pajdo R, Ptak-Belowska A, Kwiecien S, Brzozowski T. Experimental healing of preexisting gastric ulcers induced by hormones controlling food intake ghrelin, orexin-A and nesfatin-1 is impaired under diabetic conditions. A key to understanding the diabetic gastropathy? *J Physiol Pharmacol.* 2013;**64**(5):625–637.
81. Zhang AQ, Li XL, Jiang CY, Lin L, Shi RH, Chen JD, Oomura Y. Expression of nesfatin-1/NUCB2 in rodent digestive system. *World J Gastroenterol.* 2010;**16**(14):1735–1741.
82. Mohan H, Unniappan S. Ontogenic pattern of nucleobindin-2/nesfatin-1 expression in the gastroenteropancreatic tissues and serum of sprague dawley rats. *Regul Pept.* 2012;**175**(1–3):61–69.
83. Foo KS, Brauner H, Ostenson CG, Broberger C. Nucleobindin-2/nesfatin in the endocrine pancreas: distribution and relationship to glycaemic state. *J Endocrinol.* 2010;**204**(3):255–263.
84. Morton KA, Hargreaves L, Mortazavi S, Weber LP, Blanco AM, Unniappan S. Tissue-specific expression and circulating concentrations of nesfatin-1 in domestic animals. *Domest Anim Endocrinol.* 2018;**65**:56–66.
85. Riva M, Nitert MD, Voss U, Sathanoori R, Lindqvist A, Ling C, Wierup N. Nesfatin-1 stimulates glucagon and insulin secretion and beta cell NUCB2 is reduced in human type 2 diabetic subjects. *Cell Tissue Res.* 2011;**346**(3):393–405.

86. Gonzalez R, Reingold BK, Gao X, Gaidhu MP, Tsushima RG, Unniappan S. Nesfatin-1 exerts a direct, glucose-dependent insulinotropic action on mouse islet beta- and MIN6 cells. *J Endocrinol.* 2011; **208**(3):R9–R16.
87. Nakata M, Manaka K, Yamamoto S, Mori M, Yada T. Nesfatin-1 enhances glucose-induced insulin secretion by promoting Ca(2+) influx through L-type channels in mouse islet beta-cells. *Endocr J.* 2011; **58**(4):305–313.
88. Maejima Y, Horita S, Kobayashi D, Aoki M, O'Hashi R, Imai R, Sakamoto K, Mori M, Takasu K, Ogawa K, Takenoshita S, Zhao S, Hazama A, Shimomura K. Nesfatin-1 inhibits voltage gated K(+) channels in pancreatic beta cells. *Peptides.* 2017; **95**:10–15.
89. Su Y, Zhang J, Tang Y, Bi F, Liu JN. The novel function of nesfatin-1: anti-hyperglycemia. *Biochem Biophys Res Commun.* 2010; **391**(1):1039–1042.
90. Gonzalez R, Perry RL, Gao X, Gaidhu MP, Tsushima RG, Ceddia RB, Unniappan S. Nutrient responsive nesfatin-1 regulates energy balance and induces glucose-stimulated insulin secretion in rats. *Endocrinology.* 2011; **152**(10):3628–3637.
91. Ramesh N, Mortazavi S, Unniappan S. Nesfatin-1 stimulates glucagon-like peptide-1 and glucose-dependent insulinotropic polypeptide secretion from STC-1 cells in vitro. *Biochem Biophys Res Commun.* 2015; **462**(2):124–130.
92. Li Z, Gao L, Tang H, Yin Y, Xiang X, Li Y, Zhao J, Mulholland M, Zhang W. Peripheral effects of nesfatin-1 on glucose homeostasis. *PLoS One.* 2013; **8**(8):e71513.
93. Dong J, Xu H, Xu H, Wang PF, Cai GJ, Song HF, Wang CC, Dong ZT, Ju YJ, Jiang ZY. Nesfatin-1 stimulates fatty-acid oxidation by activating AMP-activated protein kinase in STZ-induced type 2 diabetic mice. *PLoS One.* 2013; **8**(12):e83397.
94. Wu D, Yang M, Chen Y, Jia Y, Ma ZA, Boden G, Li L, Yang G. Hypothalamic nesfatin-1/NUCB2 knockdown augments hepatic gluconeogenesis that is correlated with inhibition of mTOR-STAT3 signaling pathway in rats. *Diabetes.* 2014; **63**(4):1234–1247.
95. Yang M, Zhang Z, Wang C, Li K, Li S, Boden G, Li L, Yang G. Nesfatin-1 action in the brain increases insulin sensitivity through Akt/AMPK/TORC2 pathway in diet-induced insulin resistance. *Diabetes.* 2012; **61**(8):1959–1968.
96. Bonnet MS, Djelloul M, Tillement V, Tardivel C, Mounien L, Trouslard J, Troadec JD, Dallaporta M. Central NUCB2/nesfatin-1-expressing neurones belong to the hypothalamic-brainstem circuitry activated by hypoglycaemia. *J Neuroendocrinol.* 2013; **25**(1):1–13.
97. Dong J, Guan HZ, Jiang ZY, Chen X. Nesfatin-1 influences the excitability of glucosensing neurons in the dorsal vagal complex and inhibits food intake. *PLoS One.* 2014; **9**(6):e98967.
98. Gantulga D, Maejima Y, Nakata M, Yada T. Glucose and insulin induce Ca2+ signaling in nesfatin-1 neurons in the hypothalamic paraventricular nucleus. *Biochem Biophys Res Commun.* 2012; **420**(4):811–815.
99. Mimee A, Ferguson AV. Glycemic state regulates melanocortin, but not nesfatin-1, responsiveness of glucose-sensing neurons in the nucleus of the solitary tract. *Am J Physiol Regul Integr Comp Physiol.* 2015; **308**(8):R690–R699.
100. Zhang Z, Li L, Yang M, Liu H, Boden G, Yang G. Increased plasma levels of nesfatin-1 in patients with newly diagnosed type 2 diabetes mellitus. *Exp Clin Endocrinol Diabetes.* 2012; **120**(2):91–95.
101. Algul S, Ozkan Y, Ozcelik O. Serum nesfatin-1 levels in patients with different glucose tolerance levels. *Physiol Res.* 2016; **65**(6):979–985.
102. Zhai T, Li SZ, Fan XT, Tian Z, Lu XQ, Dong J. Circulating nesfatin-1 levels and type 2 diabetes: a systematic review and meta-analysis. *J Diabetes Res.* 2017; **2017**:7687098.
103. Chen K, Zhuo T, Wang J, Mei Q. Saxagliptin upregulates nesfatin-1 secretion and ameliorates insulin resistance and metabolic profiles in type 2 diabetes mellitus. *Metab Syndr Relat Disord.* 2018; **16**(7):336–341.
104. Wang C, Wang Y, Hu W. Association of the polymorphism in NUCB2 gene and the risk of type 2 diabetes. *Diabetol Metab Syndr.* 2017; **9**(1):39.
105. Wang R, Wang J, Wan X. Association of the polymorphism in nucleobindin 2 gene and the risk of metabolic syndrome. *Genet Test Mol Biomarkers.* 2016; **20**(1):49–52.
106. Zhang Y, Lu JH, Zheng SY, Yan JH, Chen L, Liu X, Wu WZ, Wang F. Serum levels of nesfatin-1 are increased in gestational diabetes mellitus. *Gynecol Endocrinol.* 2017; **33**(8):621–624.
107. Kucukler FK, Gorkem U, Simsek Y, Kocabas R, Gulen S, Guler S. Low level of nesfatin-1 is associated with gestational diabetes mellitus. *Gynecol Endocrinol.* 2016; **32**(9):759–761.
108. Ademoglu EN, Gorar S, Keskin M, Carlioglu A, Ucler R, Erdamar H, Culha C, Aral Y. Serum nesfatin-1 levels are decreased in pregnant women newly diagnosed with gestational diabetes. *Arch Endocrinol Metab.* 2017; **61**(5):455–459.

109. Chen YC, Inui A, Chang ES, Chen SC, Lee WJ, Chen CY. Comparison of gut hormones and adipokines stimulated by glucagon test among patients with type II diabetes mellitus after metabolic surgery. *Neuropeptides*. 2016;**55**:39–45.
110. Dore R, Levata L, Gachkar S, Jöhren O, Mittag J, Lehnert H, Schulz C. The thermogenic effect of nesfatin-1 requires recruitment of the melanocortin system. *J Endocrinol*. 2017;**235**(2):111–122.
111. Verdegue F, Soustek MS, Hatting M, Blättler SM, McDonald D, Barrow JJ, Puigserver P. Brown adipose YY1 deficiency activates expression of secreted proteins linked to energy expenditure and prevents diet-induced obesity. *Mol Cell Biol*. 2015;**36**(1):184–196.
112. Wang Y, Li Z, Zhang X, Xiang X, Li Y, Mulholland MW, Zhang W. Nesfatin-1 promotes brown adipocyte phenotype. *Sci Rep*. 2016;**6**(1):34747.
113. Yin Y, Li Z, Gao L, Li Y, Zhao J, Zhang W. AMPK-dependent modulation of hepatic lipid metabolism by nesfatin-1. *Mol Cell Endocrinol*. 2015;**417**:20–26.
114. Başar O, Akbal E, Köklü S, Koçak E, Tuna Y, Ekiz F, Gültuna S, Yılmaz FM, Aydoğan T. A novel appetite peptide, nesfatin-1 in patients with non-alcoholic fatty liver disease. *Scand J Clin Lab Invest*. 2012;**72**(6):479–483.
115. Merali Z, Cayer C, Kent P, Anisman H. Nesfatin-1 increases anxiety- and fear-related behaviors in the rat. *Psychopharmacology (Berl)*. 2008;**201**(1):115–123.
116. Ge JF, Xu YY, Qin G, Pan XY, Cheng JQ, Chen FH. Nesfatin-1, a potent anorexic agent, decreases exploration and induces anxiety-like behavior in rats without altering learning or memory. *Brain Res*. 2015;**1629**:171–181.
117. Jing FC, Zhang J, Feng C, Nian YY, Wang JH, Hu H, Yang BD, Sun XM, Zheng JY, Yin XR. Potential rat model of anxiety-like gastric hypersensitivity induced by sequential stress. *World J Gastroenterol*. 2017;**23**(42):7594–7608.
118. Hofmann T, Ahnis A, Elbelt U, Rose M, Klapp BF, Stengel A. NUCB2/nesfatin-1 is associated with elevated levels of anxiety in anorexia nervosa. *PLoS One*. 2015;**10**(7):e0132058.
119. Hofmann T, Stengel A, Ahnis A, Busse P, Elbelt U, Klapp BF. NUCB2/nesfatin-1 is associated with elevated scores of anxiety in female obese patients. *Psychoneuroendocrinology*. 2013;**38**(11):2502–2510.
120. Hofmann T, Elbelt U, Ahnis A, Rose M, Klapp BF, Stengel A. Sex-specific regulation of NUCB2/nesfatin-1: differential implication in anxiety in obese men and women. *Psychoneuroendocrinology*. 2015;**60**:130–137.
121. Hofmann T, Weibert E, Ahnis A, Obbarius A, Elbelt U, Rose M, Klapp BF, Stengel A. Alterations of circulating NUCB2/nesfatin-1 during short term therapeutic improvement of anxiety in obese inpatients. *Psychoneuroendocrinology*. 2017;**79**:107–115.
122. Ari M, Ozturk OH, Bez Y, Oktar S, Erduran D. High plasma nesfatin-1 level in patients with major depressive disorder. *Prog Neuropsychopharmacol Biol Psychiatry*. 2011;**35**(2):497–500.
123. Algul S, Ozelik O. Evaluating the levels of nesfatin-1 and ghrelin hormones in patients with moderate and severe major depressive disorders. *Psychiatry Investig*. 2018;**15**(2):214–218.
124. Xiao MM, Li JB, Jiang LL, Shao H, Wang BL. Plasma nesfatin-1 level is associated with severity of depression in Chinese depressive patients. *BMC Psychiatry*. 2018;**18**(1):88.
125. Xia QR, Liang J, Cao Y, Shan F, Liu Y, Xu YY. Increased plasma nesfatin-1 levels may be associated with corticosterone, IL-6, and CRP levels in patients with major depressive disorder. *Clin Chim Acta*. 2018;**480**:107–111.
126. Xu YY, Liang J, Cao Y, Shan F, Liu Y, Xia QR. High levels of nesfatin-1 in relation to the dysfunction of the hypothalamic-pituitary-adrenal and hypothalamus-pituitary-thyroid axes in depressed patients with subclinical hypothyroidism. *Neuropsychiatr Dis Treat*. 2017;**13**:1647–1653.
127. Zhang N, Li J, Wang H, Xiao L, Wei Y, He J, Wang G. The level of nesfatin-1 in a mouse gastric cancer model and its role in gastric cancer comorbid with depression. *Shanghai Arch Psychiatry*. 2018;**30**(2):119–126.
128. Bloem B, Xu L, Morava E, Faludi G, Palkovits M, Roubos EW, Kozicz T. Sex-specific differences in the dynamics of cocaine- and amphetamine-regulated transcript and nesfatin-1 expressions in the midbrain of depressed suicide victims vs. controls. *Neuropharmacology*. 2012;**62**(1):297–303.
129. Tanida M, Mori M. Nesfatin-1 stimulates renal sympathetic nerve activity in rats. *Neuroreport*. 2011;**22**(6):309–312.
130. Yosten GL, Samson WK. Nesfatin-1 exerts cardiovascular actions in brain: possible interaction with the central melanocortin system. *Am J Physiol Regul Integr Comp Physiol*. 2009;**297**(2):R330–R336.
131. Tanida M, Gotoh H, Yamamoto N, Wang M, Kuda Y, Kurata Y, Mori M, Shibamoto T. Hypothalamic nesfatin-1 stimulates sympathetic nerve activity via hypothalamic ERK signaling. *Diabetes*. 2015;**64**(11):3725–3736.

132. Aydin B, Guvenc G, Altinbas B, Niaz N, Yalcin M. Modulation of nesfatin-1–induced cardiovascular effects by the central cholinergic system. *Neuropeptides*. 2018;**70**:9–15.
133. Brailoiu GC, Deliu E, Tica AA, Rabinowitz JE, Tilley DG, Benamar K, Koch WJ, Brailoiu E. Nesfatin-1 activates cardiac vagal neurons of nucleus ambiguus and elicits bradycardia in conscious rats. *J Neurochem*. 2013;**126**(6):739–748.
134. Yilmaz MS, Altinbas B, Guvenc G, Erkan LG, Avsar O, Savci V, Kucuksen DU, Arican I, Yalcin M. The role of centrally injected nesfatin-1 on cardiovascular regulation in normotensive and hypotensive rats. *Auton Neurosci*. 2015;**193**:63–68.
135. Yosten GL, Samson WK. Neural circuitry underlying the central hypertensive action of nesfatin-1: melanocortins, corticotropin-releasing hormone, and oxytocin. *Am J Physiol Regul Integr Comp Physiol*. 2014;**306**(10):R722–R727.
136. Mimeo A, Smith PM, Ferguson AV. Nesfatin-1 influences the excitability of neurons in the nucleus of the solitary tract and regulates cardiovascular function. *Am J Physiol Regul Integr Comp Physiol*. 2012;**302**(11):R1297–R1304.
137. Mori Y, Shimizu H, Kushima H, Hiromura M, Terasaki M, Tanaka M, Osaki A, Hirano T. Increased blood pressure in nesfatin/nucleobindin-2–transgenic mice. *Hypertens Res*. 2017;**40**(10):861–867.
138. Zhang B, Nakata M, Lu M, Nakae J, Okada T, Ogawa W, Yada T. Protective role of AgRP neuron's PDK1 against salt-induced hypertension. *Biochem Biophys Res Commun*. 2018;**500**(4):910–916.
139. Osaki A, Shimizu H. Peripheral administration of nesfatin-1 increases blood pressure in mice. *Hypertens Res*. 2014;**37**(2):185–186.
140. Ayada C, Turgut G, Turgut S, Güçlü Z. The effect of chronic peripheral nesfatin-1 application on blood pressure in normal and chronic restraint stressed rats: related with circulating level of blood pressure regulators. *Gen Physiol Biophys*. 2015;**34**(1):81–88.
141. Yamawaki H, Takahashi M, Mukohda M, Morita T, Okada M, Hara Y. A novel adipocytokine, nesfatin-1 modulates peripheral arterial contractility and blood pressure in rats. *Biochem Biophys Res Commun*. 2012;**418**(4):676–681.
142. Lu QB, Wang HP, Tang ZH, Cheng H, Du Q, Wang YB, Feng WB, Li KX, Cai WW, Qiu LY, Sun HJ. Nesfatin-1 functions as a switch for phenotype transformation and proliferation of VSMCs in hypertensive vascular remodeling. *Biochim Biophys Acta*. 2018;**1864**(6 Pt A):2154–2168.
143. Zhang JR, Lu QB, Feng WB, Wang HP, Tang ZH, Cheng H, Du Q, Wang YB, Li KX, Sun HJ. Nesfatin-1 promotes VSMC migration and neointimal hyperplasia by upregulating matrix metalloproteinases and downregulating PPARgamma. *Biomed Pharmacother*. 2018;**102**:711–717.
144. Kovalyova O, Ashcheulova T, Demydenko A, Vizir M, Kochubiei O. Nesfatin-1 activity in patients with essential hypertension and prediabetes, type 2 diabetes. *Georgian Med News*. 2017;**263**:44–49.
145. Zhao Y, Ma X, Wang Q, Zhou Y, Zhang Y, Wu L, Ji H, Qin G, Lu J, Bi Y, Ning G. Nesfatin-1 correlates with hypertension in overweight or obese Han Chinese population. *Clin Exp Hypertens*. 2015;**37**(1): 51–56.
146. Sahin FK, Sahin SB, Ural UM, Cure MC, Senturk S, Tekin YB, Balik G, Cure E, Yuce S, Kirbas A. Nesfatin-1 and vitamin D levels may be associated with systolic and diastolic blood pressure values and hearth rate in polycystic ovary syndrome. *Bosn J Basic Med Sci*. 2015;**15**(3):57–63.
147. Angelone T, Filice E, Pasqua T, Amodio N, Galluccio M, Montesanti G, Quintieri AM, Cerra MC. Nesfatin-1 as a novel cardiac peptide: identification, functional characterization, and protection against ischemia/reperfusion injury. *Cell Mol Life Sci*. 2013;**70**(3):495–509.
148. Nair N, Gerger C, Hatf A, Weber LP, Unniappan S. Ultrasonography reveals in vivo dose-dependent inhibition of end systolic and diastolic volumes, heart rate and cardiac output by nesfatin-1 in zebrafish. *Gen Comp Endocrinol*. 2016;**234**:142–150.
149. Barutcigil A, Tasatargil A. Effects of nesfatin-1 on atrial contractility and thoracic aorta reactivity in male rats. *Clin Exp Hypertens*. 2018;**40**(5):414–420.
150. Mazza R, Gattuso A, Filice M, Cantafio P, Cerra MC, Angelone T, Imbrogno S. Nesfatin-1 as a new positive inotrope in the goldfish (*Carassius auratus*) heart. *Gen Comp Endocrinol*. 2015;**224**:160–167.
151. Tasatargil A, Kusu N, Dalaklioglu S, Adiguzel D, Celik-Ozenci C, Ozdem S, Barutcigil A, Ozdem S. Cardioprotective effect of nesfatin-1 against isoproterenol-induced myocardial infarction in rats: role of the Akt/GSK-3beta pathway. *Peptides*. 2017;**95**:1–9.
152. Feijóo-Bandín S, Rodríguez-Penas D, García-Rúa V, Mosquera-Leal A, Abu-Assi E, Portolés M, Roselló-Lletí E, Rivera M, Diéguez C, González-Juanatey JR, Lago F. 24 h nesfatin-1 treatment promotes apoptosis in cardiomyocytes. *Endocrine*. 2016;**51**(3):551–555.
153. Robinson C, Tsang L, Solomon A, Woodiwiss AJ, Gunter S, Mer M, Hsu HC, Gomes M, Norton GR, Millen AME, Dessein PH. Nesfatin-1 and visfatin expression is associated with reduced atherosclerotic disease risk in patients with rheumatoid arthritis. *Peptides*. 2018;**102**:31–37.

154. Nozawa S, Kimura T, Kurishima M, Mimura K, Saeki K, Miki Y, Oda H, Mori A, Momota Y, Azakami D, Ishioka K. Analyses of a satiety factor NUCB2/nesfatin-1; gene expressions and modulation by different dietary components in dogs. *J Vet Med Sci.* 2016;**78**(3):411–417.
155. Banerjee S, Chaturvedi CM. Nesfatin-1: localization and expression in avian gonads and its modulation by temporal phase relation of neural oscillations in female Japanese quail, *Coturnix japonica*. *Gen Comp Endocrinol.* 2015;**224**:205–215.
156. García-Galiano D, Navarro VM, Roa J, Ruiz-Pino F, Sánchez-Garrido MA, Pineda R, Castellano JM, Romero M, Aguilar E, Gaytán F, Diéguez C, Pinilla L, Tena-Sempere M. The anorexigenic neuropeptide, nesfatin-1, is indispensable for normal puberty onset in the female rat. *J Neurosci.* 2010;**30**(23):7783–7792.
157. Çatlı G, Anık A, Küme T, Çalan ÖG, Dündar BN, Böber E, Abacı A. Serum nesfatin-1 and leptin levels in non-obese girls with premature thelarche. *J Endocrinol Invest.* 2015;**38**(8):909–913.
158. Gao X, Zhang K, Song M, Li X, Luo L, Tian Y, Zhang Y, Li Y, Zhang X, Ling Y, Fang F, Liu Y. Role of nesfatin-1 in the reproductive axis of male rat. *Sci Rep.* 2016;**6**(1):32877.
159. Hatel A, Unniappan S. Gonadotropin-releasing hormone, kisspeptin, and gonadal steroids directly modulate nucleobindin-2/nesfatin-1 in murine hypothalamic gonadotropin-releasing hormone neurons and gonadotropes. *Biol Reprod.* 2017;**96**(3):635–651.
160. Seon S, Jeon D, Kim H, Chung Y, Choi N, Yang H. Testosterone regulates NUCB2 mRNA expression in male mouse hypothalamus and pituitary gland. *Dev Reprod.* 2017;**21**(1):71–78.
161. Legg-St Pierre CB, Mackova M, Miskiewicz EL, Hemmings DG, Unniappan S, MacPhee DJ. Insulinotropic nucleobindin-2/nesfatin-1 is dynamically expressed in the haemochorial mouse and human placenta. *Reprod Fertil Dev.* 2018;**30**(3):519–532.
162. Chung Y, Kim H, Seon S, Yang H. Serum cytokine levels are related to nesfatin-1/NUCB2 expression in the implantation sites of spontaneous abortion model of CBA/j x DBA/2 mice. *Dev Reprod.* 2017;**21**(1):35–46.