


Glycogenosis is common in nonalcoholic fatty liver disease and is independently associated with ballooning, but lower steatosis and lower fibrosis

Daniela S Allende¹ | Samer Gawrieh² | Oscar W Cummings³ | Patricia Belt⁴ |
 Laura Wilson⁴ | Mark Van Natta⁴ | Cynthia A Behling⁵ | Danielle Carpenter⁶ |
 Ryan M Gill⁷ | David E Kleiner⁸  | Mathew M Yeh⁹ | Naga Chalasani² |
 Cynthia D Guy¹⁰  | NASH Clinical Research Network

¹Department of Pathology, Cleveland Clinic, Cleveland, OH, USA

²Department of Medicine, Indiana University School of Medicine, Indianapolis, IN, USA

³Department of Pathology, Indiana University School of Medicine, Indianapolis, IN, USA

⁴Center for Clinical Trials, The Johns Hopkins University, Bloomberg School of Public Health, Baltimore, MD, USA

⁵Department of Pathology, Pacific Rim Pathology, San Diego, CA, USA

⁶Department of Pathology, Saint Louis University, St Louis, MI, USA

⁷Department of Pathology, University of California San Francisco, San Francisco, CA, USA

⁸Laboratory of Pathology, National Cancer Institute, Bethesda, MD, USA

⁹Department of Pathology, University of Washington School of Medicine, Seattle, WA, USA

¹⁰Department of Pathology, Duke University, Durham, NC, USA

Correspondence

Cynthia D Guy, MD, Duke University Health System, Department of Pathology, Duke South, Room 327B, 40 Duke Medicine Drive, Durham, NC 27710, USA.
 Email: cynthia.guy@duke.edu

Abstract

Background/Aims: Glycogen synthesis and storage are normal hepatocyte functions. However, glycogenosis, defined as excess hepatocyte glycogen visible by routine H&E light microscopy, has not been well characterized in nonalcoholic fatty liver disease (NAFLD).

Methods: Glycogenosis in NAFLD liver biopsies was graded as “none”, “focal” (in <50% of hepatocytes), or “diffuse” (in ≥50% of hepatocytes). Clinical and pathological variables associated with glycogenosis were assessed. 2047 liver biopsies were prospectively analysed.

Results: In adults and children, any glycogenosis was present in 54% of cases; diffuse glycogenosis was noted in approximately 1/3 of cases. On multiple logistic regression analysis, adults with glycogenosis tended to be older ($P = .003$), female ($P = .04$), have higher serum glucose ($P = .01$), and use insulin ($P = .02$). Adults tended to have lower steatosis scores ($P = .006$) and lower fibrosis stages ($P = .005$); however, unexpectedly, they also tended to have more hepatocyte injury including ballooning ($P = .003$). On multiple logistic regression analysis, paediatric patients with glycogenosis were more likely to be Hispanic ($P = .03$), have lower body weight ($P = .002$), elevated triglycerides ($P = .001$), and a higher fasting glucose ($P = .007$). Paediatric patients with glycogenosis also had less steatosis ($P < .001$) than those without.

Conclusions: Glycogenosis is common in adult and paediatric NAFLD, and is associated with clinical features of insulin resistance. Glycogenosis is important to recognize histologically because it may be misinterpreted as ballooning, and when

Abbreviations: AIC, akaike's information criteria; ALT, alanine aminotransferase; AST, aspartate aminotransferase; BMI, body mass index (weight in kg/height in metres²); H&E, haematoxylin and eosin; HbA1c, haemoglobin A1c; HDL, high density lipoprotein cholesterol; HOMA-IR, homeostatic model assessment of insulin resistance; IQR, inter-quartile range; NAFL, nonalcoholic fatty liver (“isolated steatosis”); NAFLD, nonalcoholic fatty liver disease; NAS, NAFLD activity score; NASH CRN, nonalcoholic steatohepatitis clinical research network; NASH, nonalcoholic steatohepatitis; OR, odds ratio; PAS, periodic acid-Schiff; PASD, periodic acid-Schiff with diastase; PPP1R3B, protein phosphatase 1 regulatory subunit 3B. Daniela S Allende and Samer Gawrieh are co first authors.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2020 The Authors. *Liver International* published by John Wiley & Sons Ltd.

Funding information

The Nonalcoholic Steatohepatitis Clinical Research Network (NASH CRN) is supported by the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) (grants U01DK061718, U01DK061728, U01DK061731, U01DK061732, U01DK061734, U01DK061737, U01DK061738, U01DK061730, U01DK061713). Additional support is received from the National Center for Advancing Translational Sciences (NCATS) (grants UL1TR000439, UL1TR000077, UL1TR000436, UL1TR000150, UL1TR000424, UL1TR000006, UL1TR000448, UL1TR000040, UL1TR000100, UL1TR000004, UL1TR000423, UL1TR000058, and UL1TR000454). This research was supported in part by the Intramural Research Program of the NIH, National Cancer Institute

Handling Editor: Utpal Pajvani

diffuse, confusion with glycogen storage disorders or glycogenic hepatopathy must be avoided. The newly observed dichotomous relationship between glycogenesis and increased liver cell injury but decreased steatosis and fibrosis requires further study.

KEYWORDS

adults, children, hepatic glycogen, pathology, steatohepatitis

1 | INTRODUCTION

The liver is central to lipid and carbohydrate metabolism in the healthy state, but these become dysregulated in the setting of obesity and metabolic syndrome, and contribute to the pathogenesis of nonalcoholic fatty liver disease (NAFLD).¹ Steatosis, hepatocellular lipid accumulation, is a hallmark histopathologic lesion in NAFLD,² and the histological, laboratory, radiographic and clinical associations of steatosis have been widely studied.^{3,4} However, relatively little is known about hepatocellular glycogen processing in NAFLD.

Hepatocytes normally contain glycogen in the unfasted state, but the hepatocellular glycogen is not distinctly visible by light microscopy using routine haematoxylin & eosin (H&E). In contrast, glycogen is distinctly visible by routine methods in pathological conditions such as glycogen storage diseases and glycogenic hepatopathy.⁵ This abnormal glycogen accumulation visible on routine H&E, termed glycogenesis, is characterized by faint grey to light pink cytoplasmic pallor and rarefaction.

The observation by our group that some hepatocytes in NAFLD contained glycogenesis, led us to a systematic investigation. Our aim was to describe the prevalence of hepatic glycogenesis in adults and children enrolled in the National Institutes of Health-sponsored Nonalcoholic Steatohepatitis Clinical Research Network (NASH CRN) and to evaluate the associated clinical, laboratory and histopathological variables.

2 | MATERIALS AND METHODS

Haematoxylin & eosin, Masson trichrome and Prussian blue stained slides from the NASH CRN Database Study, a noninterventonal

registry, were prospectively reviewed and scored by the NASH CRN Pathology Committee in a consensus manner. The NASH CRN Pathology Committee is comprised of nine expert liver pathologists who practice at US-based academic medical centres. All cases in this study were scored during group review, with a minimal number of three pathologists in attendance at any given scoring session. Final scores were determined by discussion and a majority opinion. Approval from the institutional review boards at each participating center and the data coordinating center had been obtained. Each case was evaluated according to the published and validated NASH CRN scoring system.⁶ In addition, glycogenesis was defined as a faint grey to light pink cytoplasmic pallor seen on routine H&E using light microscopy (Figure 1A,B). The degree of glycogenesis was scored as "none", "focal" or "diffuse". "None" was defined as complete absence of glycogenesis. "Focal" was defined as glycogenesis involving less than 50% of hepatocytes. "Diffuse" was defined as glycogenesis involving equal to or more than 50% of hepatocytes. Because Periodic acid-Schiff (PAS) and Periodic acid-Schiff with diastase (PASD) histochemistry is not included among the standard NASH CRN stains, the presence of glycogen was confirmed with PAS and PASD in a subset of cases (n = 4). Similarly, electron microscopy was used on 1 case to confirm the presence of glycogen.

Demographics and clinical data were collected at the time of patient entry into the NASH CRN. Laboratory values were collected within 6 months prior to or up to one month after the date of the biopsy.

The two primary comparisons were the presence of any glycogenesis vs none, and focal vs diffuse glycogenesis. Unadjusted comparisons were stratified by adults and children (age < 18 years

of age). Distributions were summarized using proportions for categorical variables and means and standard deviations for normally distributed variables, or medians and inter-quartile ranges [IQR] for non-normal data. *P*-values for the two primary comparisons were derived using the chi-square test for categorical variables and *t*-tests for normally distributed variables or Wilcoxon rank sum test for non-normally distributed variables. Multiple logistic regression was used to test for independent effects on any vs no glycogenesis and diffuse vs focal glycogenesis. Adults and children were combined for the latter two analyses. Results from stratified analyses were similar to the combined results (data not shown). Akaike's information criteria (AIC) were used to select the set of variables that maximized the amount of information from a candidate list of all variables in Table 1. *P*-values were two-sided, nominal and not adjusted for multiple comparisons. Analyses were performed using SAS version 9.3 (SAS Institute) and Stata version 14 (StataCorp).

3 | RESULTS

A total of 2047 liver biopsies were analysed, including 1348 from adults and 699 from children. Glycogenesis, focal or diffuse, was observed in 54% of adult cases and 53.5% of paediatric cases. Diffuse glycogenesis was identified in 28% of adults and 34% of children.

Glycogen was confirmed by PAS and PASD histochemistry in a subset of cases ($n = 4$, Figure 1C,D). Electron microscopy from a formalin fixed paraffin embedded liver biopsy was performed on one representative case (Figure 2).

3.1 | Clinical and laboratory features associated with glycogenesis in adults

Compared to those without glycogenesis, adults with any glycogenesis were older (mean age in years was 51.6 ± 11.7 vs 48.9 ± 12.3 , $P < .0001$), predominantly female (68% vs 59%, $P = .0008$) and more frequently had diabetes mellitus (45% vs 39%, $P = .05$). There were only four adult subjects with type 1 diabetes mellitus, and all were in the glycogenesis group. BMI was not different between the two groups ($P = .51$). Patients in this cohort (both those with and without glycogenesis) were mostly White, and Hispanic ethnicity was reported in a small subset of cases in both groups (14% in those with glycogenesis vs 11% in those without glycogenesis, $P = .09$) (Table 1).

Patients with any glycogenesis demonstrated a higher serum glucose ($115 \text{ mg/dL} \pm 42$ vs 108 ± 39 , $P = .002$) and HbA1c ($6.4\% \pm 1.3\%$ vs $6.2\% \pm 1.1$, $P = .008$) in comparison to patients without glycogenesis. There were no significant differences in ALT, AST, alkaline phosphatase, insulin, HOMA-IR or lipoprotein levels between those with or without glycogenesis.

Patients with diffuse glycogenesis as opposed to those with focal glycogenesis were more frequently women (75% vs 60%, $P < .0001$), of Hispanic ethnicity (17% vs 11%, $P = .03$), and had lower ALT levels

Layman Summary


- The global "epidemic" of obesity continues to be an important and growing health concern. Obesity and related conditions including hypertension, dyslipidaemia and diabetes mellitus work synergistically to damage the liver. This is called nonalcoholic fatty liver disease (NAFLD). NAFLD has a mild subtype (NAFL) and a serious subtype called nonalcoholic steatohepatitis (NASH) that can lead to liver scarring (cirrhosis), liver cancer and liver failure. Distinguishing NAFL from NASH currently requires a liver biopsy. Pathologists review the liver tissue under a microscope to look for fat, inflammation, liver cell death, and fibrosis, and use what they see to make a diagnosis.
- A group of NIH-sponsored expert liver pathologists reviewed a large number of NAFLD liver biopsies from adults and children, and for the first time described that not only were the liver cells stuffed with lipid, but many cases (more than half) also showed liver cells stuffed with glycogen ("glycogenesis"). Statistical analysis showed that adult NAFLD cases with glycogen were associated with older age, women, high blood levels of glucose and insulin. Furthermore, although cases with glycogenesis were associated with more cell injury and cell death, there was relatively less fat in the cells and importantly less scarring (fibrosis). This is a new insight; in NAFLD not only do the liver cells have abnormal lipid metabolism and storage, but they also have abnormal glycogen metabolism and storage. Our findings suggest the possibility that shunting of substrates away from lipid droplets towards glycogen deposition may protect the liver from scarring, and thus in the long term may protect from liver failure. More studies are needed so that we can better understand the interplay between lipid and glycogen metabolism inside the NAFLD liver cells, this may broaden our attempts to find good therapies for NAFLD.

($64 \text{ U/L} \pm 47$ vs 74 ± 49 , $P = .009$), and higher HDL levels ($46 \text{ mg/dL} \pm 13$ vs 43 ± 12 , $P = .02$), but there were no differences in glucose, insulin, HOMA-IR or other lipoprotein levels.

3.2 | Clinical and laboratory features associated with glycogenesis in children

Compared to those without glycogenesis, children with any glycogenesis were younger ($12.2 \text{ years} \pm 2.8$ vs 12.8 ± 2.8 , $P = .008$) had a lower BMI ($31.3 \text{ kg/m}^2 \pm 6.0$ vs 32.6 ± 6.8 , $P = .008$), were less frequently White (59% vs 74%, $P < .0001$) and mainly Hispanic

FIGURE 1 Glycogenosis as seen on routine light microscopy. (A) Medium to low power H&E showing diffuse cytoplasmic glycogen accumulation in hepatocytes (200x). (B) Higher power magnification shows cytoplasmic pale grey to faint pink glycogen collections (600x). (C) PAS histochemistry highlights the presence of diffuse cytoplasmic glycogen (200x). (D) PAS with diastase shows complete glycogen removal



(77% vs 66%, $P = .001$). A similar and small subset of patients in both groups carried a diagnosis of type 2 diabetes mellitus (6% vs 6%, $P = .75$). There were three children with type 1 diabetes mellitus, all of whom had diffuse glycogenosis (Table 2).

Paediatric subjects with any glycogenosis had higher levels of alkaline phosphatase ($232 \text{ U/L} \pm 101$ vs 216 ± 110 , $P = .04$), glucose ($92 \text{ mg/dl} \pm 30$ vs 87 ± 11), HbA1c ($5.6\% \pm 1.1$ vs 5.4 ± 0.4 , $P = .002$) and triglycerides ($159 \text{ mg/dL} \pm 88$ vs 143 ± 73 , $P = .009$). ALT, AST, insulin, HOMA-IR and other lipoprotein levels were not statistically different between the two groups.

After stratifying patients with glycogenosis into “focal” and “diffuse” groups, most of the clinical and laboratory variables were not significantly different except for age and alkaline phosphatase. Children with diffuse glycogenosis were slightly younger ($12.0 \text{ years} \pm 2.7$ vs 12.7 ± 2.8 , $P = .02$) and had higher levels of alkaline phosphatase ($241 \text{ U/L} \pm 102$ vs 217 ± 98 , $P = .02$).

3.3 | Histological associations of glycogenosis in adults

Compared to those without glycogenosis, patients with any glycogenosis had lower steatosis grades (1.69 ± 0.89 vs 1.88 ± 0.90 , $P = .0001$), higher ballooning score (1.07 ± 0.84 vs 0.90 ± 0.85 , $P = 0.0002$), and a higher proportion of acidophil bodies (49% vs 43%, $P = .03$), megamitochondria (33% vs 23%, $P < .0001$) and Mallory-Denk bodies (36% vs 29%, $P = .009$). Those with any glycogenosis were more frequently diagnosed with definite steatohepatitis (61% vs 51%, $P = .0007$) compared to borderline NASH or nonalcoholic fatty liver (NAFL, isolated steatosis). Even though many histologic features of severe hepatocyte injury were more frequent in those

with any glycogenosis vs those with none, the NAFLD Activity Score (NAS) and fibrosis stage were not significantly different between the two groups. Furthermore, no significant differences were noted between those with any glycogenosis and those with no glycogenosis in lobular inflammation, portal inflammation, microvesicular steatosis, glycogenic nuclei and iron deposition (Table 1).

When compared to focal glycogenosis, cases with diffuse glycogenosis were characterized by a milder steatosis grade (1.44 ± 0.85 vs 1.96 ± 0.85 , $P < .0001$), less frequent microvesicular steatosis (9% vs 15%, $P = .01$) and lower lobular inflammation score (1.49 ± 0.70 vs 1.62 ± 0.72 , $P = .02$). As a consequence, those with diffuse glycogenosis had lower NAS (4.0 ± 1.7 vs 4.7 ± 1.7 , $P < .0001$) and lower frequency of definite steatohepatitis (58% vs 63%, $P = .0003$) in comparison to focal glycogenosis cases.

3.4 | Histological associations of glycogenosis in children

Similar to adults, paediatric patients with any glycogenosis also had lower steatosis grades (1.89 ± 0.95 vs 2.23 ± 0.87 , $P < .0001$). Unlike adults, there was no difference in the diagnostic classification between glycogenosis cases and those without ($P = .07$). The NAS was slightly lower in children with glycogenosis vs those with none (3.8 ± 1.6 vs 4.2 ± 1.5 , $P = .0001$). No significant differences were observed between children with and without glycogenosis in lobular inflammation, portal inflammation, microvesicular steatosis, megamitochondria, ballooning grade, the presence of Mallory-Denk bodies, acidophil bodies or iron deposition. The fibrosis stage was not different in those children with glycogenosis compared to those without glycogenosis (Table 2).

TABLE 1 Demographics, laboratory data and histologic features by glycogenesis in 1348 adults

| | None (n=620) | Any (n=728) | None vs Any P ^a | Focal (n=348) | Diffuse (n=380) | Focal vs Diffuse P ^a | None vs Diffuse P ^a |
|---------------------------------------|------------------|------------------|----------------------------|------------------|------------------|---------------------------------|--------------------------------|
| | N (%) or Mean±SD | N (%) or Mean±SD | | N (%) or Mean±SD | N (%) or Mean±SD | | |
| Demographics | | | | | | | |
| Age at biopsy -yrs | 48.9±12.3 | 51.6±11.7 | <.0001 | 51.1±11.9 | 52.1±11.6 | .28 | .0001 |
| Body mass index-kg/m ² | 34.1±6.4 | 34.3±6.6 | .51 | 34.3±6.4 | 34.4±6.9 | .84 | .51 |
| Male Sex | 254 (41) | 233 (32) | .0008 | 139 (40) | 94 (25) | <.0001 | <.0001 |
| Race | | | .52 | | | .54 | .32 |
| White | 497 (80) | 579 (80) | | 281 (81) | 298 (78) | | |
| Black | 29 (5) | 27 (4) | | 14 (4) | 13 (3) | | |
| Other | 94 (15) | 122 (17) | | 53 (15) | 69 (18) | | |
| Ethnicity | | | | | | | |
| Hispanic | 69 (11) | 104 (14) | .09 | 39 (11) | 65 (17) | .03 | .007 |
| Diabetes Mellitus (any) | 244 (39) | 325 (45) | .05 | 152 (44) | 173 (46) | .65 | .05 |
| Type 2 | 243 (39) | 321 (44) | .08 | 151 (43) | 170 (45) | .77 | .08 |
| Laboratory measures | | | | | | | |
| AST, U/L | 50±35 | 50±32 | .86 | 53±33 | 48±32 | .07 | .42 |
| ALT, U/L | 68±54 | 69±48 | .90 | 74±49 | 64±47 | .009 | .22 |
| Alkaline phosphatase, U/L | 83±52 | 83±33 | .84 | 81±33 | 85±32 | .06 | .38 |
| Glucose, mg/dL | 108±39 | 115±42 | .002 | 116±42 | 114±42 | .66 | .02 |
| Insulin, μU/mL | 24±30 | 27±29 | .06 | 26±23 | 29±33 | .13 | .03 |
| HOMA-IR, μU/mL ^b mg/dL/405 | 7.1±12.2 | 8.2±10.1 | .07 | 7.8±8.4 | 8.7±11.5 | .22 | .05 |
| HbA1c, % | 6.2±1.1 | 6.4±1.3 | .008 | 6.5±1.3 | 6.4±1.2 | .41 | .08 |
| Total cholesterol, mg/dL | 191±48 | 189±42 | .57 | 191±44 | 188±41 | .29 | .31 |
| Triglycerides, mg/dL | 178±154 | 180±158 | .81 | 187±193 | 172±117 | .21 | .58 |
| HDL, mg/dL | 43±12 | 45±13 | .13 | 43±12 | 46±13 | .02 | .01 |
| LDL, mg/dL | 115±40 | 112±37 | .25 | 114±37 | 110±36 | .12 | .07 |
| Histological features | | | | | | | |
| Biopsy Length, mm (range) | 20.4±10.0 (1-70) | 20.2±9.3 (3-64) | .70 | 20.3±9.7 (4-64) | 20.1±8.8 (3-58) | .81 | .65 |
| Steatosis Grade | 1.88±0.90 | 1.69±0.89 | .0001 | 1.96±0.85 | 1.44±0.85 | <.0001 | <.0001 |
| 0 (<5%) | 32 (5) | 56 (8) | .001 | 10 (3) | 46 (12) | <.0001 | <.0001 |
| 1 (5-33%) | 198 (32) | 268 (37) | | 103 (30) | 165 (43) | | |
| 2 (33-67%) | 205 (33) | 251 (34) | | 126 (36) | 125 (33) | | |
| 3 (>67%) | 185 (30) | 153 (21) | | 109 (31) | 44 (12) | | |
| Steatosis location^b | | | | | | | |
| Zone 3 Predominant | 316 (51) | 334 (46) | .005 | 155 (45) | 179 (48) | .04 | .0004 |
| Zone 1 Predominant | 4 (1) | 8 (1) | | 3 (1) | 5 (1) | | |

(Continues)

TABLE 1 (Continued)

| | None (n=620) | Any (n=728) | None vs Any P ^a | Focal (n=348) | Diffuse (n=380) | Focal vs Diffuse P ^a | None vs Diffuse P ^a |
|----------------------------------|------------------|------------------|----------------------------|------------------|------------------|---------------------------------|--------------------------------|
| | N (%) or Mean±SD | N (%) or Mean±SD | | N (%) or Mean±SD | N (%) or Mean±SD | | |
| Azonal | 160 (26) | 248 (34) | | 112 (32) | 136 (36) | | |
| Panacinar | 137 (22) | 131 (18) | | 78 (22) | 53 (14) | | |
| Steatosis macrovesicular type | | | .002 | | | .05 | .0002 |
| Large droplet | 444 (72) | 576 (80) | | 265 (76) | 311 (83) | | |
| Mixed large and small droplet | 157 (25) | 126 (17) | | 71 (20) | 55 (15) | | |
| Small droplet | 16 (3) | 19 (3) | | 12 (3) | 7 (2) | | |
| Microvesicular Steatosis | 69 (11) | 84 (12) | .86 | 51 (15) | 33 (9) | .01 | .22 |
| Lobular inflammation score | 1.53±0.70 | 1.55±0.71 | .53 | 1.62±0.72 | 1.49±0.70 | .02 | .45 |
| 0 (no foci) | 11 (2) | 13 (2) | .93 | 3 (1) | 10 (3) | .08 | .78 |
| 1 (<2 foci/20x hpf) | 334 (54) | 380 (52) | | 172 (49) | 208 (55) | | |
| 2 (2-4 foci/20x hpf) | 211 (34) | 254 (35) | | 128 (37) | 126 (33) | | |
| 3 (>4 foci/20x hpf) | 66 (10) | 81 (11) | | 45 (13) | 36 (9) | | |
| Portal inflammation score | 1.12±0.58 | 1.18±0.61 | .07 | 1.16±0.61 | 1.19±0.61 | .45 | .06 |
| 0-None | 72 (12) | 83 (11) | .07 | 41 (12) | 42 (11) | .72 | .06 |
| 1-Mild | 402 (65) | 434 (60) | | 211 (61) | 223 (59) | | |
| 2- More than mild | 145 (23) | 211 (29) | | 96 (28) | 115 (30) | | |
| Ballooning injury score | 0.90±0.85 | 1.07±0.84 | .0002 | 1.11±0.84 | 1.03±0.84 | .19 | .02 |
| 0-None | 259 (42) | 236 (32) | .0009 | 106 (30) | 130 (34) | .41 | .05 |
| Few | 167 (27) | 208 (29) | | 98 (28) | 110 (29) | | |
| 2-Many | 194 (31) | 284 (39) | | 144 (41) | 140 (37) | | |
| Classic ballooning ^c | 161 (26) | 230 (32) | .02 | 114 (33) | 116 (31) | .52 | .12 |
| Acidophil Bodies | 268 (43) | 358 (49) | .03 | 183 (53) | 175 (46) | .09 | .38 |
| Megamitochondria | 143 (23) | 239 (33) | <.0001 | 104 (30) | 135 (36) | .11 | <.0001 |
| Mallory Bodies | 182 (29) | 263 (36) | .009 | 125 (36) | 138 (36) | .94 | .02 |
| Fibrosis stage | 1.55±1.25 | 1.59±1.30 | .57 | 1.62±1.28 | 1.57±1.32 | .58 | .86 |
| 0: None | 152 (25) | 192 (26) | .54 | 88 (25) | 104 (27) | .14 | .19 |
| 1A: Mild perisinusoidal only | 71 (11) | 80 (11) | | 33 (10) | 47 (12) | | |
| 1B: Moderate perisinuosoidal | 82 (13) | 85 (12) | | 42 (12) | 43 (11) | | |
| 1C: Periportal only | 26 (4) | 23 (3) | | 8 (2) | 15 (4) | | |
| 2: Periportal and perisinusoidal | 125 (20) | 128 (18) | | 75 (22) | 53 (14) | | |

(Continues)

TABLE 1 (Continued)

| | None (n=620) | Any (n=728) | None vs Any P ^a | Focal (n=348) | Diffuse (n=380) | Focal vs Diffuse P ^a | None vs Diffuse P ^a |
|----------------------------|------------------|------------------|----------------------------|------------------|------------------|---------------------------------|--------------------------------|
| | N (%) or Mean±SD | N (%) or Mean±SD | | N (%) or Mean±SD | N (%) or Mean±SD | | |
| 3: Bridging fibrosis | 119 (19) | 161 (22) | | 74 (21) | 87 (23) | | |
| 4: Cirrhosis | 44 (7) | 58 (8) | | 27 (8) | 31 (8) | | |
| Diagnostic classification | | | | | | | |
| Not NAFLD | 28 (5) | 34 (5) | .0007 | 4 (1) | 30 (8) | .0003 | .004 |
| NAFLD, Not steatohepatitis | 146 (24) | 144 (20) | | 76 (22) | 68 (18) | | |
| Borderline, Zone 3 pattern | 123 (20) | 95 (13) | | 41 (12) | 54 (14) | | |
| Borderline, Zone 1 pattern | 6 (1) | 14 (2) | | 8 (2) | 6 (2) | | |
| Definite steatohepatitis | 317 (51) | 441 (61) | | 219 (63) | 222 (58) | | |
| NAFLD activity score | | | | | | | |
| 0 | 5 (1) | 9 (1) | .33 | 1 (0) | 8 (2) | <.0001 | .02 |
| 1 | 26 (4) | 25 (3) | | 4 (1) | 21 (6) | | |
| 2 | 79 (13) | 93 (13) | | 33 (9) | 60 (16) | | |
| 3 | 93 (15) | 119 (16) | | 63 (18) | 56 (15) | | |
| 4 | 147 (24) | 141 (19) | | 57 (16) | 84 (22) | | |
| 5 | 107 (17) | 143 (20) | | 69 (20) | 74 (19) | | |
| 6 | 83 (13) | 112 (15) | | 61 (18) | 51 (13) | | |
| 7 | 58 (9) | 71 (10) | | 47 (14) | 24 (6) | | |
| 8 | 22 (4) | 15 (2) | | 13 (4) | 2 (1) | | |
| Mean±SD | 4.3±1.8 | 4.3±1.8 | .94 | 4.7±1.7 | 4.0±1.7 | <.0001 | .003 |

^aDerived from chi-squared test for nominal variables and t-test with unequal variance for continuous variables

^b10 cases had no steatosis; 166 cases are missing microgranulomas, large lipogranulomas, pigmented macrophages and glycogen nuclei—no longer scored

^cSevere ballooning and presence of Mallory bodies

Compared to children with focal glycogenosis, those with diffuse glycogenosis demonstrated a lower steatosis grade (1.80 ± 0.93 vs 2.04 ± 0.96 , $P = .02$), a higher frequency of megamitochondria (14% vs 7%, $P = .03$) and a slightly lower NAS (3.6 ± 1.6 vs 4.0 ± 1.6 , $P = .04$).

3.5 | Covariates of glycogenosis (none, any, focal, diffuse) on multiple logistic regression analyses in adults

Older age at biopsy (OR 1.012, 95% CI 1.006, 1.027, $P = .003$), female sex (OR 1.31, 95% CI 1.1, 1.71, $P = .04$), high glucose (OR 1.005, 95% CI 1.001, 1.009, $P = .01$), higher insulin (OR 1.013, 95% CI 1.002, 1.025, $P = .02$), and lower HOMA-IR (OR 0.97, 95% CI 0.94, 1.02, $P = .06$) significantly increased the risk for any glycogenosis. Female sex (OR 2.04, 95% CI 1.41, 2.95, $P < .001$) and use of insulin (OR 1.009, 95% CI 1.001, 1.017, $P = .02$) were associated with diffuse glycogenosis (Table 3).

Of the histological features, lower steatosis grade (OR 0.82, 95% CI 0.71, 0.94, $P = .006$), lower fibrosis stage (OR 0.84, CI 95% 0.75, 0.95, $P = .005$) were associated with higher risk for glycogenosis. The presences of megamitochondria (OR 1.48, 95% CI 1.13, 1.95, $P = .004$), acidophil bodies (OR 1.32, 95% CI 1.03, 1.69, $P = .03$), and higher ballooning grade (OR 1.30, 95% CI 1.09, 1.55, $P = .003$) were associated with higher odds of finding glycogenosis on histological exam. Diffuse glycogenosis compared to focal glycogenosis was associated with lower steatosis (OR 0.53, 95% CI 0.43, 0.64, $P < .001$) and less microvesicular steatosis (OR 0.56, 95% CI 0.33, 0.97, $P = .04$).

3.6 | Covariates of glycogenosis (none, any, focal, diffuse) on multiple logistic regression analyses in children

Hispanic ethnicity (OR 1.53, 95% CI 1.03, 2.25, $P = .03$), elevated triglycerides (OR 1.004, 95% CI 1.002, 1.007, $P = .001$) and high




FIGURE 2 Glycogenosis as seen on electron microscopy. (A) Toluidine stained thick section reveals cytoplasmic glycogen (400x). (B) Ultrastructure of the same hepatocyte confirms cytoplasmic glycogen aggregates (1000x)

glucose (OR 1.017, 95% CI 1.005, 1.030, $P = .007$) increased the odds of finding any glycogenosis. In contrast, White race (OR 0.46, 95% CI 0.32, 0.67, $P < .001$), higher weight (OR 0.989, 95% CI 0.982, 0.996, $P = .002$), higher AST (OR 0.997, 95% CI 0.994, 1.000, $P = .04$), and higher steatosis grade (OR 0.65, 95% CI 0.54, 0.79, $P < .001$) reduced the odds of finding any glycogenosis. The diagnosis of diabetes (OR 0.2, 95% CI 0.05, 0.72, $P = .01$), and more acidophil bodies (OR 0.63, 95% CI 0.39, 1.00, $P = .05$) reduced the odds of having diffuse glycogenosis in children (Table 4).

4 | DISCUSSION

Glycogen is a branched polymer of glucose stored primarily in the liver and to a lesser extent in skeletal muscle.⁷ It provides a readily available and easy to mobilize source of glucose. Hepatocellular

glycogen can accumulate when hepatic carbohydrate metabolism becomes dysregulated in NAFLD.¹ As glycogen accumulates, the hepatocyte cytoplasm can take on a glassy, pale grey to light pink appearance, termed glycogenosis. To the best of our knowledge, this is the first study to systematically and prospectively characterize glycogenosis in the setting of NAFLD on a large scale in adults and children. We demonstrate that glycogenosis is common, occurring in more than half of both adults and children with NAFLD. Furthermore, diffuse glycogenosis is also common and was identified in approximately one third of both adults and children (28% and 34% respectively).

In adults, the presence of glycogenosis was associated with older age, female sex, higher serum glucose level, and higher insulin level. The histological variables associated with the presence of glycogenosis were lower steatosis grade, but greater hepatocellular injury (higher ballooning grade, the presence of acidophil bodies, Mallory-Denk bodies and megamitochondria).

Based on multiple logistic regression analysis in adults, a lower steatosis score and histologic features indicative of hepatocellular injury such as ballooning, megamitochondria and acidophil bodies were positively associated with glycogenosis in adults. Interestingly, however, despite increased hepatocyte injury, a lower fibrosis stage was associated with the presence of any glycogenosis. This suggests that glycogenosis may have a protective effect on disease progression.

Our study illuminated some curious observations. In some instances of glycogenosis, individual cells may become swollen and take on cytoplasmic pallor, and in this context the enlarged, pale, glycogen-filled hepatocytes may be confused with ballooned hepatocytes (Figure 1A,B). Accurate identification of hepatocyte ballooning is critical as ballooned hepatocytes are a hallmark histopathologic finding for the diagnosis of steatohepatitis and help distinguish the severe form of NAFLD, steatohepatitis, from nonalcoholic fatty liver.^{6,8,9} Cytoskeletal alterations⁹ and fat droplet accumulation in ballooned hepatocytes have been documented,¹⁰ but to our knowledge abundant glycogen accumulation in ballooned hepatocytes has not previously been described. We have observed that ballooned hepatocytes may rarely contain abundant visible glycogen (Figure 3); the large majority of ballooned hepatocytes in our experience, however, do not contain visible glycogen. Glycogenosis in NAFLD can be diffuse, involving essentially every hepatocyte, or it can be focal, occurring in small groups of or individual hepatocytes. Accurate determination of the NAFLD hepatocellular ballooning grade can be challenging^{9,11,12} and the presence of glycogenosis may further add to the difficulty. A raised awareness that glycogenosis is common in NAFLD may be advantageous for correct histopathological classification.

Glycogenosis in NAFLD is a distinct entity and should not be confused with glycogen storage diseases and glycogenic hepatopathy. Glycogenic hepatopathy has been best characterized in patients with type 1 diabetes mellitus and poor glycaemic control, and in patients following high dose corticosteroid therapy.^{5,13,14} Glycogenic hepatopathy has rarely been reported in patients with type 2 diabetes

TABLE 2 Demographics, laboratory data and histologic features by glycogenesis in 699 children

| | None (n=325) | | Any (n=374) | | None vs Any P ^a | | Diffuse (n=236) | | Focal vs Diffuse P ^a | | None vs Diffuse P ^a |
|---------------------------------------|------------------|---------|-----------------|---------|----------------------------|----------------|-----------------|---------|---------------------------------|----------------|--------------------------------|
| | N (%) | Mean±SD | N (%) | Mean±SD | N (%) | P ^a | N (%) | Mean±SD | N (%) | P ^a | |
| Demographics | | | | | | | | | | | |
| Age at biopsy-yr | 12.8±2.8 | | 12.2±2.8 | | .008 | | 12.7±2.8 | | .02 | | .0005 |
| Body mass index-kg/m ² | 32.6±6.8 | | 31.3±6.0 | | .008 | | 31.6±6.4 | | .40 | | .006 |
| Male Sex | 233 (72) | | 257 (69) | | .41 | | 96 (70) | | .82 | | .37 |
| Race | | | | | <.0001 | | | | .89 | | .0006 |
| White | 241 (74) | | 219 (59) | | | | 80 (58) | | | | |
| Black | 9 (3) | | 14 (4) | | | | 6 (4) | | | | |
| Other | 75 (23) | | 141 (38) | | | | 52 (38) | | | | |
| Ethnicity | | | | | | | | | | | |
| Hispanic | 213 (66) | | 287 (77) | | .001 | | 182 (77) | | .90 | | .004 |
| Diabetes Mellitus, any | 18 (6) | | 24 (6) | | .75 | | 19 (8) | | .13 | | .24 |
| Type 2 | 18 (6) | | 21 (6) | | 1.00 | | 5 (4) | | .25 | | .54 |
| Laboratory measures | | | | | | | | | | | |
| AST, U/L | 68±63 | | 62±48 | | .16 | | 60±45 | | .54 | | .33 |
| ALT, U/L | 117±110 | | 107±94 | | .23 | | 105±87 | | .77 | | .35 |
| Alkaline phosphatase, U/L | 216±110 | | 232±101 | | .04 | | 217±98 | | .02 | | .006 |
| Glucose, mg/dL | 87±11 | | 92±30 | | .006 | | 88±19 | | .06 | | .002 |
| Insulin, μU/mL | 33±29 | | 35±46 | | .56 | | 39±68 | | .29 | | .77 |
| HOMA-IR, μU/mL ^b mg/dL/405 | 7.2±7.0 | | 8.1±11.0 | | .20 | | 8.7±14.9 | | .49 | | .41 |
| HbA1c, % | 5.4±0.4 | | 5.6±1.1 | | .002 | | 5.5±0.5 | | .08 | | .001 |
| Total cholesterol, mg/dL | 163±39 | | 166±36 | | .26 | | 163±39 | | .22 | | .11 |
| Triglycerides, mg/dL | 143±73 | | 159±88 | | .009 | | 159±93 | | .99 | | .02 |
| HDL, mg/dL | 40±10 | | 40±9 | | .25 | | 40±9 | | .75 | | .25 |
| LDL, mg/dL | 95±31 | | 96±30 | | .71 | | 93±29 | | .14 | | .33 |
| Histological features | | | | | | | | | | | |
| Biopsy Length-mm, (range) | 19.5±10.0 (4-60) | | 19.1±8.1 (3-49) | | .58 | | 19.1±8.3 (4-49) | | .93 | | .60 |
| Steatosis score | 2.23±0.87 | | 1.89±0.95 | | <.0001 | | 2.04±0.96 | | .02 | | <.0001 |
| 0 (<5%) | 12 (4) | | 24 (6) | | <.0001 | | 9 (7) | | .04 | | <.0001 |
| 1 (5-33%) | 57 (18) | | 118 (32) | | | | 33 (24) | | | | |

(Continues)

TABLE 2 (Continued)

| | None (n=325) | | Any (n=374) | | None vs Any P ^a | | Diffuse (n=236) | | Focal vs Diffuse P ^a | | None vs Diffuse P ^a | |
|---------------------------------|------------------|-----------|------------------|-----------|----------------------------|--|------------------|--|---------------------------------|--|--------------------------------|-------|
| | N (%) or Mean±SD | | N (%) or Mean±SD | | N (%) or Mean±SD | | N (%) or Mean±SD | | N (%) or Mean±SD | | N (%) or Mean±SD | |
| 2 (33-67%) | 100 (31) | 107 (29) | 39 (28) | 68 (29) | | | | | | | | |
| 3 (>67%) | 156 (48) | 125 (33) | 57 (41) | 68 (29) | | | | | | | | |
| Steatosis location ^b | | | | | | | | | | | | |
| Zone 3 Predominant | 126 (39) | 119 (32) | 58 (42) | 61 (26) | .02 | | | | .007 | | | .0004 |
| Zone 1 predominant | 54 (17) | 85 (23) | 27 (20) | 58 (25) | | | | | | | | |
| Azonal | 23 (7) | 43 (12) | 10 (7) | 33 (14) | | | | | | | | |
| Panacinar | 122 (38) | 124 (33) | 42 (31) | 82 (35) | | | | | | | | |
| Steatosis macro-vesicular type | | | | | | | | | | | | |
| Large droplet | 216 (66) | 308 (83) | 110 (80) | 198 (85) | | | | | | | | |
| Mixed large and small droplet | 101 (31) | 59 (16) | 25 (18) | 34 (15) | | | | | | | | |
| Small droplet | 8 (3) | 5 (1) | 3 (2) | 2 (1) | | | | | | | | |
| Microvesicular Steatosis | 6 (2) | 5 (1) | 3 (2) | 2 (1) | .76 | | | | .36 | | | .33 |
| Lobular inflammation score | | | | | | | | | | | | |
| 0 (no foci) | 2 (1) | 2 (1) | 0 (0) | 2 (1) | .82 | | | | .21 | | | .74 |
| 1 (<2 foci/20x hpf) | 189 (58) | 222 (59) | 75 (54) | 147 (62) | | | | | | | | |
| 2 (2-4 foci/20x hpf) | 110 (34) | 129 (34) | 56 (41) | 73 (31) | | | | | | | | |
| 3 (>4 foci/20x hpf) | 24 (7) | 21 (6) | 7 (5) | 14 (6) | | | | | | | | |
| Mean (SD) | 1.48±0.64 | 1.45±0.61 | 1.51±0.59 | 1.42±0.62 | .55 | | | | .18 | | | .26 |
| Portal inflammation score | 1.06±0.55 | 1.05±0.56 | 0.99±0.59 | 1.08±0.54 | .76 | | | | .11 | | | .64 |
| 0-None | 40 (12) | 50 (13) | 25 (18) | 25 (11) | .91 | | | | .11 | | | .82 |
| Mild | 226 (70) | 256 (69) | 89 (65) | 167 (71) | | | | | | | | |
| 2-More than mild | 59 (18) | 67 (18) | 23 (23) | 44 (19) | | | | | | | | |
| Ballooning injury score | | | | | | | | | | | | |
| 0-None | 201 (62) | 245 (66) | 89 (64) | 156 (66) | .21 | | | | .95 | | | .25 |
| Few | 84 (26) | 98 (26) | 37 (27) | 61 (26) | | | | | | | | |
| Many | 40 (12) | 31 (8) | (9) | 19 (8) | | | | | | | | |
| Mean (SD) | 0.50±0.71 | 0.43±0.64 | 0.44±0.65 | 0.42±0.64 | .13 | | | | .74 | | | .14 |
| Classic ballooning ^c | 19 (6) | 16 (4) | 7 (5) | 9 (4) | .34 | | | | .56 | | | .28 |
| Acidophil bodies | 177 (54) | 210 (56) | 84 (61) | 126 (53) | .70 | | | | .16 | | | .80 |

(Continues)

TABLE 2 (Continued)

| | None (n=325) | | Any (n=374) | | None vs Any P ^a | | Diffuse (n=236) | | Focal vs Diffuse P ^b | | None vs Diffuse P ^a | |
|----------------------------------|--------------|------------|-------------|------------|----------------------------|------------|-----------------|------------|---------------------------------|------------|--------------------------------|------------|
| | N (%) | or Mean±SD | N (%) | or Mean±SD | N (%) | or Mean±SD | N (%) | or Mean±SD | N (%) | or Mean±SD | N (%) | or Mean±SD |
| Megamitochondria | 27 (8) | | 42 (11) | | .21 | 9 (7) | 33 (14) | .03 | | | | .03 |
| Mallory bodies | 22 (7) | | 20 (5) | | .52 | 9 (7) | 11 (5) | .48 | | | | .29 |
| Fibrosis stage | 1.07±1.05 | | 0.98±0.99 | | .28 | 0.99±1.00 | 0.98±0.98 | .90 | | | | .31 |
| 0: None | 116 (36) | | 138 (37) | | .54 | 50 (36) | 88 (37) | .67 | | | | .51 |
| 1A: Mild perisinusoidal only | 28 (9) | | 24 (6) | | | 9 (7) | 15 (6) | | | | | |
| 1B: Moderate perisinusoidal | 13 (4) | | 15 (4) | | | 9 (7) | 6 (3) | | | | | |
| 1C: Periportal only | 77 (24) | | 108 (29) | | | 38 (28) | 70 (30) | | | | | |
| 2: Periportal and perisinusoidal | 45 (14) | | 50 (13) | | | 17 (12) | 33 (14) | | | | | |
| 3: Bridging fibrosis | 42 (13) | | 35 (9) | | | 13 (9) | 22 (9) | | | | | |
| 4: Cirrhosis | 3 (1) | | 4 (1) | | | 2 (1) | 2 (1) | | | | | |
| Diagnostic classification | | | | | | | | | | | | |
| Not NAFLD | 10 (3) | | 23 (6) | | .07 | 9 (7) | 14 (6) | .31 | | | | .20 |
| NAFLD, Not Steatohepatitis | 94 (29) | | 124 (33) | | | 49 (36) | 75 (32) | | | | | |
| Borderline, Zone 3 pattern | 59 (18) | | 46 (12) | | | 16 (12) | 30 (13) | | | | | |
| Borderline, Zone 1 pattern | 97 (30) | | 106 (28) | | | 31 (22) | 75 (32) | | | | | |
| Definite steatohepatitis | 65 (20) | | 75 (20) | | | 33 (24) | 42 (18) | | | | | |
| NAFLD activity score | | | | | | | | | | | | |
| 0 | 1 (0) | | 2 (1) | | .003 | 0 (0) | 2 (1) | .02 | | | | .0001 |
| 1 | 9 (3) | | 17 (5) | | | 8 (6) | 9 (4) | | | | | |
| 2 | 30 (9) | | 72 (19) | | | 23 (17) | 49 (21) | | | | | |
| 3 | 58 (18) | | 81 (22) | | | 17 (12) | 64 (27) | | | | | |
| 4 | 90 (28) | | 80 (21) | | | 35 (25) | 45 (19) | | | | | |
| 5 | 78 (24) | | 64 (17) | | | 29 (21) | 35 (15) | | | | | |
| 6 | 42 (13) | | 45 (12) | | | 22 (16) | 23 (10) | | | | | |
| 7 | 11 (3) | | 8 (2) | | | 3 (2) | 5 (2) | | | | | |
| 8 | 6 (2) | | 5 (1) | | | 1 (1) | 4 (2) | | | | | |
| Mean±SD | 4.2±1.5 | | 3.8±1.6 | | 0.001 | 4.0±1.6 | 3.6±1.6 | .04 | | | | <.0001 |

^aDerived from chi-squared test for nominal variables and t-test with unequal variance for continuous variables

^b3 cases had no steatosis; 136 cases are missing microgranulomas, large lipogranulomas, pigmented macrophages and glycogen nuclei—no longer scored

^cSevere ballooning and presence of Mallory-Denk bodies

TABLE 3 Multiple logistic regression^a of any vs. no glycogenosis and diffuse vs. focal glycogenosis on demographics, laboratory measures and histologic features in adults

| | Any vs No glycogenosis | | | Diffuse vs Focal glycogenosis | | |
|--|------------------------|--------------|---------|-------------------------------|--------------|---------|
| | Odds ratio | 95% CI | P-value | Odds ratio | 95% CI | P-value |
| Demographics | | | | | | |
| Age at biopsy-yr | 1.012 | 1.006, 1.027 | .003 | | | |
| Female sex vs male sex | 1.31 | 1.01, 1.71 | .04 | 2.04 | 1.41, 2.95 | <.001 |
| Hispanic (yes vs no) | | | | 1.58 | 0.97, 2.28 | .07 |
| Laboratory measures | | | | | | |
| Glucose-mg/dL | 1.005 | 1.001, 1.009 | .01 | | | |
| LDL-mg/dL | | | | 0.996 | 0.992, 1.001 | .10 |
| Insulin- μ U/mL | 1.013 | 1.002, 1.025 | .02 | 1.009 | 1.001, 1.017 | .02 |
| HOMA-IR- μ U/ mL ^b mg/dL/405 | 0.97 | 0.94, 1.02 | .06 | | | |
| Histologic features | | | | | | |
| Steatosis grade-score | 0.82 | 0.71, 0.94 | .006 | 0.53 | 0.43, 0.64 | <.001 |
| Steatosis type | | | .02 | | | |
| Mixed vs Large droplet | 0.67 | 0.50, 0.90 | | | | |
| Small vs Large droplet | 1.09 | 0.53, 2.24 | | | | |
| Microvesicular Steatosis (present vs absent) | | | | 0.56 | 0.33, 0.97 | .04 |
| Ballooning grade-score | 1.30 | 1.09, 1.55 | .003 | | | |
| Non-hepatocyte iron grade-score | 1.34 | 1.10, 1.62 | .003 | 1.26 | 0.97, 1.63 | .08 |
| Megamitochondria (present vs absent) | 1.48 | 1.13, 1.95 | .004 | 1.36 | 0.95, 1.94 | .10 |
| Acidophils (yes vs no) | 1.32 | 1.03, 1.69 | .03 | | | |
| Fibrosis-stage | 0.84 | 0.75, 0.95 | .005 | | | |
| Biopsy length-mm | 0.99 | 0.98, 1.00 | .14 | | | |

^aCovariates were selected using AIC criteria from two multiple logistic models regressing any (n = 672) vs no (n = 547) glycogenosis and regressing diffuse (n = 346) vs focal (n = 326) glycogenosis in adults with non-missing data on candidate set of age, gender, ethnicity, white race, BMI, weight, ALT, AST, alkaline phosphatase, triglycerides, cholesterol, HDL, LDL, glucose, insulin, HOMA-IR, HbA1c, diabetes, ballooning, lobular inflammation, portal inflammation, iron grade, Mallory bodies, steatosis grade, location and type, fibrosis stage, microvascular steatosis, megamitochondria, acidophils, non-hepatocyte iron grade and biopsy length.

mellitus.¹⁵ Glycogenic hepatopathy is associated with high transaminase elevations which may result in a liver biopsy. The diffuse hepatic glycogenosis seen in type 1 diabetes-associated glycogenic hepatopathy is rapidly reversible with good control of glucose levels. Adequate glucose control also results in reversal of hepatomegaly and return of transaminase levels to normal. Glycogenic hepatopathy typically shows diffusely pale hepatocytes and absence of histological evidence supporting of NAFLD. In this cohort, type 1 diabetes was rare (four adults and three children) and the overwhelming

majority of subjects with diabetes had type 2. Even though in our adult cases variables of insulin resistance (serum glucose levels and insulin) were associated with glycogenosis, there was no elevation of transaminases vs the nonglycogenotic (control) population. None of the subjects had a diagnosis of glycogen storage disease which would be exclusionary for entry in this NASH CRN study.

Why is glycogenosis so common in NAFLD and associated with decreased steatosis? There is a complex interplay between hepatocellular carbohydrate and lipid metabolism. These pathways are

TABLE 4 Multiple logistic regression^a of any vs no glycogenosis and diffuse vs focal glycogenosis on demographics, laboratory measures and histologic features in children

| | Any vs No glycogenosis | | | Diffuse vs Focal glycogenosis | | |
|----------------------------|------------------------|--------------|---------|-------------------------------|--------------|---------|
| | Odds ratio | 95% CI | P-value | Odds ratio | 95% CI | P-value |
| Demographics | | | | | | |
| Hispanic vs non-Hispanic | 1.53 | 1.03, 2.25 | .03 | | | |
| White vs non-white | 0.46 | 0.32, 0.67 | <.001 | | | |
| Weight-kg | 0.989 | 0.982, 0.996 | .002 | | | |
| Laboratory measures | | | | | | |
| Triglycerides-mg/dL | 1.004 | 1.002, 1.007 | .001 | | | |
| Total cholesterol-mg/dL | | | | 1.006 | 1.000, 1.023 | .07 |
| Alkaline phosphatase-U/L | | | | 1.002 | 1.000, 1.005 | .10 |
| AST-U/L | 0.997 | 0.994, 1.000 | .04 | | | |
| Glucose-mg/dL | 1.017 | 1.005, 1.030 | .007 | | | |
| Comorbidities | | | | | | |
| Diabetes (yes vs no) | | | | 0.20 | 0.05, 0.72 | .01 |
| Histologic features | | | | | | |
| Steatosis grade-score | 0.65 | 0.54, 0.79 | <.001 | 0.78 | 0.59, 1.03 | .08 |
| Steatosis type | | | <.001 | | | |
| Mixed vs large droplet | 0.40 | 0.27, 0.60 | | | | |
| Small vs large droplet | 0.39 | 0.11, 1.32 | | | | |
| Steatosis location | | | | | | .05 |
| Zone 1 vs Zone 3 | | | | 1.96 | 1.02, 3.76 | |
| Azonal vs Zone 3 | | | | 2.37 | 0.98, 5.74 | |
| Panacinar vs Zone 3 | | | | 2.06 | 1.13, 3.76 | |
| Acidophils (yes vs no) | | | | 0.63 | 0.39, 1.00 | .05 |

^aCovariates were selected using AIC criteria from two multiple logistic models regressing any (n = 355) vs no (n = 301) glycogenosis and regressing diffuse (n = 230) vs focal (n = 125) glycogenosis in children with non-missing data on candidate set of age, gender, ethnicity, white race, BMI, weight, ALT, AST, alkaline phosphatase, triglycerides, cholesterol, HDL, LDL, glucose, insulin, HOMA-IR, HbA1c, diabetes, ballooning, lobular inflammation, portal inflammation, iron grade, Mallory bodies, steatosis grade, location and type, fibrosis stage, microvascular steatosis, megamitochondria, acidophils, non-hepatocyte iron grade and biopsy length.

normally finely balanced, but they can become dysregulated and may shunt substrates from one pathway to another. For example, several studies have documented a link between high fructose diets and NAFLD.^{16,17} In humans and rodents, fructose induces glycogen synthesis to a greater degree than glucose,¹⁸⁻²⁰ and a prior adult NASH CRN study showed an inverse relationship between fructose intake and the severity of steatosis.²¹ Unfortunately no histological assessment of glycogenosis was available in that study, and data on fructose consumption was not available in the current study.

Furthermore, the inverse relationship between glycogenosis and steatosis is highlighted by a recent primate study. Baboons given a high-fat, high-simple-carbohydrate diet which induced maternal obesity during pregnancy resulted in offspring with more severe liver steatosis but a lesser degree of hepatic glycogen accumulation compared to offspring of controls.²²


The interconnection of hepatic carbohydrate and lipid metabolism, and the downstream effects on NAFLD progression risk is further illustrated by a recent important finding. In individuals at high risk of NAFLD, genetic variation (rs4841132 G < A) in protein

phosphatase 1 regulatory subunit 3B (PPP1R3B), a hepatic glycogen metabolism regulatory protein, is associated with reduced hepatic steatosis and protection against liver fibrosis, but results in increased liver glycogen.^{23,24} The authors speculate that this variant upregulates PPP1R3B levels, favouring energy storage as glycogen by shunting glucose away from glycolysis while suppressing de novo lipogenesis. The authors suggest that this variant is protective against steatosis and fibrosis.

Some of the limitations of this study are the lack of dietary information, the lack of data on prior corticosteroid use, the lack of clinical data regarding the presence or absence of hepatomegaly, the lack of imaging studies specifically addressing the presence or absence of hepatic glycogen, the lack of quantitative analysis of hepatocellular glycogen and triglyceride levels in these samples, and the lack of PPP1R3B genetic data in this cohort.

In summary, this study demonstrates that glycogenosis is commonly seen in the context of adult and paediatric NAFLD. In adults, glycogenosis is associated with older age, female sex, and higher blood glucose and insulin levels. Furthermore, we show that

FIGURE 3 Ballooned hepatocytes with cytoplasmic glycogen. (A, B). H&E stained sections of ballooned hepatocytes with prominent glycogen aggregates (200x and 400x respectively). C and D. PAS stained section of ballooned hepatocytes with cytoplasmic glycogen (200x and 400x respectively)



glycogenosis is associated with a higher degree of ballooning, but a lower steatosis grade and lower fibrosis stage. NAFLD glycogenosis should not be confused with hepatocyte ballooning, glycogenic hepatopathy or glycogen storage disorders. Although dysregulated lipid metabolism has been well documented in the setting of fatty liver disease, further studies are warranted to better understand the causes and effects of altered carbohydrate metabolism in NAFLD.

CONFLICT OF INTEREST

There are no conflicts of interest.

ETHICS APPROVAL AND PATIENT CONSENT STATEMENT

Each participating institution obtained IRB approval to enrol patients into the parent NASH CRN studies. Written consents was signed by participating patients.

PERMISSION TO REPRODUCE MATERIAL FROM OTHER SOURCES

No material is obtained/reproduced from other sources.

ORCID

David E Kleiner  <https://orcid.org/0000-0003-3442-4453>

Cynthia D Guy  <https://orcid.org/0000-0002-4959-751X>

REFERENCES

1. Bechmann LP, Hannivoort RA, Gerken G, Hotamisligil GS, Trauner M, Canbay A. The interaction of hepatic lipid and glucose metabolism in liver diseases. *J Hepatol.* 2012;56(4):952-964.
2. Schild MH, Guy CD. Nonalcoholic steatohepatitis: histopathology basics within a broader context. *Surg Pathol Clin.* 2018;11(2):267-285.
3. Neuschwander-Tetri BA, Clark JM, Bass NM, et al. Clinical, laboratory and histological associations in adults with nonalcoholic fatty liver disease. *Hepatol.* 2010;52(3):913-924.
4. Chalasani N, Wilson L, Kleiner DE, et al. Relationship of steatosis grade and zonal location to histological features of steatohepatitis in adult patients with non-alcoholic fatty liver disease. *J Hepatol.* 2008;48(5):829-834.
5. Torbenson M, Chen Y-Y, Brunt E, et al. Glycogenic hepatopathy: an underrecognized hepatic complication of diabetes mellitus. *Am J Surg Pathol.* 2006;30(4):508-513.
6. Kleiner DE, Brunt EM, Van Natta M, et al. Design and validation of a histological scoring system for nonalcoholic fatty liver disease. *Hepatol.* 2005;41(6):1313-1321.
7. Adeva-Andany MM, González-Lucán M, Donapetry-García C, Fernández-Fernández C, Ameneiros-Rodríguez E. Glycogen metabolism in humans. *BBA Clin.* 2016;5:85-100.
8. Brunt EM, Janney CG, Bisceglie AM, et al. Nonalcoholic steatohepatitis: a proposal for grading and staging the histological lesions. *Am J Gastroenterol.* 1999;94(9):2467-2474.
9. Lackner C, Gogg-Kamerer M, Zatloukal K, et al. Ballooned hepatocytes in steatohepatitis: the value of keratin immunohistochemistry for diagnosis. *J Hepatol.* 2008;48(5):821-828.
10. Caldwell S, Ikura Y, Dias D, et al. Hepatocellular ballooning in NASH. *J Hepatol.* 2010;53(4):719-723.
11. Younossi ZM, Gramlich T, Liu YC, et al. Nonalcoholic fatty liver disease: assessment of variability in pathologic interpretations. *Mod Pathol.* 1998;11(6):560-565.
12. Kleiner DE, Brunt EM, Belt P, et al. Extending the ballooning score beyond 2: A proposal for a new balloon score. *Hepatol.* 2015;157:62.
13. Mukewar S, Sharma A, Lackore KA, et al. Clinical, biochemical, and histopathology features of patients with glycogenic hepatopathy. *Clin Gastroenterol Hepatol.* 2017;15(6):927-933.
14. Iancu TC, Shiloh H, Dembo L. Hepatomegaly following short-term high-dose steroid therapy. *J Pediatr Gastroenterol Nutr.* 1986;5(1):41-46.
15. Sherigar JM, Castro JD, Yin YM, et al. Glycogenic hepatopathy: a narrative review. *World J Hepatol.* 2018;10(2):172-185.

16. Ouyang X, Cirillo P, Sautin Y, et al. Fructose consumption as a risk factor for non-alcoholic fatty liver disease. *J Hepatol*. 2008;48(6):993-999.
17. Thuy S, Ladurner R, Volynets V, et al. Nonalcoholic fatty liver disease in humans is associated with increased plasma endotoxin and plasminogen activator inhibitor 1 concentrations and with fructose intake. *J Nutr*. 2008;138(8):1452-1455.
18. Decombaz J, Jentjens R, Ith M, et al. Fructose and galactose enhance postexercise human liver glycogen synthesis. *Med Sci Sports Exerc*. 2011;43(10):1964-1971.
19. Nilsson LH, Hultman E. Liver and muscle glycogen in man after glucose and fructose infusion. *Scand J Clin Lab Invest*. 1974;33(1):5-10.
20. Nishi T, Kido Y, Ogawa A, et al. Effect of fructose on glycogen synthesis in the perfused rat liver. *Biochem Int*. 1990;20(2):329-335.
21. Abdelmalek MF, Suzuki A, Guy C, et al. Increased fructose consumption is associated with fibrosis severity in patients with nonalcoholic fatty liver disease. *Hepatol*. 2010;51(6):1961-1971.
22. Puppala S, Li C, Glenn JP, et al. Primate fetal hepatic responses to maternal obesity: epigenetic signalling pathways and lipid accumulation. *J Physiol*. 2018;596:5823-5837.
23. Kahali B, Halligan B, Speliotes E. Insights from genome-wide association analyses of non alcoholic fatty liver disease. *Sem Liver Dis*. 2015;35(4):375-391.
24. Dongiovanni P, Meroni M, Mancina RM, et al. Protein phosphatase 1 regulatory subunit 3B gene variation protects against hepatic fat accumulation and fibrosis in individuals at high risk of nonalcoholic fatty liver disease. *Hepatol Commun*. 2018;2:666-675.

How to cite this article: Allende DS, Gawrieh S, Cummings OW, et al. Glycogenesis is common in nonalcoholic fatty liver disease and is independently associated with ballooning, but lower steatosis and lower fibrosis. *Liver Int*. 2021;41:996-1011. <https://doi.org/10.1111/liv.14773>

APPENDIX

MEMBERS OF THE NONALCOHOLIC STEATOHEPATITIS CLINICAL RESEARCH NETWORK ADULT AND PEDIATRIC CLINICAL CENTERS

Baylor College of Medicine, Houston, TX: Donna Garner, CPNP; Paula Hertel, MD; Ryan Himes, MD; Alicia Lawson, BS; Tamir Miloh, MD; Nicole Triggs, CPNP.

Cincinnati Children's Hospital Medical Center, Cincinnati, OH: Kristin Bramlage, MD; April Carr, BS, CCRP; Kim Cecil, PhD; Meghan McNeill, MS; Marialena Mouzaki, MD; Andrew Trout, MD; Stavra Xanthakos, MD; Kimberlee Bernstein, BS, CCRP (2012-2018); Stephanie DeVore, MSPH (2009-2011); Rohit Kohli, MD (2009-2016); Kathleen Lake, MSW (2009-2012); Daniel Podberesky, MD (2009-2014); Alex Towbin, MD (2009-2016).

Cleveland Clinic Foundation, Cleveland, OH: Daniela Allende, MD; Annette Bellar, MSLA; Jaividhya Dasarathy, MD; Srinivasan Dasarathy, MD; Arthur J. McCullough, MD; Revathi Penumatsa, MBBS, MPH; Jaspreet S. Reen, MBBS.

Columbia University, New York, NY: Joel E. Lavine, MD, PhD; Ali Mencin, MD; Elena Reynoso, MD.

Duke University Medical Center, Durham, NC: Manal F. Abdelmalek, MD, MPH; Mustafa Bashir, MD; Stephanie Buie; Anna Mae Diehl, MD; Cynthia Guy, MD; Christopher Kigongo, MB, CHB; Mariko Kopping, MS, RD; Dawn Piercy, MS, FNP; Naglaa Tawadrous.

Emory University, Atlanta, GA: Adina Alazraki, MD; Rebecca Cleeton, MPH, CCRP; Maria Cordero, CCRP; Albert Hernandez; Saul Karpen, MD, PhD; Jessica Cruz Munos (2013-2015); Nicholas Raviele (2012-2014); Miriam Vos, MD, MSPH, FAHA.

Indiana University School of Medicine, Indianapolis, IN: Molly Bozic, MD; Laura Carr, RN; Naga Chalasani, MD; Oscar W. Cummings, MD; Samer Gawrieh, MD; Ann Klipsch, RN; Jean P. Molleston, MD; Emily Ragozzino; Linda Ragozzino, RN; Dawn Rank, RN; Girish Rao, MD; Niharika Samala, MD; Raj Vuppalanchi, MD.

Johns Hopkins Hospital, Baltimore, MD: Kimberly Kafka, RN; Ann Scheimann, MD.

NORTHWESTERN UNIVERSITY FEINBERG SCHOOL OF MEDICINE/ANN & ROBERT H. LURIE CHILDREN'S

Hospital of Chicago: Mark H. Fishbein, MD; Joy Ito, RN; Saeed Mohammad, MD; Peter F. Whittington, MD (2009-2017).

Saint Louis University, St Louis, MO: Sarah Barlow, MD (2002-2007); Elizabeth M. Brunt, MD (2002-2008); Danielle Carpenter, MD; Theresa Cattoor, RN; Jose Derdoy, MD (2007-2011); Janet Freebersyser, RN; Ajay Jain MD; Debra King, RN (2004-2015); Jinping Lai, MD (2015-2016); Brent A. Neuschwander-Tetri, MD; Joan Siegner, RN (2004-2015); Susan Stewart, RN (2004-2015); Susan Torretta; Kristina Wriston, RN (2015).

Swedish Medical Center, Seattle, WA: Maria Cardona Gonzalez; Jodie Davila; Kris V. Kowdley, MD; Nizar Mukhtar, MD; Michelle Poitevin; Brook Quist; Sherilyn Soo.

University of California San Diego, San Diego, CA: Jorge Angeles, MD; Jennifer Arin; Cynthia Behling, MD, PhD; Craig Bross; Carissa Carrier; Diana De La Pena; Janis Durelle; Mary Catherine Huckaby, MD; Joel E. Lavine, MD, PhD (2002-2010); Rohit Loomba, MD, MHS; Egbert Madamba; Michael S. Middleton, MD, PhD; Kimberly Newton, MD; Jeffrey B. Schwimmer, MD; Claude Sirlin, MD; Patricia Ugalde-Nicalo, MD.

University of California San Francisco, San Francisco, CA: Nathan M. Bass, MD, PhD (2002-2011); Danielle Brandman, MD, MAS; Jesse Courtier, MD; Ryan Gill, MD, PhD; Bilal Hameed, MD; Camille Langlois, MS; Jacquelyn Maher, MD; Emily Rothbaum Perito, MD; Philip Rosenthal, MD; Nicholas Slater, MD; Norah Terrault, MD, MPH; Patrika Tsai, MD; Ashley Ungermann, MS.

University of Washington Medical Center and Seattle Children's Hospital, Seattle, WA: Niviann Blondet, MD; Kara Cooper; Karen Murray, MD; Randolph Otto, MD; Matthew Yeh, MD, PhD; Melissa Young.

Virginia Commonwealth University, Richmond, VA: Sherry Boyett, RN, BSN; Melissa J. Contos, MD; Velimir AC Luketic, MD; Arun J. Sanyal, MD; Jolene Schlosser, RN, BSN; Mohammad S. Siddiqui, MD.

Washington University, St. Louis, MO: Elizabeth M. Brunt, MD (2008-2015); Kathryn Fowler, MD (2012-2015).

Resource Centers.

National Cancer Institute, Bethesda, MD: David E. Kleiner, MD, PhD.
National Institute of Diabetes and Digestive and Kidney Diseases,
Bethesda, MD: Edward C. Doo, MD; Sherry Hall, MS; Jay H.
Hoofnagle, MD; Patricia R. Robuck, PhD, MPH (2002-2011); Averell
H. Sherker, MD; Rebecca Torrance, RN, MS.

Data Coordinating Center, Johns Hopkins University, Bloomberg
School of Public Health, Baltimore, MD: Patricia Belt, BS; Jeanne

M. Clark, MD, MPH; John Dodge (2002-2018); Michele Donithan,
MHS (2002-2017); Milana Isaacson, BS (2002-2018); Mariana Lazo,
MD, PhD, ScM; Jill Meinert; Laura Miriel, BS; Emily P. Sharkey, MPH,
MBA; Jacqueline Smith, AA; Michael Smith, BS; Alice Sternberg,
ScM; James Tonascia, PhD; Mark L. Van Natta, MHS; Annette
Wagoner; Laura A. Wilson, ScM; Goro Yamada, PhD, MHS, MHS,
MMS; Katherine P. Yates, ScM.