

SCIENTIFIC REPORTS



OPEN

Multi-state study of *Enterobacteriaceae* harboring extended-spectrum beta-lactamase and carbapenemase genes in U.S. drinking water

Windy D. Tanner¹, James A. VanDerslice¹, Ramesh K. Goel¹, Molly K. Leecaster^{1,2}, Mark A. Fisher¹, Jeremy Olstadt³, Catherine M. Gurley⁴, Anderson G. Morris⁴, Kathryn A. Seely⁴, Leslie Chapman⁵, Michelle Korando⁵, Kalifa-Amira Shabazz⁶, Andrea Stadsholt⁷, Janice VanDeVelde⁷, Ellen Braun-Howland⁸, Christine Minihane⁸, Pamela J. Higgins⁹, Michelle Deras¹⁰, Othman Jaber¹¹, Dee Jette¹² & Adi V. Gundlapalli^{1,2}

Community-associated acquisition of extended-spectrum beta-lactamase- (ESBL) and carbapenemase-producing *Enterobacteriaceae* has significantly increased in recent years, necessitating greater inquiry into potential exposure routes, including food and water sources. In high-income countries, drinking water is often neglected as a possible source of community exposure to antibiotic-resistant organisms. We screened coliform-positive tap water samples ($n = 483$) from public and private water systems in six states of the United States for *bla*_{CTX-M}, *bla*_{SHV}, *bla*_{TEM}, *bla*_{KPC}, *bla*_{NDM}, and *bla*_{OXA-48}-type genes by multiplex PCR. Positive samples were subcultured to isolate organisms harboring ESBL or carbapenemase genes. Thirty-one samples (6.4%) were positive for *bla*_{CTX-M} ESBL-type *bla*_{SHV} or *bla*_{TEM} or *bla*_{OXA-48}-type carbapenemase genes, including at least one positive sample from each state. ESBL and *bla*_{OXA-48}-type *Enterobacteriaceae* isolates included *E. coli*, *Kluyvera*, *Providencia*, *Klebsiella*, and *Citrobacter* species. The *bla*_{OXA-48}-type genes were also found in non-fermenting Gram-negative species, including *Shewanella*, *Pseudomonas* and *Acinetobacter*. Multiple isolates were phenotypically non-susceptible to third-generation cephalosporin or carbapenem antibiotics. These findings suggest that tap water in high income countries could serve as an important source of community exposure to ESBL and carbapenemase genes, and that these genes may be disseminated by non-*Enterobacteriaceae* that are not detected as part of standard microbiological water quality testing.

Antibiotic-resistant infections are responsible for an estimated 2 million illnesses and 23,000 deaths in the United States each year¹. The rising prevalence of multidrug-resistant bacteria is especially alarming, as infections with these organisms have led to increasing use of broad spectrum antibiotics such as third and fourth generation cephalosporin and carbapenem antibiotics^{1,2}. Enzymes such as extended-spectrum beta-lactamases (ESBLs) and carbapenemases can render these antibiotics ineffective. *Enterobacteriaceae* harboring these enzymes are ranked among the most urgent antibiotic resistance threats according to the U.S. Centers for Disease Control and

¹University of Utah, Salt Lake City, UT, USA. ²VA Salt Lake City Healthcare System, Salt Lake City, UT, USA. ³Wisconsin State Laboratory of Hygiene, Madison, WI, USA. ⁴Arkansas Department of Health Public Health Laboratory, Little Rock AR, USA. ⁵Illinois Department of Public Health, Carbondale, IL, USA. ⁶Illinois Department of Public Health, Chicago, IL, USA. ⁷Illinois Department of Public Health, Springfield, IL, USA. ⁸Wadsworth Center, Albany, NY, USA. ⁹Pennsylvania Department of Environmental Protection, Harrisburg, PA, USA. ¹⁰Weber Basin Water Conservancy District, Layton, UT, USA. ¹¹Utah Public Health Laboratory, Taylorsville, UT, USA. ¹²Davis County Health Department, Clearfield, UT, USA. Correspondence and requests for materials should be addressed to W.D.T. (email: windy.tanner@hsc.utah.edu)




Figure 1. Water sample coliform screening process.

Prevention and the World Health Organization^{1,3}. Additionally, genes encoding these enzymes are often found on mobile genetic elements that can be transferred horizontally to other bacterial species⁴.

ESBL- and carbapenemase-producing bacteria are commonly associated with healthcare contact¹; however, community-associated infections have significantly increased in recent years⁵. Between 2009 and 2011, the occurrence of ESBL-producing bacteria in community-associated infections increased from 3.1% to 12.6%, while the occurrence in hospital-associated infections remained the same⁵. It is estimated that as many as two-thirds of all ESBL-producing *Enterobacteriaceae* infections are community-associated^{5,6}. In 2013, the U.S. Centers for Disease Control warned that spread of carbapenem-resistant *Enterobacteriaceae* (CRE) into the community could reasonably be expected, as experienced with ESBLs¹. A 2017 review on CRE in the community noted that the prevalence of CRE among U.S. community-associated study samples ranged from 5.6 to 10.8%⁷.

ESBL and carbapenemase genes are frequently found in *Enterobacteriaceae* such as *Klebsiella pneumoniae*, *Escherichia coli*, *Enterobacter cloacae*, and *Citrobacter* species⁴, but can also be found in non-fermenting Gram-negative species⁸. These bacteria are most commonly spread via fecal-oral transmission routes, including direct transmission (e.g. via hands) and indirect transmission (e.g. via the environment). *Enterobacteriaceae*, including ESBL- and carbapenemase-producing strains, have been reported in a number of environmental compartments including food, animals, surface waters, and drinking water^{9–11}. Studies reporting carbapenemase or ESBL genes in drinking water have largely been performed in low-income countries^{9,12}. In high income countries, reports of ESBL- and carbapenemase-producing bacteria in drinking water have been limited to single-cases or intrinsic genes in nonpathogenic environmental bacterial species^{10,13}, but comprehensive studies are lacking.

In clinical infections, ESBL and carbapenemase genes are most frequently found in *Enterobacteriaceae* species, but these organisms are typically uncommon in chlorinated public drinking water supplies in high income countries. To potentially increase detection of ESBL and carbapenemase genes in U.S. drinking water, we targeted water samples testing positive for *Enterobacteriaceae* (coliform bacteria). The objectives of our study were (1) to determine whether ESBL- and carbapenemase-producing genes are present in U.S. drinking water samples that have tested positive for *E. coli* or total coliform bacteria, (2) to describe the sample and water system characteristics associated with samples testing positive for ESBL or carbapenemase genes, and (3) to determine if the ESBL and carbapenemase genes are present in viable bacteria isolated from the coliform-positive water samples.

Methods

Water sample collection and coliform testing. Between July 2015 and November 2016, regulatory and investigational drinking water samples testing positive for *E. coli* or total coliform bacteria (*Enterobacteriaceae*) were acquired from multiple local and state public health laboratories that perform regulatory water quality testing for water utilities in their state or county. Participating laboratories included the state public health and environmental laboratories in Wisconsin (99 samples), New York (22 samples), Pennsylvania (69 samples), Illinois (125 total samples from three laboratories), Arkansas (64 samples), and Utah. Additionally, one water utility and one county health laboratory in Utah supplied coliform-positive samples (total of 104 Utah samples). Sample acquisition was opportunistic, and may not have consistently included all coliform-positive samples from each site. Private well samples were only included in the study if positive for *E. coli*, to avoid testing a high proportion of private well samples, which frequently test positive for coliform bacteria. Drinking water samples collected for regulatory water quality testing are required to be 100 milliliters in volume and must be preserved with sodium thiosulfate and tested at the laboratory within 30 hours of collection, preferably held at a temperature between 0 and 10 degrees Celsius during transport¹⁴.

Upon arrival to each public health laboratory location, drinking water samples were tested using a conventional enzyme-substrate method that indicates the presence of *E. coli* and total coliform bacteria (Fig. 1)¹⁵. The substrate reagent includes a nutrient medium that creates culture conditions in the water sample and promotes bacterial growth and has chromogenic and fluorogenic indicators to detect total coliforms and *E. coli*, respectively. Samples were tested in either a presence/absence format using the original 100-mL collection vessel, or in a quantitative format by dispensing the 100 mL sample into a multi-well tray (Quanti-Tray, IDEXX, Westbrook, ME) for quantification by the most-probable-number technique. When *E. coli* and/or coliform bacteria were detected in cultured presence/absence samples, 1 mL of the positive enriched sample was placed in a sterile cryovial with

1 mL 40% glycerol (final concentration 20% glycerol). If a sealed Quanti-Tray sample was positive, the back of the tray was disinfected and a sterile syringe was used to extract the culture liquid from positive wells. When multiple wells indicated coliforms, a composite 1 mL of enriched sample was produced by extracting liquid from several wells, and the composite was mixed with glycerol as described above. All samples were cryogenically frozen, and samples from locations other than Utah were shipped on dry ice overnight to the research laboratory in Salt Lake City, Utah.

Resistance gene detection and PCR amplicon sequencing. Upon arrival to the research laboratory, preserved samples were screened for the three ESBL genes most common in the U.S. (*bla_{SHV}*, *bla_{TEM}*, and *bla_{CTX}*) by multiplex PCR, as previously described¹⁶. Samples were also screened for the three most common carbapenemase genes (*bla_{OXA-48}*-type, *bla_{NDM}*, and *bla_{KPC}*) by a separate multiplex PCR¹⁷. Primer sequences are available in Supplemental Table 1. Four microliters of the preserved sample culture was directly used as a template for the PCR in a total reaction volume of 25 μ L. Positive controls included American Type Culture Collection isolate BAA-2146 for *bla_{SHV}*, *bla_{TEM}*, *bla_{CTX-M}*, and *bla_{NDM}* detection, a clinical KPC-producing *K. pneumoniae* for *bla_{KPC}* detection, and an OXA-48-producing *Shewanella* for *bla_{OXA-48}* detection. PCR products were run on a 1% agarose gel, and amplicon from samples putatively positive for any of the genes of interest were sequenced by the Sanger method using the forward and reverse primers used for PCR screening. If samples were positive for multiple genes, PCR was repeated with individual primer pairs in separate reactions, followed by amplicon sequencing. Sequences were searched against the Comprehensive Antibiotic Resistance Database (CARD) and the GenBank (BLASTn) database to confirm the presence of one of the six target ESBL or carbapenemase genes. SHV and TEM variants identified by these databases were compared against the Lahey Clinic designation of ESBL-type *bla_{SHV}* and *bla_{TEM}* genes¹⁸. Lack of specificity of the multiplex CTX-M primer can result in amplification of several other beta-lactamase genes, such as *bla_{OXY}*¹⁶; as a result, chromatograms from all CTX-M -positive amplicon sequences were closely inspected to see if multiple genes may have been present in the sample. In the case of mixed chromatograms, samples were tested as described above using a CTX-M group-specific multiplex PCR¹⁹ (Supplemental Table 1). Samples confirmed as positive for any of the target genes by Sanger sequencing were subsequently tested for the specific gene using primers that amplified larger segments of the genes to better identify the specific alleles^{20–23}.

Bacterial isolation and identification. Bacteria carrying the target resistance genes were isolated through selective and non-selective culture of the preserved samples as previously described^{24,25}. Samples confirmed for the presence of *bla_{TEM}*, *bla_{SHV}*, or *bla_{CTX-M}* genes were plated on CHROMagar Orientation™ agar plates (DRG International, Springfield, NJ) with and without a proprietary ESBL supplement. Approximately 50 μ L of the preserved sample was spread onto each plate, and cefotaxime (30 μ g), ceftazidime (30 μ g), and aztreonam (30 μ g) disks were placed on the non-selective CHROMagar plates. After overnight incubation, colonies on the ESBL CHROMagar plate and colonies falling within a distinct zone forming around the disks on the non-selective plate were tested by PCR for *bla_{TEM}*, *bla_{SHV}*, or *bla_{CTX-M}* genes using the primer set from the ESBL multiplex PCR.

For samples confirmed as having a *bla_{OXA-48}*-type gene, 50 μ L of preserved sample was spread onto an mSuperCARBA™ plate (DRG International, Springfield, NJ) and a non-selective CHROMagar Orientation™ plate (DRG International, Springfield, NJ) with a temocillin disk (Rosco, Taastrup, Denmark). In cases where the *bla_{OXA-48}*-type producer could not be isolated by direct plating, 20 μ L of the original culture sample was added to a Luria broth with 0.5, 1, 2, and 4 μ g/mL imipenem. Broths testing positive by PCR were plated as described above and to sheep's blood agar with 0.125 μ g/mL ertapenem. Colonies growing on the mSuperCARBA™ plate and colonies growing within the distinct zone that formed around the temocillin disk were tested for *bla_{OXA-48}*-type genes using the OXA-48 primer set from the carbapenemase gene multiplex PCR and confirmed by Sanger sequencing of the amplicon. Bacterial species identification was performed by MALDI-TOF (Biotyper, Bruker, Bellerica, MA) using the full spectral library which is composed of spectra representing roughly 2750 species of microorganisms from approximately 470 genera²⁶.

Susceptibility testing. Minimum inhibitory concentrations of isolates were determined using Thermo Scientific™ Sensititre™ Extended Spectrum Beta-lactamase plates (Trek Diagnostic Systems, Inc., Independence, OH) according to manufacturer's directions. For isolates in which the genes were lost upon subculture, fresh colonies from the original sample were used in preparation of the antimicrobial susceptibility testing inoculum, if sufficient growth was present.

Supplementary sample data. Laboratories supplying coliform-positive samples also provided limited data on sample characteristics such as water source (e.g. groundwater, [treated] surface water, or blended), sample type (regulatory or investigational), and system type (e.g. community public water system [CPWS], non-community public water system [NCPWS], or private well, as defined by the U.S. Environmental Protection Agency)²⁷. Sample results were described with respect to these characteristics. Statistical analyses to test specific hypotheses or make statements of inference would have been inappropriate because the sample collection was opportunistic.

Results

Sample characteristics. A total of 483 non-duplicate samples, representing 361 public and private water systems, were collected. Of the 483 samples, 27% were from private water systems, 31% were from CPWSs, and 42% were from NCPWSs. Spring water sources comprised 5% of the samples, while surface and ground water sources made up 9% and 78%, respectively. Approximately 6% of the samples were blended, meaning that they were from systems where both groundwater and treated surface water were supplied to consumers during periods of high demand.

PCR detection of ESBL and carbapenemase genes and amplicon sequencing. The 483 samples were screened for *bla*_{TEM}, *bla*_{SHV}, *bla*_{CTX-M}, *bla*_{KPC}, *bla*_{NDM}, and *bla*_{OXA-48}-type genes. Sixty-four samples appeared to be positive for *bla*_{CTX-M} by initial PCR and agarose gel screening; however, comparison of PCR amplicon sequences to the GenBank database revealed that the majority of the putative *bla*_{CTX-M} positives were attributed to extended-spectrum beta-lactamase genes that are typically chromosomally-encoded in various *Enterobacteriaceae* species, including *bla*_{RAHN} (*Rahnella aquatilis*), *bla*_{FONA} (*Serratia fonticola*), *bla*_{OXY} (*Klebsiella oxytoca*), *bla*_{SMO} (strain RUS)²⁸, and *Citrobacter amalonaticus* class-A beta lactamase. Because these were not target genes in the study, we did not pursue any further analysis of these samples.

Thirty-one samples (6.4%) from twenty-six (7.2%) of the water systems were positive for the target *bla*_{CTX-M}, *bla*_{OXA-48}-type, or ESBL-type *bla*_{TEM} or *bla*_{SHV} genes. Thirteen of the samples (2.7%) from twelve different water systems (3.3%) were confirmed to have *bla*_{CTX-M} genes based on amplicon sequences. Thirteen (2.7%) and ten (2.1%) samples were positive for *bla*_{SHV} and *bla*_{TEM} genes, respectively. One SHV-positive amplicon sequence most closely matched an ESBL-type *bla*_{SHV} (*bla*_{SHV-38}) and one TEM-positive amplicon sequence closely matched both ESBL- and non-ESBL-type *bla*_{TEM} genes. The carbapenemase multiplex PCR screen did not reveal any NDM- or KPC-positive samples; however, *bla*_{OXA-48}-type genes were detected in 19 samples (3.9%) from 15 (4.2%) of the water systems. The samples with *bla*_{OXA-48}-type genes were from four states and represented between 2–7% of samples from those states. Three of the samples that tested positive for *bla*_{OXA-48}-type genes were also positive for *bla*_{CTX-M} genes and were each from different states. At least one ESBL or carbapenemase gene was detected in all six states comprising between 1% and 15% of the coliform-positive samples tested from each state. Of the 185 water systems with repeat samples, six had samples that tested positive for at least one ESBL or carbapenemase gene; half of these water systems had only one positive sample and one water system had all three samples positive.

Isolation of ESBL- or carbapenemase-producing bacteria and antimicrobial susceptibility testing.

Bacteria carrying target ESBL or carbapenemase genes were isolated from 21 of the 31 positive samples (Table 1). The *bla*_{CTX-M} gene was detected in *Klebsiella oxytoca*, *Citrobacter freundii* complex, *Kluyvera ascorbata* and *Kluyvera georgiana*, the latter two being the progenitor of *bla*_{CTX-M} genes. The ESBL-type *bla*_{SHV-38} gene was present in *Klebsiella pneumoniae* and a *bla*_{TEM} variant gene exhibiting an ESBL phenotype was present in *E. coli*. Species carrying *bla*_{OXA-48}-type genes included *E. coli*, *Providencia rettgeri*, *Acinetobacter baumannii* complex, *Pseudomonas putida*, *Pseudomonas koreensis*, *Pandoraea sputorum*, *Shewanella putrefaciens*, and other *Shewanella* (n = 3) and *Pseudomonas* (n = 1) that could not be identified at the species level. The stability of ESBL and *bla*_{OXA-48}-type genes were noted based on the number of subcultures where the resistance gene could still be detected in culture. The ESBL genes were very stable and were still detected after two or more subcultures. The *bla*_{OXA-48}-type genes were typically lost after one subculture in non-*Shewanella* non-fermenting Gram-negative rods, and after two or more subcultures in *Enterobacteriaceae*, with or without selective pressure from imipenem concentrations ranging from 0.125 ug/mL to 4 ug/mL. Nucleic acid sequences of the PCR amplicon from isolates and from positive samples where an isolate could not be recovered are available in GenBank (Table 1). Antimicrobial susceptibility testing results for selected isolates are presented in Table 2.

Resistance gene presence and tap water source characteristics. The *bla*_{CTX-M}, *bla*_{OXA-48}-type, or ESBL-type *bla*_{TEM} or *bla*_{SHV} genes were found in 5.6% of samples from ground water sources and 7.0% of samples from treated surface water sources, and from 10.0% of private and 5.1% of public water systems: 8.7% of CPWS and 2.5% of NCPWS samples. Overall, 6.8% of the coliform-positive samples from public water systems also tested positive for *E. coli*; however, when considering those samples that tested positive for an ESBL or *bla*_{OXA-48}-type gene, 22.2% were positive for *E. coli*.

Discussion

Community-associated antibiotic-resistant infections caused by ESBL- and carbapenemase-producing bacteria have increased significantly in the U.S. over the past decade. Many possible transmission routes have been studied; but in high-income countries, drinking water has not been adequately assessed as a potential source. We detected ESBL genes (*bla*_{CTX-M}, *bla*_{TEM}, or *bla*_{SHV}) or *bla*_{OXA-48}-type carbapenemase genes in more than 6% of the coliform-positive U.S. drinking water samples screened. Coliform-positive drinking water samples were targeted for testing because ESBL and carbapenemase genes are most commonly found in *Enterobacteriaceae* in clinical settings.

Non-*E. coli* coliforms, such as *Klebsiella*, *Citrobacter*, *Enterobacter*, and *Serratia* species are considered to be non-pathogenic by regulatory agencies and are primarily used as indicators of potential fecal contamination or water distribution system breaches. However, it is important to consider that these species are also some of the most common carriers of ESBL and carbapenemase genes, regardless of pathogenicity²⁴. In 2015, the presence of coliform bacteria was reported in 1909 U.S. water systems serving over 10 million in total people, although audits and other compliance reports have found that underreporting of drinking water contaminants in public water systems is likely a widespread issue²⁹. Coliform bacteria are even more common in private wells, which supply approximately 15% of the U.S. population with drinking water³⁰.

This study utilized a convenience sample of coliform-positive water samples from state and local public health and environmental laboratories to increase the potential for finding ESBL- or carbapenemase-producing *Enterobacteriaceae*, not to estimate the overall frequency of occurrence of these genes in U.S. water supplies. The study design limits the conclusions that can be drawn regarding the overall prevalence of these genes in U.S. water systems and limits access to some information associated with individual samples, such as the type of water treatment used by the water systems. There is also a slight possibility that some coliform-positive samples resulted

Sample ID	Gene variant or group with highest DNA sequence similarity ^a	Gene homology with reference sequence	Protein homology with reference sequence	GenBank Nucleotide Reference (GenBank Protein reference)	Organism	<i>E. coli</i> present/absent in sample	System type ^b	Water source	GenBank Accession Number
A1 ^c	CTX-M-205	79%	86%	MG028655 (ATJ25945.1)	<i>Citrobacter freundii</i> -complex	Absent	CPWS	Ground	MG560059
A2 ^c	CTX-M-205	80%	88%	MG028655	<i>Citrobacter freundii</i> -complex	Present	CPWS	Ground	MG560060
A3a	CTX-M-40 or CTX-M-63	84%	94%	NG_048991 NG_049014	<i>Kluyvera georgiana</i>	Present	Private	Ground	MG560061
A3b	OXA-181	100%	100%	CP023897	<i>Acinetobacter baumannii</i> complex	Present	Private	Ground	MG560077
A4	CTX-M-40 or CTX-M-63	98%	99%	NG_048991 NG_049014	<i>Kluyvera georgiana</i>	Absent	CPWS	Ground	MG560062
A5	CTX-M-40 or CTX-M-63	84%	94%	NG_048991 NG_049014	<i>Kluyvera georgiana</i>	Absent	CPWS	Ground	MG560063
A6	CTX-M-9 family	86% ^d	95%	NG_049028.1	Not Isolated	Absent	CPWS	Surface	MG560055
B1	CTX-M-10 or CTX-M-34	100%	100%	NG_048898 NG_048984	<i>Kluyvera</i> species	Present	Private	Ground	MG560064
B2	CTX-M-40 or CTX-M-63	96%	95%	NG_048991 NG_049014	Not Isolated	Present	Private	Unknown	MG560056
B3a	CTX-M-12	100%	100%	DQ821704	Not Isolated	Present	Private	Ground	MG560057
B3b	OXA-48b	100%	100%	KU820807	Not Isolated	Present	Private	Ground	MG560068
B4	OXA-48b	100%	100%	KU820807	<i>E. coli</i>	Present	Private	Ground	MG560078
B5	OXA-181	85% ^d	93%	KJ620504	<i>E. coli</i>	Present	Private	Ground	MG560079
B6	OXA-48b	100%	100%	KU820804	<i>Shewanella</i> species	Present	Private	Ground	MG560080
B7	CTX-M-3	100%	100%	KU200455	<i>Klebsiella oxytoca</i>	Present	Private	Ground	MG560065
B8	OXA-181	85%	94%	KJ620504	Not Isolated	Present	Private	Ground	MG56006
B9	OXA-48b	100%	100%	KU820804	<i>Shewanella</i> species	Present	Private	Ground	MG560081
B10	OXA-252	100%	100%	CP022089	Not Isolated	Present	Private	Ground	MG560070
B11a	OXA-48b	99%	99%	KC902850	<i>Shewanella putrefaciens</i> ;	Absent	NCPWS	Ground	MG560082
B11b	OXA-48b	99%	99%	KC902850	<i>Pandoraea sputorum</i>	Absent	NCPWS	Ground	MG560083
B12	TEM-1 variant ^e	99%	99%	KU664682.1	<i>E. coli</i>	Present	Private	Ground	MG560088
B13	OXA-48b or OXA-252	100%	100%	KU820804 NG_050608.1	<i>Shewanella</i> species	Present	Private	Ground	MG560084
C1	CTX-M-3	99%	99%	Y10278	Not Isolated	Absent	CPWS	Surface	MG560058
C2	CTX-M-2-like	89%	96%	KX377894	<i>Kluyvera ascorbata</i>	Absent	CPWS	Ground	MG560066
D1 ^c	OXA-48b	100%	100%	KU820807	<i>Providencia rettgeri</i>	Absent	NCPWS	Ground	MG560085
D2 ^c	OXA-252	100%	100%	NG_050608	Not Isolated	Present	NCPWS	Ground	MG560071
D3 ^c	OXA-199	99%	99%	NG_049495	Not Isolated	Present	NCPWS	Ground	MG560072
E1a	CTX-M-36	99%	100%	NG_048986	<i>Klebsiella oxytoca</i>	Absent	CPWS	Blend ^f	MG560067
E1b	OXA-48b	100%	100%	KU820807	Not Isolated	Absent	CPWS	Blend ^f	MG560073
E2 ^c	OXA-48b	100%	100%	KU820807	Not Isolated	Absent	CPWS	Blend ^f	MG560074
E3	OXA-252	100%	100%	NG_050608	<i>Pseudomonas</i> species	Absent	CPWS	Blend ^f	MG560075
E4	OXA-48b	100%	100%	KU820802	Not Isolated	Absent	CPWS	Blend ^f	MG560076
E5 ^c	OXA-48b	100%	100%	KU820807	<i>Pseudomonas koreensis</i>	Absent	CPWS	Blend ^f	MG560086
E6 ^c	OXA-252	100%	100%	KU820800	<i>Pseudomonas putida</i>	Absent	CPWS	Blend ^f	MG560087
F1	SHV-38	100%	100%	NG_050077	<i>Klebsiella pneumoniae</i>	Present	NCPWS	Ground	MG560089

Table 1. Characteristics of sources of samples positive for *bla*_{CTX-M}, *bla*_{OXA-48}-type, and putative *bla*_{SHV} or *bla*_{TEM} genes; ^aGene variant or cluster with greatest homology in GenBank; ^bCPWS: Community public water system, NCPWS: Non-community public water system; ^cSamples collected from the same water systems: A1 and A2; D1, D2, and D3; E2, E5, and E6; ^d100% nucleotide homology to uncultured bacterium clones; ^eGene sequence is similarly homologous with both ESBL and non-ESBL *bla*_{TEM} genes, but exhibits an ESBL phenotype ^fBlended water is a composite of ground water and treated surface water.

from accidental contamination during original sample collection; however, regulatory samples are typically collected by public water system personnel trained in proper sample collection technique.

The true prevalence of the target ESBL and carbapenemase genes in these samples may be underestimated due to testing limitations. A very small proportion of the preserved water sample culture was tested, potentially missing ESBL and carbapenemase genes present in low copy numbers in the samples. Additionally, DNA was not extracted from the preserved samples prior to PCR testing, and PCR inhibitors may have been present, affecting gene detection. We were also only able to determine presence or absence of the target resistance genes and could not assess the abundance in the original water sample due to the culture step in the coliform screening process.

Sample ID	Organism (gene type)	ESBL pheno-type ^a	TAZ	FOT	FAZ	FEP	FOX	CEP	POD	AXO	IMI	MEM	GEN	AMP	CIP	P/T4
A1	<i>C. freundii</i> -complex (CTX)	No	≤0.25	≤0.25	16*	≤1	8*	>16*	1	≤1	1	≤1	≤4	≤8*	≤1	32/4
A2	<i>C. freundii</i> -complex (CTX)	No	≤0.25	0.5	>16*	≤1	8*	>16*	1	≤1	≤0.5	≤1	≤4	16*	≤1	≤4/4
A3a	<i>K. georgiana</i> ^b (CTX)	No	≤0.25	≤0.25	>16	≤1	≤4	>16	2	≤1	≤0.5	≤1	≤4	>16	≤1	≤4/4
A3b	<i>A. baumannii</i> complex (OXA-48)	Yes	8	8	>16	2	64	>16	8	8	≤0.5	≤1	≤4	16*	≤1	64/4
A4	<i>K. georgiana</i> ^b (CTX)	No	≤0.25	≤0.25	>16	≤1	≤4	>16	1	≤1	4	≤1	≤4	>16	≤1	64/4
A5	<i>K. georgiana</i> ^b (CTX)	No	0.5	0.5	>16	≤1	≤4	>16	2	2	≤0.5	≤1	≤4	>16	≤1	≤4/4
B1	<i>Kluyvera</i> species ^b (CTX)	Yes	32	2	>16	8	8	>16	16	4	≤0.5	≤1	≤4	>16	≤1	≤4/4
B4	<i>E. coli</i> (OXA-48)	Yes	8	8	16	2	64	>16	0.5	≤1	≤0.5	≤1	8	≤8	≤1	>64/4
B5	<i>E. coli</i> (OXA-48)	Yes	8	4	≤8	>16	≤4	>16	1	≤1	1	≤1	≤4	≤8	≤1	>64/4
B6	<i>Shewanella</i> species ^b (OXA-48)	No	≤0.25	2	>16	≤1	≤4	>16	≤0.25	≤1	≤0.5	>8	≤4	16	≤1	≤4/4
B7	<i>K. oxytoca</i> (CTX)	No	1	≤0.25	≤8	≤1	≤4	≤8	≤0.25	≤1	≤0.5	≤1	≤4	>16*	≤1	≤4/4
B11	<i>S. putrefaciens</i> ^b (OXA-48)	Yes	0.5	2	>16	≤1	8	≤8	≤0.25	64	≤0.5	4	≤4	16	≤1	≤4/4
B12	<i>E. coli</i> (TEM)	Yes	8	8	≤8	>16	8	>16	16	4	≤0.5	≤1	≤4	>16	≤1	>64/4
B13	<i>Shewanella</i> species ^b (OXA-48)	No	≤0.25	≤0.25	>16	≤1	≤4	>16	≤1	≤1	≤0.5	2	16	≤8	≤1	≤4/4
C2	<i>K. ascorbata</i> ^b (CTX)	No	≤0.25	≤0.25	>16	≤1	≤4	>16	1	≤1	≤0.5	≤1	≤4	≤8	≤1	≤4/4
D1	<i>P. rettgeri</i> (OXA-48)	No	0.5	≤0.25	>16*	2	64	>16*	≤0.25	≤1	≤0.5	≤1	≤4	≤8*	≤1	≤4/4
E1a	<i>K. oxytoca</i> (CTX)	No	0.5	1	≤8	≤1	≤4	>16	4	16	≤0.5	≤1	≤4	≤8*	≤1	≤4/4
E3	<i>Pseudomonas</i> species ^b (OXA)	No	1	2	>16	≤1	>64	>16	2	8	≤0.5	≤1	≤4	>16	≤1	≤4/4
F1	<i>K. pneumoniae</i> (SHV)	Yes	64	4	≤8	≤1	≤4	>16	4	≤1	≤0.5	≤1	16	>16*	≤1	>64/4

Table 2. Minimum inhibitory concentrations (MIC) (µg/mL) of isolates to selected antibiotics by broth microdilution. TAZ: ceftazidime, FOT: cefotaxime, FAZ: ceftazolin, FEP: cefepime, FOX: ceftoxitin, CEP: cephalothin, POD: cefpodoxime, AXO: ceftriaxone, IMI: imipenem, MEM: meropenem, GEN: gentamicin, AMP: ampicillin, CIP: ciprofloxacin, P/T4: piperacillin/tazobactam. Cells marked with a * indicate antibiotics to which the organism is considered intrinsically resistant according to the 2019 Clinical and Laboratory Standards Institute (CLSI) M100 ED29:2019 document³⁴; ^aAn ESBL phenotype is defined as a bacterial phenotype that exhibits a ≥3 two-fold concentration decrease in a MIC for ceftazidime or cefotaxime tested in combination with clavulanate vs the MIC of ceftazidime or cefotaxime when tested alone. Although the result is listed for all organisms in Table 2, this test is intended for *Klebsiella pneumoniae*, *Klebsiella oxytoca*, *E. coli*, and *Proteus mirabilis* per CLSI m100³⁴; ^bIndicates organisms for which an intrinsic resistance profile is not available in the CLSI m100 document.

ESBL and carbapenemase genes other than the target genes were also clearly present in these water samples. Some of these genes may have been carried by the bacterial isolates in addition to the target ESBL or carbapenemase genes that were detected, potentially contributing to any observed phenotype. The majority of these non-target genes are known to be chromosomally-located and carried by nonpathogenic bacterial species; however, some, such as *bla*_{OXY}, can be plasmid-mediated, and may still be a cause for clinical concern³¹. The *bla*_{CTX-M} and *bla*_{OXA-48}-type genes also arise from relatively nonpathogenic bacteria that are commonly found in water sources (*Kluyvera* and *Shewanella* species, respectively), but have been widely disseminated to other bacterial species via mobile genetic elements. In this study, *bla*_{OXA-48}-type genes in non-*Shewanella* species were lost in subsequent subcultures, a phenomenon also observed with non-fermenting Gram-negative rods carrying the *bla*_{NDM-1} carbapenemase gene isolated from New Delhi drinking water⁹.

Bacterial isolates were identified using the full spectral Bruker MALDI-TOF library used by diverse laboratories. Despite the broad organism coverage, it is possible that some environmental species may have limited representation in this system. In our study, all of the *bla*_{CTX-M} carriers were identified as *Enterobacteriaceae*, with approximately half being classified as *Kluyvera* species. *Enterobacteriaceae* carrying *bla*_{OXA-48}-type genes were also isolated from some water samples. The *bla*_{OXA-48}-like genes were also found in non-*Enterobacteriaceae* species, such as *Shewanella*, *Acinetobacter*, and *Pseudomonas*. The coliform species harboring resistance genes would trigger a positive coliform water test, which would theoretically be followed by attempts to remediate the contamination issue; however non-*Enterobacteriaceae* species harboring ESBL and carbapenemase genes would evade detection by the most commonly used coliform screening methods resulting in “silent” dissemination of these genes via tap water that meets all current regulatory requirements.

Studies reporting ESBL- and carbapenemase-producing bacteria in drinking water in high-income countries are extremely rare. CTX-M-producing *E. coli* was discovered in drinking water in France in a single water sample¹⁰, and carbapenemase-producing *Serratia fonticola* has previously been reported in drinking water from Portugal³², as have non-fermenting intrinsic carbapenemase-producers¹³. To our knowledge, this is the first extensive study of drinking water from multiple regions in a high income country that has revealed the geographically widespread distribution of ESBL- and carbapenemase-producing isolates of serious clinical concern. Our findings suggest that community exposure to these organisms may be more common than currently realized, and consequently, their prevalence in the general population may be underestimated.

In this era of increasing antimicrobial resistance, it is critical to determine the public health significance of antibiotic resistance genes in community drinking water regardless of country income-level classification. Public water systems provide an effective means by which pathogens and antibiotic-resistant organisms can be transmitted to large segments of the population. In high-income countries, approaches such as increased consumption of bottled water are not adequate solutions, as bottled water is often derived from the same sources as tap water, and may also contain trace levels of total coliform bacteria³³. More research is needed to better characterize the problem, understand the associated risk, and devise solutions to combat antimicrobial-resistant bacteria in tap water².

Data Availability

ESBL or carbapenemase gene sequence data that support the findings of this study have been deposited into GenBank with the accession numbers listed in Table 1. The data that support the descriptive statistical analyses of sample characteristics and resistance gene presence are available from the corresponding author upon reasonable request.

References

- Centers for Disease Control and Prevention. Antibiotic Resistance Threats in the United States, 2013, <https://www.cdc.gov/drugresistance/pdf/ar-threats-2013-508.pdf> (2013).
- World Health Organization. Antimicrobial Resistance: Global Report on Surveillance, <https://www.who.int/drugresistance/documents/surveillance-report/en/> (2014).
- World Health Organization. WHO publishes list of bacteria for which new antibiotics are urgently needed, <http://www.who.int/mediacentre/news/releases/2017/bacteria-antibiotics-needed/en/> (2017).
- Bush, K. Alarming β -lactamase-mediated resistance in multidrug-resistant Enterobacteriaceae. *Curr Opin Microbiol.* **13**, 558–564 (2010).
- Bouchillon, S. K., Badal, R. E., Hoban, D. J. & Hawser, S. P. Antimicrobial susceptibility of inpatient urinary tract isolates of gram-negative bacilli in the United States: results from the study for monitoring antimicrobial resistance trends (SMART) program: 2009–2011. *Clin Ther.* **35**, 872–877 (2013).
- Doi, Y. *et al.* Community-associated extended-spectrum beta-lactamase-producing *Escherichia coli* infection in the United States. *Clin Infect Dis.* **56**, 641–648 (2013).
- Kelly, A. M., Mathema, B. & Larson, E. L. Carbapenem-resistant Enterobacteriaceae in the community: a scoping review. *Int J Antimicrob Agents.* **50**, 127–134 (2017).
- Potron, A., Poirel, L. & Nordmann, P. Emerging broad-spectrum resistance in *Pseudomonas aeruginosa* and *Acinetobacter baumannii*: Mechanisms and epidemiology. *Int J Antimicrob Agents.* **45**, 568–585 (2015).
- Walsh, T. R., Weeks, J., Livermore, D. M. & Toleman, M. A. Dissemination of NDM-1 positive bacteria in the New Delhi environment and its implications for human health: An environmental point prevalence study. *Lancet Infect Dis.* **11**, 355–362 (2011).
- Maded, J. Y. *et al.* ST48 *Escherichia coli* carrying the blaCTX-M-1 Inc11/ST3 plasmid in drinking water, France. *Antimicrob Agents Chemother.* **60**, 6430–6432 (2016).
- Randall, L. P. *et al.* Evaluation of meat, fruit and vegetables from retail stores in five United Kingdom regions as sources of extended-spectrum beta-lactamase (ESBL)-producing and carbapenem-resistant *Escherichia coli*. *Int J Food Microbiol.* **241**, 283–290 (2016).
- Talukdar, P. K. *et al.* Antimicrobial resistance, virulence factors and genetic diversity of *Escherichia coli* isolates from household water supply in Dhaka, Bangladesh. *PLoS One.* **8**, e61090, <https://doi.org/10.1371/journal.pone.0061090> (2013).
- Henriques, I. S. *et al.* Prevalence and diversity of carbapenem-resistant bacteria in untreated drinking water in Portugal. *Microbial Drug Resist.* **18**, 531–537 (2012).
- Revisions to the Total Coliform Rule, Final Rule. 40 CFR Parts 141 and 142. United States (2013).
- American Public Health Association, American Water Works Association & Water Environment Federation. 9223 Enzyme-substrate coliform test, 21st edition. *Standard Methods for the Examination of Water and Wastewater* (American Public Health Association, 2005).
- Monstein, H. J. *et al.* Multiplex PCR amplification assay for the detection of blaSHV, blaTEM and blaCTX-M genes in Enterobacteriaceae. *APMIS.* **115**, 1400–1408 (2007).
- Poirel, L., Walsh, T. R., Cuvillier, V. & Nordmann, P. Multiplex PCR for detection of acquired carbapenemase genes. *Diagn Microbiol Infect Dis.* **70**, 119–123 (2011).
- Bush, K., Palzkill, T. & Jacoby, G. β -Lactamase Classification and Amino Acid Sequences for TEM, SHV and OXA Extended-Spectrum and Inhibitor Resistant Enzymes, www.lahey.org/studies (2018).
- Woodford, N., Fagan, E. J. & Ellington, M. J. Multiplex PCR for rapid detection of genes encoding CTX-M extended-spectrum (beta)-lactamases. *J Antimicrobial Chemother.* **57**, 154–155 (2006).
- Nuesch-Inderbinen, M. T., Hachler, H. & Kayser, F. H. Detection of genes coding for extended-spectrum SHV beta-lactamases in clinical isolates by a molecular genetic method, and comparison with the E test. *Eur J Clin Microbiol Infect Dis.* **15**, 398–402 (1996).
- Mena, A. *et al.* Characterization of a large outbreak by CTX-M-1-producing *Klebsiella pneumoniae* and mechanisms leading to *in vivo* carbapenem resistance development. *J Clin Microbiol.* **44**, 2831–2837 (2006).
- Brinas, L., Zarazaga, M., Saenz, Y., Ruiz-Larrea, F. & Torres, C. Beta-lactamases in ampicillin-resistant *Escherichia coli* isolates from foods, humans, and healthy animals. *Antimicrob Agents Chemother.* **46**, 3156–3163 (2002).
- Poirel, L., Heritier, C., Tolun, V. & Nordmann, P. Emergence of oxacillinase-mediated resistance to imipenem in *Klebsiella pneumoniae*. *Antimicrob Agents Chemother.* **48**, 15–22 (2004).
- Payne, S. J. *et al.* Molecular techniques and data integration: Investigating distribution system coliform events. *J Water Supply Res Technol.* **59**, 298–311 (2010).
- Tanner, W. D. *et al.* Development and field evaluation of a method for detecting carbapenem-resistant bacteria in drinking water. *Syst Appl Microbiol.* **38**, 351–357 (2015).
- Khot, P. D. *et al.* Optimization of Matrix-Assisted Laser Desorption Ionization–Time of Flight Mass Spectrometry Analysis for Bacterial Identification. *J Clin Microbiol.* **50**, 3845–3852 (2012).
- U.S. Environmental Protection Agency. Causes of Total Coliform-Positive Occurrences in Distribution Systems, http://www.epa.gov/ogwdw/disinfection/tcr/pdfs/issuuepaper_tcr_causes.pdf (2006).
- Lartigue, M. F. *et al.* Characterization of an extended-spectrum class A beta-lactamase from a novel enterobacterial species taxonomically related to *Rahnella* spp./*Ewingella* spp. *J Antimicrobial Chemother.* **68**, 1733–1736 (2013).
- Natural Resources Defense Council. Threats on tap: widespread violations highlight need for investment in water infrastructure and protections, <https://www.nrdc.org/sites/default/files/threats-on-tap-water-infrastructure-protections-report.pdf> (2017).
- U.S. Environmental Protection Agency. About Private Water Wells, <https://www.epa.gov/privatewells/about-private-water-wells> (2017).
- Gonzalez-Lopez, J. J. *et al.* First detection of plasmid-encoded blaOXY beta-lactamase. *Antimicrob Agents Chemother.* **53**, 3143–3146 (2009).

32. Henriques, I. *et al.* Draft Genome Sequence of *Serratia fonticola* UTAD54, a Carbapenem-Resistant Strain Isolated from Drinking Water. *Genome Announc.* 1, e00970–13, <https://doi.org/10.1128/genomeA.00970-13> (2013).
33. U.S. Food and Drug Administration. *CFR - Code of Federal Regulations Title 21*, <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm?fr=165.110> (2017).
34. Clinical and Laboratory Standards Institute (2019). Performance Standards for Antimicrobial Susceptibility Testing, 29th Edition. CLSI M100:ED29:2019, <http://em100.edaptivedocs.net/dashboard.aspx> (2019).

Acknowledgements

A portion of this work was performed in the laboratory of Catherine Loc-Carrillo, PhD at the Veterans Affairs Medical Center campus in Salt Lake City, Utah and through resources provided by the Salt Lake City VA IDEAS center (VA Center of Innovation Award #I50HX001240 from the Health Services Research and Development of the Office of Research and Development of the US Department of Veterans Affairs). Funding for this project was provided through the Health Studies Fund of the University of Utah Department of Family and Preventive Medicine and through internal University Seed Funding. We are grateful for the outreach efforts of Sarah Wright and the Association of Public Health Laboratories (APHL) in finding participating APHL member laboratories. We also appreciate the assistance of the Arkansas Department of Health Public Health Laboratory, Illinois Department of Health laboratories in Chicago, Springfield, and Carbondale, Pennsylvania Department of Environmental Protection Laboratories, Biggs Laboratory at the New York State Department of Health Wadsworth Center, Wisconsin State Laboratory of Hygiene, Utah Public Health Laboratory, Davis, Utah County Health Department Laboratory, and the Weber Basin Water Conservancy District laboratory (Utah) in providing and preserving samples for this project. We are also grateful to the Derek Warner and the University of Utah Core DNA Sequencing Laboratory for consulting on and assistance in preparations for the PCR amplicon sequencing.

Author Contributions

W.T. performed all PCR testing, bacterial isolation, and susceptibility testing. J.V., R.G. and A.G. participated in study design and manuscript preparation. M.L. performed all descriptive statistical analysis. M.F. performed MALDI-TOF testing. Authors from the Arkansas, Illinois, New York, Pennsylvania, Utah, Wisconsin, and Davis County public health laboratories and the Weber Basin Water Conservancy District assisted in preserving coliform-positive samples and compiling associated sample data. All authors were involved in compiling and reviewing data for the report and approved the final version. Authors from the Arkansas, Illinois, New York, Pennsylvania, Utah, Wisconsin, and Davis County public health laboratories and the Weber Basin Water Conservancy District all contributed equally.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-019-40420-0>.

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019