

Preplanned Studies

Analyzing the Trends and Causes of Birth Defects — Jinan City, Shandong Province, China, 2005–2022

Wei Wei^{1,&}; Wei Jiang^{2,&}; Rui Yang^{3,&}; Wenchao Cui⁴; Lihua Zhang⁵; Zhongliang Li^{6,#}

Summary

What is already known about this topic?

Numerous studies conducted in China have reported on the prevalence of birth defects (BDs). However, the limited surveillance periods in select studies curtail comprehensive trends and cause evaluation. Accordingly, the surveillance duration of BDs is extended, and a comprehensive analysis of their prevailing trends is conducted to provide a basis for government intervention and policy implementation.

What is added by this report?

There is a distinct increase in the incidence of BDs observed in Jinan. In rural areas, a pronounced upward trend was observed, and the increase was more rapid than in urban areas. BD prevalence among mothers over 35 years old and under 20 years old was substantially higher than BD prevalence rates in other maternal age brackets. Specifically, the period from 2005 to 2022 saw the prevalence of congenital heart disease surge, the fastest average annual growth rate among all birth defects.

What are the implications for public health practice?

It's essential to prioritize pregnant women in rural areas and those at both ends of the maternal age spectrum. Implementing comprehensive initiatives is crucial to address the high prevalence of congenital heart disease.

Birth defects (BDs) are a collection of structural, functional, and/or biochemical-molecular anomalies identifiable at birth (1). With advancements in economic and social sectors, mortality rates from infections and malnutrition among children and adolescents have significantly decreased. However, there's a rising trend in deaths due to BDs (2). Current estimates suggest that BD prevalence is 4.7% in developed countries, 5.6% in middle-income countries, and 6.4% in low-income countries (3). Moreover, disabilities and cancer arising from birth defects pose growing threats to children's health (4). Given these developments, BDs have become a critical

global health concern demanding urgent attention.

In this context, China has also seen a rise in the significance of BDs in public health. According to the 2012 "Birth Defect Prevention Report" published by the Ministry of Health, China's prevalence rate was 5.6 per 10,000 births in 2011 (5). Notably, 19.1% of infant deaths were attributable to BDs, and BDs were the root cause of 9.6% of disabilities. Furthermore, BDs were established as the second leading cause of infant death in China (6). Given its status as a developing nation with a large population, it is critical for China to reduce the incidence rates of birth defects.

Many Chinese studies have reported on the prevalence of BDs. Current protocols mandate reporting prenatal abnormalities in body structure, function, or metabolism. The extensive use of ultrasound technology in obstetrics has led to an increase in the detection of BDs prior to 28 weeks of gestation. Consequently, there is a compelling need to extend the surveillance period for birth defects.

In Jinan, most expectant mothers typically initiate hospital visits around the 12th week of gestation to secure a maternal record and undergo a medical examination. Therefore, starting birth defect monitoring from this gestational stage is both methodologically robust and practically feasible. As a result, a birth defect monitoring network was established across all district and county levels in Jinan in 2005. Accordingly, a birth defect monitoring network encompassing all district and county levels throughout Jinan was established in 2005. Health centers at the township level and county-level hospitals providing midwifery services were required to report relevant birth defect data. In 2011, the reach of this network was expanded to include all medical institutions offering pregnancy health and midwifery services. Additionally, an information system designed to collect precise birth defect data was implemented in the same year. All information relating to mothers and newborns, inclusive of data on birth defects, was submitted via an online reporting system, enabling accurate and reliable data.

Between 2005 and 2008, BDs were identified from the 28th week of gestation until 7 days post-birth. However, from 2008 to 2022, the period for determining BDs broadened to begin at 12 weeks gestation and extended through to 7 days after birth. Our study holds the potential to inform governmental policy-making with scientifically-based recommendations and serve as a template for developing cities to prevent BDs.

Data was aggregated from registry forms and an electronic information system. Between 2005 and 2015, paper forms were utilized for data collection. However, during 2015–2022, data compilation transitioned to an electronic information system. The collected data encompassed a range of factors such as the total number of births, maternal age, residency location (either urban or rural), and neonate gender. We accounted for 1,463,842 births and 28,537 BDs within our study.

Both external malformations and chromosomal aberrations, encompassed under BDs, were diagnosed under the International Classification of Diseases (10th edition).

To maintain data accuracy, standardized procedures were established for data handling, abstraction, and evaluation. Hospital surveillance teams verified diagnoses, reviewed data, and checked medical records following these standards. Annual surveys revealed a 0.5% underreporting rate of birth defects in Jinan. Missing reports were documented yearly.

The Cochran-Armitage test (CAT) was used to analyze the trends of BD prevalence. Differences in BDs were tested using the Chi-square test. All statistical analyses were executed using SPSS software (version 18.0; IBM Corp., Armonk, NY, USA), with a significance level of $P < 0.05$.

Between 2005 and 2022, there were 1,463,842 recorded births and 28,537 instances of BDs, yielding a prevalence of 194.93 per 10,000 births. Time trend analysis illustrated a rise in BD rates from 99.15 per 10,000 births in 2005 to 290.27 per 10,000 births in 2022. This represents an aggregate increase of 191.21% and an annual increase of 10.70% ($P < 0.05$). The peak birth defect prevalence was in 2022, at 290.27 per 10,000 births. A significant increase inflection point was identified in 2016, with prevalences exceeding 230 per 10,000 births from then on.

During the study period, urban areas had a BD rate of 234.54 per 10,000 births, while rural areas had

161.40 per 10,000 births. The prevalence in urban areas significantly surpassed that in rural areas, averaging 31.17% ($P < 0.05$). A time trend analysis indicated an increase in the incidence of BDs in both urban and rural areas, with the average annual growth rate in rural environments being faster than in urban ones (13.63% vs. 7.85%) (Table 1 and Figure 1).

Throughout our study, we observed a significantly higher prevalence of BDs in two specified age groups. Mothers under 20 had a BD rate of 272.91 per 10,000 births, and those over 35 had 236.09 per 10,000 births. These age groups showed significantly higher BD incidences compared to other groups ($P < 0.05$).

From 2005 to 2022, the proportion of pregnant women under 20 surged by 2037.50%, while those over 35 increased by 66.35% (Table 2).

From 2005 to 2022, the top 5 BDs were congenital heart disease (88.54 per 10,000 births), polydactyly (19.23 per 10,000 births), cleft lip and/or palate (10.41 per 10,000 births), hypospadias (9.84 per 10,000 births), and syndactyly (5.35 per 10,000 births).

Of the 10 major BDs analyzed, 7 showed an upward trend, and 3 declined from 2005 to 2022 ($P < 0.05$). Congenital heart disease exhibited a significant increase at 867.36%, reflecting an average annual growth of 48.19%. The occurrence of Down syndrome similarly increased by 299.14%, with an average annual increase of 16.62%. Polydactyly also rose by 223.09%, averaging an annual increase of 12.39%. On the other hand, Neural Tube Defects demonstrated a stark decrease at 90.35%, with an average annual decline of 5.02%. The rates of Congenital Hydrocephalus similarly fell by 63.81%, with an average annual decrease of 3.54%. Limb Shortening also declined by 53.19%, averaging an annual decrease of 2.96% (Table 3 and Figure 2).

DISCUSSION

The results indicated a prevalence rate of BDs in Jinan of 194.93 per 10,000 live births between 2005 and 2022. Furthermore, the prevalence of birth defects was observed to increase at an annual rate of 10.71%, surpassing the growth rates recorded in Jiangsu (7), Hunan (8), and Sichuan (9). This growing issue demands significant attention from relevant governmental sectors.

A significant shift in growth trends was identified in

TABLE 1. The prevalence of birth defects in Jinan from 2005 to 2020.

Year	Total*			Urban*			Rural*		
	Births	BDs	PRE	Births	BDs	PRE	Births	BDs	PRE
2005	47,302	469	99.15	19,467	255	130.99	27,835	214	76.88
2006	49,964	620	124.09	18,957	331	174.61	31,007	289	93.20
2007	60,206	627	104.14	25,154	375	149.08	35,052	252	71.89
2008	58,797	673	114.46	23,975	400	166.84	34,822	273	78.40
2009	66,767	690	103.34	26,603	376	141.34	40,164	314	78.18
2010	80,960	951	117.47	36,732	545	148.37	44,228	406	91.80
2011	81,422	1,431	175.75	36,700	819	223.16	44,722	612	136.85
2012	89,054	1,347	151.26	39,763	728	183.08	49,291	619	125.58
2013	77,458	1,156	149.24	35,551	600	168.77	41,907	556	132.67
2014	109,211	1,530	140.10	51,363	813	158.29	57,848	717	123.95
2015	76,440	1,420	185.77	35,959	762	211.91	40,481	658	162.55
2016	131,469	3,068	233.36	62,281	1,750	280.98	69,188	1,318	190.50
2017	102,681	2,781	270.84	49,217	1,589	322.86	53,464	1,192	222.95
2018	100,077	2,652	265.00	48,853	1,453	297.42	51,224	1,199	234.07
2019	102,254	2,718	265.81	49,126	1,500	305.34	53,128	1,218	229.26
2020	83,811	2,270	270.85	40,445	1,240	306.59	43,366	1,030	237.51
2021	73,594	2,039	276.31	36,136	1,100	304.41	37,658	939	249.35
2022	72,175	2,095	290.27	35,126	1,111	316.29	37,049	984	265.59
Total	1,463,842	28,537	194.93	671,408	15,747	234.54	792,434	12,790	161.40
AAGR			10.71			7.85			13.63

Abbreviation: BDs=birth defects; PRE=prevalence (per 10,000 births); CAT=Cochran-Armitage trend; AAGR=average annual growth rate.
* CAT $P < 0.05$.




FIGURE 1. The prevalence of birth defects between urban areas and rural areas from 2005 to 2020.

Note: Two distinct periods were demarcated based on changes in population policies: 2005–2016, preceding the implementation of the universal two-child policy; and 2017–2022 following the implementation of the universal two-child policy.

Abbreviation: BD=birth defect.

2016, potentially due to an increase in pregnancies among women of advanced maternal age. Our study

indicates a spike in pregnancies among women over 35 in 2016, reaching 25.63% from an earlier 10%. This

TABLE 2. The prevalence of BDs in different maternal age groups from 2005 to 2020 (per 10,000 births).

Year	<20 years		20–25 years		25–30 years		30–35 years		35+ years	
	Proportion, %	PRE [†]	Proportion, %	PRE [†]	Proportion, %	PRE [†]	Proportion, %	PRE [†]	Proportion, %	PRE [†]
2005	0.08	277.78	22.58	108.62	52.78	84.11	19.65	112.96	4.91	172.12
2006	0.11	357.14	26.82	108.96	43.70	117.24	22.43	141.90	6.94	164.41
2007	0.12	269.61	22.69	97.20	43.08	105.01	25.35	92.10	8.76	131.51
2008	0.11	292.23	24.51	98.91	40.72	122.37	24.88	104.84	9.78	126.65
2009	0.24	249.24	27.94	97.06	36.28	108.62	24.92	103.41	10.62	110.11
2010	0.47	235.60	23.22	144.19	40.26	109.21	26.66	101.45	9.39	139.49
2011	0.41	337.42	29.66	162.71	36.59	186.96	24.05	164.98	9.31	215.24
2012	0.35	317.46	24.30	143.71	39.90	152.81	26.29	150.47	9.16	193.61
2013	0.57	273.35	22.08	133.31	41.29	164.14	26.54	145.46	9.52	151.82
2014	0.49	242.54	15.41	135.46	47.81	142.31	27.14	133.60	9.15	158.16
2015	0.69	263.65	13.31	179.91	43.91	189.51	27.67	172.08	14.42	205.03
2016	0.54	252.45	9.32	163.08	32.74	203.40	31.77	136.07	25.63	213.92
2017	0.62	290.70	10.71	233.98	31.48	254.97	29.22	309.57	27.97	314.31
2018	0.64	291.02	10.07	235.62	30.13	226.98	33.84	255.27	25.32	347.89
2019	1.32	293.17	10.92	237.75	31.12	230.58	32.07	249.54	24.57	328.41
2020	1.61	296.90	11.62	239.91	30.59	249.27	29.84	215.05	26.34	295.99
2021	1.63	301.63	12.02	241.63	29.36	251.36	31.63	225.36	25.36	312.36
2022	1.71	305.32	11.68	246.36	25.65	253.69	32.69	226.39	28.27	326.69
Total	0.65	272.91*	18.27	202.76	37.63	182.21	27.59	187.97	15.86	236.09*
AAGR	113.19	0.55	-2.68	7.05	-2.86	11.20	3.69	5.58	26.43	4.99

Abbreviation: BDs=birth defects; PRE=prevalence (per 10,000 births); CAT=Cochran-Armitage trend; AAGR=average annual growth rate.

* Chi-squared test $P<0.05$.

† CAT $P<0.05$.

TABLE 3. The prevalence of major birth defects from 2005 to 2020 (per 10,000 births).

Rank	Types of BDs	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	Total	AAGR
1	Congenital heart disease*	12.47	16.61	17.51	16.22	16.63	23.34	43.97	40.87	35.89	34.06	48.01	62.10	71.20	80.32	90.63	100.23	110.36	120.63	88.54	48.19
2	Polydactyly*	8.88	13.81	9.89	11.70	11.69	15.19	15.6	16.73	15.75	15.11	18.58	21.00	25.71	26.72	27.12	27.60	28.01	28.69	19.23	12.39
3	Cleft lip and/or palate	15.45	20.22	17.52	22.23	17.23	17.42	19.49	16.05	15.49	16.3	15.56	18.74	20.25	19.81	20.33	19.63	20.12	19.68	10.41	1.52
4	Hypospadias*	4.44	8.83	7.69	5.07	6.10	6.78	12.73	4.45	6.12	4.28	7.92	10.10	12.10	11.34	12.10	10.10	12.82	11.73	9.84	9.12
5	Syndactyly	3.38	3.20	3.41	2.68	2.55	3.33	3.32	3.59	3.49	3.57	5.63	6.90	7.89	6.96	6.70	4.80	5.10	5.20	5.35	2.99
6	Down's syndrome* Other	2.33	2.10	1.78	1.50	1.20	1.61	2.21	1.68	5.03	3.85	6.93	9.28	13.93	11.66	12.9	11.40	10.80	9.30	5.05	16.62
7	malformation of external ear*	1.90	2.40	1.95	3.34	1.35	0.74	2.70	2.47	3.36	3.20	3.53	2.89	6.82	6.56	5.40	3.90	3.70	3.90	4.36	5.85
8	Congenital hydrocephalus*	9.09	8.21	5.84	6.35	4.94	5.93	6.21	5.95	4.91	5.22	4.06	5.01	5.94	5.09	4.10	3.24	4.26	3.29	4.05	-3.54
9	Neural tube defects*	10.99	11.61	9.49	9.29	8.14	4.32	4.54	3.71	4.00	4.94	4.06	4.18	4.72	4.70	3.20	1.67	1.32	1.06	2.12	-5.02
10	Limb shortening*	4.23	5.60	2.92	4.68	2.55	2.47	4.18	3.26	3.61	2.93	4.32	3.19	4.80	4.20	3.90	2.15	2.05	1.98	3.61	-2.96

Abbreviation: BD=birth defect; AAGR=average annual growth rate; CAT=Cochran-Armitage trend.

* CAT $P<0.05$.

change likely stems from China's "universal two-child policy", allowing all women to have two children. Moreover, the traditional Chinese belief, deeming numerous sons as a source of abundant blessings,

deeply entrenched in the society, also pose an effect. Consequently, following the policy's introduction, many women, particularly those of advanced maternal age, chose to have children. It is widely established




FIGURE 2. The prevalence of (A) congenital heart disease, (B) polydactyly, (C) Down's syndrome, (D) neural tube defects, (E) congenital hydrocephalus, and (F) limb shortening from 2005 to 2020. Abbreviation: BD=birth defect.

through numerous studies that an increased maternal age poses risk factors for BDs (10), highlighting 2016 as a significant turning point in the rising birth defect trends.

Our findings revealed a higher prevalence of BDs in urban areas compared to rural areas, potentially due to increased work-related stress and environmental impacts in urban areas. Furthermore, birth defects grew faster in rural areas, a trend rarely seen in other studies. This observation could potentially be due to the inadequate awareness and application of preventive measures amongst pregnant women in rural areas, compounded by suboptimal lifestyle habits (11).

Our data shows a distinct “U” shaped pattern in the maternal age distribution of the prevalence of BDs, particularly among women under 20 and above 35. This trend mirrors earlier studies (12) and may be attributable to unhealthy lifestyles and reproductive challenges. We also highlight a rapidly growing proportion of pregnancies within these age groups, particularly among women under 20 — a demographic subgroup that has previously garnered little research attention. In China, pregnancies in women under 20 are commonly unintended and frequently result in induced abortion, which poses substantial health risks.

Our time trend analysis examined the prevalence of key BDs, revealing a notable increase in the prevalence of congenital heart disease (CHD), aligning with the findings of numerous studies (13). It is well

understood that CHD originates from complex, multifactorial causes. The noted escalation in prevalence can be attributed to advancements in diagnostic and screening techniques, lifestyle behaviors of parents, genetic and environmental contributors to BDs, or a combination thereof. The widespread implementation of ultrasound technology in obstetrics should not be dismissed. A significant growth inflection point was observed for CHDs in our study in 2011. This corresponds with the introduction of free neonatal screenings for CHD in Jinan that year. The percentage of neonatal screenings soared from an average of 35.67% before 2011 to 85.36% thereafter, leading to the detection of previously undiagnosed cases, such as mild VSD or ASD. Similar observations have been reported in other scholarly articles (14).

Our research also indicates a notable decrease in the prevalence of neural tube defects, which can be attributed to the widespread promotion of folic acid programs in Jinan. The protective effect of folic acid is well-supported by multiple studies (15). Following the introduction of free folic acid for expectant mothers into public health initiatives in 2009, nearly 95% of them began a regular intake of folic acid during their pregnancies. This is likely associated with the marked decrease in the prevalence of neural tube defects in Jinan post-2010.

Our study has certain limitations. First, the birth defect data was hospital-based. Although the

monitoring phase extended up to 12 gestation weeks, any birth defects occurring prior to this period or outside of hospital settings may have been overlooked. Second, environmental hazards and unhealthy lifestyles play a significant role in the incidence of birth defects. However, due to data limitations, we couldn't fully assess the impact of these factors on Jinan's birth defect rates, indicating the need for additional research. Third, the annual variation in the monitoring time duration for birth defects could potentially adversely affect the trend analysis for birth defects. Lastly, given the extended monitoring period, changes in diagnostic criteria or improvements in diagnostic methods were inevitable, meaning that the prevalence of birth defects could be underestimated.

In summary, there has been a notable escalation in the prevalence of BDs in Jinan, with the growth rate in rural areas surpassing that in urban regions. Data indicates that mothers over 35 and under 20 exhibit a pronounced BD prevalence relative to other maternal age groups, and representation within these age groups is also dramatically rising. Furthermore, there is a marked increase in the prevalence of congenital heart disease, while the prevalence of neural tube defects shows a decrease. Targeted interventions for expectant mothers in rural territories, as well as those in the aforementioned age groups, are imperative.

Acknowledgements: Appreciation is expressed to all the participants for their contribution.

doi: 10.46234/ccdcw2023.184

* Corresponding author: Zhongliang Li, lizhongliang903@163.com.

¹ Department of General management, Jinan Maternity and Child Care Hospital Affiliated to Shandong First Medical University, Jinan City, Shandong Province, China; ² Department of Medical Service, Jinan Maternity and Child Care Hospital Affiliated to Shandong First Medical University, Jinan City, Shandong Province, China; ³ Department of Outpatient, Jinan Maternity and Child Care Hospital Affiliated to Shandong First Medical University, Jinan City, Shandong Province, China; ⁴ Department of Neonatology, Jinan Maternity and Child Care Hospital Affiliated to Shandong First Medical University, Jinan City, Shandong Province, China; ⁵ Department of Medicine, Jinan Maternity and Child Care Hospital Affiliated to Shandong First Medical University, Jinan City, Shandong Province, China; ⁶ Department of Women Healthcare, Jinan Maternity and Child Care Hospital Affiliated to Shandong First Medical University, Jinan City, Shandong Province, China.

† Joint first authors.

Submitted: August 24, 2023; Accepted: September 11, 2023

REFERENCES

- Perin J, Mai CT, De Costa A, Strong K, Diaz T, Blencowe H, et al. Systematic estimates of the global, regional and national under-5 mortality burden attributable to birth defects in 2000–2019: a summary of findings from the 2020 WHO estimates. *BMJ Open* 2023; 13(1):e067033. <http://dx.doi.org/10.1136/bmjopen-2022-067033>.
- Perin J, Mulick A, Yeung D, Villavicencio F, Lopez G, Strong KL, et al. Global, regional, and national causes of under-5 mortality in 2000-19: an updated systematic analysis with implications for the Sustainable Development Goals. *Lancet Child Adolesc Health* 2022;6(2):106 – 15. [http://dx.doi.org/10.1016/S2352-4642\(21\)00311-4](http://dx.doi.org/10.1016/S2352-4642(21)00311-4).
- World Health Organization. The global burden of disease: 2004 update. Geneva: World Health Organization; 2008. <https://www.who.int/publications/i/item/9789241563710>.
- Spector LG, Kochilas L. Major birth defects and cancer. *BMJ* 2020;371:m4464. <http://dx.doi.org/10.1136/bmj.m4464>.
- Qin H, Zhu J. Birth defect prevention report: report on birth defects prevention and treatment in China (2012). 2013. <https://www.gov.cn/govweb/gzdt/att/att/site1/20120912/1c6f6506c7f811bacf9301.pdf>.
- Chen GX, Zeng GZ, Chen XZ, Wu LS. Meta analysis for surveillance of mortality among children under 5 years old in China from 2001 to 2010. *Chin J Healthy Birth Child Care* 2013;19(8):619-21. <https://d.wanfangdata.com.cn/periodical/zgysyy201308004>. (In Chinese).
- Zhou Y, Mao XQ, Zhou H, Wang L, Qin ZQ, Cai ZM, et al. Birth defects data from population-based birth defects surveillance system in a district of southern Jiangsu, China, 2014–2018. *Front Public Health* 2020;8:378. <http://dx.doi.org/10.3389/fpubh.2020.00378>.
- Xie DH, Yang TB, Liu ZY, Wang H. Epidemiology of birth defects based on a birth defect surveillance system from 2005 to 2014 in Hunan Province, China. *PLoS One* 2016;11(1):e0147280. <http://dx.doi.org/10.1371/journal.pone.0147280>.
- Zhou TJ, Liu WX, Zhao ZL, Zhang G, Liu JT, He LK, et al. Changes in the incidence of birth defects in Sichuan Province, southwest China from 2006 to 2019: a retrospective analysis based on hospital birth defect monitoring. *Res Square* 2021;2(23):271811. <http://dx.doi.org/10.21203/rs.3.rs-271811/v1>.
- Zhang XG, Su L, Wu SQT, Hao XJ, Guo SY, Suzuki K, et al. Prevalence of birth defects and risk-factor analysis from a population-based survey in Inner Mongolia, China. *BMC Pediatr* 2012;12(1):125. <http://dx.doi.org/10.1186/1471-2431-12-125>.
- Cui MM, Lu XL, Lu YY, Wang F, Xie XL, Cheng XY, et al. Knowledge and intake of folic acid to prevent neural tube defects among pregnant women in urban China: a cross-sectional study. *BMC Pregnancy Childbirth* 2021;21(1):432. <http://dx.doi.org/10.1186/s12884-021-03893-4>.
- Luo YL, Li CY, Gao XH, Tan SQ, Li JM, Wang W, et al. Maternal age, parity and isolated birth defects: a population-based case-control study in Shenzhen, China. *PLoS One* 2013;8(11):e81369. <http://dx.doi.org/10.1371/journal.pone.0081369>.
- van der Linde D, Konings EEM, Slager MA, Witsenburg M, Helbing WA, Takkenberg JJM, et al. Birth prevalence of congenital heart disease worldwide: a systematic review and meta-analysis. *J Am Coll Cardiol* 2011;58(21):2241 – 7. <http://dx.doi.org/10.1016/j.jacc.2011.08.025>.
- GBD 2017 Congenital Heart Disease Collaborators. Global, regional, and national burden of congenital heart disease, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet Child Adolesc Health* 2020;4(3):185 – 200. [http://dx.doi.org/10.1016/S2352-4642\(19\)30402-X](http://dx.doi.org/10.1016/S2352-4642(19)30402-X).
- De Wals P, Tairou F, Van Allen MI, Uh SH, Lowry RB, Sibbald B, et al. Reduction in neural-tube defects after folic acid fortification in Canada. *N Engl J Med* 2007;357(2):135 – 42. <http://dx.doi.org/10.1056/NEJMoa067103>.