


# Intra-Articular Injection of Rebamipide-Loaded Nanoparticles Attenuate Disease Progression and Joint Destruction in Osteoarthritis Rat Model: A Pilot Study

CARTILAGE  
January-March 2022: 1–13  
© The Author(s) 2022  
DOI: 10.1177/19476035211069250  
journals.sagepub.com/home/CAR  


Sung Eun Kim<sup>1</sup>, Sung Jae Choi<sup>2,3</sup>, Kyeongsoon Park<sup>4</sup>, Hak-Jun Kim<sup>1,2</sup>,  
Gwan Gyu Song<sup>2,5</sup>, and Jae Hyun Jung<sup>2,3</sup> 

## Abstract

**Objective.** Rebamipide has antioxidant effects and is a drug with a local rather than systemic mechanism of action. Oxidative stress and inflammation in chondrocytes are the major factors contributing to the development and progression of osteoarthritis (OA). Since OA is mainly developed in weight bearing or overused joints, the locally sustained therapy is effective for targeting inflammatory component of OA. We investigated the effects of intra-articular injection of rebamipide loaded nanoparticles (NPs) in OA rat model. **Design.** We fabricated rebamipide-loaded methoxy poly(ethylene glycol)-b-poly(D,L-lactide) (mPEG-PDLLA) and poly(D, L-lactide-co-glycolide) (PLGA) NPs that allow the sustained release of rebamipide. **In vitro**, chondrocytes from rat were used to investigate the cytotoxicity and anti-inflammatory effect of rebamipide-loaded NPs. **In vivo**, monosodium iodoacetate (MIA)-induced OA rats were divided into 7 groups, consisting of healthy control rats and rats injected with MIA alone or in combination with NPs, rebamipide (1 mg)/NPs, rebamipide (10 mg)/NPs, rebamipide (10 mg) solution, or oral administration. **Results.** *In vitro*, rebamipide/NPs dose-dependently suppressed the mRNA levels of pro-inflammatory mediators, including interleukin (IL)-1 $\beta$ , IL-6, tumor necrosis factor- $\alpha$ , matrix metalloproteinase (MMP)-3, MMP-13, and cyclo-oxygenase-2. *In vivo*, the mRNA levels of pro-inflammatory components most markedly decreased in the intra-articularly injected rebamipide (10 mg)/NP group compared to other groups. Macroscopic, radiographic, and histological evaluations showed that the intra-articular injection of rebamipide/NPs inhibited cartilage degeneration more than rebamipide solution or rebamipide administration. **Conclusions.** Using a chemically induced rat model of OA, intra-articular delivery of rebamipide was associated with decreased local and systemic inflammatory response decreased joint degradation and arthritic progression.

## Keywords

rebamipide, methoxy poly(ethylene glycol)-b-poly(D,L-lactide), poly(D, L-lactide-co-glycolide), intra-articular injection, osteoarthritis

## Introduction

Osteoarthritis (OA) is the most common chronic degenerative disease in the elderly and various factors affect the development and progression of OA. Continuous oxidative stress in chondrocytes is an important factor contributing to the development of OA.<sup>1</sup> Oxidative stress is produced by the imbalance between the production and clearance of reactive oxygen species (ROS), which is high in OA cartilage, causing chronic inflammation.<sup>2</sup> One of the oxidants, nitric oxide (NO), and ROS, plays a

role in chondrocyte insensitivity to the anabolic actions of insulin-like growth factor-1, which leads to the pathological degradation of the cartilage extracellular matrix. ROS cause cellular damage, and excessive production of oxidants is associated with the apoptosis of cartilage chondrocytes.<sup>3</sup> Apoptotic cell death is associated with senescence, and oxidative stress-induced cell death is observed in OA chondrocytes.<sup>4</sup> In addition to aging, various inflammatory cytokines affect the development of OA, and one of the key cytokines, interleukin-1 beta (IL-1 $\beta$ ), also stimulates the production of ROS.<sup>5</sup> Inflammatory



cytokines, such as IL-1 $\beta$ , IL-6, and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) are highly upregulated in OA joints and expression of matrix degrading proteases leads to cartilage extracellular matrix degradation.<sup>2</sup> ROS and inflammation are considered as possible targets for the treatment of OA.

Oxidative stress is caused by an imbalance of ROS and antioxidant defense, and attempts have been made to suppress ROS production in OA.<sup>6</sup> Tenoxicam has an antioxidant effect and improves OA,<sup>7</sup> and celecoxib or hyaluronan can reduce nitrite and NO.<sup>7,8</sup> In an animal study, statin were shown to be effective in improving OA through antioxidant action.<sup>9</sup> Rebamipide protects the gastrointestinal mucosa by inhibiting inflammation through an antioxidant mechanism that scavenges hydroxyl radicals and suppresses superoxide production.<sup>10</sup> In addition, rebamipide inhibits the signal transducer and activator of transcription factor 3, suppresses the nuclear factor (NF) kappa-B transcription factor, and inhibits the expression of the receptor activator of nuclear factor kappa-B ligand, preventing the progression of arthritis.<sup>10-12</sup> NF-kappa B activation is inhibited in the presence of antioxidants.<sup>13</sup> The antioxidant and anti-inflammatory effects of rebamipide have been applied to various diseases, such as liver injury, dry eye, and Behçet's disease.<sup>3,11</sup> Rebamipide also attenuates subchondral trabecular bone resorption and produces proteoglycans.<sup>14,15</sup> In previous animal studies, systemic administration of rebamipide was found to be effective in improving arthritis.<sup>3,16</sup> However, rebamipide is considered to have a local rather than systemic effect with respect to the mechanism of action.<sup>17</sup> Thus, topical rebamipide is used as a topical agent for the treatment of ocular dryness.<sup>18</sup>

Currently, intra-articular injections are mainly administered with steroids and hyaluronic acid. Rebamipide may directly affect directly joints presenting OA *via* intra-articular injection. However, the half-life of rebamipide is approximately 2 h;<sup>19</sup> therefore, a repetitive intra-articular injection of rebamipide may be necessary. For the long-term sustained release of the drug, nanoparticulate drug carriers have been developed as advanced drug delivery systems. Methoxy poly(ethylene glycol)-*b*-poly (D,L-lactide) (mPEG-PDLLA) is an amphiphilic diblock copolymer synthesized by ring-opening polymerization and is

widely used because of its biodegradability and non-cytotoxicity.<sup>20</sup> Poly(D,L-lactic-*co*-glycolic acid) (PLGA) is a polymer that is widely used because of its favorable degradation characteristics and good biocompatibility compared to various other types of nanoparticles (NPs).<sup>21</sup> In this study, we examine the effect of sustained intra-articular delivery of rebamipide on cartilage degeneration *in vivo* using a chemically induced model of rat OA.

## Methods

### Materials

The mPEG-PDLLA (molecular weight [MW]: 5,000–20,000 Da) was purchased from Akina Inc. (West Lafayette, IN, USA). PLGA (50:50; Resomer<sup>®</sup> RG505) was supplied by Boehringer Ingelheim (Ingelheim, Germany). Rebamipide was purchased from the Tokyo Chemical Industry Co., Ltd. (Tokyo, Japan). Dimethyl sulfoxide (DMSO) and lipopolysaccharide (LPS) were obtained from Sigma-Aldrich (St. Louis, MO, USA). A cellulose-ester dialysis membrane (MW cut-off [MWCO]: 6–8 kDa) was obtained from Spectrum Laboratories (CA, USA). Dulbecco's modified Eagle's medium (DMEM), fetal bovine serum, phosphate-buffered saline (PBS), IL-1 $\beta$  and penicillin-streptomycin were obtained from Gibco BRL (Rockville, MD, USA). Chondrocytes were purchased from Lonza Ltd. (Walkersville, MD, USA).

### Fabrication of mPEG-PDLLA and PLGA NPs and Rebamipide-Loaded NPs

To fabricate the rebamipide-loaded NPs, mPEG-PDLLA (40 mg mL<sup>-1</sup>) and PLGA (160 mg mL<sup>-1</sup>) were dissolved in DMSO solution and slowly stirred for 30 min at room temperature, followed by the addition of rebamipide (0.2 or 2 mg for 1 and 10 mg [w/w] of the total polymer) and stirred for 24 h. The resulting solution was dialyzed against distilled water (DW) and lyophilized. The mPEG-PDLLA and PLGA NPs without rebamipide were similarly fabricated as the mPEG-PDLLA and PLGA NPs with rebamipide. The mPEG-PDLLA and PLGA NPs loaded with rebamipide (1 or 10 mg) are hereafter referred to as rebamipide

<sup>1</sup>Department of Orthopedic Surgery and Rare Diseases Institute, Korea University Guro Hospital, Seoul, Republic of Korea

<sup>2</sup>Korea University College of Medicine, Seoul, Republic of Korea

<sup>3</sup>Division of Rheumatology, Department of Internal Medicine, Korea University Ansan Hospital, Ansan-si, Republic of Korea

<sup>4</sup>Department of Systems Biotechnology, Chung-Ang University, Anseong-si, Republic of Korea

<sup>5</sup>Division of Rheumatology, Department of Internal Medicine, Korea University Guro Hospital, Seoul, Republic of Korea

Supplementary material for this article is available on the Cartilage website at <http://cart.sagepub.com/supplemental>.

### Corresponding Author:

Jae Hyun Jung, Division of Rheumatology, Department of Internal Medicine, Korea University Ansan Hospital, 123 Jeokgeum-ro, Danwon-gu, Ansan-si 15355, Gyeonggi-do, Republic of Korea.  
Email: [mcfriend82@naver.com](mailto:mcfriend82@naver.com)

(1 mg)/NPs and rebamipide (10 mg)/NPs. In addition, mPEG-PDLLA and PLGA NPs without rebamipide are hereafter referred to as NPs.

### Characterization of Bare NPs and Modified NPs

We confirmed the shapes of NPs with or without rebamipide using a JEM-F2100 (JEOL Ltd., Japan) field emission-transmission electron microscope (FE-TEM). Before the FE-TEM exhibition, each specimen (100 µg) was placed in a tube with EtOH (1 mL) and scattered by a bath-type sonicator (Powersonic 405; 40 kHz, power: 350 W, Hwashin Tech Co., Ltd. Korea) for 1 h at 4°C. Each NP (10 µg) was carefully moved on to a TEM grid (CF200-Cu, Electron Microscopy Sciences, Hatfield, PA, USA), and the shape of each NP was monitored using FE-TEM at 200 kV. To perform particle range tests using dynamic light scattering (DLS), NPs with or without rebamipide were dispersed using a bath-type sonicator containing DW for 1 h at 4°C. The size distribution and zeta potential values were determined using a DLS instrument (Malvern Zetasizer 3000, Malvern, UK) equipped with a helium-neon laser at a wavelength of 633 nm. The zeta potential, the electrostatic potential at the slipping plane, is used to exam colloid-electrolyte interactions, and the basic principle is that particles with opposite charge surfaces are attracted, while particles of similar charge are repelled.<sup>22,23</sup> Zeta potential is one of the main properties that can affect the stability and cell immersion of NPs by measuring the surface charge in suspension through electrophoretic light scattering. The higher zeta potential, the higher is the repulsion force between the particles and the lower the probability for a collision with subsequent aggregating.<sup>24</sup> To investigate the amount of rebamipide loaded on NPs, 10 mg of rebamipide (1 mg)/NPs and 10 mg of rebamipide (10 mg)/NPs were dissolved in 2 mL of dichloromethane. Then, PBS (2 mL) was added to the rebamipide/NPs solution. The rebamipide concentration was determined from the standard curve of various rebamipide solution using the following standard curve equation:

$$Y = 0.1655 + 0.00317, R^2 : 0.99$$

The drug concentration in each group was measured using a Flash Multimode Reader (Varioskan™, Thermo Scientific, USA) at 315 nm.

### Rebamipide Release from Rebamipide/NPs

To investigate the release of rebamipide from rebamipide (1 mg)/NPs and rebamipide (10 mg)/NPs, each concentration of rebamipide/NPs, distributed in 1 mL of PBS solution (pH 7.4), was added to a dialysis bag (MWCO 6–8 kDa). Each bag containing the substrate was carefully transferred into a 50 mL tube with 5 mL of PBS solution (pH 7.4),

followed by gentle shaking (100 rpm) in a water bath at 37°C. At predetermined time intervals points (1, 5, 9, and 12 h, and 1, 3, 5, 7, 14, 21, and 28 days), the supernatant of each rebamipide/NPs was harvested and replaced with fresh PBS. The amount of rebamipide from the NPs was investigated by measuring the absorbance at 315 nm using a multimode reader.

### Cytotoxicity Test

The cytotoxicity of NPs, rebamipide (1 mg)/NPs, rebamipide (10 mg)/NPs, and rebamipide (10 mg) solution was tested against chondrocyte on days 1 and 3. Cells ( $5 \times 10^4$  cells/well) were seeded into 96-well culture plates and allowed to adhere for 24 h, followed by the addition of DMEM in the presence or absence of 100 µg/mL NPs with or without rebamipide. The cell were treated with rebamipide solution as positive control. On the first and third day of incubation, the cells were washed with PBS and cell counting kit-8 (CCK-8) reagents (Dojindo, Tokyo, Japan) were added, followed by incubation for 1 h. The optical density was monitored at 450 nm using a Flash Multimode Reader (Varioskan™, Thermo Scientific).

### In Vitro Anti-Inflammatory Effects on Inflamed Chondrocytes

To examine the effect of rebamipide/NPs on the mRNA levels of pro-inflammatory cytokines in LPS + IL-1β-promoted chondrocytes, cells ( $1 \times 10^5$  cells/well) were seeded in 24-well plates and treated with LPS (10 ng/mL) and IL-1β (10 ng/mL) solutions and with NPs, rebamipide (1 mg)/NPs, and rebamipide (10 mg)/NPs (at a concentration of 100 µg/mL) or rebamipide (100 µg/mL) and incubated at 37°C in a humidified 5% CO<sub>2</sub> atmosphere. LPS- and IL-1β-stimulated cells were used as positive controls, and non-stimulated cells were used as negative controls. The expression of IL-1β, IL-6, TNF-α, matrix metalloproteinase-3 (MMP-3), MMP-13, and cyclo-oxygenase-2 (COX-2) was measured. At 1- and 3-day time-points, cells were harvested for total RNA isolation. Total RNA was isolated using the RNeasy Plus Mini Kit (Qiagen, Valencia, CA, USA). For the reverse transcription of the total RNA (1 µg) into complementary deoxyribonucleic acid (cDNA), AccuPower RT PreMix (Bioneer, Daejeon, Korea) was used. All PCR amplifications were performed using AccuPower PCR PreMix (Bioneer, Daejeon, Korea). The primer sequences of the pro-inflammatory genes are described in the Supplementary Materials (Table 1). PCR amplification and detection were conducted using an ABI7300 Real-Time Thermal Cycler (Applied Biosystems, Foster City, CA, USA). The relative mRNA levels of inflammatory markers were normalized to those of glyceraldehyde 3-phosphate dehydrogenase (GAPDH).

**Table 1.** All Primers of the Pro- and Anti-Inflammatory Component Target Genes Subjected to Real-Time PCR *in Vitro* and *in Vivo*.

| Gene              | Primer sequence                    |
|-------------------|------------------------------------|
| IL-1 (F)          | 5'-CCA CCT CCA GGG ACA GGA TA-3'   |
| IL-1 (R)          | 5'-AAC ACG CAG GAC AGG TAC AG -3'  |
| IL-6 (F)          | 5'- CCG TTT CTA CCT GGA GTT TG -3' |
| IL-6 (R)          | 5'- GTT TGC CGA GTA GAC CTC AT -3' |
| TNF- $\alpha$ (F) | 5'- CTC CCA GAA AAG CAA GCA AC -3' |
| TNF- $\alpha$ (R) | 5'- CGA GCA GGA ATG AGA AGA GG-3'  |
| MMP-3 (F)         | 5'-ACC TGT CCC TCC AGA ACC TG-3'   |
| MMP-3 (R)         | 5'-AAC TTC ATA TGC GGC ATC CA-3'   |
| MMP-13 (F)        | 5'-AAG GAG CAT GGC GAC TTC TA-3'   |
| MMP-13 (R)        | 5'-GGT CCT TGG AGT GGT CAA GA-3'   |
| COX-2 (F)         | 5'-CAG CCA TAC TAT GCC TCG GA-3'   |
| COX-2 (R)         | 5'-GGA TGT CTT GCT CGT CGT TC-3'   |

PCR = polymerase chain reaction; IL = interleukin; F = forward; C = cytosine; A = adenine; T = thymine; G = guanine; R = reverse; TNF = tumor necrosis factor; MMP = matrix metalloproteinase.

### OA-Induced Animal Model and Experimental Plan

For the *in vivo* study, 28 male Sprague-Dawley 8-week-old rats were supplied by DooYeal Biotech (Seoul, Korea). Among these 28 rats, 4 rats without treatment were used as the negative control. MIA-induced model was used for OA model. MIA is an inhibitor of glyceraldehyde-3-phosphate dehydrogenase, which causes chondrocyte death, cartilage degeneration, osteophyte. In addition MIA-induced model elevated the expression changes of MMP and pro-inflammatory cytokines such as IL-1 $\beta$  and TNF- $\alpha$ .<sup>25</sup> The intra-articular cavity of the remaining 24 rats was administered 50  $\mu$ L of MIA (10 mg/mL in PBS [pH 7.4]) using a 1-cm<sup>3</sup> syringe. After the injection of MIA, four rats were maintained as positive controls and given no NPs with or without rebamipide. NPs, rebamipide (1 mg)/NPs and rebamipide (10 mg)/NPs were mixed with 1 mL of 2% carboxymethyl cellulose (CMC) solution. One week after MIA injection, the remaining rats, excluded from the positive control, were treated with 200  $\mu$ L of CMC solution containing NPs, rebamipide (1 mg)/NPs or rebamipide (10 mg)/NPs. As a drug control, 200  $\mu$ L rebamipide solution (10 mg/mL) was intra-articular injected into rats. As another drug control, rebamipide (10 mg/kg/day) was administered orally for 8 weeks. The number of rats per each treated group was four. Figure 1 showed the overview of this study. The actual treatment dosages of rebamipide were 1.55  $\mu$ g/each rat for rebamipide (1 mg)/NPs and 13.82  $\mu$ g/each rat for rebamipide (10 mg)/NPs. The actual dose of oral rebamipide was 2.11 mg/kg/day for each rat, respectively. The total dose of rebamipide orally administered for 4 or 8 weeks was 59.08 or 118.16 mg/kg for each rat, respectively. At the end of the


animal experiments, all rats were euthanized to demonstrate the therapeutic effect of rebamipide/NPs. The knee joints of the euthanized rats were analyzed using micro-computed tomography (micro-CT) and safranin-O/fast green staining. The whole blood of the rats was isolated 4 and 8 weeks following the administration of the NPs with or without rebamipide, rebamipide solution, and oral rebamipide to confirm the anti-inflammatory effect of rebamipide/NPs. The *in vivo* study was approved by the Institutional Animal Care and Use Committee of the Korea University Medical Center (KOREA-2018-0112).

### Histological Examination and Micro-CT

For *ex vivo* characterization, the right knee of the rats was fixed for 24 h in 3.7% formaldehyde and subsequently decalcified in 10% formic acid for 4 weeks. The decalcified tissue was embedded in paraffin and cross-sectioned to a thickness of 8  $\mu$ m in a longitudinal parallel direction using a microtome machine (HM 355S Automatic Microtomes, Thermo Scientific Inc., USA). The sectioned tissues were then stained with hematoxylin and eosin (H&E) and safranin-O/fast green staining. The stained tissues were observed under a light microscope (CX31RTSF; Olympus, Tokyo, Japan). The histology of specimen was assessed by a pathologist. To confirm the changes in the subchondral bone, each sample was subjected to micro-CT (Albira II Imaging System, Carestream Health). The micro-CT system was operated at 40 kV, a current of 250  $\mu$ A, and a nominal resolution of 9  $\mu$ m/pixel. The result of micro-CT was assessed by two authors independently, and validated by another author.

### In Vivo Anti-Inflammatory Effects

Total RNA was extracted from the whole blood of rats using the QIAamp RNA Blood Mini Kit (Qiagen, Valencia, CA, USA) according to the manufacturer's instructions at 4 and 8 weeks after the administration of NPs with or without rebamipide, rebamipide solution, and oral rebamipide. Then, 1  $\mu$ g of total RNA was used for cDNA synthesis *via* reverse transcription with AccuPower RT PreMix (Bioneer, Daejeon, Korea) in accordance with the manufacturer's instructions. The primer sequences described above for the *in vitro* anti-inflammatory study were used as templates for PCR. PCR amplification and detection were performed using an ABI7300 Real-Time Thermal Cycler (Applied Biosystems) with the triplication of each cDNA. The levels of these pro-inflammatory factors were normalized to those of GAPDH. All experiment process including RNA extraction and PCR analysis were performed simultaneously on a single batch with the same library and time after collecting the samples from the experiment.



**Figure 1.** The experimental overview of the study. NPs = nanoparticles.


### Statistical Analysis

All tests were repeated 3 times. Data are expressed as mean  $\pm$  standard deviation, and statistical significance was determined using one-way ANOVA in SigmaPlot (Systat Software, Inc., IL, USA) *via* the Holm-Sidak *t*-test for multiple comparisons. *P*-values were compared among all rebamipide members. Differences were considered statistically significant at *P*-values  $< 0.05$  and  $< 0.01$ .

## Results


### Characteristics of NPs with or without Rebamipide

The morphology of the NPs, rebamipide (1 mg)/NPs, and rebamipide (10 mg)/NPs were confirmed by FE-TEM. Each NP group exhibited by TEM had a spherical form and was nanosized. The average NP sizes and polydispersity indices were  $241.70 \pm 38.29$  nm and 0.312 for NPs,  $231.00 \pm 40.80$  nm and 0.294 for rebamipide (1 mg)/NPs, and  $238.00 \pm 34.07$  nm and 0.307 for rebamipide (10 mg)/NPs (Fig. 2), respectively. The demonstrated zeta potential values were  $-14.20 \pm 2.17$  mV for NPs,  $-19.73 \pm 1.11$  mV for rebamipide (1 mg)/NPs and  $-33.97 \pm 2.61$  mV for rebamipide (10 mg)/NPs (Fig. 3). The loading amount (percentage) of rebamipide in the rebamipide (1 mg)/NPs



**Figure 2.** Particle size and distribution of (A) NPs, (B) rebamipide (1 mg)/NPs, and (C) rebamipide (10 mg)/NPs. Transmission electron microscope scale bar = 0.5  $\mu$ m. NPs = Nanoparticles.

and rebamipide (10 mg)/NPs was  $7.73 \pm 0.54$   $\mu$ g ( $77.33 \pm 5.40\%$ ) and  $69.08 \pm 3.34$   $\mu$ g ( $69.08 \pm 3.34\%$ ), respectively.




**Figure 3.** Zeta potential of NPs, rebamipide (1 mg)/NPs, and rebamipide (10 mg)/NPs. NPs = nanoparticles.

Dialysis membrane method was used to confirm the release kinetics of rebamipide from NPs. A dialysis membrane method set-up is divided into two compartments, 1 consisting of a dialysis device surrounded by a NP formulation or free drug solution, and the other consisting of a sink transmitter.<sup>26,27</sup> The release kinetics of NPs was measured by combining results of drug diffuse in a sink transmitter and the drug releasing from NPs the medium inside the dialysis device. The release kinetics of rebamipide from rebamipide (1 mg)/NPs and rebamipide (10 mg)/NPs for up to 28 days are shown in Figure 4A. On day 1, the amounts of the released rebamipide (percentage) were  $4.49 \pm 0.03 \mu\text{g}$  ( $58.16 \pm 0.39\%$ ) for rebamipide (1 mg)/NPs and  $41.18 \pm 0.36 \mu\text{g}$  ( $60.27 \pm 0.52\%$ ) for rebamipide (10 mg)/NPs. After 28 days, rebamipide (1 mg)/NPs and rebamipide (10 mg)/NPs released  $7.22 \pm 0.06 \mu\text{g}$  ( $93.45 \pm 0.82\%$ ) and  $62.17 \pm 0.75 \mu\text{g}$  ( $90.65 \pm 1.09\%$ ) of rebamipide, respectively. Figure 4B shows the cytotoxicity results for each group at 1 and 3 days of incubation with the chondrocytes. Cell viability after treatment with each NP substance was over 95% for 3 days compared with that of the control.

### Gene Levels of Cytokines in IL-1 & LPS-Promoted Chondrocytes after Treatment with Rebamipide/NPs

To determine whether rebamipide/NPs remarkably suppressed the IL-1- and LPS-promoted inflammatory response in chondrocytes, we conducted real-time PCR to estimate the mRNA levels of pro-inflammatory elements (i.e., IL-1 $\beta$ , IL-6, TNF- $\alpha$ , MMP-3, MMP-13, and COX-2) in chondrocytes treated with NPs, rebamipide (1 mg)/NPs, rebamipide (10 mg)/NPs or rebamipide for 1 or 3 days (Fig. 5 and Suppl. Table S1). Compared with cells without IL-1 and




**Figure 4.** (A) *In vitro* cumulative release profiles of rebamipide release from rebamipide (1 mg)/NPs and rebamipide (10 mg)/NPs. (B) Cytotoxicity assays following treatments of chondrocytes with nanoparticles (NPs), rebamipide (1 mg)/NPs, rebamipide (10 mg)/NPs, and rebamipide (10 mg) on days 1 and 3. NPs = nanoparticles.

LPS induction, IL-1- and LPS-induced cells exhibited an upregulation of pro-inflammatory elements for 3 days. The mRNA levels of IL-1 $\beta$ , IL-6, TNF- $\alpha$ , MMP-3, MMP-13, and COX-2 in cells treated with NPs were similar to those in cells treated with IL-1 and LPS, whereas the expression levels of pro-inflammatory elements in IL-1- and LPS-induced chondrocytes cultured with rebamipide/NPs decreased in a time- and concentration-dependent manner. Moreover, IL-1 $\beta$ , IL-6, TNF- $\alpha$ , MMP-3, MMP-13, and COX-2 gene levels in IL-1- and LPS-promoted cells treated with rebamipide solution were prominently suppressed compared with those in all other groups.

### Changes in Pro-Inflammatory Factors *in Vivo*

Four and 8 weeks after the administration of the NPs with or without rebamipide treatment for MIA-induced OA in rats, we performed real-time PCR to analyze the changes in



**Figure 5.** The mRNA expression levels of pro-inflammatory component, including (A) IL-1 $\beta$ , (B) IL-6, (C) tumor necrosis factor-alpha, (D) MMP-3, (E) MMP-13, and (F) cyclo-oxygenase-2 in IL-1- and LPS-stimulated chondrocytes treated with NPs, rebamipide (1 mg)/NPs, rebamipide (10 mg)/NPs, and rebamipide (10 mg) after 1 and 3 days of incubation. *P*-value was indicated by using one-way ANOVA via the Holm-Sidak multiple comparison test. IL = interleukin; MMP = matrix metalloproteinase; LPS = lipopolysaccharide; NPs = nanoparticles; GAPDH = glyceraldehyde 3-phosphate dehydrogenase; TNF = tumor necrosis factor; COX = cyclo-oxygenase. \*\**P* < 0.01.

the mRNA levels of the pro-inflammatory elements (IL-1 $\beta$ , IL-6, TNF- $\alpha$ , MMP-3, MMP-13, and COX-2) using whole blood samples. As shown in Figure 6 and Supplementary Table S2, the mRNA levels of pro-inflammatory components in the NPs-only group improved in time-dependent manner, similar to those in the MIA-induced group. At 4 weeks, the levels of the 6 genes in the rats treated with the rebamipide/NPs at both rebamipide concentrations decreased greatly compared to those in the rats treated with NPs. Moreover, differences in the mRNA levels of IL-1 $\beta$ , IL-6, TNF- $\alpha$ , MMP-3, MMP-13, and COX-2 were substantially reduced in the whole blood of rats between the oral rebamipide-treated group and the NP-treated group, with similar reduction in difference observed between the rebamipide solution-treated group and the NP-treated group. At 8 weeks, the IL-1 $\beta$ , IL-6, TNF- $\alpha$ , MMP-3, MMP-13, and COX-2 gene levels were markedly reduced in the rebamipide/NP-treated group in a time-dependent manner, whereas they increased slightly in both the rebamipide-treated group and rebamipide solution-treated group in a time-dependent manner. In particular, at 8 weeks after MIA injection, the mRNA expression levels of IL-6 were increased by approximately 3.7 times compared to the control. While after the intra-articular injection of rebamipide (10 mg)/NPs, the mRNA expression levels of IL-6 were increased by approximately 2.5 times. This indicated that the mRNA expression level of IL-6 between MIA only and rebamipide (10 mg)/NPs injection groups was reduced by approximately 34%. In the case of the mRNA expression level of TNF- $\alpha$ , it was increased by approximately 3.5 times compared to those of control at 8 weeks after MIA injection. While after the intra-articular injection of rebamipide (10 mg)/NPs, the mRNA expression level of TNF- $\alpha$  was increased by approximately 2.4 times, which was reduced by approximately 31% compared to those of MIA injection only.


### Gross, Radiologic, and Histological Determinations

Eight weeks after treatment with rebamipide oral administration, rebamipide solution, rebamipide (1 mg)/NPs or rebamipide (10 mg)/NP injections, gross and radiological examinations were conducted to confirm cartilage alterations and amelioration of OA progression. As described in Figure 7A, rats treated with only MIA or with MIA and NPs displayed severe joint surface irregularities, space narrowing, articular cartilage erosions, subchondral bone sclerosis, and femoral condyle demolition. The rats in the oral rebamipide administration or rebamipide solution injection groups showed irregular surfaces and erosions of cartilage compared to the rats in the MIA-only or MIA + NP treatment groups. Intra-articular treatment with NPs containing rebamipide (1 or 10 mg) in rats with MIA-promoted OA


remarkably decreased the erratic cartilage surface and subchondral sediment. In particular, the rebamipide (10 mg)/NP-injected group exhibited an even articular surface, which was analogous to that seen in the control group. Gross examinations were also performed to further demonstrate cartilage transformation during the progression of OA (Fig. 7B). MIA-only and MIA with NP injection groups revealed whole spaces of cartilage destruction with irregular surface fibrillation. Treatment with oral rebamipide or rebamipide solution resulted in partial degenerative changes. Treatment of OA with rebamipide/NP resulted in less degradation of cartilage surfaces compared to the rebamipide oral administration or rebamipide solution treatments. The injection of rebamipide (10 mg)/NPs in joints with OA resulted in the lowest degradation of cartilage surfaces compared to the other treatment groups.

The results of H&E staining showed a clear alteration in the cartilage lesion between all rebamipide treatment groups (Fig. 7C). MIA-only and NP-treated groups showed roughness and destruction of the cartilage and meniscus. The rebamipide solution-injected group and the orally administered rebamipide group showed surface roughness, but mild cartilage destruction compared with the MIA-only and NP groups. Rats treated with NPs with different rebamipide concentrations (1 or 10 mg) exhibited little cartilage breakdown and surface roughness. Owing to the anti-inflammatory activity of the rebamipide released from the NPs, the rebamipide (10 mg)/NP-treated group displayed very little surface roughness as well as a minimum degree of cartilage destruction among the treated groups. In H&E staining, the cytoplasm and nucleus are mainly stained. In the control and rebamipide (10 mg)/NPs groups, the cytoplasm of the cartilage were intact and purple nuclei were visible, whereas in the MIA group, the cytoplasm and nuclei of the cartilage were hardly visible and in rebamipide oral administration group, the cytoplasm reduced and nuclei were almost invisible. The results of safranin-O staining allowed the determination of proteoglycan content during the progression of OA. In the MIA-only and NP treatment groups, a loss of proteoglycan to the deep area of the cartilage layers, subchondral bone exposure, and erosion and fibrillation of cartilage were observed (Fig. 7D). The rebamipide solution-injected group or rebamipide orally administered group showed a loss of proteoglycan in the superficial and intermediate layers of the cartilage layer. A slight loss of proteoglycan in the superficial zone of cartilage layers was observed in the rebamipide (1 mg)/NP group. However, rebamipide (10 mg)/NP-treated group showed a minor loss of proteoglycan in the superficial area of the cartilage layer, and most of the proteoglycan in the cartilage was retained. Osteoarthritis Research Society International (OARSI) scoring was conducted to further confirm the *in vivo* restraint effect of rebamipide/NPs on the progression of MIA-induced OA (Fig. 8). Significant differences were






**Figure 6.** The mRNA expression levels of pro-inflammatory components, including (A) IL-1 $\beta$ , (B) IL-6, (C) tumor necrosis factor- $\alpha$ , (D) MMP-3, (E) MMP-13, and (F) cyclo-oxygenase-2 in whole blood samples from the MIA-induced OA rats in each group. Whole blood was obtained at baseline and 4th and 8th weeks after treatment of rebamipide. Each mRNA expression level was evaluated using real-time PCR analysis. The number (n) of animals per group was 4 and the data are expressed as the mean  $\pm$  standard deviation. *P*-value was indicated by using one-way ANOVA via the Holm-Sidak multiple comparison test. IL = interleukin; MMP = matrix metalloproteinase; MIA = monosodium iodoacetate; OA = osteoarthritis; PCR = polymerase chain reaction; NPs = nanoparticles; GAPDH = glyceraldehyde 3-phosphate dehydrogenase; TNF = tumor necrosis factor; COX = cyclo-oxygenase. \**P* < 0.05 and \*\**P* < 0.01.



**Figure 7.** (A) Micro-computed tomography, (B) gross morphology, (C) hematoxylin and eosin staining results, and (D) safranin-O staining results of the rat knee joint at the ninth week of MIA injection (eighth week of rebamipide treatment). The treated groups were divided into the (a) control, (b) MIA, (c) MIA + NPs, (d) MIA + rebamipide (1 mg)/NPs, (e) MIA + rebamipide (10 mg)/NPs, (f) oral administration, and (g) rebamipide solution groups. Arrows indicate erosions. The number (n) of animals per group was 4. MIA = monosodium iodoacetate; NPs = nanoparticles.



**Figure 8.** Osteoarthritis Research Society International scores of the rat knee joints during at the ninth week of monosodium iodoacetate injection (eighth week of rebamipide treatment). The score was measured by averaging the values of 4 rats belonging to each of the 7 groups. MIA = monosodium iodoacetate; NPs = nanoparticles; OARSI = Osteoarthritis Research Society International.

observed between the OARSI scores obtained after rebamipide oral administration, rebamipide solution, and rebamipide/NP injection and those obtained after MIA induction and NP injection (\*\* $P < 0.01$ ). In addition, the OARSI scores in the rebamipide (10 mg)/NP-treated group were considerably lower than those in the MIA induction, NP alone, rebamipide oral administration, rebamipide solution, and rebamipide (1 mg)/NPs (\*\* $P < 0.01$ ) groups.

## Discussion

OA initiates cartilage damage, causes synovial inflammation and sclerosis of subchondral bones, and damages bones resulting in the formation of osteophytes.<sup>28</sup> Disease-modifying OA drugs (DMOADs) have been developed to inhibit the progression of OA by targeting each process. Phase 3 studies on inducible NO synthase inhibitors, gene therapy, and existing possible DMOADs, such as diacerein, calcitonin, and doxycycline, are being conducted, and drugs that inhibit various mechanisms that affect the development of OA are being studied.<sup>29</sup> Oxidative stress plays a critical role in the development and progression of OA, and drugs that inhibit oxidative stress are fundamental DMOADs. Rebamipide is an oxygen radical scavenger that inhibits oxidative stress and osteoclastogenesis, and is considered a drug that can be used as a DMOAD.<sup>10</sup> In this study, rebamipide-loaded mPEG-PDLLA and PLGA NPs released rebamipide in a sustained manner for up to approximately 2 months without cytotoxic effects against synoviocytes. In addition, rebamipide released from NPs in the joint cavity

significantly reduced the mRNA levels of pro-inflammatory cytokines in the blood and inhibited the progression of OA by protecting against cartilage degradation. In particular, a single intra-articular injection of rebamipide (10 mg)/NPs was more effective locally and systemically in inhibiting OA progression than daily rebamipide consumption.

Both mPEG-PDLLA and PLGA are NPs that have been used for cancer treatment, with tested efficacy and stability. When rebamipide was loaded onto the NPs of these 2 substances, the product maintained the nanosize and reduced inflammation in synoviocytes and chondrocytes without cytotoxicity. The MIA + NP-treated group showed no significant elevation in pro-inflammatory cytokines in the early or late stage compared to the MIA-treated group; therefore, the mPEG-PDLLA and PLGA NPs did not worsen arthritis. In addition, approximately 60% of rebamipide was released from the mPEG-PDLLA and PLGA NPs in 2 days, and rebamipide was gradually released continuously, which may suppress the elevation of IL-1 $\beta$  and TNF- $\alpha$  levels in synoviocytes. Although evaluated *in vitro* for up to 7 weeks, the slope of the cumulative release of rebamipide over time gradually decreased for both rebamipide (1 mg)/NPs and rebamipide (10 mg)/NPs, but the total cumulative dose of rebamipide continued to increase.

OA progresses by increasing ROS levels due to oxidative stress caused by several factors, such as aging. ROS increase the secretion of pro-inflammatory cytokines, such as IL-1 and TNF- $\alpha$ , in chondrocytes and increase the expression of enzymes related to the progression of OA, such as MMP-1 and COX-2.<sup>28</sup> IL-1 and TNF- $\alpha$  further activate the cells to promote the secretion of other inflammatory or catabolic enzymes, such as NO synthase-2, COX-2, MMP-1, MMP-3, and MMP-13.<sup>30</sup> IL-1 is a powerful inducer of cartilage degradation and induces the mRNA expression of disease-relevant proteases, such as MMP13 and ADAMTS5.<sup>31</sup> TNF- $\alpha$  upregulates MMP gene expression and induces cartilage degeneration.<sup>32</sup> In *in vitro* studies, rebamipide/NPs were less effective in suppressing the mRNA levels of pro-inflammatory cytokines and enzymes than rebamipide alone, possibly because of the relatively low drug concentrations maintained by the sustained release of the drug following the slow degradation of mPEG-PDLLA and PLGA NPs. These findings are similar to the results of the previous study that diacerein, which is considered as a DMOAD, also inhibited the expression of pro-inflammatory cytokines in LPS + IL-1 $\beta$ -promoted synoviocytes when the drugs were loaded to PLGA NPs.<sup>30</sup>

Since rebamipide has been developed as a gastrointestinal protective drug, it is unclear whether rebamipide has the role of inhibiting joint inflammation and OA progression after intra-articular injection *in vivo* even if inhibition of OA-related inflammatory mediators was detected *in vitro*. To test the efficacy of rebamipide/NPs *in vivo*, MIA-induced

OA rat models were established. MIA induces inflammation, subchondral bone damage, and chondrocyte death by inhibiting GAPDH, resulting in the development of OA.<sup>33</sup> A single intra-articular injection of rebamipide solution showed anti-inflammatory effect at 4 weeks; however, the anti-inflammatory effect decreased at 8 weeks, which may be due to the short half-life of rebamipide and the short retention time of the drug in the joint cavity. Moreover, rebamipide/NPs significantly reduced the levels of pro-inflammatory cytokines and enzymes in whole blood compared with the daily oral administration of rebamipide. Furthermore, macroscopic and histological evaluations showed that in rats intra-articularly injected with rebamipide (10 mg)/NPs, the prevention of cartilage degradation was more pronounced than that observed in the rats given a daily oral administration of rebamipide. However, there are limitations as a pilot study due to its small sample size in this study; thus, a caution is required in interpretation of this study and a larger-scale study will be required for a further validation study. Several drugs that can reduce ROS, such as vitamin C, coenzyme Q10, and hyaluronic acid, are expected to prevent structural changes in OA cartilage; however, there is no consistent evidence that these antioxidants are effective in treating OA.<sup>28</sup> Rebamipide, which has an antioxidant effect, can protect cartilage via systemic (oral administration) and focal dissemination (intra-articular injection) in OA.<sup>3,11</sup> Upon comparing the clinical treatment dosage of rebamipide (10 mg)/NPs (13.82 µg/rat) and the daily oral administration dosage of rebamipide (118.16 mg/rat), the intra-articular injection of rebamipide/NPs emerges as the more effective treatment approach for protecting the cartilage and suppressing the expression of inflammatory cytokines. In addition, rebamipide shows different aqueous solubility depending on the pH.<sup>34</sup> When rebamipide was loaded on mPEG-PDLLA and PLGA NPs, it was continuously released in the environment, including acidic environments in the inflamed synovial fluid, and the released rebamipide worked effectively on the cartilage affected by OA.

In conclusions, the sustained drug delivery system involving rebamipide-loaded mPEG-PDLLA and PLGA NPs suppressed the mRNA levels of pro-inflammatory elements in inflamed synoviocytes stimulated by LPS and IL-1. A single intra-articular injection of rebamipide/NPs also suppressed inflammatory reactions and appeared to mitigate cartilage degradation in MIA-induced OA. Considering the actual treatment dosage and the degree of anti-inflammatory effect, rebamipide/NP injection appeared to be more effective in the treatment of OA than the intra-articular injection or daily oral administration of rebamipide alone. The preliminary results of this pilot study support further adequately powered *in vivo* study of the effect of intra-articular administration of rebamipide NPs for the treatment of OA; ultimately optimization of doses and dose schedules are also warranted.

## Acknowledgments and Funding

We would like to thank Nano-based Disease Control Institute for assistance of the experiment.

The research was supported by a Korea University Ansan Hospital Grant (grant number: O1904051) and the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (grant number: 2018R1D1A1B07050687).

## Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Ethical Approval

The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Institutional Animal Care and Use Committee of the Korea University Medical Center (KOREA-2018-0112).

## ORCID iD

Jae Hyun Jung  <https://orcid.org/0000-0002-3835-3587>

## References

- Alcaraz MJ, Megias J, Garcia-Arnanadis I, Clerigues V, Guillen MI. New molecular targets for the treatment of osteoarthritis. *Biochem Pharmacol.* 2010;80(1):13-21.
- Ansari MY, Ahmad N, Haqqi TM. Oxidative stress and inflammation in osteoarthritis pathogenesis: role of polyphenols. *Biomed Pharmacother.* 2020;129:110452.
- Moon SJ, Woo YJ, Jeong JH, Park MK, Oh HJ, Park JS, *et al.* Rebamipide attenuates pain severity and cartilage degeneration in a rat model of osteoarthritis by downregulating oxidative damage and catabolic activity in chondrocytes. *Osteoarthritis Cartilage.* 2012;20(11):1426-38.
- Carlo MD Jr, Loeser RF. Increased oxidative stress with aging reduces chondrocyte survival: correlation with intracellular glutathione levels. *Arthritis Rheum.* 2003;48(12):3419-30.
- Wojdasiewicz P, Poniatowski LA, Szukiewicz D. The role of inflammatory and anti-inflammatory cytokines in the pathogenesis of osteoarthritis. *Mediators Inflamm.* 2014;2014:561459.
- Pazdzior M, Kielczykowska M, Kurzepa J, Luchowska-Kocot D, Kocot J, Musik I. The oxidative stress in knee osteoarthritis patients. An attempt of evaluation of possible compensatory effects occurring in the disease development. *Medicina (Kaunas).* 2019;55(5):150.
- Ozgoemen S, Ardicoglu O, Erdogan H, Fadillioglu E, Gudul H. In vivo effect of celecoxib and tenoxicam on oxidant/anti-oxidant status of patients with knee osteoarthritis. *Ann Clin Lab Sci.* 2005;35(2):137-43.
- Kalaci A, Yilmaz HR, Aslan B, Sogut S, Yanat AN, Uz E. Effects of hyaluronan on nitric oxide levels and superoxide dismutase activities in synovial fluid in knee osteoarthritis. *Clin Rheumatol.* 2007;26(8):1306-11.
- Pathak NN, Balaganur V, Lingaraju MC, Kant V, Kumar D, Kumar D, *et al.* Effect of atorvastatin, a HMG-CoA reductase inhibitor in monosodium iodoacetate-induced osteoarthritic

- pain: implication for osteoarthritis therapy. *Pharmacol Rep.* 2015;67(3):513-9.
10. Nanke Y, Kobashigawa T, Yago T, Kawamoto M, Yamanaka H, Kotake S. Rebamipide, an amino acid analog of 2(1H)-quinolinone, inhibits the formation of human osteoclasts. *Biomed Res Int.* 2016;2016:6824719.
  11. Suzuki Y, Hasegawa M, Matsui Y, Unno H, Iino T, Yoshida T, et al. Intra-articular injection of rebamipide prevents articular cartilage degeneration in murine post-traumatic osteoarthritis models. *Mod Rheumatol.* 2020;30(4):765-72.
  12. Min HK, Kim JK, Lee SY, Kim EK, Lee SH, Lee J, et al. Rebamipide prevents peripheral arthritis and intestinal inflammation by reciprocally regulating Th17/Treg cell imbalance in mice with curdlan-induced spondyloarthritis. *J Transl Med.* 2016;14(1):190.
  13. Beg AA, Finco TS, Nantermet PV, Baldwin AS Jr. Tumor necrosis factor and interleukin-1 lead to phosphorylation and loss of I kappa B alpha: a mechanism for NF-kappa B activation. *Mol Cell Biol.* 1993;13(6):3301-10.
  14. Izawa T, Hutami IR, Tanaka E. Potential role of rebamipide in osteoclast differentiation and mandibular condylar cartilage homeostasis. *Curr Rheumatol Rev.* 2018;14(1):62-9.
  15. Pujari-Palmer M, Pujari-Palmer S, Engqvist H, Karlsson Ott M. Rebamipide delivered by brushite cement enhances osteoblast and macrophage proliferation. *PLoS One.* 2015;10(5):e0128324.
  16. Moon SJ, Park JS, Woo YJ, Lim MA, Kim SM, Lee SY, et al. Rebamipide suppresses collagen-induced arthritis through reciprocal regulation of th17/treg cell differentiation and heme oxygenase 1 induction. *Arthritis Rheumatol.* 2014;66(4):874-85.
  17. Arakawa T, Higuchi K, Fujiwara Y, Watanabe T, Tominaga K, Sasaki E, et al. 15th anniversary of rebamipide: looking ahead to the new mechanisms and new applications. *Dig Dis Sci.* 2005;50(Suppl 1):S3-11.
  18. Ueda K, Matsumiya W, Otsuka K, Maeda Y, Nagai T, Nakamura M. Effectiveness and relevant factors of 2% rebamipide ophthalmic suspension treatment in dry eye. *BMC Ophthalmol.* 2015;15:58.
  19. Hasegawa S, Sekino H, Matsuoka O, Saito K, Sekino H, Morikawa A, et al. Bioequivalence of rebamipide granules and tablets in healthy adult male volunteers. *Clin Drug Investig.* 2003;23(12):771-9.
  20. Tian Y, Shen Y, Jv M. Synthesis, characterization and evaluation of tinidazole-loaded mPEG-PDLLA (10/90) in situ gel forming system for periodontitis treatment. *Drug Deliv.* 2016;23(8):2726-35.
  21. Swider E, Koshkina O, Tel J, Cruz LJ, de Vries IJM, Srinivas M. Customizing poly(lactic-co-glycolic acid) particles for biomedical applications. *Acta Biomater.* 2018;73:38-51.
  22. Davis ME, Chen ZG, Shin DM. Nanoparticle therapeutics: an emerging treatment modality for cancer. *Nat Rev Drug Discov.* 2008;7(9):771-82.
  23. Vogel R, Pal AK, Jambhrunkar S, Patel P, Thakur SS, Reátegui E, et al. High-resolution single particle zeta potential characterisation of biological nanoparticles using tunable resistive pulse sensing. *Sci Rep.* 2017;7(1):17479.
  24. Doane TL, Chuang CH, Hill RJ, Burda C. Nanoparticle  $\zeta$ -potentials. *Acc Chem Res.* 2012;45(3):317-26.
  25. Xu J, Yan L, Yan B, Zhou L, Tong P, Shan L. Osteoarthritis pain model induced by intra-articular injection of mono-iodoacetate in rats. *J Vis Exp.* 2020;159. doi:10.3791/60649
  26. Zambito Y, Pedreschi E, Di Colo G. Is dialysis a reliable method for studying drug release from nanoparticulate systems? A case study. *Int J Pharm.* 2012;434(1-2):28-34.
  27. Modi S, Anderson BD. Determination of drug release kinetics from nanoparticles: overcoming pitfalls of the dynamic dialysis method. *Mol Pharm.* 2013;10(8):3076-89.
  28. Lepetsos P, Papavassiliou AG. ROS/oxidative stress signaling in osteoarthritis. *Biochim Biophys Acta.* 2016;1862(4):576-91.
  29. Oo WM, Yu SP, Daniel MS, Hunter DJ. Disease-modifying drugs in osteoarthritis: current understanding and future therapeutics. *Expert Opin Emerg Drugs.* 2018;23(4):331-47.
  30. Jung JH, Kim SE, Kim HJ, Park K, Song GG, Choi SJ. A comparative pilot study of oral diacerein and locally treated diacerein-loaded nanoparticles in a model of osteoarthritis. *Int J Pharm.* 2020;581:119249.
  31. Vincent TL. IL-1 in osteoarthritis: time for a critical review of the literature. *F1000Res.* 2019;8. doi:10.12688/f1000research.18831.1
  32. Fernandes JC, Martel-Pelletier J, Pelletier JP. The role of cytokines in osteoarthritis pathophysiology. *Biorheology.* 2002;39(1-2):237-46.
  33. Pitcher T, Sousa-Valente J, Malcangio M. The moniodoacetate model of osteoarthritis pain in the mouse. *J Vis Exp.* 2016;111:53746.
  34. Shin BS, Kim CH, Jun YS, Yoon CH, Rho JI, Lee KC, et al. Oral absorption and pharmacokinetics of rebamipide and rebamipide lysinate in rats. *Drug Dev Ind Pharm.* 2004;30(8):869-76.