

Anti-apoptotic proteins and catalase-dependent apoptosis resistance in nickel chloride-transformed human lung epithelial cells

YU-XIU YANG¹, XIU-LING LI¹, LEI WANG², SHUANG-YIN HAN¹, YAN-RUI ZHANG¹,
POYIL PRATHEESHKUMAR², XIN WANG², JIAN LU², YUAN-QIN YIN², LI-JUAN SUN²,
AMIT BUDHRAJA², ANDREW J. HITRON² and SONG-ZE DING^{1,2}

¹Department of Internal Medicine, Henan Provincial People's Hospital, Zhengzhou University, Zhengzhou, Henan 450000, P.R. China; ²Graduate Center for Toxicology, College of Medicine, University of Kentucky, Lexington, KY 40536, USA

Received April 15, 2013; Accepted May 29, 2013

DOI: 10.3892/ijo.2013.2004

Abstract. Chronic exposure to nickel compounds is associated with increased incidence of certain types of human cancer, including lung and nasal cancers. Despite intensive investigation, the oncogenic processes remain poorly understood. Apoptosis resistance is a key feature for tumor cells to escape physiological surveillance and acquire growth advantage over normal cells. Although NiCl₂ exposure induces transformation of human lung epithelial cells, little information is available with regard to its molecular mechanisms, it is also not clear if the transformed cells are apoptosis resistant and tumorigenic. We explored the apoptosis resistance properties of nickel chloride-transformed human lung epithelial cells and the underlying mechanisms. The results showed that transformed BEAS-2B human lung epithelial cells are resistant to NiCl₂-induced apoptosis. They have increased Bcl-2, Bcl-xL and catalase protein levels over the passage matched non-transformed counterparts. The mechanisms of apoptosis resistance are mitochondria-mediated and caspase-dependent. Forced overexpression of Bcl-2, Bcl-xL and catalase proteins reduced NiCl₂-induced cell death; siRNA-mediated knockdown of their expression sensitized the cells to nickel-induced apoptosis, suggesting that Bcl-2, Bcl-xL and catalase protein expression plays a critical role in apoptosis resistance. Akt also participates in this process, as its overexpression increases Bcl-xL protein expression levels and attenuates NiCl₂-induced apoptosis. Furthermore, transformed cells are tumorigenic in a xenograft model. Together, these results demonstrate that nickel-transformed cells are apoptosis-resistant and

tumorigenic. Increased expression of Bcl-2, Bcl-xL and catalase proteins are important mechanisms contributing to transformed cell oncogenic properties.

Introduction

Nickel is a ubiquitous environmental metal, widely used in industrial processes. Nickel compounds are important human carcinogens. Epidemiologically, chronic and professional exposure to nickel compounds is associated with increased incidence of certain human cancer, including lung and nasal cancers (1,2). Both the soluble and insoluble forms of nickel compounds are detrimental to health with the insoluble form possesses higher risk, such as nickel oxide and nickel subsulfide (3,4). In 1990, nickel is classified as established carcinogen to humans (group 1) by International Agency for Research on Cancer (IARC) (3,4).

Several mechanisms have been proposed for nickel-induced carcinogenesis. These include production of reactive oxygen species (ROS), induction of DNA damage (1,4), alteration of epigenetic changes such as histone modifications (5,6), disruption of cellular iron homeostasis (7), and activation of oncogenic pathways (8,9). Despite intensive investigation, the molecular features of nickel-induced oncogenic transformation, which are essential components in tumor initiation and progression, remain elusive.

Apoptosis (programmed cell death) is a key mechanism to maintain normal cell homeostasis, it removes cells that carry abnormal genetic information and keeps the functional integrity of organ or cell populations (10). Aberrant expression of anti-apoptotic protein is a common feature to many types of malignancy including lung, breast, prostate cancer, melanoma and leukemia. Their overexpression confers the cancer cells a growth advantage to escape from physiological surveillance and promote tumor formation (10,11). Recent studies also suggest that activation of oncogenic pathways mediate nickel chloride (NiCl₂)-induced epithelial cell oncogenic transformation accompanied by enhanced Bcl-2, Bcl-xL protein expression (9).

Correspondence to: Dr Song-Ze Ding, Department of Internal Medicine, Henan Provincial People's Hospital, Zhengzhou University, no. 7 Wei Wu Road, Zhengzhou, Henan 450000, P.R. China
E-mail: dingsongze@hotmail.com

Key words: Bcl-2, Bcl-xL, apoptosis, nickel, BEAS-2B cell, oxidative stress, carcinogenesis, tumor, lung cancer

Bcl-2, and Bcl-xL protein expression has been implicated in cancer and cell transformation (10,12), therefore, we hypothesized that their overexpression in transformed cells may lead to apoptosis resistance contributing to nickel-induced carcinogenesis. However, it is not clear if a metal compounds such as nickel might induce apoptosis resistance in transformed cells and the underlying molecular mechanisms, neither is it known if these cells are tumorigenic. Therefore, we investigated the role of Bcl-2, Bcl-xL and ROS scavenging enzyme in nickel-induced apoptosis resistance and the molecular mechanisms, which are necessary steps to understand nickel-induced carcinogenesis.

We demonstrate in this study that increased Bcl-2, Bcl-xL and ROS scavenging enzyme expression are important in conferring NiCl₂-transformed BEAS-2B cell apoptosis resistance. The mechanisms are mitochondria-mediated and caspase-dependent. Transformed cells are also tumorigenic with accompanied oncogenic protein expression. The results characterize the oncogenic properties of nickel-transformed human lung epithelial cells and provide insight into nickel-induced carcinogenesis.

Materials and methods

Cell culture and reagents. Immortalized normal human bronchial epithelial cell line, BEAS-2B was purchased from American Type Culture Collection (ATCC, Manassas, VA). NiCl₂-transformed BEAS-2B cells were isolated from soft agar by low dose NiCl₂ exposure with BEAS-2B cells for 6 months (9). These cells showed anchorage-independent growth, proliferated faster, but were morphologically similar to normal BEAS-2B cells (9). BEAS-2B cells that stably express catalase (OriGene, Rockville, MD), Akt and dominant negative-Akt (DN-Akt) which is a kinase-dead (K179M) mutant, were generated by integration of catalase, wild-type Akt (Akt-C), DN-Akt expression vectors into BEAS-2B cells and selected with G418. Characterization of these cells has been described before (13,14). Tissue culture reagents were purchased from Gibco (Invitrogen, Carlsbad, CA). Cells were maintained in DMEM medium supplemented with 10% fetal bovine serum and antibiotics at 37°C in a humidified 10% CO₂ incubator.

Nickel chloride (NiCl₂) was purchased from Sigma Chemical Company (St. Louis, MO, USA). Polyclonal and monoclonal antibodies against phospho-Akt at Ser473 (#9275), phospho-Stat3 (#9145), Bcl-xL (#2764), catalase (#8841), HIF-1 α (#3434), cleaved caspase-3 (#9661), cleaved caspase-7 (#9491), cleaved PARP (#9542), peroxiredoxin 1 (#8499) were purchased from Cell Signaling Technology (Danvers, MA). Bcl-2 was from Dako (Carpinteria, CA). SOD1 (sc-11407), SOD2 (sc-130345), β -actin (sc-47778), lamin A/C (sc-6215), Akt1 (sc-5298) and NF- κ B p50 (sc-7178) were purchased from Santa Cruz Biotechnology (Santa Cruz, CA).

Assays for apoptosis. Apoptosis was determined by Annexin V/propidium iodide (PI) staining kit (BD Pharmingen, San Jose, CA). In some experiments, due to the expression of green fluorescence protein (GFP) interfering with Annexin V-FITC reading, a double stain apoptosis detection kit using Hoechst 33342/PI staining (GenScript, Piscataway, NJ) was chosen to replace Annexin V-FITC. Briefly, cells (1x10⁶) treated with or without NiCl₂ were collected and washed once with

cold PBS, they were stained with 5 μ l of Annexin V-FITC or Hoechst 33342 followed by 10 μ l of PI (5 μ g/ml) in binding buffer [10 mmol/l HEPES (pH 7.4), 140 mmol/l NaOH, and 2.5 mmol/l CaCl₂] for 15 min at room temperature in dark. The apoptotic cells were determined using a Becton-Dickinson FACScan cytofluorometer. Both early apoptotic (Annexin V or Hoechst 33342-positive and PI-negative) and late apoptotic (Annexin V or Hoechst 33342-positive and PI-positive) cells were included in cell death determinations.

Western blot analyses. Cell lysates were collected in lysis buffer (50 mM Tris-HCl, pH 7.4, 1 mM EDTA, 150 mM NaCl, 1% NP-40, 0.25% Na-deoxycholate, and 1 μ g/ml of aprotinin, leupeptin and pepstatin). Proteins (20 μ g) were separated on 10% SDS polyacrylamide gels. Proteins were subsequently transferred from gel onto nitrocellulose membrane (Bio-Rad, Hercules, CA). Membranes were blocked for 1 h at room temperature in Tris-buffered saline plus 0.01% Tween-20 (TBS-T) in 5% non-fat dry milk (pH 7.4). Various antibodies were diluted at 1:1,000 in TBS-T with 5% non-fat dry milk solution, incubated with membrane at 4°C overnight and washed three times with TBS-T. The secondary antibody, horseradish peroxidase (HRP)-conjugated antibodies, diluted at 1:2,500 in TBS-T with 5% non-fat dry milk were incubated with membrane at room temperature for 2-3 h. Signal was detected with an enhanced chemiluminescence detection kit (Perkin Elmer Life Sciences, Boston, MA). In each experiment, either anti- β -actin or anti-lamin A/C antibody was reprobed to monitor protein loading. Image quantification was processed by ImageJ software (NIH, Rockville, MD). In some experiments, mean densitometry values were adjusted with β -actin and expressed as fold changes over the appropriate controls.

Quantitative RT-PCR for gene expression. Total RNA from BEAS-2B cells was extracted using TRIzol reagents as described previously (15). Reverse transcription of 2 μ g of total cellular RNA was performed in a final volume of 20 μ l containing 5X first strand buffer (Invitrogen), 1 mM of each dNTP, 20 units of placental RNase inhibitor, 5 μ M random hexamer and 9 units of Moloney murine leukemia virus reverse transcriptase (Invitrogen). After incubation at 37°C for 45 min, the samples were heated for 5 min at 92°C to end the reaction, diluted at 1:4 and stored at -20°C until PCR. cDNA (2 μ l) was subjected to real-time quantitative PCR using the real-time PCR detection systems (Bio-Rad, Hercules, CA) with SYBR-Green I (Molecular Probes, Eugene, OR) as a fluorescent reporter. Threshold cycle number of duplicate reactions was determined using PCR software and levels of selected gene mRNA expression were normalized to hypoxanthine phosphoribosyltransferase (HPRT) levels using the formula $2^{(Rt-Et)}$; where *Rt* is the mean threshold cycle for the reference gene HPRT and *Et* is the mean threshold cycle for experimental gene. Data are expressed as arbitrary units and adjusted to fold changes over the non-stimulated control cells. Primer sequences are provided in Table I.

Measurement of mitochondrial membrane potential. Mitochondrial membrane potential ($\Delta\Psi$ m) was monitored using MitoProbe JC-1 assay kit (Invitrogen). 5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolyl-carbocyanine iodide (JC-1),

Table I. PCR primers.

Gene name	For RT-PCR (5'→3')
<i>Bcl-2</i>	Fwd: GCCTTCTTTGAGTTCGGTGG Rev: ATCTCCCGGTTGACGCTCT
<i>Bcl-xL</i>	Fwd: GAGCTGGTGGTTGACTTTCTC Rev: TCCATCTCCGATTACAGTCCCT
<i>Hprt</i>	Fwd: TTGGAAAGGGTGTATTCTCA Rev: TCCAGCAGGTCAGCAAAGAA
<i>Hprt</i> , hypoxanthine phosphoribosyltransferase.	

a lipophilic cationic fluorescence dye, is capable of selectively entering mitochondria, where it forms monomers, and emits green fluorescence (FL-1) when $\Delta\Psi_m$ is relatively low. At high $\Delta\Psi_m$, JC-1 aggregates and gives red fluorescence (FL-2) (16). Thus the red and green fluorescence of JC-1 reflect the change of $\Delta\Psi_m$ of the mitochondrial membrane. Briefly, BEAS-2B or transformed BEAS-2B cells (5×10^5) were seeded into 60-mm culture dishes and treated with NiCl_2 for 16 h. Cells were trypsinized, washed in ice-cold PBS and incubated with $2 \mu\text{M}$ JC-1 at 37°C for 20 min. Cells were washed once with PBS and analyzed by FACScan cytofluorometer.

Caspase activity assay. Caspase activity was assessed using the luminescent Caspase-Glo[®] 3/7 assay system (Promega, Madison, WI) following manufacturer's instructions. Briefly, BEAS-2B or transformed BEAS-2B cells were treated with or without NiCl_2 at 1.5 mM for 16 h. Caspase-Glo 3/7 reagent (100 μl) was added into 96-well plates and incubated for 1 h at room temperature. The luminescence was measured using a Glomax[™] 96 microplate luminometer (Promega). Z-VAD-FMK (carbobenzoxy-valyl-alanyl-aspartyl-[O-methyl]-fluoromethylketone), a cell-permeant pan-caspase inhibitor that irreversibly binds to the catalytic site of caspase proteases that inhibit induction of apoptosis was added 30 min before nickel stimulation at $20 \mu\text{M}$ solution in DMSO (Promega). Data were collected and expressed as fold change over the control without NiCl_2 treatment.

Small interfering RNA transfection. Transfection procedures followed manufacturer's recommended protocol. Small interference RNA (siRNA) for *Bcl-2* (sc-29214), *Bcl-xL* (sc-43630) were purchased from Santa Cruz Biotechnology (Santa Cruz, CA). Scrambled control and siRNA for catalase (s-2443) were obtained from Ambion (Austin, TX). Briefly, siRNA for *Bcl-2*, *Bcl-xL*, catalase and controls were incubated with Lipofectamine[™] RNAiMAX in Opti-MEM I medium (Invitrogen) for 20 min at room temperature. They were then added to cell culture media without antibiotics. Media were replaced 24 h post-transfection. Cells were treated with or without NiCl_2 for another 24 h. Cell lysates or cells were collected either for western blot or apoptosis analysis as mentioned above.

Retrovirus infection. GFP labeled p-MIG, p-MIG-*Bcl-2*, pMIG-*Bcl-xL* retroviral vector (17) were kind gifts from Dr Stanley Korsmeyer, Dana-Farber Cancer Institute, USA

(Addgene, #9044, #3541, #3544). The plasmids were transfected into 293-GPG cell using Lipofectamine 2000 reagent, viruses were produced in 293-GPG cells with subsequent centrifugation, filtration and stored at -20°C . BEAS-2B or T-BEAS-2B cells were then infected with control, *Bcl-2*, *Bcl-xL* retroviruses by adding virus solution into the cells with polybrene for 48 h before they were used for apoptosis and western blot analysis, virus infection in cells was monitored by checking GFP under fluorescence microscope.

Tumor xenograft model. Athymic nude (nu/nu) female mice, 5 weeks old and weighing about 18 to 20 g were purchased from Jackson Laboratory (Bar Harbor, MI). Mice were handled following the University of Kentucky guidelines for care and use of laboratory animals and approved by the ethical committee. BEAS-2B or transformed BEAS-2B cells (2×10^6) were mixed with Matrigel and injected in both side of flank region in 50 μl volume. Five animals in each group gave 10 injection sites. Mice were fed normally, the body weight, and tumor growth was monitored twice weekly until the tumor size reached $1 \times 1 \times 1 \text{ cm}^3$ in size or 8 weeks post inoculation.

Statistical analysis. All values are expressed as mean \pm standard error (SEM). Student's t-test was used to compare the difference between various controls and NiCl_2 -treated groups. P-value less than 0.05 was considered statistically significant.

Results

Transformed BEAS-2B cells have increased *Bcl-2*, *Bcl-xL* protein expression and are resistant to nickel-induced apoptosis. To evaluate *Bcl-2*, *Bcl-xL* protein expression and their role in nickel-induced apoptosis, BEAS-2B and transformed BEAS-2B (T-BEAS-2B) cells were first treated with or without 0.5, 1 and 1.5 mM of NiCl_2 for 24 h for apoptosis induction. At 0.5, and 1 mM doses, only BEAS-2B cells showed apoptosis induction in 24 h, T-BEAS-2B cell showed little or no apoptosis (data not shown), therefore, 1.5 mM of NiCl_2 was selected for apoptosis induction in both types of cells.

BEAS-2B and T-BEAS-2B cells were treated with or without 1.5 mM of NiCl_2 for 24 h to analysis apoptosis. The results indicated that T-BEAS-2B cells had significantly lower level of apoptosis (Fig. 1A), but increased *Bcl-2*, and *Bcl-xL* protein expression (Fig. 1B and C) when compared with control BEAS-2B cells. Following NiCl_2 exposure, the *Bcl-2*, and *Bcl-xL* protein expression declined in both type of cells, but control BEAS-2B cells had greater level of reduction than T-BEAS-2B cells (Fig. 1B and C). To understand if *Bcl-2*, and *Bcl-xL* protein reduction might be due to NiCl_2 -induced transcriptional repression, quantitative RT-PCR assay was performed in both types of cells. Fig. 1D shows that *Bcl-2* and *Bcl-xL* mRNA expression was repressed in BEAS-2B cells following NiCl_2 -treatment. Similar pattern of mRNA expression was also noted in T-BEAS-2B cells (data not shown). These results suggested a mechanism of NiCl_2 -induced gene transcription repression (Fig. 1D).

We tested if the levels of *Bcl-2* and *Bcl-xL* protein expression in BEAS-2B and T-BEAS-2B cells might affect NiCl_2 -induced apoptosis. Forced overexpression of *Bcl-2*, and *Bcl-xL* protein with plasmids in BEAS-2B cells reduced NiCl_2 -induced cell




Figure 1. Transformed BEAS-2B cells are resistant to nickel-induced apoptosis. BEAS-2B or transformed BEAS-2B (T-BEAS-2B) cells (1x10⁶) were treated with or without 1.5 mM of NiCl₂ for 24 h. (A) Apoptosis was analyzed from the treated cells by Annexin V/PI staining followed by flow cytometry analysis. (B) Cell lysates were collected and resolved in 10% SDS-PAGE to detect respective protein expression, arrows indicate the specific band. (C) Scanned densitometry data were expressed as arbitrary unit and adjusted as fold changes over the control. (D) Bcl-2 and Bcl-xL mRNA expression from the treated BEAS-2B cells were determined by real-time PCR as stated in Materials and methods. Data are mean ± SEM from 3-4 separate experiments. *P<0.05 when compared with control cells.




Figure 2. Effects of Bcl-2, and Bcl-xL expression in NiCl₂-induced apoptosis. BEAS-2B cells were infected with control empty vector, Bcl-2, and Bcl-xL retrovirus (control, Bcl-2, Bcl-xL), transformed BEAS-2B cells (T-BEAS-2B) were transfected with control, Bcl-2, and Bcl-xL siRNA (siRNA). Cells were treated with or without NiCl₂ at 0.75 and 1.5 mM for 24 h. To determine Bcl-2, and Bcl-xL specific protein expression, cell lysates were collected from the retrovirus infected and siRNA transfected cells (A). Protein (20 μg) was resolved in 10% SDS-PAGE to detect the respective protein expression. Blots are representative of 2-3 separate experiments with similar results, arrows indicate the specific band. Apoptosis was assayed using either Annexin V/PI (B and C) or Hoechst 33342/PI staining (D) followed by flow cytometry analysis. Data are mean ± SEM from three separate experiments, *P<0.05 when compared with controls.




Figure 3. Effects of catalase in NiCl₂-induced apoptosis. BEAS-2B (B-2B), transformed BEAS-2B (T-BEAS-2B, T-B-2B) (1x10⁶) were treated with or without NiCl₂ at 1.5 mM for 24 h. Cell lysates were collected and resolved in 10% SDS-PAGE to detect catalase and other ROS scavenger proteins expression (A-C), blots are representative of 2-3 separate experiments with similar results. Arrows indicate the specific band. Densitometry data from catalase were scanned as arbitrary unit and adjusted as fold changes over the control (B). To determine the role of catalase in NiCl₂-induced apoptosis (D), BEAS-2B (B-2B) cells were transfected with or without catalase siRNA (B-2B-Cat-siRNA) for 48 h. Cells were treated with or without NiCl₂ at 1.5 mM for another 24 h. BEAS-2B catalase-stable expressing cells (Cat-stable) also received the same treatment. Apoptosis was assayed by Annexin V/PI staining followed by flow cytometry analysis (D). Data are mean ± SEM from three separate experiments, *P<0.05 when compared with control cells.

apoptosis in a dose-dependent manner (Fig. 2A-C). In contrast, siRNA knockdown of Bcl-2, and Bcl-xL protein expression in T-BEAS-2B cells enhanced the apoptosis rate (Fig. 2A and D). These results suggested that the relative levels of Bcl-2, and Bcl-xL protein are important for apoptosis induction/resistance in both types of cells.

Effects of catalase in NiCl₂-induced apoptosis. After cell transformation, ROS generation is reduced in transformed cells, a mechanism that attributed to increased ROS scavenging enzyme expression (18). We examined if the apoptosis resistance in NiCl₂-transformed cells are accompanied by higher ROS scavenging proteins expression. The results showed upregulation of catalase in nickel-transformed cells over the controls (Fig. 3A and B). Overexpression of catalase by using a stable cell line reduced NiCl₂-induced apoptosis; in contrast, using catalase siRNA to reduce its expression (data not shown) enhanced apoptosis (Fig. 3C). These results suggested that catalase or generation of ROS is important mechanism in NiCl₂-induced apoptosis. Western blot analysis also indicated that other ROS scavenging enzymes such as SOD2, and Prx-1 protein levels were increased in T-BEAS-2B cells when compared with control non-transformed BEAS-2B cells (Fig. 3D), while SOD1, and Phox47 protein expression showed no difference (data not shown). Thus, these results provide mechanistic explanation of

reduced ROS level in T-BEAS-2B cells as reported previously (9), and these mechanisms may contribute to the apoptosis resistance in transformed cells.

Transformed cells are resistant to NiCl₂-induced mitochondria damage. To investigate whether transformed cells are resistant to mitochondria damage. We evaluated NiCl₂-induced mitochondria damage in both types of cells and the role of Bcl-2 and Bcl-xL in these processes. The results showed that T-BEAS-2B cells are more resistant to NiCl₂-induced mitochondrial damage compared with non-transformed BEAS-2B cells (Fig. 4A and B). Forced overexpression of Bcl-2, Bcl-xL and catalase largely abolished NiCl₂-induced mitochondria damage in BEAS-2B cells (Fig. 4C and D). Significant difference was noted when the quantitative data were compared with control cells (*P<0.05). These results indicated that enhanced Bcl-2, Bcl-xL and catalase protein expression are important mechanisms for reduced mitochondria damage in transformed cells.

Caspase activation in nickel-transformed cells. The above results demonstrated that T-BEAS-2B cells differ from BEAS-2B cells in NiCl₂-induced apoptosis resistance and mitochondria damage; thus we further studied the underlying molecular mechanisms. We compared caspase activation in both types of cells. The




Figure 4. Transformed cells are resistant to NiCl₂-induced mitochondria damage. To evaluate NiCl₂-induced mitochondria damage, BEAS-2B, transformed BEAS-2B (T-BEAS-2B), control empty vector, Bcl-2, and Bcl-xL-retrovirus infected BEAS-2B (control, Bcl-2, Bcl-xL) and BEAS-2B catalase-stable expressing cells (Cat-stable) (1x10⁶) were treated with or without NiCl₂ at 1.5 mM for 16 h. Cells were stained with JC-1 dye followed by flow cytometry analysis. (A and C) Representative images are shown; (B and D) quantification of three separate experiments are expressed as mean ± SEM. (B and D) *P<0.05 when compared with controls.




Figure 5. Caspase activation in nickel-transformed cells. BEAS-2B, transformed BEAS-2B (T-BEAS-2B), control empty vector, Bcl-2, and Bcl-xL-retrovirus infected BEAS-2B (Bcl-2, Bcl-xL) and BEAS-2B catalase-stable expressing cells (Cat-stable) (1×10^6) were treated with or without NiCl₂ at 1.5 mM for 24 h. Cell lysate was collected and 20 μ g of protein resolved in 10% SDS-PAGE to detect respective protein expression (A, C, E-G). Blots are representative from 3-4 separate experiments with similar results, arrows indicate the specific band. To determine caspase 3/7 enzymatic activity, BEAS-2B and T-BEAS-2B cells were treated with or without NiCl₂ at 1.5 mM for 16 h and enzymatic activity was assayed as described in Materials and methods (B). For apoptosis assay, BEAS-2B, transformed BEAS-2B (T-BEAS-2B) cells were treated with or without NiCl₂ at 1.5 mM and Z-VAD for 24 h, apoptosis was assayed using the Annexin V/PI staining followed by flow cytometry (D). Quantitative data are expressed as mean \pm SEM from three separate experiments, *P<0.05 when compared with respective controls.

result in Fig. 5 clearly indicates that upon NiCl₂ stimulation, BEAS-2B cells had much stronger caspase activation, presented as increased c-caspase 3, 7 and c-PARP protein expression when compared with T-BEAS-2B cells (Fig. 5A). This was accompa-

nied by increased caspase 3/7 enzymatic activity, while Z-VAD, a pan-caspase inhibitor, blocked the caspase 3/7 enzymatic activity (Fig. 5B), reduced c-caspase 3, 7 expression (Fig. 5C), and attenuated apoptosis (Fig. 5D).




Figure 6. Effects of Akt expression in NiCl₂-induced apoptosis. BEAS-2B and BEAS-2B stably expressing Akt (Akt-C), dominant negative-Akt (DN-Akt) cells were treated with or without NiCl₂ at 1.5 mM for 24 h. Cell lysates were collected and 20 μ g of protein was resolved in 10% SDS-PAGE to detect protein expression. Blots are representative of three separate experiments with similar results. (A) Arrows indicate the specific band. (B) Scanned densitometry data are expressed as arbitrary units and adjusted as fold changes over the control. (C) Apoptosis was determined by Annexin V/PI staining followed by flow cytometry analysis. Data are mean \pm SEM from three separate experiments, *P<0.05 when compared with control cells.

In addition, we evaluated if Bcl-2, Bcl-xL and catalase proteins that are overexpressed in T-BEAS-2B cell may antagonize the NiCl₂-induced caspase activation. The results indicated that overexpression of Bcl-2, Bcl-xL and catalase protein effectively reduced the c-caspase 3, 7 and c-PARP protein expression (Fig. 5E-G), respectively, correlating with reduced mitochondria damage and apoptosis. Therefore, these results demonstrated that upregulation of Bcl-2, Bcl-xL and catalase in T-BEAS-2B suppresses caspase activation, which are essential mechanisms underlying the adapted apoptosis resistance in these cells.

Role of p-Akt in NiCl₂ induced-apoptosis. p-Akt expression was upregulated in T-BEAS-2B cells in a previous study (9). To understand its role in NiCl₂-induced apoptosis, we used Akt overexpression (Akt-C) and dominant negative Akt (DN-Akt) stable cell lines to determine the effects of Akt on NiCl₂-induced apoptosis. The results indicated that Akt stable cells had persistently enhanced p-Akt level (Fig. 6A), and increased Bcl-xL protein expression, but not Bcl-2 expression; while in DN-Akt cells, Bcl-2, and Bcl-xL protein expression remain at the basal control level (Fig. 6A and B). In addition, NiCl₂ challenge induced marginally higher levels of apoptosis in control and DN-Akt cells over the Akt stable cells (Fig. 6C),

pointing out the protective roles of Akt in NiCl₂-induced BEAS-2B cell apoptosis.

Tumorigenic properties of NiCl₂ transformed cells. To evaluate if the transformed cells are tumorigenic, we injected both control BEAS-2B and T-BEAS-2B cells into nude mice and observed the tumor growth. The results indicated that among five injected mice in each group, only T-BEAS-2B cells induced tumor growth, and tumors occurred at multiple injection sites, grew into various sizes within 8 weeks (Fig. 7A and B). Furthermore, western blot analysis revealed increased HIF-1 α NF- κ B-p50 subunit and p-Stat3 protein expression in T-BEAS-2B cells when compared with non-transformed cells, underlying the oncogenic features of the NiCl₂-transformed cells (Fig. 7C).

Discussion

Nickel is a known human carcinogen and induces genotoxic stress. A number of mechanisms have been proposed for its carcinogenic effects; such as generation of ROS, induction of DNA damage, and activation of oncogenic pathways (3). However, little information is available with regard to the mechanisms of apoptosis resistance in nickel-transformed




Figure 7. Tumorigenic properties of NiCl₂-transformed cells. BEAS-2B or transformed BEAS-2B (T-BEAS-2B) cells (2x10⁶) were mixed with matrigel and injected subcutaneously into athymic nude mice on both sides of flank region. (A and B) Five animals were injected in each group, tumor growth were monitored twice weekly until the tumor size reach 1 x 1 x 1 cm³ in size or by 8 weeks post inoculation. (C) To monitor the oncogenic protein expression in transformed BEAS-2B cells, cell lysates from each type of cell were collected and resolved in 10% SDS-PAGE to detect p-Stat3, NF-κB-p50 subunit and HIF-1α protein expression. Blots are representative of 3-4 separate experiments with similar results. Arrows indicate the specific band.

cells. In this study, we focused on apoptosis resistant and tumorigenic property of NiCl₂-transformed human lung epithelial cells, and explored the potential molecular mechanisms. The results demonstrate that NiCl₂ transformed BEAS-2B cells are resistant to apoptosis and tumorigenic. Higher levels of Bcl-2, Bcl-xL and catalase protein expression are important mechanisms contributing to the transformed cell oncogenic properties.

Previously *in vitro* studies have shown that water-insoluble nickel compounds such as nickel oxide, nickel subsulfide are more readily phagocytized than the water-soluble nickel compounds such as nickel chloride. They can reach the nucleus of the cell in greater amounts than that of water-soluble nickel compounds and induce more cell damage (19,20). In the current study, NiCl₂ was selected due to the BEAS-2B cells being transformed by NiCl₂ exposure (9). One important characteristic of nickel compounds is the generation of ROS in host cells, and ROS appear critical in nickel-induced cell damage, such as apoptosis and oncogenic transformation (1,4). Increasing evidence has indicated that ROS plays an important role in inducing apoptosis and carcinogenesis following diverse exposure to environmental stimuli (21). However, ROS generation in transformed cells is reduced (9,18), the mechanism is not clear. In the current study, we confirm that antioxidant enzymes are increased in transformed cells including catalase, SOD2 and Prx-1 (22), suggesting that they are responsible for the reduced ROS generation in transformed cells (9,18). Higher

ROS scavenging enzyme expression may also contribute to NiCl₂-induced apoptosis resistance in transformed cells by attenuating the detrimental effects from ROS generated by NiCl₂ exposure. However, the biological implications of these changes during transformation have yet to be explored, as well as the molecular mechanisms that result in their increased expression in transformed cells. The data provide evidence that transformation involves multiple cellular signal alternations, which deserve further investigation.

Bcl-2 and Bcl-xL proteins are overexpressed in a variety of human cancers (23), and are important members of anti-apoptotic proteins family. The mechanisms of Bcl-2 and Bcl-xL protect cells from apoptosis through either heterodimerization with pro-apoptotic proteins, or their direct pore-blocking effects on the outer membrane of mitochondria (11,23). Mitochondria play a critical role in apoptosis induction, which involves both caspase-dependent and -independent pathways. We noted that NiCl₂-induced apoptosis can be affected by both Bcl-2 and Bcl-xL expressing vector and siRNA transfection, indicating that their relative levels regulate cell apoptosis resistance in T-BEAS-2B cells. The question of how transformed cell evolve into higher level of Bcl-2, and Bcl-xL protein expression remain to be investigated, we hypothesis that either genetic or epigenetic modifications may be responsible during the transformation and selection processes.

In our study, NiCl₂ treatment reduced the cellular level of Bcl-2 and Bcl-xL proteins in both transformed and non-trans-

formed cells (Fig. 1A and B), but the level of reduction in transformed cells is less dramatic, which is associated with low apoptotic rate. This effect also correlates with reduced mitochondrial stress, c-caspase 3, 7 protein level, caspase enzymatic activity and PARP cleavage. Together, they suggest that transformed cells evolve a mechanism of increased Bcl-2/Bcl-xL protein expression, reduced mitochondrial stress and alleviated caspase activation upon NiCl₂ exposure, therefore, confer these cells the apoptosis resistant property.

Several signaling pathways such as PI3K, NF- κ B, Akt, and Sonic hedgehog have been shown to regulate Bcl-2 and Bcl-xL protein expression and apoptosis in different cell types (12,24,25). Induction of cyclooxygenase 2 is also important in nickel induced apoptosis (26). In the current study, we focus on the effects of Bcl-2, Bcl-xL, and catalase by examining their roles in apoptosis resistance in transformed cells, as they are either the effector proteins downstream of the above mentioned signal pathways or involved in ROS generation. Akt is a serine/threonine kinase, also known as protein kinase B (PKB). It is involved in regulating multiple cellular functions, including cell growth, proliferation, survival, glucose metabolism and genome stability (27,28). Aberrant Akt activation has been noted in breast, prostate, lung, pancreatic, liver, ovarian and colorectal cancers, and in malignant transformation (27-30). Akt overexpression enhances Bcl-xL expression, which is correlated with the fact that Akt-stable expressing cells are more resistant to NiCl₂-induced apoptosis. The results uncovered anti-apoptotic protein-dependent mechanism in nickel-induced apoptosis resistance in transformed cells.

It is worth noting that overexpression of Akt could not completely block the nickel-induced apoptosis (Fig. 6), and even this effect was marginal. Since there are two major apoptotic pathways, the death-receptor pathway and the mitochondrial pathway, we consider other signal pathways might also contribute to the nickel-induced apoptosis. It has been reported that increased FasL expression, cell cycle alteration, activation of c-Myc through ERK pathway, caspase-8/AIF-mediated pathways, and activation of NF- κ B/Cox2 pathway all participate in nickel-induced apoptosis (26,31-34). Therefore, each may initiate and contribute to the pathway or cell type-specific apoptosis processes in addition to the mechanisms discussed in the present study. However, increased Bcl-2, Bcl-xL and catalase protein expression, reduced mitochondria damage and caspase activation described here appear to be the essential mechanisms involved in NiCl₂-mediated cell death resistance program in transformed BEAS-2B cells. Future investigations will be required to clarify if other mechanisms might also be involved in transformed cell resistance to apoptosis.

One important feature of transformed cells is their tumorigenic property. These cells also have enhanced p-Stat3, NF- κ B-p50 subunit, and HIF-1 α protein expression. The altered oncogene expression probably reflects extensive genetic or epigenetic rearrangements during the transformation processes which allow transformed cells to adopt and grow in a new environment. It is also possible that they acquire cancer stem cell trait to allow cell resistant apoptosis. Future studies focusing on nickel-induced apoptosis resistance and cell transformation are of great interest to understand nickel-induced carcinogenic mechanisms and to provide options for prevention.

Acknowledgements

This research was supported in part by NIH grants (1R01CA119028) and a Henan provincial government biomedical key research grant (132102310444).

References

- Costa M: Molecular mechanisms of nickel carcinogenesis. *Annu Rev Pharmacol Toxicol* 31: 321-337, 1991.
- Doll R, Morgan LG and Speizer FE: Cancers of the lung and nasal sinuses in nickel workers. *Br J Cancer* 24: 623-632, 1970.
- Costa M, Davidson TL, Chen H, *et al*: Nickel carcinogenesis: epigenetics and hypoxia signaling. *Mutat Res* 592: 79-88, 2005.
- Lu H, Shi X, Costa M and Huang C: Carcinogenic effect of nickel compounds. *Mol Cell Biochem* 279: 45-67, 2005.
- Salnikow K and Zhitkovich A: Genetic and epigenetic mechanisms in metal carcinogenesis and cocarcinogenesis: nickel, arsenic, and chromium. *Chem Res Toxicol* 21: 28-44, 2008.
- Salnikow K and Costa M: Epigenetic mechanisms of nickel carcinogenesis. *J Environ Pathol Toxicol Oncol* 19: 307-318, 2000.
- Davidson T, Chen H, Garrick MD, D'Angelo G and Costa M: Soluble nickel interferes with cellular iron homeostasis. *Mol Cell Biochem* 279: 157-162, 2005.
- Pan J, Chang Q, Wang X, *et al*: Reactive oxygen species-activated Akt/ASK1/p38 signaling pathway in nickel compound-induced apoptosis in BEAS 2B cells. *Chem Res Toxicol* 23: 568-577, 2010.
- Pan JJ, Chang QS, Wang X, *et al*: Activation of Akt/GSK3 β and Akt/Bcl-2 signaling pathways in nickel-transformed BEAS-2B cells. *Int J Oncol* 39: 1285-1294, 2011.
- Cory S and Adams JM: The Bcl2 family: regulators of the cellular life-or-death switch. *Nat Rev Cancer* 2: 647-656, 2002.
- Gross A, McDonnell JM and Korsmeyer SJ: BCL-2 family members and the mitochondria in apoptosis. *Genes Dev* 13: 1899-1911, 1999.
- Pugazhenth S, Nesterova A, Sable C, *et al*: Akt/protein kinase B up-regulates Bcl-2 expression through cAMP-response element-binding protein. *J Biol Chem* 275: 10761-10766, 2000.
- Ding SZ, Yang YX, Li XL, *et al*: Epithelial-mesenchymal transition during oncogenic transformation induced by hexavalent chromium involves reactive oxygen species-dependent mechanism in lung epithelial cells. *Toxicol Appl Pharmacol* 269: 61-71, 2013.
- Zhou BP, Hu MC, Miller SA, *et al*: HER-2/neu blocks tumor necrosis factor-induced apoptosis via the Akt/NF-kappaB pathway. *J Biol Chem* 275: 8027-8031, 2000.
- Ding SZ, Fischer W, Kaparakis-Liaskos M, *et al*: *Helicobacter pylori*-induced histone modification, associated gene expression in gastric epithelial cells, and its implication in pathogenesis. *PLoS One* 5: e9875, 2010.
- Cossarizza A, Baccarani-Contri M, Kalashnikova G and Franceschi C: A new method for the cytofluorimetric analysis of mitochondrial membrane potential using the J-aggregate forming lipophilic cation 5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolcarbocyanine iodide (JC-1). *Biochem Biophys Res Commun* 197: 40-45, 1993.
- Cheng EH, Wei MC, Weiler S, *et al*: BCL-2, BCL-X(L) sequester BH3 domain-only molecules preventing BAX- and BAK-mediated mitochondrial apoptosis. *Mol Cell* 8: 705-711, 2001.
- Chang Q, Pan J, Wang X, Zhang Z, Chen F and Shi X: Reduced reactive oxygen species-generating capacity contributes to the enhanced cell growth of arsenic-transformed epithelial cells. *Cancer Res* 70: 5127-5135, 2010.
- Dunnick JK, Elwell MR, Radovsky AE, *et al*: Comparative carcinogenic effects of nickel subsulfide, nickel oxide, or nickel sulfate hexahydrate chronic exposures in the lung. *Cancer Res* 55: 5251-5256, 1995.
- Oller AR, Costa M and Oberdorster G: Carcinogenicity assessment of selected nickel compounds. *Toxicol Appl Pharmacol* 143: 152-166, 1997.
- Zhao J, Shi X, Castranova V and Ding M: Occupational toxicology of nickel and nickel compounds. *J Environ Pathol Toxicol Oncol* 28: 177-208, 2009.
- Kim YJ, Ahn JY, Liang P, Ip C, Zhang Y and Park YM: Human prx1 gene is a target of Nrf2 and is up-regulated by hypoxia/reoxygenation: implication to tumor biology. *Cancer Res* 67: 546-554, 2007.

23. Yip KW and Reed JC: Bcl-2 family proteins and cancer. *Oncogene* 27: 6398-6406, 2008.
24. Catz SD and Johnson JL: Transcriptional regulation of bcl-2 by nuclear factor kappa B and its significance in prostate cancer. *Oncogene* 20: 7342-7351, 2001.
25. Bigelow RL, Chari NS, Uden AB, *et al*: Transcriptional regulation of bcl-2 mediated by the sonic hedgehog signaling pathway through gli-1. *J Biol Chem* 279: 1197-1205, 2004.
26. Ding J, Zhang X, Li J, *et al*: Nickel compounds render anti-apoptotic effect to human bronchial epithelial BEAS-2B cells by induction of cyclooxygenase-2 through an IKKbeta/p65-dependent and IKKalpha- and p50-independent pathway. *J Biol Chem* 281: 39022-39032, 2006.
27. Lawlor MA and Alessi DR: PKB/Akt: a key mediator of cell proliferation, survival and insulin responses? *J Cell Sci* 114: 2903-2910, 2001.
28. Morley S, Wagner J, Kauppinen K, Sherman M and Manor D: Requirement for Akt-mediated survival in cell transformation by the dbl oncogene. *Cell Signal* 19: 211-218, 2007.
29. Bellacosa A, Kumar CC, Di Cristofano A and Testa JR: Activation of AKT kinases in cancer: implications for therapeutic targeting. *Adv Cancer Res* 94: 29-86, 2005.
30. Manning BD and Cantley LC: AKT/PKB signaling: navigating downstream. *Cell* 129: 1261-1274, 2007.
31. Li Q, Suen TC, Sun H, Arita A and Costa M: Nickel compounds induce apoptosis in human bronchial epithelial Beas-2B cells by activation of c-Myc through ERK pathway. *Toxicol Appl Pharmacol* 235: 191-198, 2009.
32. Ding J, He G, Gong W, *et al*: Effects of nickel on cyclin expression, cell cycle progression and cell proliferation in human pulmonary cells. *Cancer Epidemiol Biomarkers Prev* 18: 1720-1729, 2009.
33. Kim K, Lee SH, Seo YR, Perkins SN and Kasprzak KS: Nickel(II)-induced apoptosis in murine T cell hybridoma cells is associated with increased fas ligand expression. *Toxicol Appl Pharmacol* 185: 41-47, 2002.
34. Zhao J, Bowman L, Zhang X, *et al*: Metallic nickel nano- and fine particles induce JB6 cell apoptosis through a caspase-8/AIF mediated cytochrome c-independent pathway. *J Nanobiotechnol* 7: 2, 2009.