

## Research Article

# The Effect of Teach-Back Combined with King Interactive Standard Mode on the Life of COPD Patients

Jiayi Rang <sup>1</sup>, Liming Peng,<sup>2</sup> Long Wen,<sup>3</sup> Zhiguo Zhou,<sup>3</sup> You Xia,<sup>1</sup> Chaoying Xie,<sup>4</sup> Ting Xie,<sup>3</sup> and Jing Tan<sup>3</sup>

<sup>1</sup>Nursing Department, Changsha City First Hospital, Changsha 410000, Hunan, China

<sup>2</sup>Lao Gan Activity Center, Changsha City First Hospital, Changsha 410000, Hunan, China

<sup>3</sup>The Department of Respiratory and Critical Care Medicine, Changsha City First Hospital, Changsha 410000, Hunan, China

<sup>4</sup>Outpatient Office, Changsha City First Hospital, Changsha 410000, Hunan, China

Correspondence should be addressed to Jiayi Rang; 2016120220@jou.edu.cn

Received 12 July 2022; Revised 2 August 2022; Accepted 16 August 2022; Published 20 September 2022

Academic Editor: Yuvaraja Teekaraman

Copyright © 2022 Jiayi Rang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Background.** COPD is a common clinical chronic airway inflammatory disease that occurs mostly in middle-aged and older adults over the age of 40. The incidence of COPD is increasing year by year and the onset of age is gradually becoming younger. **Objective.** To observe the effect of teach-back combined with king interaction on the life of patients with chronic obstructive pulmonary disease (COPD). **Methods.** A total of 100 COPD patients admitted to our hospital from Jan 2021 to Jan 2022 were retrospectively selected to be divided into 50 cases in the control group and 50 cases in the observation group according to the nursing methods. The control group was treated with routine nursing intervention, while the observation group was treated with teach-back combined with king interactive standard mode intervention. The differences in Self-Care Ability Assessment Scale (ESCA) score, St. George's Respiratory Questionnaire (SGRQ) score, Mental State Assessment Scale (MSSNS) score, 6-minute walking distance (6MWD), and pulmonary function indexes were compared between the two groups before and after the intervention. The success rate and patient compliance of each index in the groups were also recorded. **Results.** After 3 months and 6 months of intervention, the total SGRO score and its factor scores of self-care skills, self-care responsibility, self-concept, health knowledge level in them were all higher than those before the intervention, while the total SGRO score and its factor scores of respiratory symptoms, activity limitation, disease influence, and so on were all decreased compared with those before the intervention. The ESCA score of the observation group was significantly higher than that of the control one after 3 months and 6 months of intervention, while the SGRQ score was significantly lower than that of the control one, with statistical significance ( $P < 0.05$ ). After 3 months of intervention, the total score of MSSNS and the scores of anxiety, depression, loneliness, and other factors in both groups were decreased compared with those before intervention. After 6 months of intervention, the total score of MSSNS and scores of each factor in both groups were decreased compared with those before intervention, and the MSSNS scores in the observation group were significantly lower than those in the control group after the intervention, which was statistically significant ( $P < 0.05$ ). After 3 months and 6 months of intervention, 6MWD, forced vital capacity (FVC), forced expiratory value in 1 second (FEV1), and FVC/FEV1 in them were all higher than those before intervention, and 6MWD and pulmonary function were significantly higher in the observation group than in the control group after 3 and 6 months of intervention, which was statistically significant ( $P < 0.05$ ). The ESCA score, SGRQ score, MSSNS score, pulmonary function compliance rate, and compliance rate in the observation group were significantly higher than those in the control group, which was statistically significant ( $P < 0.05$ ). **Conclusion.** The use of teach-back combined with king interactive standard mode in COPD patients can improve the patient's self-care ability, reduce psychological negative emotions, and improve the quality of life.

## 1. Introduction

COPD is an airway inflammatory disease characterized by incompletely reversible airflow limitation, with clinical manifestations of chronic cough, sputum, and dyspnea. The symptoms of acute exacerbation are aggravated, and the patients can die from respiratory failure [1, 2]. The main goals of current clinical treatment of COPD are controlling the disease and preventing acute exacerbations. However, the risk of rehospitalization within one year after discharge increases due to the lack of awareness about COPD-related knowledge. The burden of COPD is heavy. It is expected to become the third leading cause of human death by 2030 [3]. Therefore, it is of great importance to improve the self-management ability and life of COPD patients.

Teach-back is a scientific method of communication in which nursing staff provides health education to patients and instructs patients to repeat the knowledge points in their own words to ensure that they really master health knowledge [4]. King interaction standard mode is a mutual pointing interaction model, which deepens the communication and interaction between nurses and patients through the process of perception, judgment, behavior, and response, so as to achieve a good communication effect [5]. The teach-back and king interaction mode have been used in the nursing of respiratory diseases, while the research was not sufficient. In order to improve the awareness rate of COPD-related knowledge and improve the self-management ability of patients and their life quality, the teach-back combined with king interactive compliance model was applied to COPD patients in this study. The report is as follows.

*1.1. Core Tips.* In this study, the scientific communication model teach-back combined with king interactive compliance model was used to intervene in patients with COPD, and it was found that it could improve the self-care ability of patients, reduce psychological distress, and improve their quality of life.

## 2. Data and Methods

*2.1. General Information.* A total of 100 patients with COPD admitted to our hospital from Jan 2021 to Jan 2022 were retrospectively selected, of whom 59 were males and 41 were females, aged from 40 to 75 years old, with an average of  $(60.44 \pm 8.85)$  years. The course of disease was 2~17 years, with an average of  $(8.96 \pm 1.77)$  years. They were divided into control group (50 cases) and observation group (50 cases) according to the nursing methods. There was no significant difference in specific general data between them ( $P > 0.05$ ).

*2.2. Case Selection Criteria.* Inclusion criteria: (1) in line with the criteria for stable COPD in the Guidelines for the Diagnosis and Treatment of Chronic Obstructive Pulmonary Disease; (2) ages from 18 to 75 years old; (3) having basic verbal communication skills and being able to cooperate with the intervention and assessment of relevant scores; (4) with complete clinical and follow-up data.

Exclusion criteria: (1) with severe mental diseases; (2) presence of in situ or metastatic malignancy; (3) poor overall physical condition with an expected survival of less than six months; (4) inability to perform the 6-minute walk test due to the presence of physical dysfunction; (5) with functional impairment of important organs; (6) long-term bed rest; (7) with hematopoietic system diseases.

*2.3. Method.* In the control group, conventional nursing interventions were used to instruct patients to maintain healthy habits, follow medical prescriptions, perform respiratory function exercises, and pay attention to the prevention of cold, infection, and inhalation of harmful fumes.

The observation group was intervened by teach-back combined with king interactive compliance model. The intervention group of teach-back combined with king interactive compliance model was established, with the nurse manager as the group leader and the charge nurse as the group member. The group was organized to learn COPD rehabilitation training, teach-back and king interactive compliance concept, and other related knowledge. Nursing staff was allowed to join the group after passing the examination. Intervention: (1) Evaluation: One-to-one communication was used on the day of admission or the next day to know the patients' understanding of COPD-related knowledge, self-management ability, willingness, and attitude to participate in interaction. (2) Plan: Develop an intervention plan based on the assessment results by reviewing the relevant literature. The teach-back questionnaire was formulated to carry out the scenario simulation exercise. After that, the existing problems were analyzed and corrective measures were proposed. (3) Implementation: Nursing staff used easy-to-understand language to explain COPD-related knowledge according to the patient's understanding of them, guiding patients to establish a healthy lifestyle, follow medical prescriptions, perform respiratory function exercises, and pay attention to the prevention of cold, infection, and inhalation of harmful fumes. They should ask patients according to the teach-back questionnaire: Did you understand what I just said? What do you agree with? Why is it important to establish a healthy lifestyle, take medication as prescribed, and perform respiratory function exercises? How many times a day do you do respiratory function exercises? How long do you do them each time? Can you show me how to do them? Is there anything else that is unclear to you? If the patient cannot repeat them correctly, the nursing staff should repeat the explanation again. (4) Evaluation: Assess the achievement of the single target value of the plan, analyze the reasons for the failure, and timely adjust the intervention measures. After the patients were discharged from the hospital, the interventions were continued up to 6 months by outpatient follow-up, telephone follow-up or WeChat follow-up, family visits, and other forms.

*2.4. Detection Method.* Exercise endurance was measured by 6MWD before intervention, 3 months and 6 months after intervention. Patients walked as fast as they could on a flat

surface after 15 min of rest, and the walking distance within 6 min was recorded with the unit of meter. The forced vital capacity (FVC) forced expiratory volume in one second (FEV1) with the unit of ml and FVC/FEV1 were measured by German Yager pulmonary function detector.

**2.5. Score Standard.** Self-care ability assessment scale (ESCA) score [6]: evaluation was conducted before intervention, 3 months and 6 months after intervention, including four aspects of self-care skills, sense of responsibility, self-concept, and health knowledge level, with a total of 43 pieces and a single score of 0 to 4. The high ~ low ESCA score indicates the high ~ low self-care ability.

Saint George's Respiratory Questionnaire (SGRQ) score [7]: 43 questions were assessed before intervention, 3 and 6 months after intervention, including respiratory symptoms, activity limitation, and disease impact, with 43 questions and a total score range of 0 to 100. The high-low SGRQ score represented the low-high quality of life.

Mental state assessment scale (MSSNS) score [8]: The evaluation was conducted before intervention, 3 months and 6 months after intervention, including four aspects of anxiety, depression, anger, and loneliness, with a total of 38 pieces and a single score of 0 to 4. The high ~ low MSSNS score indicates the high ~ low of negative psychology.

Compliance rate: the number of indicators reaching the standard/total number of cases  $\times$  100%.

Compliance rate: the self-developed score of this study was used, which covered respiratory function exercise, medication, lifestyle, etc., with a total score range of 0~100, of which 90 or more points were defined as complete compliance, 60~89 points as basic compliance, 60 points below as noncompliance. Complete and basic compliance were defined as compliance.

**2.6. Statistical Method.** The data were processed by SPSS26.0. The normality and homogeneity of the measurement data were detected by the K-S method test and Levene's method test, respectively. The measurement data conforming to the standard were described by ( $\chi \pm s$ ). The *t*-test was used for comparison.  $\chi^2$  test was applied to compare the count data by the four-compartment table method or the  $\chi^2$  test with row  $\times$  list, and  $P < 0.05$  was statistically significant.

### 3. Results

**3.1. Analysis of Two Groups of General Data.** There was no significant difference in the general data of residence, pulmonary function classification, education level, age, gender, and combined diseases between them ( $P > 0.05$ ). See Table 1.

**3.2. Comparison of ESCA Scores between the Two Groups.** Before intervention, there was no significant difference in ESCA score between them ( $P > 0.05$ ). After 3 months and 6 months of intervention, the total SGRO score and its factor scores of self-care skills, self-care responsibility, self-concept, health knowledge level in them were all higher than before,

and the ESCA score of the observation group was significantly higher than that of the control one after 3 months and 6 months of intervention, with statistical significance ( $P < 0.05$ ). See Table 2.

**3.3. Comparison of SGRO Scores between the Two Groups.** Before intervention, there was no significant difference in SGRO score between them ( $P > 0.05$ ); the total SGRO score and its factor scores of respiratory symptoms, activity limitation, disease influence, and so on were all decreased compared with those before the intervention, and the SGRO score was significantly lower than that of the control one, with statistical significance ( $P < 0.05$ ). See Table 3.

**3.4. Comparison of MSSNS Scores between the Two Groups.** Before intervention, there was no significant difference in MSSNS score between them ( $P > 0.05$ ). After 3 months and 6 months of intervention, the total score of MSSNS score and the scores of anxiety, depression, anger (except for 3 months of intervention), and loneliness in them were lower than those before intervention, and the MSSNS score and its factor score (except for anger at 3 months of intervention) in the observation one were significantly lower than those in the control one, with statistical significance ( $P < 0.05$ ). See Table 4.

**3.5. Comparison of 6MWD and Pulmonary Function between Two Groups.** Before intervention, there was no significant difference in 6MWD and pulmonary function indexes between them ( $P > 0.05$ ). After 3 months and 6 months of intervention, the 6MWD, FVC, FEV1, and FVC/FEV1 of them were higher than those before intervention, and the 6MWD and pulmonary function of the observation one were significantly higher than those of the control one after 3 months and 6 months of intervention, with statistical significance ( $P < 0.05$ ). See Figure 1.

**3.6. Comparison of Two Groups of Indicators Compliance Rate.** The ESCA score, SGRQ score, MSSNS score, and lung function compliance rate in the observation one were 90.00% (45/50), 84.00% (42/50), 88.00% (44/50), and 82.00% (82/50), respectively, which were significantly higher than those in the control one with 74.00% (37/50), 66.00% (33/50), 72.00% (36/50), and 64.00% (32/50), respectively, with statistical significance ( $P < 0.05$ ). See Table 5.

**3.7. Compliance Comparison between Two Groups.** The compliance rate of the observation group was 88.00% (44/50), which was significantly higher than 68.00% (34/50) of the control one, with statistical significance ( $P < 0.05$ ). See Table 6.

### 4. Discussion

Dyspnea can not only lead to a decline in exercise endurance and reduce the life quality of patients, but also cause greater economic burden on patients and cause psychological

TABLE 1: Analysis of the general data of the two groups.

Normal information	Control one ( $n = 50$ )	Observation one ( $n = 50$ )	$\chi^2$ or $t$	$P$
Gender [ $n(\%)$ ]				
Male	27 (54.00)	32 (64.00)	1.034	0.309
Female	23 (46.00)	18 (36.00)		
Age [ $(\bar{x} \pm s)$ , age]	61.42 $\pm$ 9.43	59.86 $\pm$ 10.04	0.801	0.425
Body mass index [ $(\bar{x} \pm s)$ , kg/m <sup>2</sup> ]	24.51 $\pm$ 2.12	24.39 $\pm$ 2.25	0.274	0.784
Course of disease [ $(\bar{x} \pm s)$ , year]	8.67 $\pm$ 1.85	9.05 $\pm$ 1.77	1.049	0.297
Place of residence [ $n(\%)$ ]				
City	28 (56.00)	26 (52.00)	0.161	0.688
Rural	22 (44.00)	24 (48.00)		
Pulmonary function class [ $n(\%)$ ]				
Class I	28 (56.00)	25 (50.00)	0.486	0.784
Class II	16 (32.00)	17 (34.00)		
Class III	6 (12.00)	8 (16.00)		
Educational level [ $n(\%)$ ]				
Junior high school and below	18 (36.00)	15 (30.00)	0.963	0.618
High school and secondary school	22 (44.00)	21 (42.00)		
College and above	10 (20.00)	14 (28.00)		
Concomitant disease [ $n(\%)$ ]				
Hypertension	17 (34.00)	14 (28.00)	0.421	0.517
Diabetes	8 (16.00)	12 (24.00)	1.000	0.317
Cerebrovascular disease	14 (28.00)	15 (30.00)	0.049	0.826
Hyperlipidemia	9 (18.00)	11 (22.00)	0.250	0.617

TABLE 2: Comparison of ESCA scores between the two groups [ $(\bar{x} \pm s)$ , fraction].

Factor	Time	Control one $n = 50$	Observation one $n = 50$	$t$	$P$
Self-care skills	Before intervention	20.55 $\pm$ 4.78	20.46 $\pm$ 4.83	0.094	0.926
	Intervention for 3 months	26.88 $\pm$ 5.11*	30.13 $\pm$ 5.15*	3.168	0.002
	Intervention for 6 months	32.25 $\pm$ 6.62*	39.25 $\pm$ 7.14*	5.084	<0.001
Self-care responsibility	Before intervention	14.25 $\pm$ 3.28	14.09 $\pm$ 3.31	0.243	0.809
	Intervention for 3 months	16.88 $\pm$ 3.33*	18.96 $\pm$ 3.29*	3.142	0.002
	Intervention for 6 months	19.54 $\pm$ 3.17*	22.45 $\pm$ 3.69*	4.230	<0.001
Self-concept	Before intervention	17.44 $\pm$ 3.13	17.52 $\pm$ 2.97	0.131	0.896
	Intervention for 3 months	19.05 $\pm$ 3.05*	21.53 $\pm$ 3.39*	3.846	<0.001
	Intervention for 6 months	22.96 $\pm$ 3.45*	25.85 $\pm$ 3.59*	4.104	<0.001
Health knowledge level	Before intervention	41.08 $\pm$ 8.22	40.73 $\pm$ 8.69	0.207	0.837
	Intervention for 3 months	45.45 $\pm$ 6.78*	49.63 $\pm$ 8.85*	2.900	0.005
	Intervention for 6 months	50.22 $\pm$ 7.89*	57.14 $\pm$ 7.61*	4.464	<0.001
ESCA total score	Before intervention	93.32 $\pm$ 9.04	92.87 $\pm$ 10.11	0.235	0.815
	Intervention for 3 months	108.26 $\pm$ 12.78*	120.25 $\pm$ 13.69*	4.527	<0.001
	Intervention for 6 months	124.97 $\pm$ 15.47*	144.69 $\pm$ 18.55*	5.773	<0.001

\* $P < 0.05$  indicates the comparison with the control group.

TABLE 3: Comparison of SGRQ scores between the two groups [ $(\bar{x} \pm s)$ , fraction].

Factor	Time	Control one $n = 50$	Observation one $n = 50$	$t$	$P$
Respiratory symptoms	Before intervention	63.25 $\pm$ 6.85	64.32 $\pm$ 6.72	0.788	0.432
	Intervention for 3 months	53.77 $\pm$ 5.16*	49.08 $\pm$ 5.46*	4.414	<0.001
	Intervention for 6 months	46.26 $\pm$ 5.58*	43.56 $\pm$ 5.05*	2.537	0.013
Limited activity	Before intervention	60.74 $\pm$ 7.88	59.65 $\pm$ 7.96	0.688	0.493
	Intervention for 3 months	52.77 $\pm$ 7.45*	46.32 $\pm$ 6.39*	4.647	<0.001
	Intervention for 6 months	45.36 $\pm$ 5.97*	40.45 $\pm$ 4.82*	4.525	<0.001
Disease impact	Before intervention	61.78 $\pm$ 7.12	62.67 $\pm$ 6.31	0.661	0.510
	Intervention for 3 months	54.55 $\pm$ 6.85*	49.44 $\pm$ 6.66*	3.782	<0.001
	Intervention for 6 months	44.89 $\pm$ 5.34*	41.28 $\pm$ 5.13*	3.447	0.001
Overall SGRQ score	Before intervention	61.92 $\pm$ 5.88	62.21 $\pm$ 6.03	0.243	0.808
	Intervention for 3 months	53.69 $\pm$ 5.41*	48.28 $\pm$ 5.11*	5.140	<0.001
	Intervention for 6 months	45.50 $\pm$ 4.97*	41.76 $\pm$ 4.57*	3.917	<0.001

\* $P < 0.05$  indicates the comparison with the control group.

TABLE 4: Comparison of MSSNS scores between the two groups  $[(\chi \pm s), \text{fraction}]$ .

Factor	Time	Control one $n = 50$	Observation one $n = 50$	$t$	$P$
Anxiety	Before intervention	24.21 $\pm$ 3.77	24.08 $\pm$ 4.56	0.155	0.877
	Intervention for 3 months	22.74 $\pm$ 3.12*	21.51 $\pm$ 2.98*	2.016	0.047
	Intervention for 6 months	20.39 $\pm$ 2.63*	19.23 $\pm$ 3.12*	2.010	0.047
Depression	Before intervention	15.88 $\pm$ 2.12	15.79 $\pm$ 2.26	0.205	0.838
	Intervention for 3 months	13.73 $\pm$ 2.05*	12.26 $\pm$ 1.79*	3.819	<0.001
	Intervention for 6 months	11.64 $\pm$ 1.85*	10.25 $\pm$ 1.74*	3.870	<0.001
Anger	Before intervention	10.56 $\pm$ 2.44	10.48 $\pm$ 2.36	0.167	0.868
	Intervention for 3 months	9.87 $\pm$ 2.11	9.71 $\pm$ 1.98	0.391	<0.001
	Intervention for 6 months	9.46 $\pm$ 1.89*	8.34 $\pm$ 1.45*	3.325	0.001
Lonely	Before intervention	13.66 $\pm$ 2.54	13.74 $\pm$ 2.48	0.159	0.874
	Intervention for 3 months	12.06 $\pm$ 2.13*	10.58 $\pm$ 1.91*	3.658	<0.001
	Intervention for 6 months	9.65 $\pm$ 1.82*	8.77 $\pm$ 1.65*	2.533	0.013
MSSNS total score	Before intervention	64.31 $\pm$ 5.22	64.09 $\pm$ 4.93	0.217	0.829
	Intervention for 3 months	58.41 $\pm$ 4.53*	54.06 $\pm$ 4.11*	5.029	<0.001
	Intervention for 6 months	51.14 $\pm$ 4.15*	48.87 $\pm$ 4.03*	2.775	0.007

\* $P < 0.05$  indicates the comparison with the control group.




FIGURE 1: The FVC, FEV1, FVD/FEV1, and 6MWD of two groups before intervention, 3 months after intervention and 6 months after intervention.

disorders [9, 10]. A survey found that Chinese residents have low awareness of COPD-related knowledge, resulting in about 43% of COPD patients needing hospitalization at least once a year, and more than half of the first-time hospitalized patients died within 3.6 years [11, 12]. Therefore, special

attention should be paid to the development and improvement of patients' self-management skills in interventions for COPD to reduce acute COPD exacerbation, which can not only reduce the cost of hospitalization, but also delay the disease and improve the prognosis [13].

TABLE 5: Comparison of the compliance rate of each index between the two groups [ $n(\%)$ ].

Group	$n$	ESCA score compliance rate	SGRQ score compliance rate	MSSNS score compliance rate	Pulmonary function compliance rate
Control one	50	37 (74.00)	33 (66.00)	36 (72.00)	32 (64.00)
Observation one	50	45 (90.00)	42 (84.00)	44 (88.00)	41 (82.00)
$\chi^2$		4.336	4.320	4.000	4.110
$P$		0.037	0.038	0.046	0.043

TABLE 6: Comparison of compliance between the two groups [ $n(\%)$ ].

Group	$n$	Full compliance	Basic compliance	Noncompliance	Compliance rate
Control one	50	14 (28.00)	20 (40.00)	16 (32.00)	34 (68.00)
Observation one	50	32 (64.00)	12 (24.00)	6 (12.00)	44 (88.00)
$\chi^2$					5.828
$P$					0.016

Teach-back method is a simple and practical way of health education, which ensures patients to complete health education better and improve their disease-related knowledge level through explaining, answering, retelling, correcting, and other steps, especially for people with poor health knowledge [4, 14]. At present, teach-back method has been widely used in postoperative rehabilitation, chronic diseases, teaching, and other fields [15, 16]. King interactive standard model believes that the interaction between nurses and patients can help nurses to fully understand the psychological and physiological changes of patients, formulate targeted nursing intervention, and it can stimulate the potential of patients, giving full play to their subjective initiative, which has been applied in obstetrics, surgery, internal medicine, and other fields [17, 18].

In this study, the teach-back combined with king interactive compliance model was applied to the intervention of COPD patients. It was found that the total score of SGRO score and its self-care skills, self-care responsibility, self-concept, health knowledge level of patients after 3 months and 6 months of intervention were higher than those of patients with conventional intervention. The total score of SGRQ score and its respiratory symptoms, activity limitation, and disease impact were lower than those of patients with conventional intervention. Teach-back combined with king interactive compliance model for COPD patients can improve their self-care ability and respiratory function. The results of 6MWD and pulmonary function test suggested that the 6MWD, FVC, FEV1, and FVC/FEV1 of the intervention group using teach-back combined with king interactive compliance model after 3 and 6 months of intervention were higher than those of the conventional intervention one, and the ESCA score, SGRQ score, MSSNS score, and pulmonary function compliance rate were significantly higher than those of the conventional intervention one. The results suggested that teach-back combined with king interactive compliance model for COPD patients could improve exercise tolerance and pulmonary ventilation function, which was due to the fact that the teach-back method eliminated the one-way information transmission mode of routine health education and evaluated and provided feedback on patients' knowledge

mastery through retelling to help patients better grasp and understand health education information [19, 20]. In the process of king interactive standard intervention, it focuses on the interaction between nurses and patients and strengthens health knowledge through the spiral process of continuous cognition, error correction, and recognition, so as to improve the self-management ability of patients, stimulate their subjective initiative, and better implement the compliance behavior [21–23]. In the king interactive compliance model, nurses participate in the whole process of interaction, which can discover the problems in the plan and adjust the intervention scheme in time to ensure the continuity of intervention [24, 25].

COPD patients have decreased motor function due to somatic symptoms, and some patients even lose their ability to work and self-care in daily life, which causes negative emotions [26–28]. Severe negative emotion can affect patients' sleep quality and their treatment compliance, which is detrimental to the prognosis [29–31]. The study found that the total MSSNS score and its factor scores at 3 months and 6 months of the teach-back combined with king interactive compliance model were lower than those who used the conventional intervention, and the compliance rate was higher than that of the conventional intervention. It suggests that teach-back combined with king interactive compliance model for COPD patients can reduce the negative emotions of patients and improve treatment compliance. This is due to the fact that patients in teach-back combined with king interactive compliance model have better mastery of disease-related knowledge and pay more attention to disease treatment and compliance behavior, resulting in an increased compliance. Patients communicate with nursing staff throughout the process in this model, so as to find out their negative emotions and get timely relief, thus reducing their negative emotions [32, 33].

There were some limitations of this study. This study was a retrospective study, and the selection bias was inevitable. Furthermore, the sample of this study was only 100 patients. And finally, this study was performed in only one hospital. Thus, the results should be confirmed by further multicenter randomized controlled clinical trials.

In conclusion, teach-back combined with king interactive compliance model for COPD patients can improve their self-care ability, reduce psychological distress, and improve the quality of life.

### Data Availability

The datasets used and analyzed during the current study can be obtained from the corresponding author upon reasonable request.

### Conflicts of Interest

The authors declare that they have no conflicts of interest.

### Authors' Contributions

Jiayi Rang, Liming Peng, and Long Wen have contributed equally to this work and share first authorship.

### Acknowledgments

This work was supported by the influence of diversified care on the rehabilitation and psychological status of Novel Coronavirus pneumonia patients in the study (kq2001010) and the effect of the Teach-back combined with king interaction-compliance mode on the quality of life of patients COPD (202114010379).

### References

- [1] B. Zarrabian and M. Mirsaedi, "A trend analysis of chronic obstructive pulmonary disease mortality in the United States by race and sex," *Annals of the American Thoracic Society*, vol. 18, no. 7, pp. 1138–1146, 2021.
- [2] J. S. Li, "Guidelines for Chinese medicine rehabilitation of chronic obstructive pulmonary disease," *World J Tradit Chin Med*, vol. 6, no. 3, pp. 295–306, 2020.
- [3] K. M. Suhas, G. K. Alaparathi, S. K. Krishnan, and K. C. Bairapareddy, "Upper limb extremity muscle-dysfunction in chronic obstructive pulmonary disease: a narrative review," *Current Respiratory Medicine Reviews*, vol. 16, no. 1, pp. 11–20, 2020.
- [4] S. Ahmadidarrehsima, E. A. Bidmeshki, M. Rahnama, K. Babaei, M. Afshari, and B. Khandani, "The effect of self-management education by the teach-back method on uncertainty of patients with breast cancer: a quasi-experimental study," *Journal of Cancer Education*, vol. 35, no. 2, pp. 366–372, 2020.
- [5] D. Boges, M. Agus, R. Sicut, P. J. Magistretti, M. Hadwiger, and C. Cali, "Virtual reality framework for editing and exploring medial axis representations of nanometric scale neural structures," *Computers & Graphics*, vol. 91, no. 10, pp. 12–24, 2020.
- [6] L. Zixu, "Effect of graphic media authorized education method on ESCA score and quality of life in patients with ischemic stroke at convalescence," *Gansu Medicine*, vol. 38, no. 12, pp. 1111–1113, 2019.
- [7] Application of Liu Tao, "Cai Boros.St. George's respiratory questionnaire in quality of life assessment of patients with chronic obstructive pulmonary disease," in *Proceedings of the 2008 Beijing Annual Conference Papers of Respiratory Professionals*, pp. 206–207, 2008.
- [8] G. Cuiying, X. Su, and Y. Wang, "Establishment and application of psychological nursing mode in general hospitals," *Chinese Journal of Practical Nursing*, vol. 31, no. 21, pp. 1609–1611, 2015.
- [9] D. Lee, R. Lee, A. Tran et al., "Hemiarthroplasty versus total hip arthroplasty for femoral neck fractures in patients with chronic obstructive pulmonary disease," *European Journal of Trauma and Emergency Surgery*, vol. 47, no. 2, pp. 547–555, 2021.
- [10] A. L. Young, F. J. S. Bragman, B. Rangelov et al., "Disease progression modeling in chronic obstructive pulmonary disease," *American Journal of Respiratory and Critical Care Medicine*, vol. 201, no. 3, pp. 294–302, 2020.
- [11] R. Duan, H. Niu, T. Yu et al., "Adverse effects of short-term personal exposure to fine particulate matter on the lung function of patients with chronic obstructive pulmonary disease and asthma: a longitudinal panel study in Beijing, China," *Environmental Science and Pollution Research*, vol. 28, no. 34, pp. 47463–47473, 2021.
- [12] Y. Yang, Z. Mao, Q. Li, and J. Mao, "Fatigue and health-related quality of life among patients with chronic obstructive pulmonary disease in China," *The Clinical Respiratory Journal*, vol. 14, no. 2, pp. 109–115, 2020.
- [13] S. Bchir, S. Boumiza, H. ben Nasr et al., "Impact of cathepsin D activity and C224T polymorphism (rs17571) on chronic obstructive pulmonary disease: correlations with oxidative and inflammatory markers," *Clinical and Experimental Medicine*, vol. 21, no. 3, pp. 457–465, 2021.
- [14] A. N. Miller, R. Zraick, S. Atmakuri et al., "Characteristics of teach-back as practiced in a university health center, and its association with patient understanding, self-efficacy, and satisfaction," *Patient Education and Counseling*, vol. 104, no. 11, pp. 2700–2705, 2021.
- [15] H. Eloi, "Implementing teach-back during patient discharge education," *Nursing Forum*, vol. 56, no. 3, pp. 766–771, 2021.
- [16] Y. R. Hong, J. Huo, A. Jo, M. Cardel, and A. G. Mainous, "Association of patient-provider teach-back communication with diabetic outcomes: a cohort study," *The Journal of the American Board of Family Medicine*, vol. 33, no. 6, pp. 903–912, 2020.
- [17] M. A. Nas, Y. Cayir, and A. Bilen, "The impact of teach-back educational method on diabetes knowledge level and clinical parameters in type 2 diabetes patients undergoing insulin therapy," *International Journal of Clinical Practice*, vol. 75, no. 4, p. e13921, 2021.
- [18] E. G. Oh, H. J. Lee, Y. L. Yang, and Y. M. Kim, "Effectiveness of discharge education with the teach-back method on 30-day readmission: a systematic review," *Journal of Patient Safety*, vol. 17, no. 4, pp. 305–310, 2021.
- [19] P. Saiklang, R. Puntumetakul, M. Swangnetr Neubert, R. Boucaut, and N. Swangnetr, "The immediate effect of the abdominal drawing-in maneuver technique on stature change in seated sedentary workers with chronic low back pain," *Ergonomics*, vol. 64, no. 1, pp. 55–68, 2021.
- [20] K. S. Lokesh, S. K. Chaya, B. S. Jayaraj et al., "Vitamin D deficiency is associated with chronic obstructive pulmonary disease and exacerbation of COPD," *The Clinical Respiratory Journal*, vol. 15, no. 4, pp. 389–399, 2021.
- [21] M. K. Momeni, S. A. J. Moosavi, B. Zamani et al., "The prevalence of restless legs syndrome in chronic obstructive pulmonary disease (COPD) patients," *Sleep and Biological Rhythms*, vol. 19, no. 2, pp. 201–204, 2021.
- [22] M. Kaminska, K. P. Rimmer, D. Morrison et al., "Long-term non-invasive ventilation in patients with chronic obstructive

- pulmonary disease (COPD): 2021 Canadian Thoracic Society Clinical Practice Guideline update,” *Canadian Journal of Respiratory, Critical Care, and Sleep Medicine*.vol. 5, no. 3, pp. 160–183, 2021.
- [23] G. Hahn, M. H. Cho, E. K. Silverman, C. Lange, S. M. Lutz, and N. Laha, “A fast and efficient smoothing approach to Lasso regression and an application in statistical genetics: polygenic risk scores for chronic obstructive pulmonary disease (COPD),” *Statistics and Computing*, vol. 31, no. 3, pp. 35–3511, 2021.
- [24] S. Peckens, M. M. Adelman, D. King, A. M. Ashcraft, J. Xiang, and B. Sheppard, “Improving chronic obstructive pulmonary disease (COPD) symptoms using a team-based approach,” *The Journal of the American Board of Family Medicine*, vol. 33, no. 6, pp. 978–985, 2020.
- [25] J. Xie and C. Wu, “Ultrasensitive quantification of extracellular vesicles through dual signal amplification for the early diagnosis and prognosis of chronic obstructive pulmonary disease (COPD),” *Analytical Methods*, vol. 13, no. 38, pp. 4437–4441, 2021.
- [26] H. Anees, A. Mohamed Azmi, A. Muhammad Sohail et al., “The economic burden of chronic obstructive pulmonary disease (COPD) in the USA, Europe, and Asia: results from a systematic review of the literature,” *Expert Review of Pharmacoeconomics & Outcomes Research*, vol. 20, no. 6, pp. 661–672, 2020.
- [27] F. Pishgar, M. Shabani, T. Quinaglia A C Silva et al., “Quantitative analysis of adipose depots by using chest CT and associations with all-cause mortality in chronic obstructive pulmonary disease: longitudinal Analysis from MESArthritis ancillary study,” *Radiology*, vol. 299, no. 3, pp. 703–711, 2021.
- [28] M. E. Wilson, C. C. Dobler, A. S. Morrow et al., “Association of home noninvasive positive pressure ventilation with clinical outcomes in chronic obstructive pulmonary disease,” *JAMA: The Journal of the American Medical Association*, vol. 323, no. 5, pp. 455–465, 2020.
- [29] S. A. Alsubheen, A. Wshah, R. Goldstein, and D. Brooks, “Psychometric properties of patient-reported outcome measures assessing self-efficacy in patients with Chronic Obstructive Pulmonary Disease (COPD): a systematic review,” *COPD: Journal of Chronic Obstructive Pulmonary Disease*, vol. 18, no. 2, pp. 254–263, 2021.
- [30] S. Keogan, T. Alonso, S. Sunday et al., “Lung function changes in patients with chronic obstructive pulmonary disease (COPD) and asthma exposed to secondhand smoke in outdoor areas,” *Journal of Asthma*, vol. 58, no. 9, pp. 1169–1175, 2021.
- [31] K. Fujikura, A. Albin, R. G. Barr et al., “Aortic enlargement in chronic obstructive pulmonary disease (COPD) and emphysema: the Multi-Ethnic Study of Atherosclerosis (MESA) COPD study,” *International Journal of Cardiology*, vol. 331, no. 1, pp. 214–220, 2021.
- [32] C. Tang, H. Mistry, N. Bayman et al., “Outcomes of curative-intent radiotherapy in non-small cell lung cancer (NSCLC) patients with chronic obstructive pulmonary disease (COPD) and interstitial lung disease (ILD),” *Radiotherapy and Oncology*, vol. 160, no. 60, pp. 78–81, 2021.
- [33] S. Alshafie, G. O. El-Helw, A. M. Fayoud et al., “Efficacy of dietary nitrate-rich beetroot juice supplementation in patients with chronic obstructive pulmonary disease (COPD): a systematic review and meta-analysis,” *Clinical Nutrition ESPEN*, vol. 42, no. 2, pp. 32–40, 2021.