

Supplementary Online Content

Park DY, Hu JR, Jamil Y, et al. Shorter dual antiplatelet therapy for older adults after percutaneous coronary intervention: a systematic review and network meta-analysis. *JAMA Netw Open*. 2024;7(3):e244000. doi:10.1001/jamanetworkopen.2024.4000

eTable 1. Search Strategy Used in This Study

eTable 2. Baseline Demographics of the Selected Trials

eTable 3. Definition of Outcomes Used in the Trials

eTable 4. Risk of Bias in the Selected Studies Using the Cochrane Risk Assessment Tool

eTable 5. Grading of Recommendations, Assessment, Development and Evaluations (GRADE) Criteria for Each Outcome

eTable 6. Heterogeneity Assessment in Network Meta-Analysis

eTable 7. Node-Splitting Analysis of Inconsistency for Each Outcome

eTable 8. Test of Inconsistency Between Designs

eTable 9. P-Scores of Each Duration of Dual Antiplatelet Therapy

eTable 10. Cumulative Event Rates at Different Durations of Dual Antiplatelet Therapy

eTable 11. Absolute Risk Differences at Shorter Durations of Dual Antiplatelet Therapy

eTable 12. Sensitivity Analysis Only Including Trials That Defined Older Adults as Age ≥ 65 Years

eTable 13. Sensitivity Analysis Only Including Trials That Reported Major Bleeding in Older Adults

eFigure. Node-Splitting Analysis of Inconsistency for Each Outcome

eReferences.

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable 1. Search Strategy Used in This Study

Database	Search Strategy
Cochrane Library	#1 ((duration or length or timing or time or months or reduce* or alterat*) near/3 (dose* or therapy* or treatment* or medication* or DAPT or "dual anti-platelet")):ti,ab OR (de-escalat* or "prescribing practice*" or short-term or abbreviat*):ti,ab OR (duration or length or timing or time or month*):ti #2 ((antiplatelet or anti-platelet) near/3 (dual or combination) near/3 (intervention* or therap*)):ti,ab or ((Aspirin or "Acetylsalicylic acid") and (clopidogrel or plavix or prasugrel or effient or ticagrelor or brilinta or "P2Y12 inhibitor*")):ti,ab or (DAPT):ti,ab #3 ("percutaneous coronary" near/3 (interven* or revascular*)):ti,ab or (PCI):ti,ab or ((balloon or stent*) near/3 angioplast*):ti,ab or (("drug coated" or "drug eluting" or "drug releasing") near/3 stent*):ti,ab #4 #1 and #2 and #3
Ovid Embase	1 exp percutaneous coronary intervention/ 2 exp drug eluting stent/ 3 (percutaneous coronary adj3 (interven* or revascular*)):tw,kf. 4 PCI.ti,ab. 5 ((balloon or stent*) adj3 angioplast*):tw,kf. 6 ((drug coated or drug eluting or drug releasing) adj3 stent*):tw,kf. 7 1 or 2 or 3 or 4 or 5 or 6 8 exp dual antiplatelet therapy/ 9 ((antiplatelet or anti-platelet) adj3 (dual or combination) adj3 (intervention* or therap*)):tw,kf. 10 DAPT.ti,ab. 11 ((Aspirin or Acetylsalicylic acid) and (clopidogrel or plavix or prasugrel or effient or ticagrelor or brilinta or P2Y12 inhibitor*)):tw,kf. 12 8 or 9 or 10 or 11 13 7 and 12 14 exp randomized controlled trial/ 15 exp single blind procedure/ 16 double blind procedure/ 17 crossover procedure/ 18 (random* or factorial* or crossover* or placebo* or assign* or allocat* or volunteer*):tw,kf. 19 (cross adj1 over):tw,kf. 20 ((double or triple or single) adj1 (mask* or blind*)):tw,kf. 21 14 or 15 or 16 or 17 or 18 or 19 or 20 22 13 and 21 23 treatment duration/ 24 ((duration or length or timing or time or months or reduce* or alterat*) adj3 (dose* or therap* or treatment* or medication* or DAPT or dual anti-platelet)):tw,kf. 25 (duration or length or timing or time or month*):ti. 26 (de-escalat* or prescribing practice* or short-term or abbreviat*):tw,kf. 27 prescribing practice/ 28 23 or 24 or 25 or 26 or 27 29 22 and 28
Ovid MEDLINE	1 exp Percutaneous Coronary Intervention/ 2 exp drug eluting stent/ 3 (percutaneous coronary adj3 (interven* or revascular*)):tw,kf.

	4 PCI.ti,ab. 5 ((balloon or stent*) adj3 angioplast*).tw,kf. 6 (((drug coated or drug eluting or drug releasing) adj3 stent*).tw,kf. 7 1 or 2 or 3 or 4 or 5 or 6 8 ((antiplatelet or anti-platelet) adj3 (dual or combination) adj3 (intervention* or therap*)).tw,kf. 9 DAPT.ti,ab. 10 ((Aspirin or Acetylsalicylic acid) and (clopidogrel or plavix or prasugrel or effient or ticagrelor or brilinta or P2Y12 inhibitor*)).tw,kf. 11 8 or 9 or 10 12 7 and 11 5188 13 "duration of therapy"/ 14 ((duration or length or timing or time or months or reduce* or alterat*) adj3 (dose* or therap* or treatment* or medication* or DAPT or dual anti-platelet)).tw,kf. 15 (duration or length or timing or time or month*).ti. 16 (de-escalat* or prescribing practice* or short-term or abbreviat*).tw,kf. 17 13 or 14 or 15 or 16 18 12 and 17 19 exp randomized controlled trial/ 20 single-blind method/ 21 Double-Blind Method/ 22 cross-over studies/ 23 (random* or factorial* or crossover* or placebo* or assign* or allocat* or volunteer*).tw,kf. 24 (cross adj1 over).tw,kf. 25 ((double or triple or single) adj1 (mask* or blind*)).tw,kf. 26 19 or 20 or 21 or 22 or 23 or 24 or 25 27 18 and 26
PubMed	((((random*[Title/Abstract] OR factorial*[Title/Abstract] OR crossover*[Title/Abstract] OR placebo*[Title/Abstract] OR assign*[Title/Abstract] OR allocat*[Title/Abstract] OR volunteer*[Title/Abstract])) OR (double mask*[Title/Abstract] OR double blind*[Title/Abstract] OR single mask*[Title/Abstract] OR single blind*[Title/Abstract] OR triple blind*[Title/Abstract] OR triple mask*[Title/Abstract] OR cross over[Title/Abstract])) OR (randomized controlled trial[MeSH Terms])) AND (((de-escalat*[Title/Abstract] OR prescribing practice*[Title/Abstract] OR short-term[Title/Abstract] OR abbreviat*[Title/Abstract] OR duration[Title/Abstract] OR length[Title/Abstract] OR timing[Title/Abstract] OR time[Title/Abstract] OR month*[Title/Abstract]) AND (((Aspirin[Title/Abstract] OR Acetylsalicylic acid[Title/Abstract]) AND (clopidogrel[Title/Abstract] OR plavix[Title/Abstract] OR prasugrel[Title/Abstract] OR effient[Title/Abstract] OR ticagrelor[Title/Abstract] OR brilinta[Title/Abstract] OR P2Y12 inhibitor*[Title/Abstract])))) OR (DAPT[Title/Abstract] OR dual antiplatelet[Title/Abstract] OR dual anti-platelet[Title/Abstract])) AND (percutaneous coronary interven*[Title/Abstract] OR percutaneous coronary revascular*[Title/Abstract] OR PCI[Title/Abstract] OR balloon angioplast*[Title/Abstract] OR stent* angioplast*[Title/Abstract] OR drug coated stent*[Title/Abstract] OR drug eluting stent*[Title/Abstract] OR drug releasing stent*[Title/Abstract]))
Scopus	(TITLE-ABS-KEY((antiplatelet OR anti-platelet) W/3 (dual OR combination) W/3 (intervention* OR therap*)) OR TITLE-ABS-KEY ((aspirin OR "Acetylsalicylic acid") AND (clopidogrel OR plavix OR prasugrel OR effient OR ticagrelor OR brilinta OR "P2Y12 inhibitor*")) OR TITLE-ABS-KEY (dapt)) AND (TITLE-ABS-KEY ("percutaneous coronary" W/3 (interven* OR revascular*)) OR TITLE-ABS-KEY (pci) OR TITLE-ABS-KEY ((balloon OR stent*) W/3 angioplast*) OR TITLE-ABS-KEY (("drug coated" OR "drug eluting" OR "drug releasing") W/3 stent*)) AND (TITLE-ABS-KEY ((duration OR length OR timing OR time OR months OR reduce* OR alterat*) W/3 (dose* OR therap* OR treatment* OR medication* OR dapt OR "dual anti-platelet")) OR TITLE-ABS-KEY (de-escalat* OR "prescribing practice*" OR short-term OR abbreviat*) OR TITLE (duration OR length OR timing OR time OR month*)) AND (TITLE-ABS-KEY

	(random* OR factorial* OR crossover* OR placebo* OR assign* OR allocat* OR volunteer*) OR TITLE-ABS-KEY (cross W/1 over) OR TITLE-ABS-KEY ((double OR triple OR single) W/1 (mask* OR blind*)))
Web of Science Core Collection	#1 TS=("percutaneous coronary" near/3 (interven* or revascular*)) or TS=(PCI) or TS=((balloon or stent*) near/3 angioplast*) or TS(("drug coated" or "drug eluting" or "drug releasing") near/3 stent*) #2 TS=((antiplatelet or anti-platelet) near/3 (dual or combination) near/3 (intervention* or therap*)) or TS=((Aspirin or "Acetylsalicylic acid") and (clopidogrel or plavix or prasugrel or effient or ticagrelor or brilinta or "P2Y12 inhibitor*")) or TS=(DAPT) #3 TS=((duration or length or timing or time or months or reduce* or alterat*) near/3 (dose* or therap* or treatment* or medication* or DAPT or "dual anti-platelet")) OR TS=(de-escalat* or "prescribing practice*" or short-term or abbreviat*) OR TI=(duration or length or timing or time or month*) #4 TS=(random* or factorial* or crossover* or placebo* or assign* or allocat* or volunteer*) OR TS=(cross near/1 over) OR TS=((double or triple or single) near/1 (mask* or blind*)) #5 #1 and #2 and #3 and #4

eTable 2. Baseline Demographics of the Selected Trials

Abbreviated/Standard ^a , %	HOST-IDEA	MASTER DAPT	TICO	SMART-CHOICE	TWILIGHT	STOPDAPT-2	REDUCE
Age, year, mean	65.6/65.9	76.1/76.0	61.0/61.0	64.6/64.4	65.2/65.1	68.1/69.1	61.0/60.0
Female	27.0/25.2	30.7/30.8	21.0/20.0	27.3/25.8	23.8/23.9	21.1/23.5	17.4/22.7
BMI, kg/m ² , mean	NA	27.3/27.4	24.9/24.9	24..5/24.7	28.6/28.5	24.4/24.2	NA
Diabetes mellitus	40.5/37.4	32.9/34.3	27.0/27.0	38.2/36.8	9.4/10.5	39.0/38.0	21.6/19.5
Hypertension	73.3/73.5	76.9/78.2	50.0/51.0	61.6/61.3	72.6/72.2	73.7/74.0	50.7/50.7
Dyslipidemia	81.2/80.1	67.2/68.1	61.0/60.0	45.1/45.5	60.7/60.2	74.4/74.8	46.3/44.9
Current smoking	NA	10.0/8.1	NA	28.4/24.5	20.4/23.1	26.6/20.6	42.1/42.7
Family history	NA	NA	NA	NA	NA	NA	35.0/36.0
Chronic kidney disease	11.0/10.5	18.2/20.1	19.0/22.0	2.9/3.5	16.8/16.7	5.5/5.6	NA
Peripheral vascular disease	1.9/1.9	NA	NA	NA	6.9/6.8	6.4/6.6	NA
Heart failure	NA	NA	NA	NA	NA	7.7/7.1	NA
Prior myocardial infarction	4.5/4.2	18.9/18.8	4.0/3.0	4.1/4.3	28.7/28.6	13.8/13.2	NA
Prior PCI	13.2/14.1	25.9/26.0	NA	11.5/11.8	42.3/42.0	33.5/35.1	11.7/9.8
Prior CABG		7.4/7.5	1.0/1.0	NA	10.2/9.8	1.1/2.8	2.8/2.8
Prior stroke	6.9/6.3	NA	4.0/4.0	6.6/6.8	NA	NA	1.5/2.0
Prior bleeding	NA	7.2/6.8	NA	NA	0.9/0.9	1.3/1.9	NA
LVEF, %, mean	58.2/58.6	53.5/53.0	NA	60.0/59.9	NA	59.8/59.7	NA
Multivessel disease	51.8/51.7	NA	55.0/56.0	50.1/49.0	63.9/61.6	NA	NA
Clinical presentation							
Silent ischemia	43.4/46.3	10.7/12.0	NA	NA	6.6/6.3	NA	NA
Stable angina		40.2/40.6	NA	41.8/41.8	29.5/28.0	62.3/61.4	NA
Unstable angina	35.7/34.7	11.3/11.4	29.0/32.0	31.2/32.8	35.1/34.9	12.9/14.2	15.2/13.8
NSTEMI	20.9/19.0	25.9/24.4	35.0/32.0	16.0/15.4	28.8/30.8	5.4/6.6	35.6/41.0
STEMI	NA	11.9/11.6	36.0/36.0	11.0/10.0	NA	19.4/17.9	49.3/45.2

eTable 2. Baseline Demographics of the Selected Trials (continued)

Abbreviated/Standard ^a , %	GLOBAL LEADERS	SMART-DATE	IVUS-XPL	ISAR-SAFE	I-LOVE-IT 2	RESET	EXCELLENT
Age, year, mean	64.9/64.8	62.0/62.2	63.0/64.0	67.2/67.2	60.4/60.0	62.4/62.4	63.0/62.4
Female	24.0/23.5	25.1/24.1	33.0/30.0	19.3/19.5	32.8/31.3	35.6/37.1	34.9/36.1
BMI, kg/m ² , mean	NA	24.3/24.5	24.8/24.6	27.2/27.5	25.1/25.3	25.0/24.9	24.9/25.1
Diabetes mellitus	24.3/23.7	26.9/28.1	36.0/37.0	24.8/24.2	23.2/22.1	29.8/28.8	37.7/38.6
Hypertension	72.5/72.3	49.9/48.7	NA	90.1/91.5	61.0/64.8	62.3/61.4	72.7/73.8
Dyslipidemia	63.3/65.3	24.2/25.2	68.0/65.0	87.5/87.4	25.3/23.4	57.7/59.9	75.2/76.3
Current smoking	2.6/2.6	38.0/40.1	25.0/24.0	14.6/15.3	36.6/38.3	25.2/22.8	27.4/25.8
Family history	NA	NA	NA	NA	6.3/5.1	NA	NA
Chronic kidney disease	13.4/13.1	1.0/0.5	NA	NA	NA	NA	0.8/1.2
Peripheral vascular disease	6.7/7.9	NA	NA	NA	1.4/1.1	NA	NA
Heart failure	NA	NA	NA	NA	NA	11.3/11.8	0.6/0.7
Prior myocardial infarction	22.9/23.6	2.3/1.7	5.0/4.0	25.9/24.5	17.2/15.8	1.8/1.6	6.5/3.7
Prior PCI	32.6/33.9	4.9/3.9	10.0/10.0	NA	8.5/6.5	3.5/3.0	9.3/8.6
Prior CABG	NA	NA	3.0/2.0	7.7/7.5	0.4/0.4	0.2/0.6	NA
Prior stroke	NA	3.9/4.4	NA	NA	9.2/9.5	NA	6.5/6.7
Prior bleeding	0.7/0.6	NA	NA	NA	NA	NA	NA
LVEF, %, mean	55.1/55.3	55.5/55.4	62.3/63.1	NA	60.8/60.3	64.2/63.9	61.0/61.6
Multivessel disease	NA	43.6/46.6	NA	NA	NA	43.1/42.9	NA
Clinical presentation							
Silent ischemia	NA	NA	NA	10.9/11.3	3.0/4.0	NA	48.9/48.0
Stable angina	48.9/49.9	NA	51.0/51.0	48.6/47.8	14.3/15.1	44.5/46.3	
Unstable angina	12.9/13.2	31.0/3.7	34.0/33.0	21.5/21.9	58.0/56.5	40.8/39.9	48.5/48.4
NSTEMI	20.0/19.4	31.5/31.4	15/16	10.4/10.1	13.4/13.7	14.7/13.8	
STEMI	18.2/17.5	37.5/37.9		7.9/8.3	11.3/10.7		

^aThe data entry on the left (“abbreviated”) refers to the experimental group in which the duration of dual antiplatelet therapy was shortened. The data entry on the right (“standard”) refers to the control group in which the duration of dual antiplatelet therapy was continued for standard duration according to current guidelines.

Abbreviations: BMI, body mass index; CABG, coronary artery bypass graft; LVEF, left ventricular ejection fraction; NSTEMI, non-ST-elevation myocardial infarction; PCI, percutaneous coronary intervention; STEMI, ST-elevation myocardial infarction

eTable 3. Definition of Outcomes Used in the Trials

Trial	Net Adverse Clinical Events	Major Adverse Cardiovascular Events	Bleeding
HOST-IDEA	Cardiac death, target vessel MI, clinically driven TLR, stent thrombosis, BARC type 3 or 5 bleeding	NA ^a	NA ^a
MASTER DAPT	MACE, BARC type 3 or 5 bleeding	All-cause mortality, MI, stroke	Major or nonmajor clinically relevant bleeding
TICO	MACE, TIMI major bleeding	All-cause mortality, MI, stent thrombosis, stroke, TVR	TIMI major bleeding
SMART-CHOICE	NA ^a	All-cause mortality, MI, stroke	BARC type 2-5 bleeding
TWILIGHT	NA ^a	All-cause mortality, MI, stroke	BARC type 2, 3, or 5 bleeding
STOPDAPT-2	Cardiovascular death, MI, stent thrombosis, stroke, any bleeding	NA ^a	NA ^a
REDUCE	All-cause mortality, MI, stent thrombosis, stroke, TVR, BARC type 2, 3, or 5 bleeding	NA ^a	NA ^a
GLOBAL LEADERS	NA ^a	All-cause mortality, MI, stroke, urgent TVR	BARC type 3 or 5 bleeding
SMART-DATE	NA ^a	All-cause mortality, MI, stroke	NA ^a
IVUS-XPL	Cardiac death, MI, stroke, TIMI major bleeding	Cardiac death, MI, revascularization	TIMI major bleeding
ISAR-SAFE	All-cause mortality, MI, stent thrombosis, stroke, TIMI major bleeding	NA ^a	NA ^a
I-LOVE-IT 2	All-cause mortality, MI, stroke, BARC type 3-5 bleeding	Cardiac death, target vessel MI, clinically indicated TLR	NA ^a
RESET	Cardiac death, MI, stent thrombosis, ischemia-driven TVR, TIMI major bleeding	NA ^a	NA ^a
EXCELLENT	NA ^a	All-cause mortality, MI, stroke, revascularization	NA ^a

^aNot applicable because this outcome was not reported in the older adult population
Abbreviations: BARC, Bleeding Academic Research Consortium; MI, myocardial infarction; TIMI, Thrombolysis In Myocardial Infarction; TLR, target lesion revascularization; TVR, target vessel revascularization

eTable 4. Risk of Bias in the Selected Studies Using the Cochrane Risk Assessment Tool

Trial	Random Sequence Generation (Selection Bias)	Allocation Concealment (Selection Bias)	Blinding of Participants and Personnel (Performance Bias)	Blinding of Outcome Assessment (Detection Bias)	Incomplete Outcome Data (Attrition Bias)	Selective Reporting (Reporting Bias)	Other Bias
HOST-IDEA	Low	Low	High	Low	Low	Low	Unclear
MASTER DAPT	Low	Low	Low	Low	Low	Low	Unclear
TICO	Low	Low	High	Low	Low	Low	Unclear
SMART CHOICE	Low	Low	High	Low	Low	Low	Unclear
TWILIGHT	Low	Low	Low	Low	Low	Low	Unclear
STOPDAPT-2	Low	Unclear	High	Low	Low	Low	Unclear
REDUCE	Low	Low	High	Low	Low	Low	High
GLOBAL LEADERS	Low	Low	Low	Low	Low	Low	Unclear
SMART-DATE	Low	Low	High	Low	Low	Low	Unclear
IVUS-XPL	Low	Low	High	Low	Low	Low	High
ISAR-SAFE	Low	Low	Low	Low	Low	Low	Unclear
I-LOVE-IT 2	Low	Unclear	Unclear	Unclear	Low	Low	High
RESET	Low	Low	High	Unclear	Low	Low	Unclear
EXCELLENT	Low	Low	High	Low	Low	Low	High
Overall Bias	Low	Low	High	Low	Low	Low	Unclear

eTable 5. Grading of Recommendations, Assessment, Development and Evaluations (GRADE) Criteria for Each Outcome

Outcome	Trials	Risk of Bias	Imprecision	Inconsistency	Indirectness	Publication Bias	Certainty
Net adverse clinical events	9	Low	High	Low	Low	Low	⊕⊕⊕○
Major adverse cardiovascular events	9	Low	Moderate	Low	Low	Low	⊕⊕⊕⊕
Bleeding	6	Low	Low	Moderate	Low	Low	⊕⊕⊕⊕

eTable 6. Heterogeneity Assessment in Network Meta-Analysis

Outcome	τ^2	I^2
Net adverse clinical events	0.0782	46.7%
Major adverse cardiovascular events	0.0015	2.6%
Bleeding	0	0%

eTable 7. Node-Splitting Analysis of Inconsistency for Each Outcome

Outcome	Comparison	<i>K</i> ^a	Prop ^b	<i>NMA</i> ^c	Direct ^d	Indirect ^e	<i>Z</i> -value ^f	<i>P</i> -value ^g
Net adverse clinical events	12 months versus 1 month	1	0.42	0.5164	0.8085	0.3077	0.83	0.4084
	3 months versus 1 month	0	0	0.2561	NA	0.2561	NA	NA
	6 months versus 1 month	1	0.75	0.2359	0.1092	0.6099	-0.83	0.4084
	12 months versus 3 months	4	1.00	0.2603	0.2603	NA	NA	NA
	12 months versus 6 months	3	0.84	0.2805	0.1985	0.6993	-0.83	0.4084
	3 months versus 6 months	0	0	0.0202	NA	0.0202	NA	NA
Major adverse cardiovascular events	12 months versus 1 month	1	0.63	0.2408	0.2739	0.1844	0.34	0.7322
	3 months versus 1 month	0	0	0.1756	NA	0.1756	NA	NA
	6 months versus 1 month	1	0.70	0.1236	0.0969	0.1863	-0.34	0.7322
	12 months versus 3 months	3	1.00	0.0652	0.0652	NA	NA	NA
	12 months versus 6 months	4	0.67	0.1172	0.0875	0.1770	-0.34	0.7322
	3 months versus 6 months	0	0	0.0520	NA	0.0520	NA	NA
Bleeding	12 months versus 1 month	1	0.86	0.2462	0.1861	0.6217	-0.68	0.4949
	3 months versus 1 month	0	0	-0.3207	NA	-0.3207	NA	NA
	6 months versus 1 month	1	0.96	0.3807	0.3961	-0.0395	0.68	0.4949
	12 months versus 3 months	3	1.00	0.5669	0.5669	NA	NA	NA
	12 months versus 6 months	1	0.17	-0.1345	0.2256	-0.2100	0.68	0.4949
	3 months versus 6 months	0	0	-0.7014	NA	-0.7014	NA	NA

^aNumber of studies providing direct evidence

^bProportion of direct evidence

^cEstimated treatment effect (risk ratio) in network meta-analysis

^dEstimated treatment effect (risk ratio) derived from direct evidence

^eEstimated treatment effect (risk ratio) derived from indirect evidence

^f*Z*-value of test for disagreement between direct and indirect evidence

^g*P*-value of test for disagreement between direct and indirect evidence

Abbreviations: NMA, network meta-analysis treatment effect; prop, proportion

eTable 8. Test of Inconsistency Between Designs

Outcome	<i>P</i> -value		
	Within Designs	Between Designs	Total
Net adverse clinical events	0.0891	0.1904	0.0806
Major adverse cardiovascular events	0.3032	0.7223	0.4058
Bleeding	0.4737	0.4949	0.5807

eTable 9. P-Scores of Each Duration of Dual Antiplatelet Therapy

Outcome	1 month	3 months	6 months	12 months
Net adverse clinical events	0.8445	0.5373	0.5355	0.0827
Major adverse cardiovascular events	0.8866	0.5323	0.4147	0.1664
Bleeding	0.6553	0.9658	0.2809	0.098

eTable 10. Cumulative Event Rates at Different Durations of Dual Antiplatelet Therapy

Outcome	12 months	6 months	3 months	1 month
Net adverse clinical events	5.40%	6.03%	4.82%	6.40%
Major adverse cardiovascular events	5.76%	5.96%	3.98%	7.19%
Bleeding	5.84%	8.67%	3.55%	6.16%

eTable 11. Absolute Risk Differences at Shorter Durations of Dual Antiplatelet Therapy

	6 months	3 months	1 month
Net adverse clinical events	+0.63%	-0.58%	+1.00%
Major adverse cardiovascular events	+0.20%	-1.78%	+1.43%
Bleeding	+2.83%	-2.29%	+0.32%

eTable 12. Sensitivity Analysis Only Including Trials That Defined Older Adults as Age ≥65 Years

Net adverse clinical events ^a (6 trials included)	1 month			
	NA	3 months		
	NA	0.76 (0.46-1.26)	6 months	
	NA	0.64 (0.46-0.91)	0.85 (0.58-1.24)	12 months
Major adverse cardiovascular events ^a (7 trials included)	1 month			
	NA	3 months		
	NA	1.01 (0.66-1.56)	6 months	
	NA	0.92 (0.69-1.22)	0.91 (0.66-1.26)	12 months
Bleeding ^a (4 trials included)	1 month			
	NA	3 months		
	NA	0.71 (0.22-2.27)	6 months	
	NA	0.57 (0.45-0.71)	0.80 (0.26-2.49)	12 months


^aThis sensitivity analysis included trials that provided results in older adults ≥65 years-old, as well as one trial using a cutoff of >67.2 years (ISAR-SAFE¹) and one trial using a cutoff of >65 years (HOST-IDEA²). The duration of dual antiplatelet therapy at the rightmost column serves as the reference group for the respective column. Values are not available for 1 month of dual antiplatelet therapy because of the absence of trial that set its experimental group as 1 month of dual antiplatelet therapy and defined older adults as age ≥65 years.


eTable 13. Sensitivity Analysis Only Including Trials That Reported Major Bleeding in Older Adults

Major bleeding ^a (3 trials included)	1 month			
	1.74 (0.77-3.94)	3 months		
	1.04 (0.30-3.55)	0.60 (0.16-2.25)	6 months	
	0.83 (0.52-1.32)	0.48 (0.24-0.94)	0.80 (0.26-2.49)	12 months

^aThe duration of dual antiplatelet therapy at the rightmost column serves as the reference group for the respective column.

eFigure. Node-Splitting Analysis of Inconsistency for Each Outcome





Abbreviation: CI, confidence interval; RR, risk ratio

eReferences.

1. Schulz-Schupke S, Byrne RA, Ten Berg JM, et al. ISAR-SAFE: a randomized, double-blind, placebo-controlled trial of 6 vs. 12 months of clopidogrel therapy after drug-eluting stenting. *Eur Heart J*. May 21 2015;36(20):1252-63. doi:10.1093/eurheartj/ehu523
2. Han JK, Hwang D, Yang S, et al. Comparison of 3- to 6-Month Versus 12-Month Dual Antiplatelet Therapy After Coronary Intervention Using the Contemporary Drug-Eluting Stents With Ultrathin Struts: The HOST-IDEA Randomized Clinical Trial. *Circulation*. May 2 2023;147(18):1358-1368. doi:10.1161/CIRCULATIONAHA.123.064264