

# BMJ Open Differential intervention effectiveness of a universal school-based resilience intervention in reducing adolescent substance use within student subgroups: exploratory assessment within a cluster-randomised controlled trial

Rebecca Kate Hodder,<sup>1,2,3</sup> Megan Freund,<sup>2,3</sup> Jenny Bowman,<sup>3,4</sup> Luke Wolfenden,<sup>1,2,3</sup> Elizabeth Campbell,<sup>1,3</sup> Julia Dray,<sup>1,3,4</sup> Christophe Lecathelinais,<sup>1,3</sup> Christopher Oldmeadow,<sup>5</sup> John Attia,<sup>5</sup> John Wiggers<sup>1,2,3</sup>

**To cite:** Hodder RK, Freund M, Bowman J, *et al*. Differential intervention effectiveness of a universal school-based resilience intervention in reducing adolescent substance use within student subgroups: exploratory assessment within a cluster-randomised controlled trial. *BMJ Open* 2018;**8**:e021047. doi:10.1136/bmjopen-2017-021047

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2017-021047>).

Received 7 December 2017  
Revised 8 June 2018  
Accepted 6 July 2018



© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr Rebecca Kate Hodder;  
[rebecca.hodder@hnehealth.nsw.gov.au](mailto:rebecca.hodder@hnehealth.nsw.gov.au)

## ABSTRACT

**Objectives** Interventions addressing the individual and environmental protective factors of adolescents are suggested to have potential for reducing adolescent substance use. While universally delivered school-based substance use prevention interventions are common, previous studies have suggested variable effectiveness by subgroups of students. An exploratory study was undertaken to examine the differential effectiveness of a universal school-based resilience intervention on adolescent substance use and protective factors according to their sociodemographic and previous substance use.

**Design** Secondary analysis of data from a cluster-randomised controlled trial.

**Setting** 32 Australian secondary schools.

**Participants** Cohort of grade 7 students (n=3155) followed up in grade 10 (aged 15–16 years; 2014; n=2105).

**Intervention** Three-year universal school-based intervention implemented by school staff that targeted a range of student resilience protective factors (2012–2014).

**Measurements** Primary outcomes included: tobacco (recent, number of cigarettes) and alcohol (recent, ‘risk’ and number of drinks) use, and secondary outcomes included: marijuana (recent) and other illicit substance (recent) use, and aggregate individual and environmental protective factor scores. Generalised and linear mixed models examined interactions between treatment and student subgroups (gender; socioeconomic disadvantage (low/high); geographic location (major city/inner regional/outer regional-remote); and previous substance use (non-user/user)) at follow-up (36 models).

**Results** Analysis of student follow-up data showed no differential intervention effect for any substance use or protective factor outcome for any subgroup, with the exception of one differential effect found by socioeconomic status for the outcome of mean number of cigarettes smoked by recent smokers (p=0.003). There was no

## Strengths and limitations of this study

- The cluster-randomised controlled study design and the real-world pragmatic intervention delivered by existing school staff was a strength of the study.
- A further strength of the study was the use of best practice statistical methods for subgroup analyses.
- While typical for school-based research, less than 50% of students completed both the baseline and follow-up surveys, which is a limitation of the study.
- The study was limited by its exploratory nature as it was not designed to be powered to detect differences in subgroup analyses.

evidence of an intervention effect within the low (mean difference (MD) –12.89, 95% CI –26.00 to 0.23) or high (MD 16.36, 95% CI –1.03 to 33.76) socioeconomic subgroups.

**Conclusions** No evidence of an intervention effect on substance use and protective factors was found according to student subgroups defined by sociodemographic characteristics or previous substance use.

**Trial registration number** ACTRN12611000606987.

## BACKGROUND

Initiation of tobacco, alcohol and illicit substance use in high-income countries generally occurs during adolescence,<sup>1–3</sup> with earlier use associated with greater dependence in adulthood and a range of negative health outcomes.<sup>4</sup> Despite declining trends in adolescent use of tobacco, alcohol and illicit substances internationally,<sup>3 5</sup> a considerable proportion of adolescents from high-income

countries continue to report such use: 23%<sup>6</sup> and 45%<sup>5</sup> having smoked a cigarette, 43%<sup>6</sup> and 71%<sup>5</sup> having consumed an alcoholic drink and 15%<sup>7</sup> and 40%<sup>6</sup> having used an illicit substance.<sup>3 5 6</sup> Similar levels of adolescent substance use are reported in Australia, with 19%,<sup>3</sup> 68%<sup>7</sup> and 16%<sup>7</sup> of adolescents respectively reporting having ever smoked a cigarette, had an alcoholic drink or used an illicit substance. Additionally within Australia, the prevalence of tobacco, alcohol and illicit substance use is reported to differ by gender,<sup>7</sup> socioeconomic disadvantage<sup>8</sup> and geographic location.<sup>8</sup>

Evidence from cross-sectional and longitudinal studies suggests that a range of individual and environmental factors are associated with a decreased likelihood, and considered protective factors, of adolescent tobacco, alcohol and illicit substance use.<sup>9–22</sup> Individual factors found to be associated with a decreased likelihood of adolescent substance use include self-efficacy, problem solving and self-awareness, whereas environmental factors include caring relationships with adults and peers, and meaningful participation in home, school and community settings.<sup>9–22</sup> These protective factors are also sometimes described as contributing to ‘resilience’.<sup>23–25</sup> Most broadly, resilience has been defined as a ‘process, capacity or outcome of successfully adapting to challenging or threatening circumstances’.<sup>26</sup> In disadvantaged populations in particular, ‘resilience’ or the presence of these protective factors has been found to characterise students with good health and life outcomes despite greater risk status.<sup>26</sup> This is consistent with meta-analysis results from a recent systematic review that reported universal school-based interventions that address individual and environmental resilience protective factors to be effective in reducing illicit substance use by adolescents.<sup>27</sup> Only one controlled trial has examined the effectiveness of an intervention approach focused solely on resilience protective factors in reducing adolescent substance use.<sup>28</sup> While the study reported significant results for tobacco midintervention, no effect on tobacco, alcohol or marijuana use outcomes were evident at follow-up.<sup>29</sup> In contrast, a non-controlled pre-post study of a 3-year universal intervention focused solely on individual and environmental protective factors reported significant reductions in the use of tobacco (50% to 27%), alcohol (34% to 17%) and marijuana (16% to 7%) use among cross-sections of students.<sup>30</sup> Such evidence suggests the potential of school-based resilience interventions that address these individual and environmental protective factors as a means to reducing adolescent substance use.<sup>9 11–22</sup>

School-based substance use prevention interventions that are delivered to all students in a school or classroom regardless of risk; that is, universal interventions,<sup>31 32</sup> are recommended and commonly implemented by governments worldwide.<sup>33–36</sup> It has however been suggested that not all students within a population may benefit equally from universally implemented substance use

prevention interventions, with certain subgroups of students either benefiting more or less than others.<sup>37</sup> As a result, investigation of the generalisability of universal substance use intervention effects across subgroups of students has been recommended.<sup>38</sup> This recommendation is consistent with the standards of evidence for effective programmes and policies developed by the Society for Prevention Research.<sup>39</sup> For interventions found to be effective overall, investigation of the generalisability of effect across subgroups provides guidance for how to enhance intervention effectiveness for all student subgroups. For interventions that have a null effect overall, such investigation can provide guidance to hypothesise whether an intervention may be effective for particular subgroups and identify opportunities for future studies to test such hypotheses.

The only universally implemented school-based intervention focused solely on targeting resilience protective factors has not investigated any variable patterns of effect by participant subgroups. However, of those studies that have implemented school-based interventions that address resilience protective factors among other factors as part of a broader intervention approach, a variable pattern of effect by participant subgroup has been reported. Such variability has been reported to occur between students defined by sociodemographic and previous substance use characteristics.<sup>40–43</sup> For example, studies have reported differential intervention effects on tobacco use by gender, such as reductions in tobacco use for either females<sup>40</sup> or males,<sup>42 43</sup> whereas other studies have reported differential effects by socioeconomic level, such as reductions in alcohol use for students of low socioeconomic schools but no effect in schools of medium or high socioeconomic level.<sup>41</sup> No universally delivered school-based studies addressing protective factors could be found that examined differential intervention effects by subgroups of students defined by geographic location. Previous studies have also examined differential effectiveness of school-based substance use interventions in terms of students classified by risk of substance use, most often defined as substance use initiation prior to intervention. Such studies report mixed results,<sup>41</sup> suggesting such interventions are more effective for existing substance users than non-users, more effective for existing non-users than users or no differential effect according to previous substance use.<sup>41 44–46</sup>

A cluster-randomised control study was undertaken to investigate the overall effectiveness of a universally delivered school-based resilience intervention in reducing substance use by adolescents (20 intervention; 12 control schools).<sup>47</sup> As previously reported, the study found no effect on the primary (tobacco and alcohol use; mental health problems) or secondary (illicit substance use, individual and environmental resilience protective factors) student outcomes. The study also investigated the effect of the intervention on the implementation of resilience intervention strategies in

both intervention and control schools, and of the 16 outcomes related to the implementation of resilience strategies, intervention schools were more likely than control schools to implement 9 hours of resilience curriculum.<sup>48 49</sup> In order to investigate whether any student subgroups benefited from the intervention, a study was conducted to examine the differential effectiveness of the universally delivered school-based ‘resilience’ protective factor intervention on substance use by adolescents according to their sociodemographic and previous substance use characteristics. A secondary aim was to examine the differential effectiveness of the intervention on the hypothesised mechanism of effect, student resilience protective factors.

## METHODS

### Study design and setting

The cluster-randomised controlled trial was conducted in secondary schools in one health district of New South Wales, Australia. Outcome assessments were conducted with a cohort of students at baseline in 2011 (when students were in grade 7: aged 12–13 years) and at follow-up in 2014 (when students were in grade 10).

Approximately 114 000 people aged 10–19 years reside in metropolitan, regional, rural and remote areas within the district.<sup>50 51</sup> Further study details have been reported elsewhere.<sup>47</sup>

### Participants and recruitment

#### Schools

A national schools database<sup>52</sup> identified 172 schools with secondary enrolments within the study area. Schools were eligible if they: were a government or Catholic secondary school located within a socioeconomically disadvantaged local government area (defined by the Socio-Economic Indexes or Areas (SEIFA) Index of Relative Socio-economic Advantage/Disadvantage),<sup>53</sup> had enrolments in grades 7–10 (aged 12–16 years) and had more than 400 total student enrolments. Schools were ineligible if they were: single gender, independent (private), special needs, selective, central (for students aged 5–18 years) or boarding schools.

#### Randomisation of schools

Eligible schools were randomly ordered using a random number function in Excel and approached in that order until a quota of 32 schools consented. The 32 consenting schools were then stratified according to participation in a government disadvantaged schools initiative (yes/no)<sup>54</sup> and school size (medium 400–800; large >800). Schools were then randomly allocated to intervention or control in a 20:12 block design ratio (based on stakeholder request to increase the reach of anticipated intervention benefit) by an independent statistician using a random number function in Microsoft Excel prior to baseline data collection.

### Students

All students enrolled in grade 7 (first year at secondary school) were eligible to participate (n=4589), and active parental consent for student participation in data collection was sought via a mailed study information pack. After 2 weeks, non-responding parents were prompted via telephone by school-affiliated staff who were blind to group allocation. A toll-free number was provided for parents who wished to decline the telephone prompt.

### Intervention

A 3-year universal (‘whole of school’) intervention was delivered by school staff to all students in the cohort during grades 8–10. The intervention, based on a pilot study,<sup>30</sup> involved 16 broad strategy areas (see [box 1](#)) seeking to build the protective factors of students implemented across all three domains of the Health Promoting Schools framework<sup>55</sup> ([box 1](#)). Each broad intervention strategy addressed at least one individual (self-efficacy, problem solving, cooperation/communication, self-awareness, empathy and goals/aspirations) or environmental protective factor (school support, school meaningful participation, community support, community meaningful participation, home support, home meaningful participation, peer-caring relationships and prosocial peers). Such protective factors align with a ‘resilience’ approach.<sup>23–25 56</sup> Schools were provided with details of existing available resources and programmes targeting the protective factors identified by researchers. While schools were required to implement all strategies, they were given the flexibility to select the order in which they were implemented and which resources or programmes they used when doing so.

To ensure implementation of intervention strategies, schools were provided with a comprehensive range of support strategies, including an embedded implementation support officer, and other strategies that have been previously reported to facilitate implementation of interventions ([box 1](#)).<sup>57–64</sup>

Control schools implemented usual school curricula that may have included protective factor strategies and resources similar to or the same as those systematically provided to the intervention schools and were not provided with programme resources or implementation support. A report describing school-level student substance use and protective factor characteristics at baseline was provided to control schools.

### Data collection procedures

#### Student demographic and protective factor characteristics and substance use

Students completed a confidential web-based survey in class time prior to intervention commencement (baseline: August–November 2011) and immediately following intervention completion (follow-up: July–November 2014). Neither the school staff nor researchers were blind to group allocation.

## Box 1 Intervention and implementation support strategies

### Intervention strategies by Health Promoting Schools domain

#### Curriculum, teaching and learning

1. Age-appropriate lessons (9 hours) on individual protective factors across school subjects (eg, MindMatters<sup>78</sup> or school-developed curriculum resources).<sup>l</sup>
2. Non-curriculum programmes (9 hours) targeting protective factors (eg, the Resourceful Adolescent Program).<sup>79l, E</sup>
3. Additional programme targeting protective factors for Aboriginal students.<sup>\*l, E</sup>

#### Ethos and environment

4. Rewards and recognition programme.<sup>l, E</sup>
5. Peer support/peer mentoring programmes.<sup>l, E</sup>
6. Antibullying programmes.<sup>l, E</sup>
7. Empowerment/leadership programmes.<sup>l</sup>
8. Additional empowerment/leadership/mentoring programmes for Aboriginal students.<sup>\*l, E</sup>
9. Aboriginal cultural awareness strategies.<sup>l, E</sup>

#### Partnerships and services

10. Promotion/engagement of local community organisations/groups/clubs in school (eg, charity organisations).<sup>#</sup>
11. Additional/enhanced consultation activities with Aboriginal community groups.<sup>\*l, E</sup>
12. Promotion/engagement of health, community and youth services in the school.<sup>l, E</sup>
13. Additional/enhanced Aboriginal community organisations promoted or engaged.<sup>\*l, E</sup>
14. Referral pathways to health, community and youth services developed and promoted.<sup>l, E</sup>
15. Strategies to increase parental involvement in school (eg, school events).<sup>E</sup>
16. Information regarding student protective factors provided to parents via school newsletter.<sup>E</sup>

#### Implementation support strategies

1. Engagement with school community including presentations at school staff meetings regarding planned intervention.<sup>†</sup>
2. Embedded staff support:
  - School intervention officer 1 day a week to support programme implementation.
  - Project coordinator to liaise with school sectors and support school intervention officers.<sup>‡</sup>
3. School intervention team formed (new team or realignment of existing team, inclusive of school intervention officer and school executive member) to implement intervention.
4. Structured planning process to prioritise and select appropriate resources/programmes:
  - Needs assessment of student protective factors (when study sample in grade 7).
  - Two school community planning workshops and one strategy review workshop.<sup>‡</sup>
  - School plan to address intervention strategies endorsed by the school executive.
5. Intervention implementation guide that described the intervention, planning process, available resources and programmes, tools and templates for intervention implementation.
6. Staff mental health training (minimum of 1 hour per school during staff meetings)

Continued

## Box 1 Continued

7. \$A2000 per year each for:
  - Teacher release time for intervention implementation or professional development.
  - Strategies specifically for Aboriginal students.\*
8. Feedback reports regarding student substance use and protective factors, and intervention implementation (termly).<sup>‡</sup>
9. An Aboriginal Cultural Steering group with representation from local Aboriginal community organisations was formed to provide Aboriginal cultural advice and direction regarding the study design, implementation, evaluation and dissemination.

Following publication of the study protocol<sup>47</sup> and based on advice received from an Aboriginal Cultural Steering Group intervention strategies 3, 8, 11 and 13 were added.

\*Implemented in years 2 and 3 only; †year 1 only; ‡years 1 and 2 only.

<sup>l</sup>To target individual protective factors; <sup>E</sup>to target environmental protective factors.

## Measures

### Student demographic characteristics

The student survey addressed: age, gender, residential postcode, Aboriginal and/or Torres Strait Islander status, ethnicity and non-English speaking background.

### Student substance use characteristics

Substance use data were collected using items from an ongoing Australian triennial survey of school students' health behaviours.<sup>3</sup> Outcomes included recent tobacco, recent alcohol, 'risk' alcohol, recent marijuana and other illicit substance use as well as the number of cigarettes and alcoholic drinks consumed in the last week (table 1).

### Student individual and environmental protective factors

The Resilience and Youth Development module of the California Healthy Kids Survey was used to measure individual and environmental protective factors.<sup>65</sup> Items for six individual and three environmental factor subscales (table 1) were selected. Aggregate individual and environmental protective factor scores were used as outcome measures. Consistent with a previous study of the survey tool,<sup>65</sup> analysis of baseline responses showed the subscales were reasonably internally consistent and valid (Cronbach alpha coefficients: individual 0.55–0.81; environmental 0.77–0.88). Confirmatory factor analysis<sup>65</sup> demonstrated the subscale factor structure to be a good model fit ( $X^2(1133)=6573$ ,  $p<0.0001$ ; comparative fit index 0.92, root mean square error of approximation 0.04).

## Statistical analysis

### Student sociodemographic subgroups

Student-reported residential postcode was used to calculate student socioeconomic status<sup>53</sup> and remoteness of residential location.<sup>66</sup> Students were classified into the following subgroups based on their baseline survey characteristics: gender (males, females), socioeconomic status (as defined by SEIFA Index of Relative Socio-economic advantage/disadvantage; low: scores of <990 (most disadvantaged), high: scores  $\geq 990$ <sup>53</sup> and geographic location



**Table 1** Substance use and protective factor outcome measures at follow-up

	Survey item	Response options
<b>Primary outcomes</b>		
Tobacco use: recent	Have you smoked a cigarette in the last week?	Yes/no
Number cigarettes: last week*	If yes, starting from yesterday, please record the number of cigarettes that you smoked on each day of last week. <sup>3</sup>	0–99
Alcohol use: recent	Have you had any alcoholic drinks, such as beer, wine or alcopops/premix drinks in the last week? (do not count sips or tastes)	Yes/no
Number alcoholic drinks: last week*	If yes, starting from yesterday, please record the number of alcoholic drinks that you had on each day of last week. <sup>3</sup>	0–99
Alcohol use: ‘risk’	In the last 4 weeks, how many times have you had five or more alcoholic drinks in a row? <sup>3</sup>	None/once/twice/3–6 times/7 or more times
<b>Secondary outcomes</b>		
Marijuana use	How many times in the last 4 weeks have you smoked or used marijuana/cannabis (grass, hash, dope, weed, mull, yarndi, ganga, pot, a bong and a joint)? <sup>3</sup>	None/once or twice/3–5 times/6–9 times/10–19 times/20–39 times/40 or more times
Other illicit substance use	How many times in the last 4 weeks have you used any other illegal drug or pill to get ‘high’, such as inhalants, hallucinogens (eg, LSD, acid, trips), amphetamines (eg, speed, ice), ecstasy, cocaine or heroin?	None/once or twice/3–5 times/6–9 times/10–19 times/20–39 times/40 or more times
Individual protective factors <sup>65</sup>	Cooperation and communication subscale: two items; for example, ‘I enjoy working together with other students my age’.	1: Never true, 2: true some of the time; 3: true most of the time; 4: true all of the time
	Self-efficacy subscale: four items; for example, ‘I can do most things if I try’.	As above
	Empathy subscale: three items; for example, ‘I try to understand what other people feel and think’.	As above
	Problem solving subscale: three items; for example, ‘When I need help I find someone to talk with’.	As above
	Self-awareness subscale: three items; for example, ‘I understand why I do what I do’.	As above
	Goals and aspirations subscale: three items; for example, ‘I have goals and plans for the future’.	As above
Environmental protective factors <sup>65</sup>	School support subscale: six items; for example, ‘At my school there is an adult who really cares about me’.	As above
	School meaningful participation subscale: three items; for example, ‘At my school, I help decide things like class activities or rules’.	As above
	Peer caring relationships subscale: three items; for example, ‘I have a friend who helps me when I’m having a hard time’.	As above

\*At baseline, students were asked whether they had ever smoked a cigarette/consumed an alcoholic drink.

(as defined by the Accessibility/Remoteness Index of Australia; major city: 0–0.2, inner regional: >0.2–2.4, outer regional/remote: >2.4–15).

#### Previous substance use subgroups

Ever use of tobacco was used to define baseline tobacco use, and ever use of alcohol was used to define baseline alcohol use (user and non-user). Use of marijuana in the last 4 weeks was used to define baseline marijuana use, and use of other illicit substances in the last 4 weeks was used to define baseline other illicit substance use (user and non-user). Baseline use of any substance was defined as use of at least one

substance derived from baseline use of tobacco, alcohol, marijuana or illicit substances (user and non-user).

#### Primary outcomes: student alcohol and tobacco use

Recent tobacco use was defined as having smoked at least one cigarette in the last week, and recent alcohol use was defined as at least one alcoholic drink in the last week (yes/no). An average number of cigarettes and alcoholic drinks consumed in the last week was calculated from the responses for daily consumption. The response options for ‘risk alcohol use’ were dichotomised as ‘none’ or ‘use’ (‘once’/‘twice’/‘3–6 times’/‘7 or more times’).

### Secondary outcomes: student illicit substance use

The response options for both marijuana and other illicit substance use were dichotomised as 'none' or 'use' ('once or twice'/'3–5 times'/'6–9 times'/'10–19 times'/'20–39 times'/'40 or more times').

### Secondary outcomes: student individual and environmental protective factor scores

Student protective factor subscale scores were calculated by averaging the responses to all items in each subscale. Aggregate individual and environmental protective factor scores were calculated by averaging all relevant subscale scores for each student.<sup>65</sup> Mean scores ranged from 1 to 4, with higher scores more favourable.

### Subgroup analyses

Best practice principles for subgroup analysis specify that such analyses should be: exploratory; limited to primary outcomes with a small number of predefined subgroups; analysis by formal statistical tests of interaction; and analysis within subgroups conducted only if an interaction is statistically significant.<sup>67 68</sup> As such, comparisons between treatment groups for each dichotomous (five outcomes) and continuous (four outcomes) outcome at follow-up for the cohort grade 10 students in intervention and control schools by each of the four subgroups was undertaken to determine the effectiveness of the intervention using generalised linear mixed models (binomial distribution with a logit link) (20 models) and linear mixed models (16 models), respectively. All models included a fixed effect for treatment group (intervention vs control), a random effect for each school to account for clustering of responses within schools and an interaction term (treatment × subgroup) to determine differential intervention effect. ORs with 95% Wald CIs were calculated for each subgroup category. Where an interaction term was significant, comparisons between treatment groups within each of the subgroups was undertaken using the same modelling approach to determine the effectiveness of the intervention within each individual subgroup.

A criterion for statistical significance of  $p \leq 0.01$  was used due to multiple testing.<sup>69</sup> All analyses were undertaken using SAS Software V.9.4.<sup>70</sup>

### Patient and public involvement

School staff from intervention schools were involved in implementing and selecting which resources and strategies to implement as part of the intervention. The deidentified and aggregate overall results of the study were disseminated to the principals of participating schools at the conclusion of the study.

## RESULTS

### Sample

#### Schools

Forty-four of the 47 eligible schools were approached prior to obtaining the quota of 32 participating schools (73%

consent rate). Of those 28 were government and four Catholic schools, and 21 were medium and 11 were large-sized schools (see [figure 1](#)). No schools withdrew following allocation.

### Students

At baseline, parental consent was provided for 3530 grade seven students (76.9% of enrolled students), of which 3115 students participated in the baseline survey (67.9% of enrolled students; 88.2% of students with parental consent). Follow-up data were collected from 2149 of the students who completed the baseline survey (retention rate 69.0%; intervention 67.3%, control 71.6%; 46.8% of students enrolled at baseline). There was no differential loss to follow-up overall between intervention and control groups ( $p=0.1$ ). Students who moved between schools ( $n=30$ ) and those who participated but did not answer substance use items at baseline ( $n=14$ ) were excluded resulting in a cohort of 2105 students for the primary analysis. The demographic characteristics of students who completed the baseline and follow-up survey are shown in [table 2](#).

### Substance use


[Table 2](#) shows the characteristics of the subgroups at follow-up. Online supplementary appendix A shows the proportion of, or mean score for, students in each subgroup reporting each outcome at baseline and follow-up, respectively. There was no difference between intervention and control students for any measure of substance use at follow-up for students overall ([table 3](#)). Similarly, there was no differential effect for any of the seven substance use outcomes between subgroups defined by gender, geographic location or previous substance use ([table 3](#)). A differential effect was found for one of the seven outcomes for subgroups defined by socioeconomic status; there was a significant interaction ( $p=0.003$ ) between treatment and socioeconomic status on the mean number of cigarettes smoked (by students who were recent smokers). The CIs for the estimates within the both low (mean difference (MD)  $-12.89$ , 95% CI  $-26.00$  to  $0.23$ ) and high (MD  $16.36$ , 95% CI  $-1.03$  to  $33.76$ ) socioeconomic subgroups included the null value ([table 3](#)).

### Student individual and environmental protective factors

At follow-up, there was no difference in mean individual or environmental protective factor scores between intervention and control students ([table 3](#)). Similarly, there was no differential effect for any of the protective factor outcomes by any subgroup ([table 3](#)).

## DISCUSSION

This study sought to examine the differential effectiveness of a universally delivered school-based 'resilience' protective factor intervention on substance use by adolescents according to their baseline sociodemographic characteristics and previous substance use. The study found negligible evidence (1 of 36 tests) of differential intervention



**Figure 1** Study flow diagram.

effectiveness between student subgroups. A differential intervention effect was found for number of cigarettes smoked by recent smokers, by socioeconomic status which, on further examination within subgroups, showed no significant difference in the consumption of cigarettes in intervention compared with control students residing within either subgroup. These findings, coupled with the primary study finding of no intervention effect on any measure of adolescent substance use or protective factors for the overall trial sample,<sup>49</sup> suggests that a pragmatic approach to the universal delivery of a resilience-focused substance use intervention was not effective at any level.

The findings of the study with respect to subgroups defined by sociodemographic characteristics are generally

consistent with previous universal school-based studies that targeted protective factors as part of a broader intervention approach. With respect to subgroups defined by gender, previous studies have reported equivocal support for such interventions being effective for alcohol or illicit substance use for males only,<sup>42</sup> for females only<sup>71</sup> or effective for both males and females<sup>40</sup> in comparison with the current study that found no differential effect. Only one previous study was identified that investigated differential effectiveness of such an intervention approach by socioeconomic status, which reported an intervention effect for drunkenness for students of low but not high socioeconomic status.<sup>41</sup> No previous studies were identified that investigated differential intervention effectiveness by

**Table 2** Demographic and substance use characteristics by group at follow-up (n=2105)

Student characteristics	Intervention n (%)	Control n (%)	All students
Total students	1261	844	2105
Age, M (SD)	15.5 (0.5)	15.5 (0.5)	15.5 (0.5)
Aboriginal/Torres Strait Islander	128 (10.2)	95 (11.3)	223 (10.6)
Gender			
Male	640 (50.8)	431 (51.1)	1071 (50.9)
Female	621 (49.3)	413 (48.9)	1034 (49.1)
Remoteness*			
Major cities	525 (41.7)	508 (60.2)	1033 (49.1)
Inner regional	612 (48.6)	262 (31.0)	874 (41.5)
Outer regional/ remote	123 (9.8)	74 (8.8)	197 (9.4)
Socioeconomic status (SES)*			
Low (<990) (most disadvantaged)	704 (55.9)	534 (63.3)	1238 (58.8)
High (≥990)	556 (44.1)	310 (36.7)	866 (41.2)
Previous substance user			
Tobacco user†	106 (8.4)	64 (7.6)	170 (8.1)
Alcohol user†	364 (28.9)	200 (23.8)	564 (26.8)
Marijuana user‡	14 (1.1)	7 (0.8)	21 (1.0)
Other illicit user‡	8 (0.6)	4 (0.5)	12 (0.6)
Any substance user§	387 (30.7)	215 (25.5)	602 (28.6)

\*SES and remoteness could not be calculated for five students as postcode was missing (four intervention, one control).

†Ever use of tobacco and alcohol were used to define baseline tobacco and alcohol use, respectively.

‡Use of marijuana and other illicit substances in the last 4 weeks was used to define baseline marijuana and other illicit substance use, respectively.

§Baseline use of any substance was defined as use of at least one substance derived from baseline use of tobacco, alcohol, marijuana or illicit substances. This was the variable used for baseline use for protective factor outcomes.

geographic location for universal school-based protective factor interventions.

The findings of previous studies of intervention effectiveness by subgroups defined by previous substance use are equivocal. For example, previous studies have found no evidence of differential intervention effect between subgroups defined by previous substance use for both tobacco<sup>45</sup> and alcohol.<sup>45 46</sup> In contrast, other studies have found results favouring students who have never used a substance<sup>44</sup> and those who are previous substance users.<sup>46</sup>

The absence of an intervention effect for the hypothesised mechanism of intervention effect, resilience protective factors, for both the overall trial sample<sup>49</sup> and across all subgroups assessed in this study precludes the ability to determine whether the enhancement of protective factors can lead to a reduction in adolescent substance use. However, the singular significant result in this

study regarding the consumption of tobacco for recent smokers by socioeconomic status suggests that despite not impacting on protective factor levels, the intervention may have a differential effect on socioeconomically disadvantaged compared with advantaged students. Such results may suggest that an alternative mechanism may be responsible for the decrease in tobacco consumption. A more likely explanation for the singular result for 1 of 36 tests undertaken is that the result was false positive due to the number of tests that were conducted.

Various aspects of the intervention design may have contributed to the null finding for resilience protective factors both overall and within the student subgroups and have been reported previously.<sup>49</sup> First, the universal delivery of the intervention without any differentiation or targeted intervention for students with lower protective factor scores or students at greater risk within particular student subgroups may have limited its ability to impact across all students. Second, the use of a pragmatic intervention approach involving school staff selection and implementation of existing readily available curriculum resources and programmes and flexibility regarding the order in which selected resources were implemented may have contributed to the null study findings as pragmatic intervention approaches have been reported to be less likely to be effective than non-pragmatic approaches.<sup>72 73</sup>

As previously reported, data regarding similar levels of implementation for 15 of 16 targeted strategies in both intervention and control groups suggests the use of programmes and resources that were also accessible to control schools may also have led to a lack of differential intervention exposure between group<sup>49</sup> or may be an indication of cross-site contamination. Third, the duration of the intervention primarily over 2 years may have resulted in insufficient exposure to impact on student protective factors. Finally, the intervention's focus on protective factors only, with no content addressing known risk factors of substance use (such as peer substance use<sup>74</sup>), and a limited focus on family and community-based protective factors of substance use<sup>75</sup> may have impacted on the findings.

Strengths of the overall study included the cluster-randomised controlled study design, the use of implementation support strategies and it being a real-world pragmatic effectiveness trial. This study adhered to criteria for a best practice analytical approach to subgroup analyses<sup>67</sup> with the exception that secondary outcomes (ie, resilience protective factor outcomes) were examined in subgroups, and the proposed subgroup analyses were not published a priori. In terms of further limitations, the number of comparisons undertaken within this study may have increased the risk of type 1 error and led to the sole significant result; however, the use of a formal statistical interaction test, rather than examination of intervention effect within individual subgroups, reduces this risk.<sup>76</sup> In the overall study, the proportion of enrolled students completing both the baseline and follow-up surveys was below 50%, and while typical for school-based research,<sup>77</sup>



**Table 3** Unadjusted intervention versus control group substance use comparisons at follow-up by all students and subgroups\*

Subgroups	Tobacco			Alcohol			Illicit substances			Protective factors			Environmental Mean difference (95% CI)
	Recent OR (95% CI)	Amount Mean difference (95% CI)†	Recent OR (95% CI)	'Risk' OR (95% CI)	Amount Mean difference (95% CI)‡	Marijuana OR (95% CI)	Other substance OR (95% CI)	Individual Mean difference (95% CI)					
<b>All students</b>													
Unadjusted	1.37 (0.85 to 2.22)	-2.30 (-11.94 to 7.69)	1.10 (0.77 to 1.57)	0.97 (0.71 to 1.33)	-0.77 (-2.80 to 1.25)	1.12 (0.74 to 1.68)	1.19 (0.67 to 2.10)	0	0	0	0	0 (-0.10 to 0.10)	
Adjusted§	1.48 (0.93 to 2.37)	-0.98 (-11.15 to 9.20)	1.10 (0.77 to 1.56)	1.03 (0.74 to 1.43)	-0.51 (-2.61 to 1.59)	1.18 (0.80 to 1.72)	1.42 (0.85 to 2.38)	-0.01 (-0.07 to 0.06)	-0.01 (-0.07 to 0.06)	-0.02 (-0.09 to 0.06)	-0.02 (-0.09 to 0.06)	-0.02 (-0.09 to 0.06)	
<b>Gender</b>													
Males	1.37 (0.77 to 2.42)	-4.56 (-19.35 to 10.23)	1.09 (0.72 to 1.66)	0.80 (0.55 to 1.16)	-1.40 (-4.29 to 1.49)	1.04 (0.65 to 1.67)	0.96 (0.51 to 1.82)	0.01 (-0.07 to 0.10)	0.01 (-0.07 to 0.10)	0.02 (-0.09 to 0.13)	0.02 (-0.09 to 0.13)	0.02 (-0.09 to 0.13)	
Females	1.38 (0.78 to 2.43)	-0.33 (-13.69 to 13.03)	1.12 (0.72 to 1.73)	1.21 (0.82 to 1.79)	0 (-3.01 to 3.00)	1.23 (0.74 to 2.04)	1.77 (0.80 to 3.92)	-0.02 (-0.11 to 0.07)	-0.02 (-0.11 to 0.07)	-0.02 (-0.13 to 0.10)	-0.02 (-0.13 to 0.10)	-0.02 (-0.13 to 0.10)	
<b>Remoteness</b>													
Major city	1.93 (0.91 to 4.07)	7.04 (-15.34 to 29.42)	1.30 (0.76 to 2.22)	1.03 (0.64 to 1.67)	-0.87 (-4.57 to 2.83)	1.49 (0.79 to 2.79)	2.09 (0.96 to 4.57)	-0.07 (-0.18 to 0.04)	-0.07 (-0.18 to 0.04)	-0.13 (-0.26 to 0.00)	-0.13 (-0.26 to 0.00)	-0.13 (-0.26 to 0.00)	
Inner regional	1.36 (0.58 to 3.21)	-7.17 (-32.99 to 18.66)	1.08 (0.59 to 1.97)	0.92 (0.53 to 1.62)	-0.78 (-4.96 to 3.39)	0.97 (0.48 to 1.99)	0.75 (0.30 to 1.87)	0	0	0.10 (-0.05 to 0.25)	0.10 (-0.05 to 0.25)	0.10 (-0.05 to 0.25)	
Outer regional/ remote	0.60 (0.18 to 2.07)	-17.91 (-56.18 to 20.36)	0.48 (0.17 to 1.33)	0.98 (0.40 to 2.39)	0.77 (-7.28 to 8.82)	0.52 (0.16 to 1.71)	0.36 (0.04 to 3.18)	0.23 (0.02 to 0.43)	0.23 (0.02 to 0.43)	0.15 (-0.11 to 0.40)	0.15 (-0.11 to 0.40)	0.15 (-0.11 to 0.40)	
<b>Socioeconomic status</b>													
Low	1.40 (0.79 to 2.48)	-12.89** (-26.00 to 0.23)	1.06 (0.70 to 1.61)	1.06 (0.72 to 1.55)	-0.91 (-3.65 to 1.84)	1.30 (0.79 to 2.15)	0.47 (0.72 to 3.01)	0.00 (-0.09 to 0.09)	0.00 (-0.09 to 0.09)	0.01 (-0.10 to 0.12)	0.01 (-0.10 to 0.12)	0.01 (-0.10 to 0.12)	
High	1.35 (0.65 to 2.81)	16.36 (-1.03 to 33.76)	1.21 (0.71 to 2.07)	0.84 (0.52 to 1.35)	-0.59 (-4.20 to 3.02)	0.85 (0.46 to 1.56)	0.82 (0.34 to 1.94)	-0.01 (-0.12 to 0.10)	-0.01 (-0.12 to 0.10)	-0.02 (-0.15 to 0.12)	-0.02 (-0.15 to 0.12)	-0.02 (-0.15 to 0.12)	
<b>Baseline use</b>													
Non-user	1.39 (0.83 to 2.31)	-5.16 (-16.55 to 6.23)	1.22 (0.76 to 1.95)	0.99 (0.69 to 1.41)	-1.05 (-3.76 to 1.67)	1.37 (0.13 to 14.23)	-¶	0.01 (-0.07 to 0.09)	0.01 (-0.07 to 0.09)	-0.01 (-0.11 to 0.09)	-0.01 (-0.11 to 0.09)	-0.01 (-0.11 to 0.09)	
User	1.34 (0.60 to 2.97)	6.83 (-11.76 to 25.43)	0.94 (0.63 to 1.42)	0.79 (0.52 to 1.21)	-0.95 (-4.1 to 2.20)	1.09 (0.69 to 1.72)	1.15 (0.58 to 2.27)	0.05 (-0.10 to 0.10)	0.05 (-0.10 to 0.10)	0.05 (-0.08 to 0.18)	0.05 (-0.08 to 0.18)	0.05 (-0.08 to 0.18)	

\*Estimates presented originate from models inclusive of interaction term (treatment × subgroup). Sample sizes for each comparison ranged from 12 to 2105 and are detailed in online supplementary appendix B.

†Of students who reported tobacco use in the last week.

‡Of students who reported alcohol use in the last week.

§Models were adjusted for a priori selected prognostic variables (age, gender, school type, school size, Aboriginal/Torres Strait Islander status, ethnicity, non-English-speaking background and socioeconomic status).

¶OR estimated as infinity due to small sample size and zero cell counts.

\*\*interaction term (treatment × subgroup) p<0.01.

it may limit the generalisability of the study results. Additionally, despite the scale of the overall study in 32 secondary schools, it was not designed to be powered to detect differences in outcomes within participant subgroups. As a result, any actual differences in intervention effectiveness by subgroup may not have been detectable. The small sample size of the exposure groups in some of the subgroup analyses, such as examination of differences in the number of cigarettes smoked, further limited the statistical power to detect differences between groups. As a result, the subgroup analyses are considered exploratory in order to generate hypotheses for future research and should be interpreted with caution. While the effect of the intervention on illicit substance use among students with low socioeconomic status was double that of students with high socioeconomic status appears promising, it is unclear whether such a result would be substantiated in a study sufficiently powered to detect difference in outcomes within participants subgroups. Finally, the study was limited by its focus on adolescent substance use and resilience outcomes, and the impact of the intervention on other student outcomes, such as alcohol-related harm, or implementation outcomes, such as assessment of teacher impact on intervention implementation, was not assessed.

## CONCLUSION

This exploratory study found negligible evidence of an intervention effect on adolescent substance use and protective factors outcomes according to student socio-demographic or substance use characteristics following implementation of a universal school-based resilience intervention. While there was some evidence of a differential intervention effect on tobacco use by smokers in subgroup analysis by socioeconomic status, which suggests there may be some potential for a resilience intervention in reducing tobacco use within this group of students, the result should be interpreted with caution and further sufficiently powered research conducted to confirm this.

### Author affiliations

<sup>1</sup>Hunter New England Population Health, Hunter New England Local Health District, Wallsend, New South Wales, Australia

<sup>2</sup>School of Medicine and Public Health, The University of Newcastle, Callaghan, New South Wales, Australia

<sup>3</sup>Hunter Medical Research Institute, Newcastle, New South Wales, Australia

<sup>4</sup>School of Psychology, The University of Newcastle, Callaghan, NSW, Australia

<sup>5</sup>Clinical Research Design, IT, and Statistical Support (CReDITSS), Hunter Medical Research Institute, Wallsend, New South Wales, Australia

**Acknowledgements** We would like to thank the staff and students from participating schools and the Healthy Schools Healthy Futures (HSHF) project team. Authors would like to acknowledge the in kind support of Principals Australia for the use of MindMatters curriculum resources and training. For the duration of the research project, a HSHF Aboriginal Cultural Steering Group made up of Aboriginal staff from local Aboriginal community organizations and Government Departments was established to provide Aboriginal cultural advice and direction regarding the design, implementation, evaluation and dissemination of all research trial elements. Similarly, an HSHF Cultural Advice Group was established consisting of Aboriginal staff from the HSHF project team to provide advice regarding the research trial.

We would like to thank the members of both the HSHF Aboriginal Cultural Steering Group and the HSHF Cultural Advice Group for their ongoing advice. Additionally, ethical approval was received from the Aboriginal Health and Medical Research Council.

**Contributors** RKH conducted the literature search, drafted the manuscript and contributed to study design, data collection, data analysis, data interpretation and coordination of the study. JW, MF, JB, LW and EC helped draft the manuscript and participated in the conception, design and coordination of the study. JD helped draft the manuscript and participated in the coordination of the study. CL, CO and JA helped draft the manuscript and conducted data analysis. All authors read and revised the manuscript critically for intellectual content and approved the final manuscript.

**Funding** The work was supported by funding from the National Health and Medical Research Council, nib Foundation, Cancer Council NSW Program Grant (PG 16-05) and Hunter New England Population Health and infrastructure support from the Hunter Medical Research Institute.

**Competing interests** None declared.

**Patient consent** Not required.

**Ethics approval** Relevant ethics committee approvals were obtained (Hunter New England Health Ref: 09/11/18/4.01; University of Newcastle Ref: H-2010-0029).

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** Requests for additional unpublished data should be forwarded to Rebecca.hodder@hnehealth.nsw.gov.au.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## REFERENCES

- Gilman SE, Abrams DB, Buka SL. Socioeconomic status over the life course and stages of cigarette use: initiation, regular use, and cessation. *J Epidemiol Community Health* 2003;57:802–8.
- Johnston LD, O'Malley PM, Bachman JG, et al. *Monitoring the future. National results on adolescent drug use: overview of key findings, 2007 (NIH Publication No. 08-6418)*. Bethesda, MD: National Institute on Drug Abuse, 2008.
- White V, Bariola E. Australian secondary schools students' use of tobacco, alcohol and over-the-counter and illicit substances in 2011. 2012 [http://www.nationaldrugstrategy.gov.au/internet/drugstrategy/Publishing.nsf/content/BCBF6B2C638E1202CA257ACD0020E35C/\\$File/National%20Report\\_FINAL\\_ASSAD\\_7.12.pdf](http://www.nationaldrugstrategy.gov.au/internet/drugstrategy/Publishing.nsf/content/BCBF6B2C638E1202CA257ACD0020E35C/$File/National%20Report_FINAL_ASSAD_7.12.pdf) (cited 19/03/2013).
- Rehm J, Taylor B, Room R. Global burden of disease from alcohol, illicit drugs and tobacco. *Drug Alcohol Rev* 2006;25:503–13.
- Centers for Disease Control and Prevention. Youth Risk Behavior Surveillance - United States, 2011. *Morbidity and Mortality Weekly Report* 2012;61:1–161.
- Henderson H, Nass L, Payne C, et al. *Smoking, drinking and drug use among young people in England 2012*. London: Health and Social Care Information Centre, 2013.
- White V, Williams T, Centre for Behavioural Research in Cancer - Cancer Council Victoria. *Australian secondary school students' use of tobacco, alcohol, and over-the-counter and illicit substances in 2014. 2016*: Drug Strategy Branch - Australian Government Department of Health, 2017.
- Australian Institute of Health and Welfare. *Drugs in Australia 2010: tobacco, alcohol and other drugs*. Canberra: AIHW, 2011. Report No.: Drug statistics series no.27. Cat. no. PHE 154.
- Carvajal SC, Granillo TM. A prospective test of distal and proximal determinants of smoking initiation in early adolescents. *Addict Behav* 2006;31:649–60.
- Hodder RK, Freund M, Bowman J, et al. Association between adolescent tobacco, alcohol and illicit drug use and individual and environmental resilience protective factors. *BMJ Open* 2016;6:e012688.
- Wiefferink CH, Peters L, Hoekstra F, et al. Clustering of health-related behaviors and their determinants: possible consequences for school health interventions. *Prev Sci* 2006;7:127–49.

12. Kavas AB. Self-esteem and health-risk behaviors among Turkish late adolescents. *Adolescence* 2009;44:187–98.
13. Fang L, Barnes-Ceeneey K, Schinke SP. Substance use behavior among early-adolescent Asian American girls: the impact of psychological and family factors. *Women Health* 2011;51:623–42.
14. Haegerich TM, Tolan PH. Core competencies and the prevention of adolescent substance use. *New Dir Child Adolesc Dev* 2008;2008:47–60.
15. Hawkins JD, Catalano RF, Miller JY. Risk and protective factors for alcohol and other drug problems in adolescence and early adulthood: Implications for substance abuse prevention. *Psychol Bull* 1992;112:64–105.
16. Bond L, Butler H, Thomas L, et al. Social and school connectedness in early secondary school as predictors of late teenage substance use, mental health, and academic outcomes. *J Adolesc Health* 2007;40:357.e9–357.e18.
17. Patton GC, Bond L, Carlin JB, et al. Promoting social inclusion in schools: a group-randomized trial of effects on student health risk behavior and well-being. *Am J Public Health* 2006;96:1582–7.
18. McLellan L, Rissel C, Donnelly N, et al. Health behaviour and the school environment in New South Wales, Australia. *Soc Sci Med* 1999;49:611–9.
19. Jessor R, Van Den Bos J, Vanderryn J, et al. Protective factors in adolescent problem behavior: moderator effects and developmental change. *Dev Psychol* 1995;31:923–33.
20. Neumark-Sztainer D, Story M, French SA, et al. Psychosocial correlates of health compromising behaviors among adolescents. *Health Educ Res* 1997;12:37–52.
21. Resnick MD, Bearman PS, Blum RW, et al. Protecting adolescents from harm. Findings from the National Longitudinal Study on Adolescent Health. *JAMA* 1997;278:823–32.
22. Vuille JC, Schenkel M. Social equalization in the health of youth. The role of the school. *Eur J Public Health* 2001;11:287–93.
23. Brownlee K, Rawana J, Franks J, et al. A systematic review of strengths and resilience outcome literature relevant to children and adolescents. *Child and Adolescent Social Work Journal* 2013;30:435–59.
24. Lee TY, Cheung CK, Kwong WM. Resilience as a positive youth development construct: a conceptual review. *ScientificWorldJournal* 2012;2012:1–9.
25. Masten AS, magic O. Resilience processes in development. *Am Psychol* 2001;56:227–38.
26. Masten AS, Best KM, Garmezy N. Resilience and development: Contributions from the study of children who overcome adversity. *Dev Psychopathol* 1990;2:425–44.
27. Hodder RK, Freund M, Wolfenden L, et al. Systematic review of universal school-based 'resilience' interventions targeting adolescent tobacco, alcohol or illicit substance use: A meta-analysis. *Prev Med* 2017;100:248–68.
28. Bond L, Patton G, Glover S, et al. The Gatehouse Project: can a multilevel school intervention affect emotional wellbeing and health risk behaviours? *J Epidemiol Community Health* 2004;58:997–1003.
29. Bond L, Thomas L, Coffey C, et al. Long-term impact of the Gatehouse Project on cannabis use of 16-year-olds in Australia. *J Sch Health* 2004;74:23–9.
30. Hodder RK, Daly J, Freund M, et al. A school-based resilience intervention to decrease tobacco, alcohol and marijuana use in high school students. *BMC Public Health* 2011;11:1–33.
31. Mrazek PJ, Haggerty RJ. *Reducing risks for mental disorders: frontiers for preventive intervention research*. Washington D.C.: Institute of Medicine, 1994.
32. Botvin GJ, Griffin KW. School-based programmes to prevent alcohol, tobacco and other drug use. *Int Rev Psychiatry* 2007;19:607–15.
33. United Nations Office for Drug Control and Crime Prevention. School-based Drug Education: A Guide for Practitioners and the Wider Community. [http://www.unicef.org/lifeskills/index\\_7242.html](http://www.unicef.org/lifeskills/index_7242.html) 2003.
34. Canadian Centre on Substance Abuse. *National framework for action to reduce the harms associated with alcohol and other drugs and substances in Canada*. Ottawa, Ontario: Government of Canada, 2005.
35. Canadian Centre on Substance Abuse. *Building on our strengths: Canadian standards for school-based youth substance abuse prevention*. Ottawa, ON: Canadian Centre on Substance Abuse, 2010.
36. Ministerial Council on Drug Strategy. *National drug strategy 2010–2015: a framework for action on alcohol, tobacco and other drugs*. Canberra, ACT: Commonwealth of Australia, 2011.
37. Spoth R, Shin C, Guyll M, et al. Universality of effects: an examination of the comparability of long-term family intervention effects on substance use across risk-related subgroups. *Prev Sci* 2006;7:209–24.
38. Oesterle S, Hawkins JD, Fagan AA, et al. Testing the universality of the effects of the communities that care prevention system for preventing adolescent drug use and delinquency. *Prev Sci* 2010;11:411–23.
39. Flay BR, Biglan A, Boruch RF, et al. Standards of evidence: criteria for efficacy, effectiveness and dissemination. *Prev Sci* 2005;6:151–75.
40. Beets MW, Flay BR, Vuchinich S, et al. Use of a social and character development program to prevent substance use, violent behaviors, and sexual activity among elementary-school students in Hawaii. *Am J Public Health* 2009;99:1438–45.
41. Caria MP, Faggiano F, Bellocco R, et al. The influence of socioeconomic environment on the effectiveness of alcohol prevention among European students: a cluster randomized controlled trial. *BMC Public Health* 2011;11:312.
42. Flay BR, Graumlich S, Segawa E, et al. Effects of 2 prevention programs on high-risk behaviors among African American youth: a randomized trial. *Arch Pediatr Adolesc Med* 2004;158:377–84.
43. Perry CL, Komro KA, Weblen-Mortenson S, et al. A randomized controlled trial of the middle and junior high school D. A.R.E. and D.A.R.E. plus programs. *Archives of Pediatrics and Adolescent Medicine* 2003;157:178–87.
44. Perry CL, Williams CL, Veblen-Mortenson S, et al. Project Northland: outcomes of a communitywide alcohol use prevention program during early adolescence. *Am J Public Health* 1996;86:956–65.
45. Spoth R, Redmond C, Shin C, et al. Substance-use outcomes at 18 months past baseline: the PROSPER Community-University Partnership Trial. *Am J Prev Med* 2007;32:395–402.
46. Spoth RL, Redmond C, Trudeau L, et al. Longitudinal substance initiation outcomes for a universal preventive intervention combining family and school programs. *Psychol Addict Behav* 2002;16:129–34.
47. Hodder RK, Freund M, Bowman J, et al. A cluster randomised trial of a school-based resilience intervention to decrease tobacco, alcohol and illicit drug use in secondary school students: study protocol. *BMC Public Health* 2012;12:1009.
48. Dray J, Bowman J, Campbell E, et al. Effectiveness of a pragmatic school-based universal intervention targeting student resilience protective factors in reducing mental health problems in adolescents. *J Adolesc* 2017;57:74–89.
49. Hodder RK, Freund M, Bowman J, et al. Effectiveness of a pragmatic school-based universal resilience intervention in reducing tobacco, alcohol and illicit substance use in a population of adolescents: cluster-randomised controlled trial. *BMJ Open* 2017;7:e016060.
50. Centre for Epidemiology and Evidence. *Australian Bureau of Statistics population estimates (SAPHaRI)*. Sydney: NSW Ministry of Health, 2012.
51. Kirk CM, Lewis RK, Lee FA, et al. The power of aspirations and expectations: the connection between educational goals and risk behaviors among African American adolescents. *J Prev Interv Community* 2011;39:320–32.
52. Australian Curriculum AaRA. *My school 2015*: Australian Curriculum, Assessment and Reporting Authority, 2011.
53. Trewin D. *Information paper census of population and housing socio-economic indexes for areas: australia 2001*. Canberra: Australian Bureau of Statistics: Commonwealth of Australia, 2003.
54. NSW Department of Education and Communities. Low socioeconomic status school communities national partnership. 2011 [http://www.lowsesschools.nsw.edu.au/wcb-content/uploads/psp/file/2\\_%20LOW\\_SES\\_InfoPackage.pdf](http://www.lowsesschools.nsw.edu.au/wcb-content/uploads/psp/file/2_%20LOW_SES_InfoPackage.pdf) (cited 5 Sep 2001).
55. World Health Organisation. *Planning meeting in health promoting schools project: background, development and strategy outline of the health promoting schools project*. Copenhagen: WHO, 1991.
56. Fergus S, Zimmerman MA. Adolescent resilience: a framework for understanding healthy development in the face of risk. *Annu Rev Public Health* 2005;26:399–419.
57. Bond L, Glover S, Godfrey C, et al. Building capacity for system-level change in schools: lessons from the Gatehouse Project. *Health Educ Behav* 2001;28:368–83.
58. Gottfredson DC, Gottfredson GD. Quality of school-based prevention programs: results from a national survey. *J Res Crime Delinq* 2002;39:3–35.
59. Hazell T. *MindMatters: evaluation of the professional development program and school-level implementation*. Hunter Institute of Mental Health: Newcastle, 2006.
60. Lezotte LW, Skaife RD, Holstead MD. *Effective schools: only you can make a difference*: All Star Publishing, 2002.
61. Luther SS, Cicchetti D, Becker B. The construct of resilience: a critical evaluation and guidelines for future work. *Child Dev* 2000;71:543–62.

62. MindMatters Evaluation Consortium. *Report of the MindMatters (National Mental Health in Schools Project) Evaluation Project, vols. 1-4*. Newcastle: Hunter Institute of Mental Health, 2000.
63. Wagner EF, Tubman JG, Gil AG. Implementing school-based substance abuse interventions: methodological dilemmas and recommended solutions. *Addiction* 2004;99(Suppl 2):106–19.
64. Wilson KD, Kurz RS. Bridging implementation and institutionalization within organizations: proposed employment of continuous quality improvement to further dissemination. *J Public Health Manag Pract* 2008;14:109–16.
65. Hanson TL, Kim JO. *Measuring resilience and youth development: the psychometric properties of the Healthy Kids Survey*. Washington, DC: U.S. Department of Education, Institute of Education Sciences, National Centre for Education Evaluation and Regional Assistance, Regional Educational Laboratory West, 2007.
66. Department of Health and Aged Care. *Measuring remoteness: Accessibility/Remoteness Index of Australia (ARIA)*. Canberra: Australian Government, 2001.
67. Assmann SF, Pocock SJ, Enos LE, *et al*. Subgroup analysis and other (mis)uses of baseline data in clinical trials. *Lancet* 2000;355:1064–9.
68. Brookes ST, Whitley E, Peters TJ, *et al*. Subgroup analyses in randomised controlled trials: quantifying the risks of false-positives and false-negatives. *Health Technol Assess* 2001;5:1-56.
69. Pagano M, Gauvreau K. *Principles of Biostatistics*. California, US: Duxbury, 2000:292–4.
70. SAS Software Version 9.3 for Windows [computer program]. Version 9.3. New York: Cary, NC, USA, 2001.
71. Abatemarco DJ, West B, Zec V, *et al*. Project northland in Croatia: a community-based adolescent alcohol prevention intervention. *J Drug Educ* 2004;34:167–78.
72. Yoong SL, Wolfenden L, Clinton-McHarg T, *et al*. Exploring the pragmatic and explanatory study design on outcomes of systematic reviews of public health interventions: a case study on obesity prevention trials. *J Public Health* 2014;36:170–6.
73. Finch M, Jones J, Yoong S, *et al*. Effectiveness of centre-based childcare interventions in increasing child physical activity: a systematic review and meta-analysis for policymakers and practitioners. *Obes Rev* 2016;17:412–28.
74. Hawkins JD, Catalano RF, Miller JY. Risk and protective factors for alcohol and other drug problems in adolescence and early adulthood: implications for substance abuse prevention. *Psychol Bull* 1992;112:64–105.
75. Cleveland MJ, Feinberg ME, Bontempo DE, *et al*. The role of risk and protective factors in substance use across adolescence. *J Adolesc Health* 2008;43:157–64.
76. Burke JF, Sussman JB, Kent DM, *et al*. Three simple rules to ensure reasonably credible subgroup analyses. *BMJ* 2015;351:h5651.
77. Tigges BB. Parental consent and adolescent risk behavior research. *J Nurs Scholarsh* 2003;35:283–9.
78. MindMatters Evaluation Consortium. *MindMatters: a mental health promotion resource for secondary schools*: Commonwealth of Australia, 2000.
79. The Queensland Government. *Social and emotional wellbeing programs*, 2008. cited 08/12/8 A.D..