



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Management of COVID-19 in patients with seizures: Mechanisms of action of potential COVID-19 drug treatments and consideration for potential drug-drug interactions with anti-seizure medications

Shreshtha Jain ^a, Heidrun Potschka ^b, P. Prarthana Chandra ^c, Manjari Tripathi ^d, Divya Vohora ^{a,*}

^a Department of Pharmacology, School of Pharmaceutical Education and Research, Jamia Hamdard, New Delhi, India

^b Institute of Pharmacology, Toxicology, and Pharmacy, Ludwig-Maximilians-University (LMU), Königinstr. 16, D-80539, Munich, Germany

^c Hamdard Institute of Medical Sciences and Research, New Delhi, India

^d Department of Neurology, All India Institute of Medical Sciences, New Delhi, India

ARTICLE INFO

Keywords:

COVID-19 candidate drugs
Seizures
Drug-drug interactions
Antiviral drugs
Immunomodulatory or anti-inflammatory drugs
NSAIDs

ABSTRACT

In regard to the global pandemic of COVID-19, it seems that persons with epilepsy (PWE) are not more vulnerable to get infected by SARS-CoV-2, nor are they more susceptible to a critical course of the disease. However, management of acute seizures in patients with COVID-19 as well as management of PWE and COVID-19 needs to consider potential drug-drug interactions between antiseizure drugs and candidate drugs currently assessed as therapeutic options for COVID-19. Repurposing of several licensed and investigational drugs is discussed for therapeutic management of COVID-19. While for none of these approaches, efficacy and tolerability has been confirmed yet in sufficiently powered and controlled clinical studies, testing is ongoing with multiple clinical trials worldwide. Here, we have summarized the possible mechanisms of action of drugs currently considered as potential therapeutic options for COVID-19 management along with possible and confirmed drug-drug interactions that should be considered for a combination of antiseizure drugs and COVID-19 candidate drugs. Our review suggests that potential drug-drug interactions should be taken into account with drugs such as chloroquine/hydroxychloroquine and lopinavir/ritonavir while remdesivir and tocilizumab may be less prone to clinically relevant interactions with ASMs.

1. Introduction

The catastrophic global pandemic of Coronavirus disease 2019 (COVID-19) has put large parts of the world on a transient standstill. COVID-19 was first discovered in December 2019 and the causative agent was a novel subtype of beta-coronaviruses, known as SARS-CoV-2 (Rothan and Byrareddy, 2020). The novel coronavirus is an enveloped non-segmented positive-sense RNA virus (Richman et al., 2016) that was transmitted to humans via zoonotic transmission, similar to its precursors, severe acute respiratory syndrome coronavirus (SARS-CoV) (Ksiazek et al., 2003) and the Middle East respiratory syndrome

coronavirus (MERS-CoV) (Zaki et al., 2012). These viruses primarily affect the respiratory system, however, clinical literature also provides evidence for neuroinvasive and neurotropic properties of SARS-CoV-2 (Carod-Artal, 2020; Li et al., 2020b; Najjar et al., 2020). Neurological symptoms reported in patients with COVID-19 include febrile seizures, status epilepticus and complications including encephalopathy, cerebral haemorrhage (Asadi-Pooya, 2020; Asadi-Pooya and Simani, 2020; Carod-Artal, 2020; Yin et al., 2020). These clinical manifestations suggest that people with neurological disorders may be more vulnerable to experience severe symptoms of COVID-19 disease. However, that does not seem to be the case. The International League Against Epilepsy

Abbreviations: ACE, angiotensin-converting enzyme inhibitors; ASM, anti-seizure medication; CBD, cannabidiol; COVID, 19- Coronavirus disease 2019; CYP, cytochrome P; EMA, European Medicines Agency; FDA, US Food and Drug Administration; G-CSF, granulocyte-colony stimulating factor; GM-CSF, granulocyte-macrophage colony-stimulating factor; HDAC2, histone deacetylases; IL, interleukin; ILAE, International League Against Epilepsy; IMP, importin; MERS-CoV, Middle East respiratory syndrome coronavirus; NIH, National Institute of Health; NSAIDs, non-steroidal anti-inflammatory drugs; PWE, persons with epilepsy; SARS-CoV, severe acute respiratory syndrome coronavirus; T-705-RTP, favipiravir ibofuranosyl-5'-triphosphate; TGA, Therapeutic Goods Administration of Australia; TMPRSS2, type 2 transmembrane serine protease; TNF, tumor necrosis factor; UGT, uridine diphosphate glucuronosyltransferase.

* Corresponding author at: Department of Pharmacology, School of Pharmaceutical Education and Research, Jamia Hamdard, New Delhi, 110062, India.

E-mail address: dvohora@jamiahamdard.ac.in (D. Vohora).

<https://doi.org/10.1016/j.epilepsyres.2021.106675>

Received 15 December 2020; Received in revised form 29 April 2021; Accepted 14 May 2021

Available online 19 May 2021

0920-1211/© 2021 Elsevier B.V. All rights reserved.

(ILAE) has informed that persons with epilepsy (PWE) are not likely to be more susceptible to get COVID-19 nor are they inclined to suffer through severe manifestations of SARS-CoV-2 infection (ILAE, 2020). Even if PWE are exposed to SARS-CoV-2, it is unlikely that the frequency of seizures increases (ILAE, 2020). Nevertheless, management of COVID-19 in PWE or with acute reactive seizures requires certain precautions and guidelines to avoid worsening of the condition. In particular, potential drug-drug interaction that may occur on concomitant administration of anti-seizure medication (ASM) along with the drugs for treatment of COVID-19 need to be taken into account.

The rapid spread of the SARS-CoV-2 pandemic with a high number of severe cases and high mortality rates has triggered almost desperate attempts to repurpose different types of drugs for treatment of patients with a severe course of COVID-19. At this point of time, it needs to be emphasized that no drug has been proven to be efficacious and safe for therapeutic management of COVID-19 based on sufficiently powered and controlled studies. This fact is also pointed out by an expert group of the National Institute of Health (<https://www.covid19treatmentguidelines.nih.gov/>). A substantial number of clinical studies are currently ongoing or planned with on today's date (15th March, 2021) 5017 clinical trials being registered at clinicaltrials.gov with a high percentage of these trials aiming to explore drug candidates for the management of COVID-19 (<https://clinicaltrials.gov/ct2/results?cond=COVID-19>). However, based on the current state-of-knowledge, only a number of drug candidates are classified by the NIH expert panel as 'therapeutic options for COVID-19 currently under investigation' along with the statement that there are 'insufficient data to either recommend for or against the use'. These drugs include immunomodulators such as interleukin (IL)-1 inhibitors (Anakinra). For other immunomodulatory drugs such as interleukin-6 inhibitors, interferons and janus kinase inhibitors along with anti-viral agents including lopinavir, ritonavir and other HIV protease inhibitors, and anti-malarial drugs such as chloroquine and hydroxychloroquine in combination with or without azithromycin, as well as for the recently reported anti-parasitic agent, ivermectin, the NIH expert panel clearly recommends against the use except in the context of a clinical trial (NIH, 2020). However, recommendations vary as per countries and these drugs are still used for COVID-19 in several other countries. The only drugs that are currently being recommended by the NIH include remdesivir and corticosteroids (dexamethasone) for the treatment of COVID-19 with a severe disease course (NIH, 2020).

For COVID-19 clinical studies, potential drug-drug interactions need to be considered for all patients with chronic diseases requiring continuous treatment. This in particular applies for PWE as several ASMs are prone to drug-drug interactions (Patsalos et al., 2018; Zaccara and Perucca, 2014). Information about drug-drug interactions is also of particular relevance for intensive care unit management of critically-ill COVID-19 patients who may develop acute seizures during a severe disease course. Here, we will discuss the mechanism of action of drugs currently under discussion as potential therapeutic options for COVID-19 management (referred to as COVID-19 candidate drugs). Further, we will also discuss the current state-of-knowledge about possible interactions between ASMs and COVID-19 candidate drugs.

2. Mechanisms of action of COVID-19 candidate drugs

The current understanding of the life cycle of the novel coronavirus SARS-CoV-2 suggests drug target candidates for the prevention and treatment of COVID-19. As it is known, the virus contains a single-stranded RNA, that upon entering the host cell unfolds and translates into polypeptides that further mitigate the synthesis of viral RNA strands via RNA-dependent viral RNA polymerase (Chen et al., 2020c). Furthermore, virus binding to ACE2 through the spike (S) protein and entering the host cell is facilitated by the host type 2 transmembrane serine protease (TMPRSS2). Based on knowledge about these viral invasion mechanisms, the targeting of viral entry is in the focus of current

efforts to develop therapeutic treatments against COVID-19 (Hoffmann et al., 2020).

In the following, we highlight some of the potential drug candidates that have been or are assessed in first clinical trials (Tu et al., 2020). In this context, it is again emphasized that for all candidate drugs listed there is a lack of sufficient evidence supporting efficacy and tolerability for management of COVID-19 at this point of time. The outcome of future studies are awaited for final conclusions about the therapeutic value of these compounds. As mentioned in the introduction, an NIH expert panel even argues against the use of some of the compounds considered here due to tolerability issues and toxicity or negative clinical efficacy data (NIH, 2020). Considering registered clinical studies, we will discuss some of these compounds despite the above recommendations by NIH as potential drug-drug interactions might be relevant for studies that are currently ongoing in different countries.

It is also emphasized that none of the following drugs has been specifically developed for management of SARS-CoV-2, and that the potential therapeutic options are so far limited to repurposing of drugs, which were originally developed and designed for different indications.

Considering the discussed candidate therapeutic options, these can be categorized as those with antiviral or anti-inflammatory potential.

2.1. Antiviral drugs

2.1.1. Remdesivir

Remdesivir is a prodrug, previously known as GS-5734. It is a monophosphate that is metabolized to an adenosine nucleoside triphosphate analogue, which is integrated into the viral RNA prior to its replication (Amirian and Levy, 2020). As a consequence, the function of RNA-dependent viral RNA polymerase and RNA synthesis is inhibited (Warren et al., 2016). Due to its antiviral action and low EC₅₀, remdesivir was explored as a possible therapeutic agent during the Ebola virus outbreak (Siegel et al., 2017). An earlier experimental study suggested an efficacy of remdesivir against various coronaviruses including SARS-CoV and MERS-CoV (Sheahan et al., 2017). As the safety of the drug was previously validated in humans and due to its broad spectrum, remdesivir became the prime candidate for clinical trials against COVID-19 (Al-Tawfiq et al., 2020). *In vitro* evaluation confirmed inhibition of SARS-CoV-2 replication at a low EC₅₀ value of 0.77 μM (Wang et al., 2020a). So far, first clinical reports and studies indicate a possible beneficial impact of remdesivir in patients with severe COVID-19 (Holshue et al., 2020) (Grein et al., 2020). Remdesivir has now been authorized by US Food and Drug Administration (FDA) for emergency use and has been approved by the European Medicines Agency (EMA) as well as Therapeutic Goods Administration of Australia (TGA) and has also been widely prescribed for patients in India, Japan and South Korea for the treatment of COVID-19.

2.1.2. Favipiravir

Favipiravir is the first oral antiviral drug that was approved by the Director General of India for the treatment of COVID-19. Apart from India, the drug has also been approved in Russia and on-going clinical trials are being conducted in the US and some parts of middle-east (Varadharajan, 2020). Favipiravir is a broad spectrum anti-viral drug that inhibits the replication of virus (Furuta et al., 2017). The pyrazine carboxamide derivative favipiravir serves as a prodrug that undergoes ribosylation and phosphorylation to form its active metabolite favipiravir ibofuranosyl-5'-triphosphate (T-705-RTP) (Furuta et al., 2017). T-705-RTP gets incorporated into the viral RNA thereby interfering with the process of viral replication and inhibiting the viral RNA-dependent RNA polymerase (Du and Chen, 2020; Furuta et al., 2017). It has already been demonstrated that T-705-RTP can limit influenza virus replication by inhibiting the RNA-dependent RNA polymerase without affecting the human DNA polymerases α, β, or γ subunit (Sissoko et al., 2016). Favipiravir has also shown its effectiveness against various other RNA viruses that cause hemorrhagic fever including arenavirus,

bunyavirus, flavivirus, and filoviruses (Du and Chen, 2020). Hence, favipiravir is being investigated globally in clinical trials as a potential therapeutic agent against SARS-CoV-2 (Karlsen et al., 2020). During an *in vitro* analysis, it was confirmed that at high maximum effective concentration ($EC_{50} = 61.88 \mu\text{M}$), favipiravir was able to inhibit SARS-CoV-2 viral infection (Wang et al., 2020a). An open-label non-randomized clinical study conducted in China, provided evidence that favipiravir exposure can result in therapeutic responses with a beneficial impact on progression of the disease and viral clearance in COVID-19 patients (Cai et al., 2020). Further clinical studies seem to confirm the promising results for favipiravir as an effective treatment against SARS-CoV-2 infection (Chen et al., 2020a; Ito et al., 2021; Noda et al., 2020). In Russia, favipiravir is marketed as AVIFAVIR for management of COVID-19 patients following first clinical studies indicating its efficacy, safety, and tolerability in phase II/III multicentric randomized clinical trials (Ivashchenko et al., 2020).

2.1.3. Lopinavir/ritonavir

The use of lopinavir/ritonavir has been discouraged recently and has also been recommended against by the NIH expert panel in their COVID-19 Treatment Guidelines, due to the adverse pharmacodynamics and no clinically beneficial outcome reported in COVID-19 patients (NIH, 2020). Lopinavir and Ritonavir are anti-retroviral agents that are used in combination against human immunodeficiency virus (Kujawski et al., 2020). They act by inhibiting the enzyme protease responsible for cleaving viral polypeptides, which are essential for viral replication. It has been described that lopinavir/ritonavir also acts on the 3CL protease enzyme involved in replication of coronaviruses (Nukoolkarn et al., 2008). Related to this effect, therapeutic effects have been previously assessed in patients with SARS-CoV or MERS-CoV infections (Yao et al., 2020a). A recent randomized clinical trial conducted by Cao and colleagues (2020) failed to confirm a benefit with lopinavir/ritonavir treatment in COVID-19 patients (Cao et al., 2020).

2.1.4. Ivermectin

Ivermectin is a broad-spectrum anti-parasitic agent that has been approved by the FDA for the treatment of internal and external parasitic infestations including tropical diseases such as onchocerciasis, helminthiasis, and scabies (Crump and Omura, 2011; Omura and Crump, 2014). Ivermectin has also exhibited anti-viral potential *in vitro* with efficacy against RNA viruses such as dengue virus (DENV), Zika virus, HIV and influenza (Barrows et al., 2016; Götz et al., 2016; Wagstaff et al., 2012). Ivermectin inhibits the intracellular transportation of the virus in the host cells by targeting the host importin (IMP) α/β nuclear transport proteins (Wagstaff et al., 2012; Yang et al., 2020b). Based on corresponding evidence that $IMP\alpha/\beta$ seems to possess a role during SARS-CoV infection, an *in vitro* study was conducted by Leon Caly and co-workers that reported a 5000-fold reduction in the RNA levels of SARS-CoV-2 at 48 h post incubation with $5 \mu\text{M}$ of ivermectin in Vero/hSLAM cells (Caly et al., 2020). However, further reports were published that warned against the use of ivermectin in COVID-19 patients as the dose required to exhibit anti-viral activity as observed *in vitro*, was 100 fold more than the clinically approved dose (Bray et al., 2020; Chaccour et al., 2020). Conversely, a retrospective cohort study conducted in South Florida, reported a lower mortality rate in patients confirmed positive for SARS-CoV-2, who were on ventilatory support or required higher inspired oxygen and were treated with ivermectin (Rajter et al., 2020). Hence, these findings are required to be followed up by controlled randomized clinical trials aiming to further evaluate the efficacy and safety of ivermectin for treating COVID-19.

2.2. Drugs with anti-inflammatory or immunomodulatory potential

2.2.1. Chloroquine/hydroxychloroquine

The US Food and Drug Administration (FDA) along with WHO and NIH has warned and recommended against the use of chloroquine or

hydroxychloroquine except for emergency use in hospitalized COVID-19 patients or in a clinical trial to avoid the risk of ventricular tachycardia and prolonged QT interval (Li et al., 2020a). However, in developing countries like India, the official guidelines recommend the use of hydroxychloroquine in combination with or without azithromycin, for treatment of mild to moderate symptomatic or asymptomatic COVID-19 patients and also as a prophylactic measure (Medina and Moncada, 2020). Chloroquine and hydroxychloroquine are anti-malarial agents that are also used as preventive treatment against chronic inflammatory diseases such as Systemic lupus erythematosus or rheumatoid arthritis (Fox, 1993). These drugs have also been repurposed for therapeutic management of viral infections. Chloroquine has been previously used against the structurally similar SARS-CoV due to its inhibitory action on the glycosylation of the ACE2 receptor (Vincent et al., 2005). Hydroxychloroquine also exhibits antiviral action by acidifying the endosomal pH and thereby blocking the fusion of virus with the host cell membrane (Savarino et al., 2003). Immunomodulatory effects are also observed in response to these drugs with a decrease in the production of pro-inflammatory cytokines, and an inhibition of autophagy and lysosomal action (Sanders et al., 2020). Recent *in vitro* studies have demonstrated inhibitory effects on SARS-CoV-2 at a half-maximal effective concentration (EC_{50}) of hydroxychloroquine and chloroquine in micromolar range ($EC_{50} = 6.14 \mu\text{M}$; $EC_{50} = 23.90 \mu\text{M}$, respectively) (Yao et al., 2020b). Few clinical trials have also validated the antiviral action of these drugs in COVID-19 patients (Chen et al., 2020b; Gao et al., 2020; Gautret et al., 2020). Tolerability concerns have been raised based on the pharmacodynamics of these compounds related to their effect on cardiac sodium channels (Frisk-Holmberg et al., 1983). Recent clinical studies exploring a potential benefit in patients with COVID-19 confirmed the occurrence of QTc interval prolongation in patients with COVID-19 (Chorin et al., 2020; van den Broek et al., 2020). Based on these findings, the need for ECG monitoring has been pointed out by the authors. The adverse effect potential has a particular relevance in the context of SARS-CoV-2 infections considering that patients with cardiovascular disease are considered patients with a high risk for a severe disease course, which therefore will also more often require pharmacotherapy.

2.2.2. Corticosteroids

According to the NIH COVID-19 treatment guidelines, the expert panel is currently recommending the use of dexamethasone, a glucocorticoid, for treating COVID-19 patients that are mechanically ventilated or require supplemental oxygen (NIH, 2020). The use of dexamethasone and other corticosteroids such as prednisone, methylprednisolone, and hydrocortisone has been advised based on their anti-inflammatory properties (Yang et al., 2020a). During the progression of SARS-CoV-2 infection, patients suffer from a "cytokine storm" that results from an abrupt release of cytokines and chemokines such as interleukins IL-6 and IL-1 β , granulocyte-colony stimulating factor (G-CSF), granulocyte-macrophage colony-stimulating factor (GM-CSF), and tumor necrosis factor (TNF) (Huang et al., 2020; Nie et al., 2020; Shi et al., 2020; Xu et al., 2020b; Zhou et al., 2020).

Glucocorticoids can limit inflammatory signaling based on an interference with gene expression. Among other effects, glucocorticoids cause activation of histone deacetylases (HDAC2) by stimulating the transcription factors CREB, AP-1, and NF- κ B (Ferner et al., 2020). This molecular mechanism is of particular interest as the SARS-CoV-2 Nsp5 protease can inhibit transport of HDAC2 into the nucleus, thereby impairing inflammation-associated cellular responses (El Baba and Herbein, 2020).

However, considering immunosuppressant effects related to the mechanism of action, the use of corticosteroids has been controversially discussed. So far there is insufficient clinical evidence, whether their administration helps to reduce the mortality of critically-ill COVID-19 patients (Singh et al., 2020). Nevertheless, application of dexamethasone has been associated with improved clinical outcomes when

administered to COVID-19 patients during the Randomized Evaluation of COVID-19 Therapy (RECOVERY) Trial. Similarly, few retrospective clinical studies have reported evidence for a beneficial impact of short-term use of the corticosteroid dexamethasone with a reduction of clinical respiratory symptoms and improvement in lung absorption (Singh et al., 2020; Wang et al., 2020b; Yang et al., 2020a). Further investigations with randomized controlled clinical trials are required to confirm the use of corticosteroids for the treatment of COVID-19 patients with a severe disease course (Nicolau and Bafadhel, 2020).

2.2.3. Tocilizumab

Tocilizumab is a monoclonal antibody that acts as an antagonist at the IL-6 receptor (Rose-John et al., 2017), and is used for therapeutic management of rheumatoid arthritis (Biggioggero et al., 2018). It inhibits the release of cytokine-storm related mediators (Borku Uysal et al., 2020). As mentioned above, there are a number of cases that report heightened levels of cytokines including IL-6 in patients with COVID-19 (Huang et al., 2020) (Zhou et al., 2020). Thus, first clinical trials have been conducted with tocilizumab. These provide limited evidence for a possible clinical improvement with reduced fever, improved respiratory functioning, and earlier discharge from the hospitals (Cao, 2020) (Xu et al., 2020a). Tocilizumab has been listed under the tentative 8th edition of the National Health Commission of the People's Republic of China COVID-19 Diagnosis and Treatment Guide, as a therapeutic agent for critically-ill patients with a severe COVID-19 course and increased levels of IL-6 (Wang, 2021; Anon, 2021b). On the contrary, the NIH expert panel has recommended against using interleukin-6 inhibitors including anti-IL-6 receptor monoclonal antibodies such as tocilizumab, sarilumab, etc. and anti-IL-6 monoclonal antibodies (siltuximab) for treating COVID-19, except in a clinical trial (NIH, 2020).

2.2.4. Anakinra

According to the NIH expert panel, there is insufficient data to recommend for or against the use of interleukin-1 inhibitors such as anakinra as a potential therapeutic option for COVID-19 management (NIH, 2020). While there seems to be no relevant clinical data supporting a benefit in patients with COVID-19, a possible use of anakinra

has been brought up considering a subgroup of COVID-19 patients developing a cytokine storm syndrome (Cron and Chatham, 2020). In these patients, both, interleukin-6 inhibitors as well as the interleukin-1 receptor antagonist anakinra are discussed. This may be of particular future interest for patients with COVID-19 and seizure development, considering that first reports during recent years have indicated that anakinra may be efficacious in the management of a febrile infection-related epilepsy syndrome with development of a super-refractory status epilepticus (Dilena et al., 2019; Kenney-Jung et al., 2016; Sa et al., 2019).

3. Potential drug-drug interactions between ASMs and COVID-19 candidate drugs

First of all, we checked the literature regarding evidence for potential pharmacodynamic or pharmacokinetic drug interactions between the repurposed candidate drugs currently assessed for COVID-19 and available ASMs. Thereby, we considered potential pharmacokinetic interactions with a focus on an impact of the drugs on CYP450 enzymes. In addition, we also considered clinical reports about a possible impact of the compounds on seizure control in PWE. A brief summary of these case reports is provided in Table 1. The causal relationship between drug use and occurrence of seizures may of course not be established in all these cases. Nevertheless, some of the reports also point to the need for a more in-depth analysis of the potential for drug-drug interactions between COVID-19 candidate drugs and ASMs.

A list of probable drug-drug interactions of COVID-19 candidate drugs is provided by the following website <https://www.covid19-druginteractions.org> (Anon, 2021a) based on an initiative by the Liverpool Drug Interactions Group at the University of Liverpool. This list includes drug-drug interactions of COVID-19 candidate drugs with several ASMs. Table 2 lists the essential classes of drugs referenced from the website that indicates different degrees of drug-drug interactions. The classification distinguishes between drugs that are not to be co-administered, drugs that require dose adjustment and/or continuous monitoring if co-administered, and drugs that have a weak interaction potential with no need of monitoring. Justification for the above classification of the drug-drug interactions is, however, not provided by the

Table 1

Summary of case reports with evidence for possible seizure induction in patients exposed to drugs that are currently assessed as COVID-19 candidate drugs.

Patients' data (Age in years/ Gender)	Dosage	Indication	Duration of treatment before onset of seizures	Type of seizures observed	Previous history of seizures	References
CNhydroxychloroquine/ Hydroxychloroquine						
Case1–35yr/ Male; Case2–19yr/ Female	Chloroquine: 100 mg/day	Prophylactic anti-malarial treatment	2 days; 8 days	Generalized tonic-clonic seizures	None	(Schiemann et al., 2000)
17yr/ Female	Hydroxychloroquine: 200 mg/day	Systemic lupus erythematosus	14 days	Tonic-clonic seizures	Clinical history of complex partial seizure	(Malcangi et al., 2000)
49yr/ Male	Chloroquine: 150 mg three times daily for the first week; 150 mg twice for the second week; 150 mg daily for the third week	Erythema nodosum leprosum (Leprosy)	9 days	Tonic-clonic seizures	None	(Ebenso, 1998)
49yr/ Female	Chloroquine: 250 mg/day	Systemic lupus erythematosus	30 days	Complex partial seizures	None	(Krzeminski et al., 2018)
14yr/ Female	Chloroquine: 500 mg/day	Systemic lupus erythematosus	21 days	Tonic-clonic seizures	None	(Tristano et al., 2004)
26yr/ Female	Hydroxychloroquine: 500 mg/day	Systemic lupus erythematosus	30 days	Generalized tonic-clonic seizures	None	(Jafri et al., 2017)
Lopinavir/ritonavir						
10yr/ Female	Second line antiretroviral therapy (zidovudine-lamivudine-lopinavir/ritonavir)	HIV	8 weeks	Generalized tonic-clonic seizures	None	(Otto et al., 2020)
54yr/ Male	Antiretroviral therapy with lopinavir/ritonavir and abacavir/lamivudine	Pseudo-HIV	5 months	Non-convulsive status epilepticus	Alcohol-induced dementia, liver cirrhosis, epilepsy and psoriatic arthritis	(Etgen et al., 2010)

Table 2

Potential drug-drug interactions between antiseizure drugs (ASMs) and COVID-19 candidate drugs*.

A) List of drug-drug interactions with documented proof of evidence-									
Antiseizure medication	COVID-19 candidate drug	Lopinavir ^b / Ritonavir	Dexamethasone						
	Relevant effect on enzymes	Inhibition of CYP3A4 ^c	Inducer/ Inhibitor of CYP3A subfamily						
<i>Enzyme Inducers</i>	Carbamazepine	Induction of CYP3A4	Decrease in concentration of LPV/RTV Increase in concentration of CBZ (83 %) (Burman and Orr, 2000) + + +	Decrease in concentration of DEX Decrease in concentration of CBZ (Clearance rate ↑ by 1.41) (Bénil and Vecht, 2015) ++ Decrease in concentration of DEX Decrease in concentration of PB (Clearance rate ↑ by 1.79) (Bénil and Vecht, 2015) ++ Decrease in concentration of DEX Decrease in concentration of PHT (50 %) (Bénil and Vecht, 2015) ++					
	Phenobarbital	Induction of CYP3A4							
	Phenytoin	Induction of CYP3A4							
<i>Enzyme Inhibitor</i>	Valproic acid		Increase in concentration of LPV (38 %) (DiCenzo et al., 2004) + +						
	Lamotrigine		Decrease in concentration of LTG (50 %) (van der Lee et al., 2006) + +						
<i>Others</i>	Midazolam		Increase in concentration of MDZ (by a factor of 25) (Greenblatt et al., 2009) + +						

B) List of drug-drug interactions with potential risk hypothesized based on literature-								
Antiseizure medication	COVID-19 candidate drug	Remdesivir ^c	Chloroquine/ Hydroxy-chloroquine ^a	Lopinavir ^b / Ritonavir	Ivermectin	Dexamethasone	Tocilizumab ^d	Anakinra ^e
	Relevant effect on enzymes		Inhibition of CYP3A4	Inhibition of CYP3A4 ^c		Inducer/ Inhibitor of CYP3A subfamily	Increase in CYP3A4 (indirect effect)	Increase in CYP3A4 (indirect effect)
<i>Enzyme Inducers</i>	Carbamazepine	Induction of CYP3A4	Decrease in concentration of RDV +	Decrease in concentration of CLQ/HCLQ + + +		Decrease in concentration of IVM ++	Decrease in concentration of CBZ +	Decrease in concentration of CBZ +
	Eslicarbazepine	Induction of CYP3A4		Decrease in concentration of CLQ/HCLQ + +	Decrease in concentration of LPV/RTV + +	Decrease in concentration of IVM ++	Decrease in concentration of DEX Decrease in concentration of ESL ++	Decrease in concentration of DEX Decrease in concentration of OXC ++
	Oxcarbazepine	Induction of CYP3A4		Decrease in concentration of CLQ/HCLQ + +	Decrease in concentration of LPV/RTV + +	Decrease in concentration of IVM ++	Decrease in concentration of DEX Decrease in concentration of OXC ++	
	Phenobarbital	Induction of CYP3A4	Decrease in concentration of RDV +	Decrease in concentration of CLQ/HCLQ + + +	Decrease in concentration of LPV/RTV + + +	Decrease in concentration of IVM ++	Decrease in concentration of PB +	Decrease in concentration of PB +
	Phenytoin	Induction of CYP3A4	Decrease in concentration of RDV +	Decrease in concentration of CLQ/HCLQ + + +	Decrease in concentration of LPV/RTV + + +	Decrease in concentration of IVM ++	Decrease in concentration of PHT +	Decrease in concentration of PHT +
	Primidone							

(continued on next page)

Table 2 (continued)

B) List of drug-drug interactions with potential risk hypothesized based on literature-								
	COVID-19 candidate drug Relevant effect on enzymes	Remdesivir ^c	Chloroquine/ Hydroxy-chloroquine ^a Inhibition of CYP3A4	Lopinavir ^b / Ritonavir Inhibition of CYP3A4 [†]	Ivermectin	Dexamethasone Inducer/ Inhibitor of CYP3A subfamily	Tocilizumab ^d Increase in CYP3A4 (indirect effect)	Anakinra ^e Increase in CYP3A4 (indirect effect)
	Induction of CYP3A4	Decrease in concentration of RDV	Decrease in concentration of CLQ/HCLQ	Decrease in concentration of LPV/RTV Decrease in concentration of PRM ^s	Decrease in concentration of IVM	Decrease in concentration of DEX Decrease in concentration of PRM	Decrease in concentration of PRM	Decrease in concentration of PRM
	Rufinamide	+	+++ Decrease in concentration of CLQ/HCLQ	+++ Decrease in concentration of LPV/RTV	++ Decrease in concentration of IVM	++ Decrease in concentration of DEX Decrease in concentration of RFN	+	+
	Cannabidiol	-	++ Increase in concentration of CLQ/HCLQ	++ Increase in concentration of CBD	++	++ Decrease concentration of corticosteroid [#]	-	-
	Clonazepam Clobazam Diazepam (BZDs)	-	++	++ Increase in concentration of BZDs	-	+	+	-
	Ethosuximide	-	-	++ Increase in concentration of ESM	-	-	-	-
Others	Felbamate	-	- Decrease in concentration of CLQ/HCLQ	++ Decrease in concentration of LPV/RTV	-	-	-	-
	Perampanel	-	++	++ Increase in concentration of PER	-	-	-	-
	Sultiame	-	-	++ Increase in concentration of STM	-	-	-	-
	Tiagabine	-	-	++ Increase in concentration of TGB	-	-	-	-
		-	-	++	-	-	-	-

Degree of drug-drug interactions:

Serious (+++)Potentially serious clinical consequences. Drugs not to be co-administered.

Moderate (+)Drugs that might require dose adjustment and periodic monitoring due to potential interaction.

Minor (+)Drugs expected to have potential weak interaction. Dose adjustment not necessary.

None (-)No clinical evidence exhibiting significant interaction.

enzymes that are relevant for a possible interaction between ASMs and COVID-19 candidate drugs based on the current state of knowledge.

For ASMs not mentioned in the above table including Gabapentin, Levetiracetam, Lorazepam, Pregabalin, Retigabine, Vigabatrin and Zonisamide, no significant evidence exists for drug-drug interactions with the COVID-19 candidate drugs.

BZD-Benzodiazepines; CLZ-Clonazepam; CBD-Cannabidiol; CBZ-Carbamazepine; CLQ/HCLQ-Chloroquine/ Hydroxychloroquine; DEX-Dexamethasone; ELS-Eslicarbazepine; ETS-Ethosuximide; IVM- Ivermectin; LPV/RTV-Lopinavir/Ritonavir; LTG-Lamotrigine; MDZ-Midazolam; OXC-Oxcarbazepine; PB-Phenobarbital; PER-Perampanel; PHT-Phenytoin; PRM-Primidone; RDV-Remdesivir; RFN- Rufinamide; STM-Sultiame; TGB-Tiagabine.

^a The information summarized in this table considered information provided by Russo and Iannone at the International League Against Epilepsy (ILAE) website (Russo, 2020) and has been updated with information released by the Liverpool Drug Interaction Group (University of Liverpool, UK, in collaboration with the University Hospital of Basel (Switzerland) and Radboud UMC (Netherlands)) (<http://www.covid19-druginteractions.org/>). Please note that we only provide information about the compound's effects on metabolic.

^a It is recommended that Chloroquine and Hydroxychloroquine are not administered in patients with recurring seizures. Both are metabolized by CYP3A4 and CYP2D6, and are also known to prolong QT interval and exhibit proarrhythmic potential and hence, their administration should in general be closely monitored.

^b Lopinavir is metabolized by CYP3A4.

^c There is only limited clinical information for the investigational drug Remdesivir. However, it might be necessary to consider evidence suggesting that remdesivir is sensitive to CYP3A4.

^d Tocilizumab causes suppression of IL-6 concentrations. IL-6 can reduce the expression of CYP3A4, CYP2C9 and CYP2C19. As a consequence disease-associated increases in IL-6 as well as pharmacological reduction and control of IL-6 concentrations can affect the rate of hepatic metabolism in the opposite direction.

^e Anakinra normalizes the increased concentration of metabolizing cytochromes (CYP450) due to inflammation, thereby it may decrease the systemic concentration of drugs metabolized by these enzymes.

^f Ritonavir is a potent inhibitor of CYP3A4.

^g Primidone is metabolized by CYP3A4 to phenobarbital, which induces CYP3A4.

^h Cannabidiol inhibits CYP3A4 and thus concomitant administration with glucocorticoids like hydrocortisone and prednisolone decrease glucocorticoid clearance, thereby increasing their systemic concentration (Wilson-Morkeah et al., 2019).

respective website. Hence, here we elaborate on the possible causes of the drug-drug interactions between commonly used ASMs and the COVID-19 candidate drugs.

3.1. Antiviral drugs

3.1.1. Remdesivir

Remdesivir is an investigational drug. Thus, only limited information is available about potential drug-drug interactions with remdesivir. It has been described that remdesivir is sensitive to CYP3A4 and hence, may act as a substrate to CYP3A4 enzyme (Streetman, 2020). However, considering the fact that it is metabolized predominantly by hydrolase *in vivo*, the risk for interaction with this nucleotide analogue compound is considered relatively low (Streetman, 2020).

Nevertheless, considering the limited clinical information, monitoring of remdesivir drug concentrations should be considered in case of co-administration of remdesivir with ASMs that induce CYP3A4 including carbamazepine, phenytoin, phenobarbital, primidone, oxcarbazepine, eslicarbazepine and rufinamide (Abou-Khalil, 2019).

3.1.2. Favipiravir

The active metabolite of favipiravir is metabolized in the liver by aldehyde oxidase and xanthine oxidase to form the inactive oxidative metabolite, T-705M1, which is then excreted *via* urine. There is insufficient to almost nil clinical data available for whether favipiravir or its active metabolite, T-705-RTP, affect hepatic metabolizing enzymes. However, pre-clinical studies have reported inhibitory effects of favipiravir on cytochrome isoenzyme, CYP2C8 (PMDA, 2014) and on aldehyde oxidase enzyme (Du and Chen, 2020). Available clinical data suggests that favipiravir has a low adverse effect potential and a favorable safety profile (Pilkington et al., 2020). Hence, even though favipiravir seems to have a low potential for drug-drug interaction, further studies are obviously required to evaluate the safety and tolerability of favipiravir, when co-administered with other drugs including ASM.

3.1.3. Lopinavir/ritonavir

Lopinavir is metabolized by the CYP3A group of cytochrome enzymes, whereas ritonavir is an inhibitor of CYP3A and, thus, is administered in combination with lopinavir to reduce the metabolism of lopinavir in plasma (Birbeck et al., 2012). This leads to a dual interaction between enzyme-inducing ASMs and lopinavir/ritonavir. The enzyme-induction property of ASMs such as carbamazepine, phenytoin, phenobarbital, primidone, oxcarbazepine, eslicarbazepine, felbamate and rufinamide, will lead to an increased metabolism of lopinavir, reducing its plasma concentration (Johannessen and Landmark, 2010). On the contrary, the enzyme inhibition property of ritonavir would block the metabolism of ASMs that are metabolized by CYP3A4, including carbamazepine, clonazepam, ethosuximide, perampanel, sultiame, and tiagabine, thereby, increasing their plasma concentration and consequently increasing their adverse effect potential (Abou-Khalil, 2019). Similarly, cannabidiol is also extensively metabolized by the cytochrome enzyme, CYP3A4, and hence, its co-administration with patients treated with ritonavir will lead to an increase in the pharmacologically active concentration of CBD in the blood plasma (Yamaori et al., 2011).

Since lopinavir/ritonavir are antiretroviral agents used in the treatment of HIV, clinical information is available for potential drug-drug interactions of these antiretroviral drugs with ASMs (Birbeck et al., 2012). A study with 24 patients receiving lamotrigine reported a 50 % decrease in the plasma concentration of lamotrigine following addition of lopinavir/ritonavir to the treatment schedule (van der Lee et al., 2006). In a cohort study, it was observed that co-administration of lopinavir/ritonavir with valproic acid, increased the plasma concentration of lopinavir by 38 % over a course of seven days (DiCenzo et al., 2004). A case report also confirmed carbamazepine toxicity due to the administration of ritonavir, whereby an 83 % dose reduction was

necessary to establish a therapeutic concentration of carbamazepine (Burman and Orr, 2000). A pharmacokinetic study conducted in healthy subjects reported increased concentrations of midazolam by a factor of 25 (400 %) in the presence of ritonavir (Greenblatt et al., 2009). The increase seems to be due to inhibition of intestinal CYP3A as a consequence of lopinavir/ritonavir co-administration (Yeh et al., 2006).

Thus, reciprocal drug-drug interactions need to be considered, when administering lopinavir and ritonavir in patients with exposure to different ASMs.

3.1.4. Ivermectin

Ivermectin has been administered in PWE as an adjuvant along with phenobarbital. The combination has even been encouraged in patients with onchocerciasis-associated epilepsy (Mandro et al., 2020).

However, ivermectin is known to be metabolized by the cytochrome CYP3A4 in human liver microsomes (Zeng et al., 1998). Hence, the plasma concentration of ivermectin should be monitored when co-administered with ASMs that induce CYP3A4 including carbamazepine, phenytoin, phenobarbital, primidone, oxcarbazepine, eslicarbazepine and rufinamide (Abou-Khalil, 2019).

3.2. Drugs with anti-inflammatory or immunomodulatory potential

3.2.1. Chloroquine/hydroxychloroquine

Both chloroquine and hydroxychloroquine are metabolized by cytochrome P450 (CYP450) enzymes, namely, CYP2C8, CYP3A4 and CYP2D6. They also act as inhibitors of CYP2D6 (Schrezenmeier and Dörner, 2020). Hence, co-administration of enzyme inducers or inhibitors will affect the plasma concentration of, both, chloroquine and hydroxychloroquine, resulting in the need for dose adjustments. ASMs such as carbamazepine, phenytoin, phenobarbital, primidone, oxcarbazepine, eslicarbazepine acetate, felbamate and rufinamide are all inducers of CYP450 enzymes including CYP3A4 (Johannessen and Landmark, 2010). Thereby, co-administration of these ASMs may lead to increased metabolism of chloroquine and hydroxychloroquine, which may require elevated dose administration of the latter. However, chloroquine and hydroxychloroquine are reported to possess a potential risk for QT prolongation, which has recently been also confirmed related to its use in patients with COVID-19 (Chorin et al., 2020; van den Broek et al., 2020). Thus, dose adjustment will be quite challenging considering respective tolerability issues.

Although the patient information leaflet states an increased risk of seizures associated with the intake of the drug, a systematic review suggests that there is no class I evidence to support this notion (Pati, 2020).

Furthermore, a critical drug-drug interaction may exist between chloroquine/hydroxychloroquine and cannabidiol (CBD), the latter being a potent inhibitor of the cytochrome P450 enzyme CYP3A4. The potential interaction might lead to increased plasma concentrations of chloroquine/hydroxychloroquine, thereby increasing the risk of arrhythmia in patients co-administered with these medications. (Browning, 2014; Yamaori et al., 2011).

In conclusion, the co-administration of ASMs including cannabidiol, carbamazepine, phenytoin, phenobarbital, primidone, oxcarbazepine, eslicarbazepine, felbamate and rufinamide, with chloroquine and hydroxychloroquine should rather be avoided. Dose adjustments would require extreme care considering the proarrhythmic potential for chloroquine and hydroxychloroquine.

3.2.2. Corticosteroids

Corticosteroids like dexamethasone can act as enzyme inducers for the CYP3A cytochrome family (Zhou, 2008). This effect is mediated by an interaction with cytosolic glucocorticoid receptors, which upon activation translocate to the nucleus, where they affect transcription levels of several target genes. Hence, the concomitant use of dexamethasone and ASMs that are metabolized by CYP3A cytochrome

(carbamazepine, phenytoin, phenobarbital, primidone, oxcarbazepine, eslicarbazepine and rufinamide) can result in decreased concentrations of the ASMs. Thus, it might be necessary to control and carefully adjust the dosing of ASMs.

Accordingly, clinical evidence indicates that co-administration of enzyme-inducing ASMs such as phenytoin (plasma concentration is decreased by 0.5), phenobarbital (clearance rate is decreased by 1.41 and 3.09) and carbamazepine (clearance rate is decreased by 1.79 and 4.42) can lead to an increase in the clearance time of corticosteroids with a subsequent decrease in their elimination half-life (Bénit and Vecht, 2015; Chalk et al., 1984; Griffiths and Taylor, 2018; Olivieri, 1986; Stjernholm and Katz, 1975). This suggests that administration of ASMs with corticosteroids, can cause a decrease in the plasma concentration due to increased metabolism of corticosteroids such as methylprednisolone, prednisolone, and dexamethasone (Bénit and Vecht, 2015).

3.2.3. Tocilizumab

Tocilizumab is a monoclonal antibody targeting cytokine IL-6 that is degraded by proteases. As expected, there is no evidence for any relevant direct interaction with cytochrome P450 enzymes. However, it may indirectly affect the expression of selected enzymes related to its effect on IL-6 function. It has previously been reported that increases in IL-6 concentrations can cause suppression of CYP450 enzymes such as CYP3A4, CYP2C19, CYP2C9 and CYP1A2 (Kim et al., 2012). As a consequence, concentrations of free drug substrates that are metabolized by these enzymes may increase (Abou-Khalil, 2019). This may also affect concentrations of ASMs such as carbamazepine, phenytoin, phenobarbital, clobazam, clonazepam, ethosuximide, tiagabine, topiramate, zonisamide. In turn, administration of tocilizumab will reduce IL-6 levels, thereby normalizing the expression of metabolizing cytochromes and the concentrations of affected drugs.

Taken together, tocilizumab possesses a weaker interaction potential than other COVID-19 candidate drugs and may not require dose adjustment or special monitoring, when co-administered with ASMs.

3.2.4. Anakinra

It has been reported that increased levels of cytokines such as IL-1 during chronic inflammation may result in reduced CYP450 enzyme activity. Thus, anakinra being an antagonist of IL-1 receptors, is theoretically expected to normalize CYP activity that was previously reduced due to release of inflammatory mediators. In view of this, dose-adjustments may be required for ASMs metabolized by CYP450 enzymes. However, we did not come across any clinical evidence for drug-drug interactions between ASMs and Anakinra (Bialer et al., 2018).

4. Potential drug interactions between ASMs and antipyretic and anti-inflammatory drugs concomitantly used in COVID-19 patients

One of the initial symptoms observed in patients with COVID-19 is fever. The first line therapy for the management of fever includes antipyretic drugs such as paracetamol (acetaminophen) (Kakodkar et al., 2020). The use of other drugs such as non-steroidal anti-inflammatory drugs (NSAIDs) has been controversially discussed in the context of COVID-19 management (Moore et al., 2020; Rinott et al., 2020). This discussion is for instance triggered by reports indicating that ibuprofen can increase the expression of ACE2 receptors (Fang et al., 2020) and can inhibit antibody production *in vitro* (Bancos et al., 2009).

However, the Food and Drug Administration later issued a statement addressing these reports and stated that there is no evidence for worsening of COVID-19 with the use of NSAIDs (FDA, 2020). Furthermore, according to the COVID-19 treatment guidelines provided by NIH, patients with COVID-19 are advised to continue taking NSAIDs as concomitant medication for any comorbid condition that has been previously directed by their physician. The expert panel has also

recommended that there is no difference in the use of anti-pyretic drugs including acetaminophen as well as NSAIDs between patients with or without COVID-19 (NIH, 2020).

NSAIDs such as ibuprofen, have been proposed to prevent the cytokine storm in patients with a severe disease course (Giollo et al., 2020). In particular, ibuprofen can reduce levels of the pro-inflammatory cytokine IL-6 (Gallelli et al., 2013). This effect is of interest as IL-6 levels can be elevated in patients with COVID-19 (Cao, 2020). Thus, effects of ibuprofen are also studied in ongoing clinical trials in patients with COVID-19 (Química Luar et al., 2021; University Hospital and Bordeaux, 2020).

Previous literature has reported various drug interactions between NSAIDs and commonly used anti-seizure drugs. A past study investigated the effects of administering antipyretic analgesics or NSAIDs including acetylsalicylic acid, phenylbutazone, paracetamol and tolfenamic acid in patients with epilepsy treated with phenytoin and carbamazepine (Neuvonen et al., 1979). Moreover, there have been several studies exploring a potential anti-convulsant action of NSAIDs given individually or in combination with anti-seizure drugs. Several of these studies pointed to a potential additive or synergistic interaction (Day et al., 2019; Srivastava and Gupta, 2001; Tandon et al., 2003; Zhu et al., 2017). This might be partly related to an impact on blood-brain barrier penetration of antiseizure drugs based on an inhibition of efflux transporter induction by NSAIDs (Potschka, 2010).

A strong drug-drug interaction potential needs to be considered, when deciding to combine an antipyretic analgesic drug or a NSAID in patients with epilepsy or acute reactive seizures.

As a consequence of paracetamol exposure, a clinical study reported a significant increase in the clearance of lamotrigine due to enhanced formation of glucuronide conjugates (Gastrup et al., 2016). The antagonistic interaction seems to be related to an interference of the drugs with uridine diphosphate glucuronosyltransferase (UGT), which catalyses the metabolism of lamotrigine and paracetamol through N-glucuronidation. Lamotrigine is metabolized by UGT enzymes, namely, UGT1A3, UGT1A4 and UGT2B7, while paracetamol is metabolized by UGT1A1, UGT1A6, UGT1A9 and UGT2B15 enzymes (Bock, 2010) (Zaccara and Perucca, 2014). Both, changes in expression and activity of enzymes or competitive inhibition may contribute to the specific interaction between paracetamol and lamotrigine. The clinical relevance was confirmed by a study reporting an enhanced potential for a decrease in lamotrigine exposure along with a detrimental effect on seizure control (Carnovale et al., 2019).

The co-administration of paracetamol with enzyme-inducing AEDs such as carbamazepine, phenobarbital, phenytoin and primidone is also not recommended due to a negative impact on paracetamol concentrations (Perucca, 2006). A drug interaction potential should also be considered for traditional NSAIDs and COX-2 inhibitors such as ibuprofen, diclofenac and celecoxib. These drugs are metabolized by the cytochrome enzyme, CYP2C9, that is induced by the above mentioned AEDs (Moore et al., 2015). Furthermore, clinical evidences suggest displacement of phenytoin, valproic acid and carbamazepine on administration of ibuprofen, thereby increasing the serum concentration of the anti-seizure drugs (Dasgupta and Volk, 1996; Nation et al., 1990). On the contrary phenobarbital is evidenced to induce metabolism of ibuprofen (Kantor, 1979).

Hence, following a careful weighing of potential beneficial and detrimental effects in individual patients, potential drug-drug interactions need to be kept in mind when co-administering NSAIDs with anti-seizure drugs in patients with epilepsy or acute reactive seizures.

5. Conclusion

Significant effort is currently underway to identify drugs that exert beneficial effects on the course and outcome of COVID-19. Considering the urgency associated with the rapid development of the pandemic, these efforts so far focus on the repurposing of licensed or investigational

drugs including compounds with possible antiviral effects and immunomodulators. While efficacy and tolerability of none of these drugs has been confirmed yet in sufficiently powered and controlled clinical studies, numerous clinical trials are currently underway worldwide. Intense care management of patients that develop seizures during a severe COVID-19 course as well as management of patients with epilepsy and SARS-CoV-2 infection requires information about possible drug-drug interactions between COVID-19 candidate drugs and ASMs. The data provided in this review suggest that potential drug-drug interactions should be taken into account when considering the combination of selected COVID-19 candidate drugs with ASMs. This in particular applies for chloroquine/hydroxychloroquine as well as lopinavir/ritonavir. In contrast, other possible COVID-19 treatment options such as tocilizumab that may be less prone to clinically relevant direct interactions with ASMs. Furthermore, remdesivir is suggested to have minor drug-drug interactions with ASMs since no clinical evidence is available that may state opposite. The outcome of ongoing and future clinical trials should be waited for final conclusions recommending for or against the drug options currently discussed for therapeutic management of COVID-19. In case of confirmation of efficacy and tolerability, depending on the compound, more research might be necessary to improve our state-of-knowledge regarding the use in patients with exposure to ASMs.

Data availability statement

This is a review article and data sharing is not applicable to this article because no new data were created or analyzed in this study.

References

- Abou-Khalil, B.W., 2019. Update on antiepileptic drugs 2019. *Contin. Lifelong Learn. Neurol.* 25, 508–536.
- Al-Tawfiq, J.A., Al-Homoud, A.H., Memish, Z.A., 2020. Remdesivir as a possible therapeutic option for the COVID-19. *Travel Med. Infect. Dis.*
- Amirian, E.S., Levy, J.K., 2020. Current knowledge about the antivirals remdesivir (GS-5734) and GS-441524 as therapeutic options for coronaviruses. *One Health* 9, 100128.
- Anon, 2021a. Interactions with Experimental COVID-19 Therapies. Liverpool Drug Interactions Group.
- Anon, 2021b. National Health Commission and State Administration of Traditional Chinese Medicine. Diagnosis and Treatment Protocol for COVID-19 Patients.
- Asadi-Pooya, A.A., 2020. Seizures associated with coronavirus infections. *Seizure* 79, 49–52.
- Asadi-Pooya, A.A., Simani, L., 2020. Central nervous system manifestations of COVID-19: a systematic review. *J. Neurol. Sci.* 413, 116832–116832.
- Bancos, S., Bernard, M.P., Topham, D.J., Phipps, R.P., 2009. Ibuprofen and other widely used non-steroidal anti-inflammatory drugs inhibit antibody production in human cells. *Cell. Immunol.* 258, 18–28.
- Barrows, N.J., Campos, R.K., Powell, S.T., Prasanth, K.R., Schott-Lerner, G., Soto-Acosta, R., Galarza-Muñoz, G., McGrath, E.L., Urrabaz-Garza, R., Gao, J., Wu, P., Menon, R., Saade, G., Fernandez-Salas, I., Rossi, S.L., Vasilakis, N., Routh, A., Bradrick, S.S., Garcia-Blanco, M.A., 2016. A screen of FDA-approved drugs for inhibitors of zika virus infection. *Cell Host Microbe* 20, 259–270.
- Bénit, C.P., Vecht, C.J., 2015. Seizures and cancer: drug interactions of anticonvulsants with chemotherapeutic agents, tyrosine kinase inhibitors and glucocorticoids. *Neurooncol. Pract.* 3, 245–260.
- Bialer, M., Johannessen, S.I., Koepp, M.J., Levy, R.H., Perucca, E., Tomson, T., White, H.S., 2018. Progress report on new antiepileptic drugs: a summary of the Fourteenth Eilat Conference on New Antiepileptic Drugs and Devices (EILAT XIV). I. Drugs in preclinical and early clinical development. *Epilepsia* 59, 1811–1841.
- Biggioggero, M., Crotti, C., Becciolini, A., Favalli, E.G., 2018. Tocilizumab in the treatment of rheumatoid arthritis: an evidence-based review and patient selection. *Drug Des. Dev. Ther.* 13, 57–70.
- Birbeck, G.L., French, J.A., Perucca, E., Simpson, D.M., Framow, H., George, J.M., Okulicz, J.F., Clifford, D.B., Hachad, H., Levy for the Quality Standards Subcommittee of the American Academy of Neurology, 2012. The ad hoc task force of the Commission on Therapeutic Strategies of the International League Against Epilepsy, R.H., 2012. Antiepileptic drug selection for people with HIV/AIDS: evidence-based guidelines from the ILAE and AAN. *Epilepsia* 53, 207–214.
- Bock, K.W., 2010. Functions and transcriptional regulation of adult human hepatic UDP-glucuronosyl-transferases (UGTs): mechanisms responsible for interindividual variation of UGT levels. *Biochem. Pharmacol.* 80, 771–777.
- Borku Uysal, B., Ikitimur, H., Yavuzer, S., Ikitimur, B., Uysal, H., Islamoglu, M.S., Ozcan, E., Aktepe, E., Yavuzer, H., Cengiz, M., 2020. Tocilizumab challenge: a series of cytokine storm therapy experiences in hospitalized COVID-19 pneumonia patients. *J. Med. Virol.* 92, 2648–2656.

- Bray, M., Rayner, C., Noël, F., Jans, D., Wagstaff, K., 2020. Ivermectin and COVID-19: a report in antiviral research, widespread interest, an FDA warning, two letters to the editor and the authors' responses. *Antiviral Res.* 178, 104805.
- Browning, D.J., 2014. *Pharmacology of Chloroquine and Hydroxychloroquine, Hydroxychloroquine and Chloroquine Retinopathy*. Springer, New York, New York, NY, pp. 35–63.
- Burman, W., Orr, L., 2000. Carbamazepine toxicity after starting combination antiretroviral therapy including ritonavir and efavirenz. *AIDS* 14.
- Cai, Q., Yang, M., Liu, D., Chen, J., Shu, D., Xia, J., Liao, X., Gu, Y., Cai, Q., Yang, Y., Shen, C., Li, X., Peng, L., Huang, D., Zhang, J., Zhang, S., Wang, F., Liu, J., Chen, L., Chen, S., Wang, Z., Zhang, Z., Cao, R., Zhong, W., Liu, Y., Liu, L., 2020. Experimental treatment with favipiravir for COVID-19: an open-label control study. *Engineering*.
- Caly, L., Druce, J.D., Catton, M.G., Jans, D.A., Wagstaff, K.M., 2020. The FDA-approved drug ivermectin inhibits the replication of SARS-CoV-2 in vitro. *Antiviral Res.* 178, 104787.
- Cao, X., 2020. COVID-19: immunopathology and its implications for therapy. *Nat. Rev. Immunol.*
- Cao, B., Wang, Y., Wen, D., Liu, W., Wang, J., Fan, G., Ruan, L., Song, B., Cai, Y., Wei, M., 2020. A trial of lopinavir-ritonavir in adults hospitalized with severe Covid-19. *N. Engl. J. Med.*
- Carnovale, C., Mosini, G., Gringeri, M., Battini, V., Mazhar, F., Pozzi, M., Clementi, E., Radice, S., 2019. Interaction between paracetamol and lamotrigine: new insights from the FDA Adverse Event Reporting System (FAERS) database. *Eur. J. Clin. Pharmacol.* 75, 1323–1325.
- Carod-Artal, F.J., 2020. Neurological complications of coronavirus and COVID-19. *Rev. Neurol.* 70, 311–322.
- Chaccour, C., Hammann, F., Ramón-García, S., Rabinovich, N.R., 2020. Ivermectin and COVID-19: keeping rigor in times of urgency. *Am. J. Trop. Med. Hyg.* 102, 1156–1157.
- Chalk, J., Ridgeway, K., Brophy, Tr., Yelland, J., Eadie, M., 1984. Phenytoin impairs the bioavailability of dexamethasone in neurological and neurosurgical patients. *J. Neurol. Neurosurg. Psychiatr.* 47, 1087–1090.
- Chen, C., Zhang, Y., Huang, J., Yin, P., Cheng, Z., Wu, J., Chen, S., Zhang, Y., Chen, B., Lu, M., Luo, Y., Ju, L., Zhang, J., Wang, X., 2020a. Favipiravir Versus Arbidol for COVID-19: A Randomized Clinical Trial. *medRxiv*, 2020.2003.2017.20037432.
- Chen, J., Liu, D., Liu, L., Liu, P., Xu, Q., Xia, L., Ling, Y., Huang, D., Song, S., Zhang, D., 2020b. A pilot study of hydroxychloroquine in treatment of patients with common coronavirus disease-19 (COVID-19). *J. Zhejiang Univ. (Med. Sci.)* 49, 0–0.
- Chen, Y., Liu, Q., Guo, D., 2020c. Emerging coronaviruses: genome structure, replication, and pathogenesis. *J. Med. Virol.* 92, 418–423.
- Chorin, E., Dai, M., Shulman, E., Wadhvani, L., Bar-Cohen, R., Barbhuiya, C., Aizer, A., Holmes, D., Bernstein, S., Spinelli, M., Park, D.S., Chinitz, L.A., Jankelson, L., 2020. The QT interval in patients with COVID-19 treated with hydroxychloroquine and azithromycin. *Nat. Med.*
- Cron, R.Q., Chatham, W.W., 2020. The rheumatologist's role in COVID-19. *J. Rheumatol.*
- Crump, A., Omura, S., 2011. Ivermectin, 'wonder drug' from Japan: the human use perspective. *Proc. Jpn. Acad. Ser. B Phys. Biol. Sci.* 87, 13–28.
- Dasgupta, A., Volk, A., 1996. Displacement of valproic acid and carbamazepine from protein binding in normal and uremic sera by tolmetin, ibuprofen, and naproxen: presence of inhibitor in uremic serum that blocks valproic acid-naproxen interactions. *Ther. Drug Monit.* 18, 284–287.
- Day, A.M., Hammill, A.M., Juhász, C., Pinto, A.L., Roach, E.S., McCulloch, C.E., Comi, A. M., Ball, K.L., Fisher, B.J., Koenig, J.I., 2019. Hypothesis: presymptomatic treatment of Sturge-Weber syndrome with aspirin and antiepileptic drugs may delay seizure onset. *Pediatr. Neurol.* 90, 8–12.
- DiCenzo, R., Peterson, D., Cruttenden, K., Morse, G., Riggs, G., Gelbard, H., Schiffling, G., 2004. Effects of valproic acid coadministration on plasma efavirenz and lopinavir concentrations in human immunodeficiency virus-infected adults. *Antimicrob. Agents Chemother.* 48, 4328–4331.
- Dilena, R., Mauri, E., Aronica, E., Bernasconi, P., Bana, C., Cappelletti, C., Carrabba, G., Ferrero, S., Giorda, R., Guez, S., Scalia Catenacci, S., Triulzi, F., Barbieri, S., Calderini, E., Vezzani, A., 2019. Therapeutic effect of Anakinra in the relapsing chronic phase of febrile infection-related epilepsy syndrome. *Epilepsia Open* 4, 344–350.
- Du, Y.-X., Chen, X.-P., 2020. Favipiravir: pharmacokinetics and concerns about clinical trials for 2019-nCoV infection. *Clin. Pharmacol. Ther.* 108, 242–247.
- Ebenso, B., 1998. Seizures following chloroquine treatment of type II lepra reaction: a case report. *Lepr. Rev.* 69, 178–181.
- El Baba, R., Herbein, G., 2020. Management of epigenomic networks entailed in coronavirus infections and COVID-19. *Clin. Epigenetics* 12, 118.
- Etgen, T., Eberl, B., Freudenberg, T., 2010. Anti-retroviral therapy-induced status epilepticus in "pseudo-HIV seroconversion". *Gen. Hosp. Psychiatry* 32 (559), e515–559 e517.
- Fang, L., Karakiulakis, G., Roth, M., 2020. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir. Med.* 8 e21–e21. FDA, 2020. FDA Advises Patients on Use of Non-steroidal Anti-inflammatory Drugs (NSAIDs) for COVID-19, p. 2020.
- Ferner, R.E., DeVito, N., Aronson, J.K., 2020. Dexamethasone. *CEBM*.
- Fox, R.I., 1993. Mechanism of action of hydroxychloroquine as an antirheumatic drug. *Semin. Arthritis Rheum.* 23, 82–91.
- Frisk-Holmberg, M., Bergqvist, Y., Englund, U., 1983. Chloroquine intoxication. *Br. J. Clin. Pharmacol.* 15, 502.
- Furuta, Y., Komeno, T., Nakamura, T., 2017. Favipiravir (T-705), a broad spectrum inhibitor of viral RNA polymerase. *Proc. Jpn. Acad. Ser. B* 93, 449–463.
- Gallelli, L., Galasso, O., Falcone, D., Southworth, S., Greco, M., Ventura, V., Romualdi, P., Corigliano, A., Terracciano, R., Savino, R., Gulletta, E., Gasparini, G., De Sarro, G., 2013. The effects of nonsteroidal anti-inflammatory drugs on clinical outcomes, synovial fluid cytokine concentration and signal transduction pathways in knee osteoarthritis. A randomized open label trial. *Osteoarthritis Cartil.* 21, 1400–1408.
- Gao, J., Tian, Z., Yang, X., 2020. Breakthrough: chloroquine phosphate has shown apparent efficacy in treatment of COVID-19 associated pneumonia in clinical studies. *Biosci. Trends.*
- Gastrup, S., Stage, T.B., Fruekilde, P.B.N., Damkier, P., 2016. Paracetamol decreases steady-state exposure to lamotrigine by induction of glucuronidation in healthy subjects. *Br. J. Clin. Pharmacol.* 81, 735–741.
- Gautret, P., Lagier, J.-C., Parola, P., Meddeb, L., Mailhe, M., Doudier, B., Courjon, J., Giordanengo, V., Vieira, V.E., Dupont, H.T., 2020. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *Int. J. Antimicrob. Agents*, 105949.
- Giollo, A., Adami, G., Gatti, D., Idolazzi, L., Rossini, M., 2020. Coronavirus disease 19 (Covid-19) and non-steroidal anti-inflammatory drugs (NSAID). *Ann. Rheum. Dis.*
- Götz, V., Magar, L., Dornfeld, D., Giese, S., Pohlmann, A., Höper, D., Kong, B.-W., Jans, D.A., Beer, M., Haller, O., 2016. Influenza A viruses escape from MxA restriction at the expense of efficient nuclear vRNP import. *Sci. Rep.* 6, 1–15.
- Greenblatt, D.J., Peters, D.E., Oleson, L.E., Harmatz, J.S., MacNab, M.W., Berkowitz, N., Zinny, M.A., Court, M.H., 2009. Inhibition of oral midazolam clearance by boosting doses of ritonavir, and by 4,4-dimethyl-benzoselenazine (ALT-2074), an experimental catalytic mimic of glutathione oxidase. *Br. J. Clin. Pharmacol.* 68, 920–927.
- Grein, J., Ohmagari, N., Shin, D., Diaz, G., Asperges, E., Castagna, A., Feldt, T., Green, G., Green, M.L., Lescure, F.-X., Nicastri, E., Oda, R., Yo, K., Quiros-Roldan, E., Studemeister, A., Redinski, J., Ahmed, S., Bernett, J., Chelliah, D., Chen, D., Chihara, S., Cohen, S.H., Cunningham, J., D'Arminio Monforte, A., Ismail, S., Kato, H., Lapadula, G., L'Her, E., Maeno, T., Majumder, S., Massari, M., Mora-Rillo, M., Mutoh, Y., Nguyen, D., Verweij, E., Zoufaly, A., Osinusi, A.O., DeZure, A., Zhao, Y., Zhong, L., Chokkalingam, A., Elboudwarej, E., Telep, L., Timbs, L., Henne, I., Sellers, S., Cao, H., Tan, S.K., Winterbourne, L., Desai, P., Mera, R., Gaggari, A., Myers, R.P., Brainard, D.M., Childs, R., Flanagan, T., 2020. Compassionate use of remdesivir for patients with severe Covid-19. *N. Engl. J. Med.*
- Griffiths, K., Taylor, T., 2018. Probable phenytoin-dexamethasone interaction: a case report of subtherapeutic phenytoin concentrations. *Can. J. Hosp. Pharm.* 52.
- Hoffmann, M., Kleine-Weber, H., Schroeder, S., Krüger, N., Herrler, T., Erichsen, S., Schiergens, T.S., Herrler, G., Wu, N.-H., Nitsche, A., 2020. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell*.
- Holshue, M.L., DeBolt, C., Lindquist, S., Lofy, K.H., Wiesman, J., Bruce, H., Spitters, C., Ericson, K., Wilkerson, S., Tural, A., 2020. First case of 2019 novel coronavirus in the United States. *N. Engl. J. Med.*
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., Cheng, Z., Yu, T., Xia, J., Wei, Y., Wu, W., Xie, X., Yin, W., Li, H., Liu, M., Xiao, Y., Gao, H., Guo, L., Xie, J., Wang, G., Jiang, R., Gao, Z., Jin, Q., Wang, J., Cao, B., 2020. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 395, 497–506.
- ILAE, 2020. FAQs About COVID-19 and Epilepsy for Clinicians in English.
- Ito, K., Yokoyama, T., Oguri, A., Kato, C., Horiuchi, M., Kato, M., Usami, I., 2021. Case Report a Case of COVID-19 Pneumonia That Did Not Worsen and Was Relieved by Early Administration of Favipiravir and Ciclesonide.
- Ivashchenko, A.A., Dmitriev, K.A., Vostokova, N.V., Azarova, V.N., Blinow, A.A., Egorova, A.N., Gordeev, I.G., Ilin, A.P., Karapetian, R.N., Kravchenko, D.V., 2020. AVIFAVIR for Treatment of Patients with Moderate COVID-19: Interim Results of a Phase II/III Multicenter Randomized Clinical Trial. *medRxiv*.
- Jafri, K., Zahed, H., Wysham, K.D., Patterson, S., Nolan, A.L., Bucknor, M.D., Chaganti, R.K., 2017. Antimalarial myopathy in a systemic lupus erythematosus patient with quadriplegia and seizures: a case-based review. *Clin. Rheumatol.* 36, 1437–1444.
- Johannessen, S.I., Landmark, C.J., 2010. Antiepileptic drug interactions - principles and clinical implications. *Curr. Neuropharmacol.* 8, 254–267.
- Kakodkar, P., Kaka, N., Baig, M.N., 2020. A comprehensive literature review on the clinical presentation, and management of the pandemic coronavirus disease 2019 (COVID-19). *Cureus* 12 e7560–e7560.
- Kantor, T.G., 1979. Ibuprofen. *Ann. Intern. Med.* 91, 877–882.
- Karlsen, A.P.H., Wiberg, S., Laigaard, J., Pedersen, C., Rokamp, K.Z., Mathiesen, O., 2020. A systematic review of trial registry entries for randomized clinical trials investigating COVID-19 medical prevention and treatment. *PLoS One* 15, e0237903.
- Kenney-Jung, D.L., Vezzani, A., Kahoud, R.J., LaFrance-Corey, R.G., Ho, M.-L., Muskardian, T.W., Wirrell, E.C., Howe, C.L., Payne, E.T., 2016. Febrile infection-related epilepsy syndrome treated with anakinra. *Ann. Neurol.* 80, 939–945.
- Kim, S., Östör, A.J., Nisar, M.K., 2012. Interleukin-6 and cytochrome-P450, reason for concern? *Rheumatol. Int.* 32, 2601–2604.
- Krzeminski, P., Lesiak, A., Narbutt, J., 2018. Seizures as a rare adverse effect of chloroquine therapy in systemic lupus erythematosus patients: a case report and literature survey. *Postepy Dermatol. Alergol.* 35, 429–430.
- Ksiazek, T.G., Erdman, D., Goldsmith, C.S., Zaki, S.R., Peret, T., Emery, S., Tong, S., Urbani, C., Comer, J.A., Lim, W., 2003. A novel coronavirus associated with severe acute respiratory syndrome. *N. Engl. J. Med.* 348, 1953–1966.
- Kujawski, S.A., Wong, K.K., Collins, J.P., Epstein, L., Killeber, M.E., Midgley, C.M., Abedi, G.R., Ahmed, N.S., Almendares, O., Alvarez, F.N., Anderson, K.N., Balter, S., Barry, V., Bartlett, K., Beer, K., Ben-Aderet, M.A., Benowitz, I., Biggs, H., Binder, A. M., Black, S.R., Bonin, B., Brown, C.M., Bruce, H., Bryant-Genevier, J., Budd, A., Buell, D., Bystritsky, R., Cates, J., Charles, E.M., Chatham-Stephens, K., Chea, N., Chiou, H., Christiansen, D., Chu, V., Cody, S., Cohen, M., Connors, E., Curns, A., Dasari, V., Dawson, P., DeSalvo, T., Diaz, G., Donahue, M., Donovan, S., Duca, L.M.,

- Erickson, K., Esona, M.D., Evans, S., Falk, J., Feldstein, L.R., Fensterseib, M., Fischer, M., Fisher, R., Foo, C., Fricchione, M.J., Friedman, O., Fry, A.M., Galang, R. R., Garcia, M.M., Gerber, S.L., Gerrard, G., Ghinai, I., Gounder, P., Grein, J., Grigg, C., Gunzenhauser, J.D., Gutkin, G.I., Haddix, M., Hall, A.J., Han, G., Harcourt, J., Harriman, K., Haupt, T., Haynes, A., Holshue, M., Hoover, C., Hunter, J. C., Jacobs, M.W., Jarashow, C., Jung, M.A., Joshi, K., Kamali, T., Kamili, S., Kim, L., Kim, M., King, J., Kirking, H.L., Kita-Yarbro, A., Klos, R., Kobayashi, M., Kocharian, A., Komatsu, K.K., Koppaka, R., Layden, J.E., Li, Y., Lindquist, S., Lindstrom, S., Link-Gelles, R., Lively, J., Livingston, M., Lo, K., Lo, J., Lu, X., Lynch, B., Madoff, L., Malapati, L., Marks, G., Marlow, M., Mathisen, G.E., McClung, N., McGovern, O., McPherson, T.D., Mehta, M., Meier, A., Mello, L., Moon, S.-s., Morgan, M., Moro, R.N., Murray, J., Murthy, R., Novosad, S., Oliver, S. E., Shea, J., Pacilli, M., Paden, C.R., Pallansal, M.A., Patel, M., Patel, S., Pedraza, I., Pillai, S.K., Pindyck, T., Pray, I., Queen, K., Quick, N., Reese, H., Rha, B., Rhodes, H., Robinson, S., Robinson, P., Rolfes, M., Routh, J., Rubin, R., Rudman, S.L., Sakthivel, S.K., Scott, S., Shepherd, C., Shetty, V., Smith, E.A., Smith, S., Stierman, B., Stoecker, W., Sunenshine, R., Sy-Santos, R., Tamin, A., Tao, Y., Terashita, D., Thornburg, N.J., Tong, S., Traub, E., Tural, A., Uehara, A., Uyeki, T.M., Vahe, G., Verani, J.R., Villarino, E., Wallace, M., Wang, L., Watson, J.T., Westercamp, M., Whitaker, B., Wilkerson, S., Woodruff, R.C., Wortham, J.M., Wu, T., Xie, A., Yousaf, A., Zahn, M., Zhang, J., 2020. First 12 Patients with Coronavirus Disease 2019 (COVID-19) in the United States. medRxiv, 2020.2003.2009.20032896.
- Li, X., Wang, Y., Agostinis, P., Rabson, A., Melino, G., Carafoli, E., Shi, Y., Sun, E., 2020a. Is hydroxychloroquine beneficial for COVID-19 patients? *Cell Death Dis.* 11, 512.
- Li, Y.C., Bai, W.Z., Hashikawa, T., 2020b. The neuroinvasive potential of SARS-CoV2 may play a role in the respiratory failure of COVID-19 patients. *J. Med. Virol.* 92, 552–555.
- Malcangi, G., Fraticelli, P., Palmieri, C., Cappelli, M., Danieli, M., 2000. Hydroxychloroquine-induced seizure in a patient with systemic lupus erythematosus. *Rheumatol. Int.* 20, 31–33.
- Mandro, M., Siewe Fodjo, J.N., Mukendi, D., Dusabimana, A., Menon, S., Haesendonckx, S., Lokonda, R., Nakato, S., Nyisi, F., Abhafule, G., Wonya' Rossi, D., Jakwong, J.M., Suykerbuyk, P., Meganck, J., Hotterbeek, A., Colebunders, R., 2020. Ivermectin as an adjuvant to anti-epileptic treatment in persons with onchocerciasis-associated epilepsy: a randomized proof-of-concept clinical trial. *PLoS Negl. Trop. Dis.* 14, e0007966.
- Medina, M.T., Moncada, S.S., 2020. Hydroxychloroquine/ chloroquine as a treatment choice or prophylaxis for Covid-19 at the primary care level in developing countries: a Primum non Nocere dilemma. *J. Neurol. Sci.* 415.
- Moore, N., Pollack, C., Butkeraif, P., 2015. Adverse drug reactions and drug-drug interactions with over-the-counter NSAIDs. *Ther. Clin. Risk Manag.* 11, 1061–1075.
- Moore, N., Carleton, B., Blin, P., Bosco-Levy, P., Droz, C., 2020. Does Ibuprofen Worsen COVID-19? *Drug Saf.* 43, 611–614.
- Najjar, S., Najjar, A., Chong, D.J., Pramanik, B.K., Kirsch, C., Kuzniacki, R.I., Pacia, S.V., Azhar, S., 2020. Central nervous system complications associated with SARS-CoV-2 infection: integrative concepts of pathophysiology and case reports. *J. Neuroinflammation* 17, 231–231.
- Nation, R.L., Evans, A.M., Milne, R.W., 1990. Pharmacokinetic drug interactions with phenytoin (Part I). *Clin. Pharmacokinet.* 18, 37–60.
- Neuvonen, P., Lehtovaara, R., Bardy, A., Elomaa, E., 1979. Antipyretic analgesics in patients on antiepileptic drug therapy. *Eur. J. Clin. Pharmacol.* 15, 263–268.
- Nicolau, D.V., Bafadhel, M., 2020. Inhaled corticosteroids in virus pandemics: a treatment for COVID-19? *Lancet Respir. Med.*
- Nie, S., Zhao, X., Zhao, K., Zhang, Z., Zhang, Z., Zhang, Z., 2020. Metabolic Disturbances and Inflammatory Dysfunction Predict Severity of Coronavirus Disease 2019 (COVID-19): A Retrospective Study. medRxiv, 2020.2003.2024.20042283.
- NIH, 2020. COVID-19 Treatment Guidelines. National Institutes of Health.
- Noda, A., Shirai, T., Nakajima, H., Oda, M., Saraya, T., Ishii, H., Takizawa, H., Shimasaki, T., Sano, A., Kurai, D., Ito, E., Tashima, Y., Hada, Y., Goto, T., Masuko, S., Katsumata, A., Nakajima, M., Ichimura, S., 2020. Two cases of COVID-19 pneumonia including use of favipiravir. *Jpn. Assoc. Infect. Dis.*
- Nukoolkarn, V., Lee, V.S., Malaisree, M., Aruksakulwong, O., Hannongbua, S., 2008. Molecular dynamic simulations analysis of ritonavir and lopinavir as SARS-CoV 3CL (pro) inhibitors. *J. Theor. Biol.* 254, 861–867.
- Olivesi, A., 1986. Modified elimination of prednisolone in epileptic patients on carbamazepine monotherapy, and in women using low-dose oral contraceptives. *Biomed. Pharmacother.* 40, 301–308.
- Omura, S., Crump, A., 2014. Ivermectin: panacea for resource-poor communities? *Trends Parasitol.* 30, 445–455.
- Otto, S.B.J., George, P.E., Mercedes, R., Nabukeera-Barungi, N., 2020. Cryptococcal meningitis and immune reconstitution inflammatory syndrome in a pediatric patient with HIV after switching to second line antiretroviral therapy: a case report. *BMC Infect. Dis.* 20, 68–68.
- Pati, S., 2020. A Systematic Review to Assess Seizure Risk with Chloroquine Therapy in Persons with Epilepsy. medRxiv, 2020.2004.2009.20056358.
- Patsalos, P.N., Spencer, E.P., Berry, D.J., 2018. Therapeutic drug monitoring of antiepileptic drugs in epilepsy: a 2018 update. *Ther. Drug Monit.* 40, 526–548.
- Perucca, E., 2006. Clinically relevant drug interactions with antiepileptic drugs. *Br. J. Clin. Pharmacol.* 61, 246–255.
- Pilkington, V., Pepperrell, T., Hill, A., 2020. A review of the safety of favipiravir - a potential treatment in the COVID-19 pandemic? *J. Virus Erad.* 6, 45–51.
- PMDA, 2014. Pharmaceuticals and Medical Devices Agency: Avigan (favipiravir). Review Report.
- Potschka, H., 2010. Modulating P-glycoprotein regulation: Future perspectives for pharmacoresistant epilepsies? *Epilepsia* 51, 1333–1347.
- Química Luar, S.R.L., National Council of, S, Technical Research, A, Centro de Excelencia en Productos y Procesos, C, 2021. Inhaled Ibuprofen to Treat COVID-19.
- Rajter, J.C., Sherman, M., Fatteh, N., Vogel, F., Sacks, J., Rajter, J.-J., 2020. ICON (Ivermectin in COVID Nineteen) study: Use of Ivermectin is Associated with Lower Mortality in Hospitalized Patients with COVID19. medRxiv, 2020.2006.2006.20124461.
- Richman, D.D., Whitley, R.J., Hayden, F.G., 2016. *Clinical Virology*. John Wiley & Sons.
- Rinott, E., Kozler, E., Shapira, Y., Bar-Haim, A., Youngster, I., 2020. Ibuprofen use and clinical outcomes in COVID-19 patients. *Clin. Microbiol. Infect.* 26 (1259), e1255–1259 e1257.
- Rose-John, S., Winthrop, K., Calabrese, L., 2017. The role of IL-6 in host defence against infections: immunobiology and clinical implications. *Nat. Rev. Rheumatol.* 13, 399–409.
- Rothan, H.A., Byrareddy, S.N., 2020. The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak. *J. Autoimmun.* 102433
- Russo, E.L.L., 2020. Clinically Relevant Drug-Drug Interaction Between AEDs and Medications Used in the Treatment of COVID-19 Patients.
- Sa, M., Singh, R., Pujar, S., D'Arco, F., Desai, N., Eltze, C., Hughes, E., Al Obaidi, M., Eleftheriou, D., Tisdall, M., Selway, R., Cross, J.H., Kaliakatos, M., Valentin, A., 2019. Centromedian thalamic nuclei deep brain stimulation and Anakinra treatment for FIRES – two different outcomes. *Eur. J. Paediatr. Neurol.* 23, 749–754.
- Sanders, J.M., Monogue, M.L., Jodowski, T.Z., Cutrell, J.B., 2020. Pharmacologic treatments for coronavirus disease 2019 (COVID-19): a review. *JAMA*.
- Savarino, A., Boelaert, J.R., Cassone, A., Majori, G., Cauda, R., 2003. Effects of chloroquine on viral infections: an old drug against today's diseases. *Lancet Infect. Dis.* 3, 722–727.
- Schiemann, R., Coulaud, J.P., Bouchaud, O., 2000. Seizures after antimalarial medication in previously healthy persons. *J. Travel Med.* 7, 155.
- Schrezenmeier, E., Dörner, T., 2020. Mechanisms of action of hydroxychloroquine and chloroquine: implications for rheumatology. *Nat. Rev. Rheumatol.* 1–12.
- Sheahan, T.P., Sims, A.C., Graham, R.L., Menachery, V.D., Gralinski, L.E., Case, J.B., Leist, S.R., Pyrc, K., Feng, J.Y., Trantcheva, I., 2017. Broad-spectrum antiviral GS-5734 inhibits both epidemic and zoonotic coronaviruses. *Sci. Transl. Med.* 9.
- Shi, Y., Tan, M., Chen, X., Liu, Y., Huang, J., Ou, J., Deng, X., 2020. Immunopathological Characteristics of Coronavirus Disease 2019 Cases in Guangzhou, China. medRxiv.
- Siegel, D., Hui, H.C., Doerfler, E., Clarke, M.O., Chun, K., Zhang, L., Neville, S., Carra, E., Lew, W., Ross, B., 2017. Discovery and synthesis of a phosphoramidate prodrug of a pyrrolo [2, 1-f]triazin-4-amino] adenine C-Nucleoside (GS-5734) for the treatment of ebola and emerging viruses. *ACS Publ.*
- Singh, A.K., Majumdar, S., Singh, R., Misra, A., 2020. Role of corticosteroid in the management of COVID-19: a systemic review and a Clinician's perspective. *Diabetes Metab. Syndr. Clin. Res. Rev.* 14, 971–978.
- Sissoko, D., Laouenan, C., Folkesson, E., M'lebing, A.-B., Beavogui, A.-H., Baize, S., Camara, A.-M., Maes, P., Shepherd, S., Danel, C., 2016. Experimental treatment with favipiravir for Ebola virus disease (the JIKI Trial): a historically controlled, single-arm proof-of-concept trial in Guinea. *PLoS Med.* 13, e1001967.
- Srivastava, A.K., Gupta, Y.K., 2001. Aspirin modulates the anticonvulsant effect of diazepam and sodium valproate in pentylenetetrazole and maximal electroshock induced seizures in mice. *Indian J. Physiol. Pharmacol.* 45, 475–480.
- Stjernholm, M.R., Katz, F.H., 1975. Effects of Diphenylhydantoin, phenobarbital, and diazepam on the metabolism of methylprednisolone and its sodium succinate. *J. Clin. Endocrinol. Metab.* 41, 887–893.
- Streetman, D.S., 2020. Drug Interaction Concerns for COVID-19 Treatments. Wolters Kluwer.
- Tandon, M., Anuradha, K., Pandhi, P., 2003. Evaluation of antiepileptic activity of aspirin in combination with newer antiepileptic lamotrigine in mice. *Methods Find. Exp. Clin. Pharmacol.* 25, 607–610.
- Tristano, A.G., Falcón, L., Willson, M., de Oca, I.M., 2004. Seizure associated with chloroquine therapy in a patient with systemic lupus erythematosus. *Rheumatol. Int.* 24, 315–316.
- Tu, Y.-F., Chien, C.-S., Yarmishyn, A.A., Lin, Y.-Y., Luo, Y.-H., Lin, Y.-T., Lai, W.-Y., Yang, D.-M., Chou, S.-J., Yang, Y.-P., Wang, M.-L., Chiou, S.-H., 2020. A review of SARS-CoV-2 and the ongoing clinical trials. *Int. J. Mol. Sci.* 21.
- University Hospital, B., Bordeaux, P., 2020. Role of Ibuprofen and Other Medicines on Severity of Coronavirus Disease, p. 2019.
- van den Broek, M.P.H., Möhlmann, J.E., Abeln, B.G.S., Liebgrens, M., van Dijk, V.F., van de Garde, E.M.W., 2020. Chloroquine-induced QTc prolongation in COVID-19 patients. *Netherlands Heart J.*
- van der Lee, M.J., Dawood, L., ter Hofstede, H.J., de Graaff-Teulen, M.J., van Ewijk-Beneken Kolmer, E.W., Caliskan-Yassen, N., Koopmans, P.P., Burger, D.M., 2006. Lopinavir/ritonavir reduces lamotrigine plasma concentrations in healthy subjects. *Clin. Pharmacol. Ther.* 80, 159–168.
- Varadharajan, P., 2020. Favipiravir Approved As Anti- COVID-19 Drug; DCGI Approves Influenza Treatment Drug for Covid-19 Treatment in India. *Inventiva*.
- Vincent, M.J., Bergeron, E., Benjannet, S., Erickson, B.R., Rollin, P.E., Ksiazek, T.G., Seidah, N.G., Nichol, S.T., 2005. Chloroquine is a potent inhibitor of SARS coronavirus infection and spread. *Virology* 339, 72–83.
- Wagstaff, K.M., Sivakumaran, H., Heaton, S.M., Harrich, D., Jans, D.A., 2012. Ivermectin is a specific inhibitor of importin α/β -mediated nuclear import able to inhibit replication of HIV-1 and dengue virus. *Biochem. J.* 443, 851–856.
- Wang, H., 2021. Diagnosis and treatment protocol for COVID-19 patients (Tentative 8th Edition). *Infect. Dis. Immun.* 1 (1), 8–16. <https://doi.org/10.1097/01.ID9.0000733564.21786.b0>.
- Wang, M., Cao, R., Zhang, L., Yang, X., Liu, J., Xu, M., Shi, Z., Hu, Z., Zhong, W., Xiao, G., 2020a. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res.* 30, 269–271.

- Wang, Y., Jiang, W., He, Q., Wang, C., Wang, B., Zhou, P., Dong, N., Tong, Q., 2020b. Early, low-dose and short-term application of corticosteroid treatment in patients with severe COVID-19 pneumonia: single-center experience from Wuhan, China. medRxiv, 2020.2003.2006.20032342.
- Warren, T.K., Jordan, R., Lo, M.K., Ray, A.S., Mackman, R.L., Soloveva, V., Siegel, D., Perron, M., Bannister, R., Hui, H.C., Larson, N., Strickley, R., Wells, J., Stuthman, K. S., Van Tongeren, S.A., Garza, N.L., Donnelly, G., Shurtleff, A.C., Retterer, C.J., Gharabeh, D., Zamani, R., Kenny, T., Eaton, B.P., Grimes, E., Welch, L.S., Gomba, L., Wilhelmson, C.L., Nichols, D.K., Nuss, J.E., Nagle, E.R., Kugelman, J.R., Palacios, G., Doerffler, E., Neville, S., Carra, E., Clarke, M.O., Zhang, L., Lew, W., Ross, B., Wang, Q., Chun, K., Wolfe, L., Babusis, D., Park, Y., Stray, K.M., Trancheva, I., Feng, J.Y., Barauskas, O., Xu, Y., Wong, P., Braun, M.R., Flint, M., McMullan, L.K., Chen, S.-S., Fearn, R., Swaminathan, S., Mayers, D.L., Spiropoulou, C.F., Lee, W.A., Nichol, S.T., Cihlar, T., Bavari, S., 2016. Therapeutic efficacy of the small molecule GS-5734 against Ebola virus in rhesus monkeys. *Nature* 531, 381–385.
- Wilson-Morkeh, H., Al-Abdulla, A., Sien, L., Mohamed, H., Youngstein, T., 2019. Important drug interactions exist between cannabidiol oil and commonly prescribed drugs in rheumatology practice. *Rheumatology* 59, 249–251.
- Xu, X., Han, M., Li, T., Sun, W., Wang, D., Fu, B., Zhou, Y., Zheng, X., Yang, Y., Li, X., 2020a. Effective treatment of severe COVID-19 patients with tocilizumab. *ChinaXiv* 202003, V1.
- Xu, Z., Shi, L., Wang, Y., Zhang, J., Huang, L., Zhang, C., Liu, S., Zhao, P., Liu, H., Zhu, L., 2020b. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir. Med.* 8, 420–422.
- Yamaori, S., Ebisawa, J., Okushima, Y., Yamamoto, I., Watanabe, K., 2011. Potent inhibition of human cytochrome P450 3A isoforms by cannabidiol: role of phenolic hydroxyl groups in the resorcinol moiety. *Life Sci.* 88, 730–736.
- Yang, J.W., Yang, L., Luo, R.G., Xu, J.F., 2020a. Corticosteroid administration for viral pneumonia: COVID-19 and beyond. *Clin. Microbiol. Infect.* 26, 1171–1177.
- Yang, S.N., Atkinson, S.C., Wang, C., Lee, A., Bogoyevitch, M.A., Borg, N.A., Jans, D.A., 2020b. The broad spectrum antiviral ivermectin targets the host nuclear transport importin $\alpha/\beta 1$ heterodimer. *Antiviral Res.* 104760
- Yao, T.T., Qian, J.D., Zhu, W.Y., Wang, Y., Wang, G.Q., 2020a. A systematic review of lopinavir therapy for SARS coronavirus and MERS coronavirus—a possible reference for coronavirus disease-19 treatment option. *J. Med. Virol.*
- Yao, X., Ye, F., Zhang, M., Cui, C., Huang, B., Niu, P., Liu, X., Zhao, L., Dong, E., Song, C., 2020b. In vitro antiviral activity and projection of optimized dosing design of hydroxychloroquine for the treatment of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). *Clin. Infect. Dis.*
- Yeh, R.F., Gaver, V.E., Patterson, K.B., Rezk, N.L., Baxter-Meheux, F., Blake, M.J., Eron Jr, J.J., Klein, C.E., Rublein, J.C., Kashuba, A.D., 2006. Lopinavir/ritonavir induces the hepatic activity of cytochrome P450 enzymes CYP2C9, CYP2C19, and CYP1A2 but inhibits the hepatic and intestinal activity of CYP3A as measured by a phenotyping drug cocktail in healthy volunteers. *JAIDS Journal of Acquired Immune Deficiency Syndromes* 42, 52–60.
- Yin, R., Feng, W., Wang, T., Chen, G., Wu, T., Chen, D., Lv, T., Xiang, D., 2020. Concomitant neurological symptoms observed in a patient diagnosed with coronavirus disease 2019. *J. Med. Virol.*
- Zaccara, G., Perucca, E., 2014. Interactions between antiepileptic drugs, and between antiepileptic drugs and other drugs. *Epileptic Disord.* 16, 409–431.
- Zaki, A.M., Van Boheemen, S., Bestebroer, T.M., Osterhaus, A.D., Fouchier, R.A., 2012. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N. Engl. J. Med.* 367, 1814–1820.
- Zeng, Z., Andrew, N.W., Arison, B.H., Luffer-Atlas, D., Wang, R.W., 1998. Identification of cytochrome P4503A4 as the major enzyme responsible for the metabolism of ivermectin by human liver microsomes. *Xenobiotica* 28, 313–321.
- Zhou, S.-F., 2008. Drugs behave as substrates, inhibitors and inducers of human cytochrome P450 3A4. *Curr. Drug Metab.* 9, 310–322.
- Zhou, F., Yu, T., Du, R., Fan, G., Liu, Y., Liu, Z., Xiang, J., Wang, Y., Song, B., Gu, X., Guan, L., Wei, Y., Li, H., Wu, X., Xu, J., Tu, S., Zhang, Y., Chen, H., Cao, B., 2020. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 395, 1054–1062.
- Zhu, K., Hu, M., Yuan, B., Liu, J.-X., Liu, Y., 2017. Aspirin attenuates spontaneous recurrent seizures in the chronically epileptic mice. *Neurol. Res.* 39, 744–757.