

Article

A Structural Equation Modeling Analysis to Explore Diabetes Self-Care Factors in a Rural Sample

Laurie Abbott ^{1,*} , Lucinda Graven ¹, Glenna Schluck ¹  and Jennifer Lemacks ² ¹ College of Nursing, Florida State University, Tallahassee, FL 32306, USA² College of Nursing and Health Professions, University of Southern Mississippi, Hattiesburg, MS 39406, USA

* Correspondence: labbott@fsu.edu

Abstract: Diabetes is a public health problem that requires management to avoid health sequelae. Little is known about the determinants that influence diabetes self-care activities among rural populations. The purpose of this analysis was to explore the relationships among diabetes self-care activities, diabetes knowledge, perceived diabetes self-management, diabetes fatalism, and social support among an underserved rural group in the southern United States. A diabetes health promotion program was tested during a cluster randomized trial that tested a disease risk reduction program among adults living with prediabetes and diabetes. A structural equation model was fit to test psychosocial factors that influence diabetes self-care activities using the Information–Motivation–Behavioral Skills Model of Diabetes Self-Care (IMB-DSC) to guide the study. Perceived diabetes self-management significantly predicted self-care behaviors, and there was also a correlation between perceived diabetes self-management and diabetes fatalism. Perceived diabetes self-management influenced diabetes self-care activities in this rural sample and had an association with diabetes fatalism. The findings of this study can facilitate clinical care and community programs targeting diabetes and advance health equity among underserved rural groups.

Keywords: rural; diabetes; health promotion; disease risk; diabetes self-management



Citation: Abbott, L.; Graven, L.; Schluck, G.; Lemacks, J. A Structural Equation Modeling Analysis to Explore Diabetes Self-Care Factors in a Rural Sample. *Healthcare* **2022**, *10*, 1536. <https://doi.org/10.3390/healthcare10081536>

Academic Editor: Ioannis Ilias

Received: 11 July 2022

Accepted: 9 August 2022

Published: 14 August 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Chronic diseases such as heart disease, stroke, cancer, and diabetes are the leading causes of death, disability, and high health care costs [1,2]. In fact, approximately 6 out of 10 adults living in the United States have been diagnosed with at least one chronic disease, and 4 out of 10 have at least two chronic diseases [1,2]. Chronic diseases have been associated with risk behaviors including tobacco and alcohol use, physical inactivity, insufficient sleep, stress, and poor nutrition [2]. However, diabetes is an underrecognized contributory risk factor for chronic diseases such as cardiovascular disease, poor health outcomes, and increased mortality [3,4]. Diabetes is prevalent among rural populations in the southern United States, where life expectancy rates are the lowest in the nation, especially among African Americans who have the highest rates of chronic disease prevalence, morbidity, and mortality [4–13].

Populations living in rural areas bear a disproportionate chronic disease burden [7,8,10,14–17]. Compared with urban groups, those living in rural geographic locations have poorer health outcomes, fewer healthcare options, higher rates of premature death, and greater chronic disease risk factors [16–18]. Resource limitations, deficits in disease knowledge, and limited access to primary and preventive care further exacerbate disease risk [4,6,8,11,12,19,20]. In addition to these factors, worse outcomes among rural populations living with poorly controlled diabetes have been associated with their lack of adherence to disease management recommendations [21–23]. Poor glycemic control is associated with higher glycosylated hemoglobin (HbA1C) levels and contributes to the development and progression of chronic conditions such as heart disease and stroke [3,4].

Health outcomes associated with diabetes among rural, southern groups can potentially be improved through health promotion and disease risk reduction interventions that promote healthier lifestyle choices, improved knowledge about diseases and ways to reduce risk, and strengthen protective psychosocial factors [9,24,25]. However, there is a gap in knowledge regarding the determinants that influence diabetes self-care and management among diverse populations living in rural, southern states [22,26–28]. The Information–Motivation–Behavioral Skills Model of Diabetes Self-Care (IMB-DSC) was the theoretical framework used to guide this study, which indicates that performing diabetes self-care activities is influenced by psychosocial factors including diabetes knowledge, perceived diabetes self-management, diabetes fatalism, and social support [29]. Diabetes fatalism is described as the emotional distress from daily living with diabetes, coping using religious and spiritual resources, and self-efficacy in managing diabetes [30]. Social support is associated with improved diabetes self-management and outcomes [31–34]. More diabetes knowledge and social support and less diabetes fatalism increase the probability that diabetes self-care behaviors linked to glycemic control would be performed [29]. The purpose of this analysis was to explore the relationships among diabetes self-care activities, diabetes knowledge, perceived diabetes self-management, diabetes fatalism, and social support.

2. Materials and Methods

This study involved an analysis of data collected in a cluster randomized trial to test a culturally relevant disease risk reduction curriculum among people living with diabetes and prediabetes (ClinicalTrials.gov: NCT04795050). The study included 12 participating churches situated in a rural area of the southeastern United States, and the primary results showed statistically significant intervention effects for diabetes knowledge and some of the measured self-care activities [35]. The study received institutional review board approval at Florida State University.

2.1. Sample and Setting

The design of the cluster randomized trial involved calculating the number of participating churches and individual participants within churches needed to account for potential intra-cluster correlation [35]. For example, individuals within a participating church may share similar characteristics or kinship bonds that could have the effect of inducing correlation among study outcomes [36]. A conservative intra-correlation value ($r = 0.008$) from previous health research [37] and a medium standardized effect size ($d = 0.50$) from a similar study [38] were used to determine the sample size. For sufficient power (80%), the study required the inclusion of at least 5 different churches and at least 71 individual participants, accounting for 10% attrition, for each of the two groups. Serving as the statistical cluster, the rural churches situated in non-metropolitan areas as classified by ZIP codes were randomized to intervention and control groups after the pastors expressed interest in study participation. Randomization involved using a randomly selected five-digit random number sequence and assigning each church to either an intervention (even number) or control (odd number) group while ensuring the parity imbalance did not exceed two. The individual study participants were recruited from within the participating church congregations and self-identified as African American, were at least 22 years old, had previously been diagnosed with either diabetes or prediabetes, and could understand and speak English. Those deemed eligible and willing to participate provided written informed consent. The participants recruited from churches allocated to the intervention group received a culturally relevant diabetes risk reduction intervention, and those in churches randomized to the control group received a patient education sheet.

2.2. Measures

Participants in the intervention group completed the measures at baseline and post-intervention following the third weekly session while participants in the control group

completed baseline measures and attended a second data collection period three weeks later. Sociodemographic information was collected at baseline, and the items included age, gender, employment status, educational attainment, previous diagnosis of either diabetes or prediabetes, diabetes management regimen, family history of heart disease or diabetes, and personal history of diabetic retinopathy. The self-reported measures for diabetes self-care activities, diabetes fatalism, perceived diabetes self-management, social support, and diabetes knowledge were collected during each data collection period.

2.2.1. Diabetes Self-Care Activities

Diabetes self-care activities were measured using the Summary of Diabetes Self-care Activities, which had adequate internal consistency ($\alpha = 0.71$) [39]. The instrument contains 15 items that measure self-care activities, such as blood sugar testing, dietary factors, medications, and foot care, using an 8-point Likert scale for the number of days (0–7) the behavior had been performed during the last week.

2.2.2. Diabetes Fatalism

Diabetes fatalism was measured using the Diabetes Fatalism Scale, a 12-item, 6-point Likert scale that had good overall internal consistency ($\alpha = 0.80$) [30]. The constructs of diabetes fatalism that were included in the instrument were the emotional distress ($\alpha = 0.86$), religious and spiritual coping ($\alpha = 0.77$), and perceived self-efficacy ($\alpha = 0.77$) subscales. Greater diabetes fatalism was indicated by higher scores (Range 12–72) on the overall instrument, including these subscales.

2.2.3. Perceived Diabetes Self-Management

Perceptions of diabetes self-management were measured using the Perceived Diabetes Self-Management Scale (PDSMS), which has eight questions and Likert scale-type answer options with total score possibilities ranging from 8 to 40 [40]. The instrument had good internal consistency ($\alpha = 0.83$), with higher scores indicating a greater level of confidence in diabetes self-management.

2.2.4. Social Support

The 20-item Medical Outcomes Study Social Support Survey was used to measure social support in this study [41]. The instrument had 1 fill-in-the-blank item to record the number of close friends and relatives and 19 Likert-scale, 5-point options. The total social support score as well as the subscales (tangible support, emotional/informational support, affectionate support, and positive social interaction) had excellent internal consistency ($\alpha = 0.91$ – 0.97). The tangible support items measured help provided by others during an illness such as providing transportation to a doctor and assisting with daily chores. Emotional/informational support measured the availability of someone else to share concerns, assist during a crisis, and provide advice. Affectionate support can be described as having someone to show displays of love and affection, and positive social interaction involves having someone to share enjoyable things with.

2.2.5. Diabetes Knowledge

The Revised Diabetes Knowledge Test had adequate reliability ($\alpha = 0.77$) and was used to measure diabetes knowledge [42]. The measure includes 23 multiple choice questions, but 3 items about insulin were not scored because the curriculum did not address insulin therapy. The 20 items were totaled, for a maximum possible score of 100 points.

2.3. Intervention

The participants in the intervention group received a manualized culturally relevant curriculum that was developed by the American Diabetes Association to improve diabetes self-management and reduce disease risk among African American adults in church settings. The program was delivered by the same advanced public health nurse in the par-

participating churches randomized to the intervention group over 3 weekly sessions that lasted approximately 90 minutes to 2 hours, depending on questions and discussion. The sessions were similar to other diabetes health promotion interventions and included information about diabetes risk and pathology, managing diabetes and prediabetes through diet and exercise, maintaining glucose and HbA1C levels within therapeutic ranges, and linkages with heart disease, kidney damage, and stroke. Interactive strategies were employed, which facilitated group discussion and engagement. The control group received a brochure about diabetes.

2.4. Data Analysis

A confirmatory factor analysis (CFA) model was fit to test the measurement model where self-care behavior, a latent variable, was predicted to load onto eight measured variables. These measured variables are general diet, specific diet related to fat intake, specific diet related to carbohydrate intake, specific diet related to produce intake, exercise, blood glucose testing, foot care, and medication management. The CFA model was nested into the full structural model testing the hypothesis that diabetes knowledge, diabetes fatalism, perceived diabetes self-management, and social support predict self-care behaviors. The hypothesized model is similar to the one proposed in a previous publication by Osborn and Egede, although the specific diet subscale is three individual subscales, and there is no smoking subscale included [29]. Additionally, perceived diabetes self-management was included as a predictor of self-care behaviors. The structural model and the CFA model were tested using MPlus version 8.5 [43]. Variables with factor loadings that were not significant were excluded from the full SEM analysis. Model fit was assessed with likelihood ratio chi-square tests and the root mean square error of approximation (RMSEA) [44,45]. Insignificant chi-square test results indicate that the data fit the hypothesized structure. RMSEA values less than 0.05 indicate a close fit, whereas values between 0.05 and 0.08 indicate a reasonable fit. RMSEA values larger than 0.1 indicate a poor fit [46].

3. Results

The sample included 12 participating churches that were randomized to the intervention ($n = 7$) and control ($n = 5$) groups. Of the 146 individual participants recruited from the randomized churches, 75 received the intervention, and 71 were in the waitlisted control group. However, a few participants ($n = 9$) from both the intervention ($n = 7$) and control ($n = 2$) groups ceased participation after providing informed consent and completing baseline measures [35]. All self-identified as African American, and there were no significant between-group differences regarding sociodemographic characteristics except that more participants in the intervention group had previously been diagnosed with diabetic retinopathy ($p = 0.003$) and were unemployed ($p = 0.02$) [35]. The average ages for participants in the intervention group were 61.8 and 61.6 for those in the control group, and there were more participants (72%) who self-identified as being unemployed in the intervention group, compared with those (49%) in the control group [35]. Additionally, women ($n = 110$; 75%) participated in the study more than men ($n = 36$; 25%), but there was no statistically significant gender difference ($p = 0.08$) between groups [35]. Similar numbers of people in both the intervention ($n = 43$) and control ($n = 44$) groups had been diagnosed with diabetes, and of those diagnosed with prediabetes, there were a few more in the intervention group ($n = 32$), compared with the control group ($n = 27$).

The CFA model is graphically depicted in Figure 1. The initial CFA model demonstrated a poor fit ($\chi^2 (20) = 54.450$, $p < 0.001$; RMSEA = 0.109) to the data; thus, model adjustments were made. First, the measured variable for specific diet related to fat intake was removed, as it was insignificant. This model also demonstrated a poor fit to the data ($\chi^2 (14) = 48.256$, $p < 0.001$; RMSEA = 0.129), and therefore, a further refinement was made by adding the correlation between blood glucose testing and medication management to the model. The adjusted model demonstrated an adequate fit ($\chi^2 (13) = 13.224$, $p = 0.4307$; RMSEA = 0.011 [90%CI: 0.000, 0.083]). Diabetes self-care behaviors loaded significantly

onto general diet (0.719, $p < 0.001$), specific diet for produce intake (0.592, $p < 0.001$), specific diet for carbohydrate intake (0.643, $p < 0.001$), exercise (0.511, $p < 0.001$), blood glucose testing (0.372, $p < 0.001$), foot care (0.374, $p < 0.001$), and medication management (0.258, $p = 0.006$).




Figure 1. Confirmatory factor analysis (CFA) model. Standardized CFA solution; *** $p < 0.001$.

The structural model that was estimated is displayed with parameter estimates for each path in Figure 2. The estimated model showed a goodness of fit ($\chi^2 (37) = 31.906, p = 0.7065$; RMSEA = 0.000 [90%CI: 0.000, 0.046]). The resulting model had no multivariate outliers or issues with collinearity. The path beginning with perceived diabetes self-management was the only significant direct path predicting self-care behaviors ($r = 0.286, p = 0.002$).

Table 1. Correlations.

	Diabetes Knowledge	Perceived Diabetes Self-Management	Diabetes Fatalism	Social Support
Diabetes Knowledge	1	-0.098	-0.066	0.155
Perceived Diabetes Self-Management		1	-0.301 ***	0.101
Diabetes Fatalism			1	-0.074

Note. *** $p < 0.001$.

Additionally, the correlation between perceived diabetes self-management and diabetes fatalism was also significant (Table 1). Model results indicated that higher perceived diabetes self-management, higher diabetes knowledge, less diabetes fatalism, and more social support predicted better self-care behaviors, with the model explaining 17.3% of the variability in diabetes self-care behaviors.




Figure 2. Structural model with parameter estimates for each path. Standardized solution: ** $p < 0.01$, *** $p < 0.001$, ^p indicates a p -value equal to 0.085. Only one correlation among exogenous variables is presented, although all were fit. See Table 1 for correlations.

4. Discussion

The IMB-DSC helped conceptualize the determinants of diabetes self-care activities among rural participants living with diabetes and prediabetes including diabetes knowledge as information, diabetes fatalism as personal motivation, social support as social motivation, and diabetes self-care as health behaviors [29]. The current study included the same components with the addition of the perceived diabetes self-management scale to represent self-efficacy in relation to diabetes self-care activities [40]. As a chronic disease, diabetes requires individual confidence, or self-efficacy, to perform the diabetes self-management activities necessary for preventing adverse consequences associated with lack of glycemic control [40]. Little is known about factors that influence diabetes self-care outcomes, and the findings of this study contribute to the literature about diabetes outcomes among underserved rural groups. In this sample, the intervention had no statistically significant impact on perceived diabetes self-management, and a possible rationale could be that the levels were already high at baseline and left little room for improvement at post-test [47]. However, in this analysis, perceived diabetes self-management had the largest impact on self-care behaviors, as greater self-efficacy in performing self-care activities was associated with the actual performance of those behaviors. Similarly, another study conducted among Chinese participants concluded that self-efficacy was an independent predictor of diabetes self-care behaviors [48].

There are a few published studies that discussed perceived diabetes self-management in other populations. For example, a study that included adults in Saudi Arabia showed a significant correlation between perceived diabetes self-management and social support [49]. Another showed that a telephone intervention delivered to Turkish adults significantly improved diabetes self-efficacy and perceived self-management [50]. Additionally, motivational interviewing had a significant impact on perceived diabetes self-management in Turkish adults [51]. An educational health technology program discussed in another study

improved the perceptions of diabetes self-care as well as glycemic control among study participants in Asia [52]. The positive associations and promising intervention delivery methods discussed in these articles provide important considerations when developing future public health interventions targeting diabetes.

In this sample, the negative correlational relationship between perceived diabetes self-management and diabetes fatalism was statistically significant. This result makes clinical sense since, with an increase in the perception of diabetes self-management, it would seem likely that people would have greater feelings of self-efficacy and fewer fatalistic beliefs about living with and managing diabetes. However, even though the perceived diabetes self-management score was reported as high in this population, diabetes fatalism was moderate, which suggests that interventions should include efforts to reduce fatalism by directly impacting psychological and behavioral factors [47,53]. This has clinical importance because other studies found that higher diabetes fatalism was directly linked to poor glycemic control, poor medication adherence, and decreased self-care [53–55]. The results of this study showed that diabetes fatalism had a small negative effect in that lower levels of fatalism were associated with a greater likelihood of performing diabetes self-care behaviors. In comparison, having more social support and increased diabetes knowledge had small positive effects on self-care activities. A future study having a larger sample size could potentially show larger effects on these outcomes in this population.

Future research can also focus on the impact of individual perceptions of diabetes self-management on self-care behaviors in similar rural groups as well as urban populations. The strategies previously discussed such as using motivational interviewing, health technology, and telephone interventions could be used to enhance diabetes health interventions and improve accessibility among underserved populations residing in any location. Poverty and poor living conditions and built environment conditions can adversely affect diabetes outcomes [56]. Further research is needed to explore the impact of social determinants of health on factors related to diabetes self-care, such as fatalism, self-efficacy, and social support in other geographic areas, and brainstorm strategies to improve health among diverse populations living with diabetes. Additionally, a future analysis can explore the impact of sociodemographic characteristics and diabetes-related complications on outcomes. For example, a study among Lebanese adults showed that fatalistic attitudes were associated with characteristics such as younger age, lower educational attainment, higher BMI, and fewer diabetes comorbidities [57]. This study had some limitations as well. First, the study was conducted among participants living in a southern region of the United States, which could limit generalizability to other populations. Second, the data were self-reported by participants, and there could have potentially been recall bias when completing the survey instruments.

5. Conclusions

Confronting inequalities requires solutions that are practical, relevant, and encompass the unique characteristics of those communities that have experienced multiple geographic and systemic disadvantages [58]. For example, people living in rural geographic locations often experience hardships related to poverty and limited access to health information, access to health care, and other resources that affect health program responses and impact outcomes. The COVID-19 pandemic has further exacerbated these challenges and will likely contribute to a resurgence of diabetes-related sequelae in addition to the consequences of other chronic diseases. Mitigating rural disparities associated with diabetes involves community-relevant strategies that address the psychological and behavioral determinants of standardized diabetes care that improve diabetes self-care activities [53]. Healthy lifestyle recommendations promote longer life expectancy and reduce chronic disease risk associated with the major causes of death associated with diabetes such as cancer, stroke, and cardiovascular disease [59–61]. The development and testing of such tailored interventions can potentially impact health outcomes, decrease chronic disease development and exacerbation, and improve the quality of life among rural dwellers.

Author Contributions: L.A. was involved in the funding acquisition, conceptualization, and implementation of the methodology, as well as data collection, writing and preparing the original draft, and further review and editing. L.G. was involved in the development of the methodology and preparation of the manuscript, including the original draft and further review and editing. G.S. was involved in the conceptualization and development of the methodology, statistical analysis, writing and preparing the original draft, and editing. J.L. was involved in the development of the methodology and reviewing and editing the original manuscript and revisions. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Florida State University FYAP grant and the Florida State University College of Nursing Infrastructure Grant.

Institutional Review Board Statement: This study was conducted in accordance with the Declaration of Helsinki and approved by the Institutional Review Board of Florida State University (HSC 2017.20379 on 3/14/17).

Informed Consent Statement: Informed consent was obtained from all subjects involved in this study.

Data Availability Statement: The data presented in this study will be considered upon request from the corresponding author. The data are not publicly available due to privacy issues.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

- Boersma, P.; Black, L.I.; Ward, B.W. Prevalence of Multiple Chronic Conditions Among US Adults, 2018. *Prev. Chronic. Dis.* **2020**, *17*, 200130. [CrossRef] [PubMed]
- CDC. National Center for Chronic Disease Prevention and Health Promotion. 2021. Available online: <https://www.cdc.gov/chronicdisease/index.htm> (accessed on 4 May 2022).
- Leon, B.M.; Maddox, T.M. Diabetes and cardiovascular disease. Epidemiology, biological mechanisms, treatment recommendations and future research. *World J. Diabetes* **2015**, *6*, 1246–1258. [CrossRef] [PubMed]
- Tsao, C.W.; Aday, A.W.; Almarzooq, Z.I.; Alonso, A.; Beaton, A.Z.; Bittencourt, M.S.; Boehme, A.K.; Buxton, A.E.; Carson, A.P.; Commodore-Mensah, Y.; et al. Heart disease and stroke statistics-2022 update: A report from the American Heart Association. *Circulation* **2022**, *145*, e153–e639. [CrossRef]
- Carnethon, M.R.; Pu, J.; Howard, G.; Albert, M.A.; Anderson, C.A.; Bertoni, A.G.; Mujahid, M.S.; Palaniappan, L.; Taylor, H.A., Jr.; Willis, M.; et al. Cardiovascular Health in African Americans: A Scientific Statement from the American Heart Association. *Circulation* **2017**, *136*, e393–e423. [CrossRef] [PubMed]
- Havranek, E.P.; Mujahid, M.S.; Barr, D.A.; Blair, I.V.; Cohen, M.S.; Cruz-Flores, S.; Davey-Smith, G.; Dennison-Himmelfarb, C.R.; Lauer, M.S.; Lockwood, D.W.; et al. Social determinants of risk and outcomes for cardiovascular disease A scientific statement From the American Heart Association. *Circulation* **2015**, *132*, 873–898. [CrossRef]
- Howard, G.; Kleindorfer, D.O.; Cushman, M.; Long, D.L.; Jasne, A.; Judd, S.E.; Higginbotham, J.C.; Howard, V.J. Contributors to the excess stroke mortality in rural areas in the United States. *Stroke* **2017**, *48*, 1773–1778. [CrossRef]
- Kulshreshtha, A.; Goyal, A.; Dabhadkar, K.; Veledar, E.; Vaccarino, V. Urban-rural differences in coronary heart disease mortality in the United States: 1999–2009. *Public Health Rep.* **2014**, *129*, 19–29. [CrossRef]
- Limdi, N.A.; Howard, V.J.; Higginbotham, J.; Parton, J.; Safford, M.M.; Howard, G. US mortality: Influence of race, geography and cardiovascular risk among participants in the population-based REGARDS cohort. *J. Racial Ethn. Health Disparities* **2016**, *3*, 599. [CrossRef] [PubMed]
- Meit, M.; Knudson, A.; Gilbert, T.; Yu, A.T.; Tanenbaum, E.; Ormson, E.; Popat, S. Rural Health Reform Policy Research Center: The 2014 Update of the Rural-Urban Chartbook. 2014. Available online: <https://ruralhealth.und.edu/projects/health-reform-policy-research-center/pdf/2014-rural-urban-chartbook-update.pdf> (accessed on 6 May 2022).
- Singh, G.K.; Daus, G.P.; Allender, M.; Ramey, C.T.; Martin, E.K.; Perry, C.; Reyes, A.A.D.L.; Vedamuthu, I.P. Social determinants of health in the United States: Addressing major health inequality trends for the nation, 1935–2016. *Int. J. MCH AIDS* **2017**, *6*, 139–164. [CrossRef] [PubMed]
- Singh, G.K.; Siahpush, M. Widening rural-urban disparities in all-cause mortality and mortality from major causes of death in the USA, 1969–2009. *J. Urban Health* **2014**, *91*, 272–292. [CrossRef] [PubMed]
- Trivedi, T.; Liu, J.; Probst, J.C.; Merchant, A.; Jones, S.; Martin, A.B. Obesity and obesity-related behaviors among rural and urban adults in the USA. *Rural. Remote Health* **2015**, *15*, 3267. [CrossRef] [PubMed]
- Coughlin, S.S.; Clary, C.; Johnson, J.A.; Berman, A.; Heboyan, V.; Benevides, T.; Moore, J.; George, V. Continuing Challenges in Rural Health in the United States. *J. Environ. Health Sci.* **2019**, *5*, 90–92.

15. Garcia, M.C.; Faul, M.; Massetti, G.; Thomas, C.C.; Hong, Y.; Bauer, U.E.; Iademarco, M.F. Reducing Potentially Excess Deaths from the Five Leading Causes of Death in the Rural United States. *MMWR Surveill. Summ.* **2017**, *66*, 1–7. [[CrossRef](#)] [[PubMed](#)]
16. Garcia, M.C.; Faul, M.; Dowling, N.F.; Thomas, C.C.; Iademarco, M.F. Bridging the Gap in Potentially Excess Deaths Between Rural and Urban Counties in the United States. *Public Health Rep.* **2020**, *135*, 177–180. [[CrossRef](#)] [[PubMed](#)]
17. Moy, E.; Garcia, M.C.; Bastian, B.; Rossen, L.M.; Ingram, D.D.; Faul, M.; Massetti, G.M.; Thomas, C.C.; Hong, Y.; Yoon, P.W.; et al. Leading Causes of Death in Nonmetropolitan and Metropolitan Areas—United States, 1999–2014. *MMWR Surveill. Summ.* **2017**, *66*, 1–8. [[CrossRef](#)] [[PubMed](#)]
18. Matthews, K.A.; Croft, J.B.; Liu, Y.; Lu, H.; Kanny, D.; Wheaton, A.G.; Cunningham, T.J.; Khan, L.K.; Caraballo, R.S.; Holt, J.B.; et al. Health-Related Behaviors by Urban-Rural County Classification—United States, 2013. *MMWR Surveill. Summ.* **2017**, *66*, 1–8. [[CrossRef](#)]
19. Logan, H.; Guo, Y.; Dodd, V.J.; Muller, K.; Riley, J. The burden of chronic diseases in a rural North Florida sample. *BMC Public Health* **2013**, *13*, 906. [[CrossRef](#)]
20. Swanoski, M.T.; Lutfiyya, M.N.; Amaro, M.L.; Akers, M.F.; Huot, K.L. Knowledge of heart attack and stroke symptomology: A cross-sectional comparison of rural and non-rural US adults. *BMC Public Health* **2012**, *12*, 283. [[CrossRef](#)]
21. Chatterjee, R.; Maruthur, N.M.; Edelman, D. Novel risk factors for type 2 diabetes in African Americans. *Curr. Diab. Rep.* **2015**, *15*, 103. [[CrossRef](#)]
22. Cummings, D.M.; Lutes, L.D.; Littlewood, K.; Solar, C.; Hambidge, B.; Gatlin, P. Impact on distress reduction on behavioral correlates and A1C in African American women with uncontrolled type 2 diabetes: Results from EMPOWER. *Ethn. Dis.* **2017**, *27*, 155–160. [[CrossRef](#)]
23. Richards Adams, I.K.; Figueroa, W.; Hatsu, I.; Odei, J.B.; Sotos-Prieto, M.; Leson, S.; Huling, J.; Joseph, J.J. An examination of demographic and psychosocial factors, barriers to health eating, and diet quality among African American adults. *Nutrients* **2019**, *11*, 519. [[CrossRef](#)] [[PubMed](#)]
24. Hill-Briggs, F.; Adler, N.E.; Berkowitz, S.A.; Chin, M.H.; Gary-Webb, T.L.; Navas-Acien, A.; Thornton, P.L.; Haire-Joshu, D. Social Determinants of Health and Diabetes: A Scientific Review. *Diabetes Care* **2020**, *44*, 258–279. [[CrossRef](#)] [[PubMed](#)]
25. Davis, A.M.; Vinci, L.M.; Okwuosa, T.M.; Chase, A.R.; Huang, E.S. Cardiovascular health disparities: A systematic review of health care interventions. *Med. Care Res. Rev.* **2007**, *64*, 29S–100S. [[CrossRef](#)]
26. Berardi, V.; Bellettiere, J.; Nativ, O.; Ladislav, S.; Hovell, M.F.; Baron-Epel, O. Fatalism, diabetes management, and the role of religiosity. *J. Relig. Health* **2016**, *55*, 602–617. [[CrossRef](#)] [[PubMed](#)]
27. Hernandez, R.; Ruggiero, L.; Riley, B.B.; Wang, Y.; Chavez, N.; Quinn, L.T.; Gerber, B.S.; Choi, Y.K. Correlates of self-care in low-income African American and Latino patients with diabetes. *Health Psychol.* **2014**, *33*, 597–607. [[CrossRef](#)] [[PubMed](#)]
28. Smith, S.A.; Ansa, B. A systematic review of lifestyle interventions for chronic diseases in rural communities. *J. Ga. Public Health Assoc.* **2016**, *5*, 304–313. [[CrossRef](#)]
29. Osborne, C.Y.; Egede, L.E. Validation of an Information-Motivation-Behavioral skills model of diabetes self-care (IMB-DSC). *Patient Educ. Couns.* **2010**, *79*, 49–54. [[CrossRef](#)] [[PubMed](#)]
30. Egede, L.E.; Ellis, C. Development and psychometric properties of the 12-item diabetes fatalism scale. *J. Gen. Intern. Med.* **2010**, *25*, 61–66. [[CrossRef](#)] [[PubMed](#)]
31. Al-Dwaikat, T.N.; Chlebowy, D.O.; Hall, L.A.; Crawford, T.N.; Yankeelov, P.A. Self-management as a mediator of the relationship between social support dimensions and health outcomes of African American adults with type 2 diabetes. *West J. Nurs. Res.* **2020**, *42*, 485–494. [[CrossRef](#)]
32. Fitzpatrick, S.L.; Hill-Briggs, F. Strategies for sustained weight management: Perspectives from African American patients with type 2 diabetes. *Diabetes Educ.* **2017**, *43*, 304–310. [[CrossRef](#)] [[PubMed](#)]
33. Hawkins, J.M.; Mitchell, J. Can social integration and social support help to explain racial disparities in health care utilization among men with diabetes? *Int. J. Men's Health* **2017**, *16*, 66–83.
34. Rutledge, P.M.; Montgomery, A.J.; Lott, S.M.; McGee, Z.M.; Burns, D.M. The relationship between family support, self-care, and health in selected African American females with type 2 diabetes. *J. Natl. Black Nurses Assoc.* **2019**, *30*, 1–9.
35. Abbott, L.S.; Slate, E.H.; Graven, L.J. Cardiovascular disease risk among rural residents living with diabetes and prediabetes: A cluster randomized trial. *Public Health Nurs.* **2020**, *37*, 16–24. [[CrossRef](#)] [[PubMed](#)]
36. Donner, A.; Klar, N. Pitfalls and controversies in cluster randomization trials. *Am. J. Public Health* **2004**, *94*, 416–422. [[CrossRef](#)] [[PubMed](#)]
37. Murray, D.M.; Blitstein, J.L. Methods to reduce the impact of intraclass correlation in group-randomized trials. *Eval. Rev.* **2003**, *27*, 79–103. [[CrossRef](#)] [[PubMed](#)]
38. Dsn, P.S.F.; Pribulick, M.; Williams, I.C.; James, G.D.; Rovynak, V.; Seibold-Simpson, S.M. Promoting heart health in rural women. *J. Rural Health* **2013**, *29*, 248–257.
39. Toobert, D.J.; Hampson, S.E.; Glasgow, R.E. The summary of diabetes self-care activities measure. *Diabetes Care* **2000**, *23*, 943–950. [[CrossRef](#)] [[PubMed](#)]
40. Wallston, K.A.; Rothman, R.L.; Cherrington, A. Psychometric properties of the perceived diabetes self-Management scale (PDSMS). *J. Behav. Med.* **2007**, *30*, 395–401. [[CrossRef](#)] [[PubMed](#)]
41. Sherbourne, C.D.; Stewart, A.L. The MOS social support survey. *Soc. Sci. Med.* **1991**, *32*, 705–714. [[CrossRef](#)]

42. Fitzgerald, J.T.; Funnell, M.M.; Anderson, R.M.; Nwankwo, R.; Stansfield, R.B.; Piatt, G.A. Validation of the revised brief Diabetes Knowledge Test (DKT2). *Diabetes Educ.* **2016**, *42*, 178–187. [[CrossRef](#)] [[PubMed](#)]
43. Muthén, L.K.; Muthén, B.O. (1998–2011). *Mplus User's Guide*, 6th ed.; Muthén & Muthén: Los Angeles, CA, USA, 1998.
44. Browne, M.W.; Cudeck, R.I. Alternative ways of assessing model fit. In *Testing Structural Equation Models*; Bollen, K., Long, K., Eds.; Sage: Newbury Park, CA, USA, 1993; pp. 136–162.
45. Steiger, J.H. Structural model evaluation and modification: An interval estimation approach. *Multivar. Behav. Res.* **1990**, *25*, 173–180. [[CrossRef](#)] [[PubMed](#)]
46. MacCallum, R.C.; Browne, M.W.; Sugawara, H.M. Power analysis and determination of sample size for covariance structure modeling. *Psychol. Methods* **1996**, *1*, 130–149. [[CrossRef](#)]
47. Abbott, L.; Slate, E.; Graven, L.; Lemacks, J.; Grant, J. Fatalism, social support and self-management perceptions among rural African Americans living with diabetes and pre-diabetes. *Nurs. Rep.* **2021**, *11*, 242–252. [[CrossRef](#)]
48. Gao, J.; Wang, J.; Zhu, Y.; Yu, J. Validation of an information-motivation-behavioral skills model of self-care among Chinese adults with type 2 diabetes. *BMC Public Health* **2013**, *13*, 100. [[CrossRef](#)]
49. Alshutwi, S.; Miligi, E.; Alhumidan, L.; Almutairi, A.F. The influence of the disclosure of diabetes on the cognitive, physical ability and diabetes self-management in diabetic employed adults in Saudi Arabia. *Nurs. Open* **2022**, *9*, 978–985. [[CrossRef](#)] [[PubMed](#)]
50. Sayin Kasar, K.; Duru Asiret, G.; Kutmec Yilmaz, C.; Canlar, Ş. The effect of model-based telephone counseling on HbA1c and self-management for individuals with type 2 diabetes: A randomized controlled trial. *Prim. Care Diabetes* **2022**, *16*, 41–48. [[CrossRef](#)]
51. Dogru, A.; Owayolu, N.; Owayolu, O. The effect of motivational interview persons with diabetes on self-management and metabolic variables. *J. Pak. Med. Assoc.* **2019**, *69*, 294–300.
52. Pai, L.W.; Chiu, S.C.; Liu, H.L.; Chen, L.L.; Peng, T. Effects of a health education technology program on long-term glycemic control and self-management ability of adults with type 2 diabetes: A randomized controlled trial. *Diabetes Res. Clin. Pract.* **2021**, *175*, 108785. [[CrossRef](#)]
53. Asuzu, C.C.; Walker, R.J.; Williams, J.S.; Egede, L.E. Pathways for the relationship between diabetes distress, depression, fatalism and glycemic control in adults with type 2 diabetes. *J. Diabetes Complicat.* **2017**, *31*, 169–174. [[CrossRef](#)]
54. Sukkarieh-Haraty, O.; Egede, L.E.; Abi Kharm, J.; Bassil, M. Diabetes fatalism and its emotional distress subscale are independent predictors of glycemic control among Lebanese patients with type 2 diabetes. *Ethn. Health* **2019**, *24*, 767–778. [[CrossRef](#)] [[PubMed](#)]
55. Walker, R.J.; Smalls, B.L.; Hernandez-Tejada, M.A.; Campbell, J.A.; Davis, K.S.; Egede, L.E. Effect of diabetes fatalism on medication adherence and self-care behaviors in adults with diabetes. *Gen. Hosp. Psychiatry* **2012**, *34*, 598–603. [[CrossRef](#)]
56. Awad, N.; Saade, R.; Bassil, M.; Sukkarieh-Haraty, O.; Egede, L.E. Relationship between social determinants of health and clinical outcomes in adults with type 2 diabetes in Lebanon. *J. Natl. Med. Assoc.* **2022**, *114*, 392–405. [[CrossRef](#)] [[PubMed](#)]
57. Sukkarieh-Haraty, O.; Egede, L.E.; Abi Kharm, J.; Bassil, M. Predictors of Diabetes Fatalism Among Arabs: A Cross-Sectional Study of Lebanese Adults with Type 2 Diabetes. *J. Relig. Health* **2018**, *57*, 858–868. [[CrossRef](#)]
58. Power, T.; Kelly, R.; Usher, K.; East, L.; Travaglia, J.; Robertson, H.; Wong, A.; Jackson, D.; Balintongog, A. Living with diabetes and disadvantage: A qualitative, geographical case study. *J. Clin. Nurs.* **2020**, *29*, 2710–2722. [[CrossRef](#)] [[PubMed](#)]
59. Li, Y.; Schoufour, J.; Wang, D.D.; Dhana, K.; Pan, A.; Liu, X.; Song, M.; Liu, G.; Shin, H.J.; Sun, Q.; et al. Healthy lifestyle and life expectancy free of cancer, cardiovascular disease, and type 2 diabetes: Prospective cohort study. *BMJ* **2020**, *368*, l6669. [[CrossRef](#)]
60. Yang, R.; Xu, H.; Pedersen, N.L.; Li, X.; Yu, J.; Bao, C.; Qi, X.; Xu, W. A healthy lifestyle mitigates the risk of heart disease related to type 2 diabetes: A prospective nested case-control study in a nationwide Swedish twin cohort. *Diabetologia* **2021**, *64*, 530–539. [[CrossRef](#)] [[PubMed](#)]
61. Zhu, M.; Wang, T.; Huang, Y.; Zhao, X.; Ding, Y.; Zhu, M.; Ji, M.; Wang, C.; Dai, J.; Yin, R.; et al. Genetic Risk for Overall Cancer and the Benefit of Adherence to a Healthy Lifestyle. *Cancer Res.* **2021**, *81*, 4618–4627. [[CrossRef](#)]