

RESEARCH

Open Access



High tumor hexokinase-2 expression promotes a pro-tumorigenic immune microenvironment by modulating CD8+/- regulatory T-cell infiltration

Sehui Kim^{1,2}, Jaemoon Koh¹, Seung Geun Song^{1,3}, Jeemin Yim¹, Miso Kim⁴, Bhumsuk Keam⁴, Young Tae Kim^{5,6}, Jihun Kim⁷, Doo Hyun Chung^{1,3} and Yoon Kyung Jeon^{1,6*}

Abstract

Background: Relationship between cancer cell glycolysis and the landscape of tumor immune microenvironment in human cancers was investigated.

Methods: Forty-one fresh lung adenocarcinoma (ADC) tissues were analyzed using flow cytometry for comprehensive immunoprofiling. Formalin-fixed tissues were immunostained for hexokinase-2 (HK2) to assess cancer cell glycolysis. For validation, formalin-fixed tissues from 375 lung ADC, 118 lung squamous cell carcinoma (SqCC), 338 colon ADC, and 78 lung cancer patients treated with anti-PD-1/PD-L1 immunotherapy were immunostained for HK2, CD8, and FOXP3.

Results: Based on immunoprofiling of lung ADC, HK2 tumor expression was associated with the composition of lymphoid cells rather than myeloid cells. High HK2 tumor expression was associated with immunosuppressive/pro-tumorigenic features, especially decreased ratio of CD8+ T-cells to Tregs ($\rho = -0.415$, $P = 0.012$). This correlation was also confirmed in four different cohorts including lung ADC and SqCC, colon ADC, and the immunotherapy cohort ($\rho = -0.175 \sim -0.335$, all $P < 0.05$). A low CD8+ T-cell to Treg ratio was associated with poor progression-free survival and overall survival in lung SqCC patients, and a shorter overall survival in the immunotherapy cohort (all, $P < 0.05$).

Conclusion: An increase in HK2 expression may contribute to shaping the immunosuppressive/pro-tumorigenic tumor microenvironment by modulating the CD8+ T-cell to Treg ratio. Targeting tumor HK2 expression might be a potential strategy for enhancing anti-tumor immunity.

Keywords: Hexokinase-2, Glycolysis, Tumor microenvironment, Tumor-infiltrating lymphocytes, CD8+ T-cell to Treg ratio, Immunotherapy

Background

Cancer immunotherapy targeting the PD-1/PD-L1 pathway is clinically beneficial and widely used in patients with a variety of cancers [1–3]. However, only a small proportion of patients show a good response to immunotherapy; thus, there have been many efforts to uncover predictive biomarkers to improve PD-1/PD-L1 blockade [4]. PD-L1 expression and the tumor mutational

*Correspondence: ykjeon@snu.ac.kr

¹ Department of Pathology, Seoul National University College of Medicine, 103 Daehak-ro, Jongno-gu, 03080 Seoul, Republic of Korea
Full list of author information is available at the end of the article



burden in lung cancer are used as predictive biomarkers in clinical practice but have limited predictability [5–7]. This may be attributed to the complexity of the tumor microenvironment (TME), since the responsiveness to immunotherapy is affected by the tumor immune microenvironment [8]. In general, a pro-inflammatory microenvironment rather than an immunosuppressive microenvironment predicts a good response to immunotherapy.

Tumor and TME crosstalk occurs via diverse genetic and non-genetic factors, which shape the tumor immune microenvironment [9]. For example, β -catenin signaling, RHOA mutation, PTEN loss, LKB1 mutation, and KRAS mutations were associated with an immunosuppressive microenvironment, which included reduced CD8+ T-cell infiltration and increased regulatory T-cell (Treg) infiltration [9–14]. In addition, the metabolic status of tumor cells can affect immune cell infiltration and function [15, 16].

Within the tumor niche, tumor cells and surrounding environmental cells, especially immune cells, compete for limited nutrients and oxygen. Therefore, tumor metabolism can affect tumor-infiltrating immune cells. Because tumor cells predominantly use glycolysis for their survival and proliferation [17], immune cells often encounter low glucose and high lactic acid levels in the TME. Previous studies demonstrated that high tumor glycolysis results in an immunosuppressive TME by reducing effector T-cell functions [15, 16]. However, the effects of tumor glycolysis on other immune cells remain unclear.

Each subset of immune cells in the TME utilizes different metabolic programs for their survival, activation, and differentiation [18, 19]. Effector T-cells are highly glycolytic, but Tregs depend on oxidative phosphorylation (OXPHOS) and fatty acid oxidation (FAO) [20–23]. Activated dendritic cells use glycolysis predominantly, and pro-inflammatory tumor-associated macrophages (TAMs) are highly glycolytic. However, tolerogenic TAMs usually use oxygen-driven metabolism, such as OXPHOS and FAO [24]. Tumor-infiltrating myeloid-derived suppressor cells (MDSCs) in mice prefer FAO over glycolysis as a primary source of energy [25]. Therefore, each immune cell subset might be uniquely affected depending on the type of tumor metabolism.

Previously, we evaluated tumor HK2 expression, as a marker of glycolysis and analyzed the relationship between tumor HK2 expression and T cell function/infiltration. Previous study showed that lung cancer cell HK2 overexpression suppressed T-cell effector functions, and that the increased HK2 expression was inversely correlated with the expression of T-cell effector molecules according to analysis of The Cancer Genome Atlas in lung cancer [16]. However, little is known about the

landscape of the tumor immune microenvironment in the context of tumor glycolysis in human cancer tissues. Thus, we comprehensively investigated the relationship between tumor glycolysis and tumor-infiltrating immune cells in human cancer tissues. We demonstrated that a higher HK2 expression which stands for higher rate of tumor glycolysis is associated with an immunosuppressive microenvironment characterized by a decrease in CD8+ T-cell infiltration relative to Treg infiltration in lung and colon cancers.

Materials and methods

Patients and samples

Fresh tumor samples from 41 lung adenocarcinoma (ADC) patients who underwent surgery at Seoul National University Hospital (SNUH) were used for flow cytometry analysis. All patients agreed to the sample collection and data analyses. Written informed consent was provided for this cohort. For immunohistochemistry (IHC) analysis, tissue microarrays were constructed from formalin-fixed paraffin-embedded (FFPE) tumor tissues from the above 41 lung ADC patients, and an additional 375 lung ADC and 118 lung squamous cell carcinoma (SqCC) patients from SNUH and 338 colon ADC patients from Asan Medical Center. Additionally, 78 non-small cell lung cancer patients treated with PD-1/PD-L1 blockade at SNUH were included for further validation. Informed consent for participation in the IHC analysis only was waived by the SNUH and Asan Medical Center institutional review boards because this was a retrospective study using archived material and did not increase patient risk. The tumor, node, and metastasis staging system was performed based on the 7th American Joint Committee on Cancer (AJCC) for lung cancer and the 8th AJCC for colon cancer [26, 27]. Clinicopathological features of patients are summarized in Supplementary Tables S1, 2, 3, 4 and 5. This study followed the World Medical Association Declaration of Helsinki recommendations and was approved by the institutional review board of SNUH (No.: H-1404-100-572 and H-1905-115-1035).

Flow cytometry analysis

To analyze tumor-infiltrating immune cell subsets, 1 g fresh lung ADC tissues were subjected to flow cytometric analysis, as described previously [28]. Data were analyzed using FlowJo v10.1 software (Treestar), and the gating strategy is shown in Supplementary Figure S1.

Immunohistochemistry (IHC)

IHC was performed using the following antibodies: rabbit anti-HK2 polyclonal antibody (Genetex, Irvine, CA, USA), rabbit anti-CD8 monoclonal antibody (clone SP16,

Thermo Fisher Scientific, Rockford, IL, USA) and rabbit anti-FOXP3 monoclonal antibody (236 A/E7, Abcam, Cambridge, UK). Immunostaining was performed using the Benchmark XT autostainer (Ventana Medical Systems, Tucson, AZ, USA). The H-score of HK2 was calculated as follows: sum of each staining intensity (0–3) · proportion (0–100%). The numbers of CD8+ and FOXP3+ tumor-infiltrating lymphocytes (TILs) per mm² were automatically counted via modified nuclear IHC algorithms using Aperio ImageScope software (Aperio Technologies, Vista, CA, USA).

Statistical analyses

The correlation between tumor HK2 expression and tumor-infiltrating immune cells was calculated using Spearman's correlation. Two-sided *P* values < 0.05 were considered statistically significant. The cutoff values of the variables were determined by receiver operating characteristic curve analysis. Survival analysis was performed using Kaplan–Meier analysis and the log-rank test. Two-sided *P* values < 0.05 were considered statistically significant in all analyses. All statistical analyses were performed using R statistical software 4.1.1 (R Foundation for Statistical Computing, Vienna, Austria). Images were created using the GraphPad Prism 7 software.

Results

Tumor glycolysis is differentially associated with each subset of immune cells

Using flow cytometry, tumor-infiltrating lymphoid cells and myeloid cells from lung ADC fresh tissues from 41 patients were comprehensively profiled (Supplementary Figure S1). IHC was used to evaluate HK2 expression in tumor cells, as an indicator of tumor glycolysis, because HK2 is the first and the rate-limiting enzyme of glycolysis. The experimental scheme is described in Fig. 1 A, and the correlations between individual immune cell subsets and tumor HK2 expression are summarized in Fig. 1 B, C; Table 1.


Total immune cell infiltration did not differ when quantified according to tumor HK2 expression. Tumor HK2 expression was more closely related to lymphoid cells compared with myeloid cells (Fig. 1 B, C). The proportion of CD3+ cells to total immune cells and CD4+ T-cells to total immune cells tended to be inversely correlated with HK2 tumor expression (spearman rho = -0.283, *P* = 0.077; spearman rho = -0.326, *P* = 0.040, respectively). The proportion of CD19+ B-cells to total immune cells was also inversely correlated with HK2 tumor expression (spearman rho = -0.377, *P* = 0.017; Table 1). Although the proportion of Tregs to total immune cells did not

significantly differ according to tumor HK2 expression, the proportion of Tregs to CD4+ T-cells showed a significant positive correlation with tumor HK2 expression (spearman rho = 0.480, *P* = 0.003). Of note, the ratio of CD8+ T-cell to Tregs was inversely correlated with tumor HK2 expression (spearman rho = -0.415, *P* = 0.012) (Fig. 1 B). Proportions of tumor-infiltrating macrophages, dendritic cells, and MDSCs did not differ according to HK2 tumor expression (Fig. 1 C). In addition, HK2 tumor expression did not affect M1 versus M2 polarization of macrophages (Table 1). Among the above differentially infiltrating immune cells, we further validated the association between tumor HK2 expression and CD8+ T-cells, Tregs, and their ratio in additional larger cohorts.


Tumor HK2 expression was inversely correlated with the ratio of CD8+ T-cells to Tregs

To validate the above findings, we evaluated tumor HK2, CD8+ cells, and Tregs (FOXP3+ cells) using IHC in 375 lung ADC, 118 lung SqCC, and 338 colon ADC cases (Fig. 2), as schematically described in Fig. 3 A. HK2 expression was correlated with some clinicopathologic parameters with marginal statistical significance (Supplementary Table S6, 7 and 8). In lung adenocarcinoma, EGFR mutant tumor tended to show lower HK2 expression than EGFR wild type tumor (*P* = 0.079). In colon adenocarcinoma, tumors with higher T stage tended to have higher HK2 expression than tumor with lower stage (*P* = 0.07) and when categorizing tumors T1-3 versus T4, T4 tumors showed statistically significant HK2 overexpression, compared to T1-3 tumors (*P* = 0.015, data not shown). The relationships of tumor HK2 expression with CD8+ T-cells, Tregs, and their ratio are summarized in Fig. 3 B–D. CD8+ T-cell infiltration was not significantly correlated with tumor HK2 expression. In contrast, Treg infiltration was positively correlated with tumoral HK2 expression in lung adenocarcinoma and squamous cell carcinoma cohort (lung ADC, spearman rho = 0.489, *P* < 0.001; lung SqCC, spearman rho = 0.306, *P* = 0.001), and tended to be positively correlated with tumoral HK2 expression in colon adenocarcinoma cohort (spearman rho = 0.111, *P* = 0.054) (Fig. 3 B–D). Of note, the ratio of CD8+ T-cells to Tregs was inversely correlated with HK2 tumor expression (lung ADC, spearman rho = -0.335, *P* < 0.001; lung SqCC, spearman rho = -0.236, *P* = 0.010; colon ADC, spearman rho = -0.175, *P* = 0.004) (Fig. 3 B–D). These findings were consistent with the flow cytometry analyses and demonstrated that an increase in tumor glycolysis as represented by HK2 tumor expression was inversely correlated with the ratio of CD8+ T-cells to Tregs in patients with lung cancer and colon cancer.

A. Experimental design of flow cytometry cohort



B. Lymphoid cells



C. Myeloid cells




Fig. 1 The relationship between tumor HK2 expression and tumor-infiltrating immune cell subsets analyzed by flow cytometry. **A** Flow cytometry experimental design. **B** The total immune cell and lymphoid cell infiltration statuses according to tumor HK2 expression are depicted. **C** Myeloid cell infiltration according to tumor HK2 expression is depicted. All P values were calculated using Spearman correlation analysis

Table 1 Correlations between tumor HK2 expression and immune cell subsets

Immune cell subsets	Correlation coefficient ^a	P value
Pan-immune cells (% of total cells)	0.030	0.855
CD3 + cells (% of ICs)	-0.283	0.077
CD4 + cells (% of CD3 + cells)	-0.225	0.162
CD4 + cells (% of ICs)	-0.326	0.040
CD8 + cells (% of CD3 + cells)	0.114	0.485
CD8 + cells (% of ICs)	-0.162	0.318
Treg (% of CD4 + cells)	0.480	0.003
Tregs (% of ICs)	0.239	0.161
CD8 + cells to Tregs ratio	-0.415	0.012
CD19 + cells (% of ICs)	-0.377	0.017
NK cells (% of ICs)	-0.134	0.411
Macrophages (% of ICs)	0.023	0.888
M0 (% of macrophages)	-0.116	0.477
M1 (% of macrophages)	0.089	0.584
M2 (% of macrophages)	-0.010	0.949
DCs (% of ICs)	0.091	0.604
MDSCs	0.028	0.901

Abbreviations: IC Immune cell, Tregs Regulatory T-cells, DC Dendritic cell, MDSC Myeloid-derived suppressor cell

^a Spearman correlation analysis

A lower CD8 + T-cell to Treg ratio was associated with poor survival in patients with lung SqCC

Whereas high CD8 + T-cell infiltration generally predicts a favorable prognosis and good response to immunotherapy in cancer patients, high Treg infiltration was not consistent in predicting the prognosis of patients [29]. Treg recruitment partially depends on the presence of CD8 + T-cells [30]. A significant positive correlation between Treg and CD8 + T-cell infiltration in lung and colon cancer cohorts was consistently observed (all $P < 0.001$) (Supplementary Figure S2).

We also evaluated the prognostic significance of the CD8 + T-cell to Treg ratio as well as the numbers of CD8 + T-cells and Tregs. In lung SqCC, a lower number of CD8 + TILs was associated with poor progression-free survival (PFS) and overall survival (OS) ($P = 0.008$ and $P = 0.004$, respectively) (Fig. 4 A). Of note, a lower ratio of CD8 + T-cells to Tregs was associated with poor PFS and OS ($P = 0.005$ and $P < 0.001$, respectively) (Fig. 4B). A lower number of CD8 + TILs was associated with poor OS in patients with advanced-stage (IIB–IIIB) cancer (Fig. 4 A) rather than early-stage (I–IIA) ($P = 0.075$) (Fig. 4 A). In contrast, a decreased ratio of CD8 + T-cells to Tregs was associated with poor OS in patients with both early and advanced stages ($P = 0.010$ and 0.002 , respectively). These findings suggest that the ratio of CD8 + T-cells to Tregs might be a more useful prognostic factor compared with CD8 + TILs. Together,






Fig. 2 Representative images of HK2 tumor expression relative to the CD8 + T-cell to Treg ratio from lung adenocarcinoma cases. **A** High tumor HK2 expression with a low CD8 + T-cell to Treg ratio and **B** low tumor HK2 expression with a high CD8 + T-cell to Treg ratio. (Original magnification: $\cdot 200$, bar = 100 μ m)


A. Experimental design of validation cohorts



B. Lung ADC



C. Lung SqCC



D. Colon ADC





Fig. 3 Correlation between HK2 tumor expression and immune cell composition including CD8 + T-cells, Tregs, and CD8 + T-cell to Treg ratio. **A** Experimental design of the validation cohorts. Results from 375 lung ADC patients (**B**), 118 lung SqCC patients (**C**), and 338 colon ADC patients (**D**) are depicted. All P values were calculated using Spearman correlation analysis

A. CD8⁺ T-cells



B. CD8⁺ T-cells/Tregs




Fig. 4 Survival analysis of patients with lung SqCC according to the number of CD8⁺T-cells and the CD8⁺T-cell to Treg ratio. **A** The progression-free survival (PFS) and overall survival (OS) according to the number of CD8⁺T-cells are displayed in total, early-stage (stage I–IIA), and advanced-stage (stage IIB–IIIIB) disease patients. **B** The PFS and OS according to the ratio of CD8⁺T-cells to Tregs are displayed in the total, early-stage (stage I–IIA), and advanced-stage (stage IIB–IIIIB) disease patients. The survival difference was plotted and analyzed using the Kaplan-Meier and log-rank test

these findings suggest that tumor HK2 expression influences patient survival indirectly via distinct immune profiling, specifically a low ratio of CD8+ T-cells to Tregs.


Implication of the CD8+ T-cell to Treg ratio in patients treated with immunotherapy

An additional validation cohort that included 78 non-small cell lung cancer patients with PD-1/PD-L1 blockade therapy was also investigated. Treg infiltration, but not CD8+ T-cell infiltration, was positively correlated with HK2 tumor expression (spearman rho=0.247, P=0.033). The CD8+ T-cell to Tregs ratio was also

inversely correlated with HK2 tumor expression (spearman rho = -0.288, P=0.013) (Fig. 5 A). The CD8+ T-cell to Tregs ratio was a better discriminator for prognosis compared with CD8+ T-cells and Tregs individually. Additionally, patients with a high CD8+ T-cell to Treg ratio tended to show prolonged PFS and OS after immunotherapy (P=0.113 and 0.049, respectively) (Fig. 5B).

Discussion

In this study, HK2 tumor expression as an indicator of glycolysis was associated with a different immune profile in the TME. Additionally, the present T-cell subsets



within the TME significantly differed according to the level of HK2 tumor expression. As described previously, each subset of immune cells relies on different metabolic pathway for their survival, activation, and differentiation.

CD4+ helper and CD8+ T-cells rely on glycolysis and mammalian target of rapamycin (mTOR) signaling for their metabolic activity. In contrast, Tregs rely on FAO and OXPHOS, and mTOR suppresses differentiation into Tregs [22, 31, 32]. Therefore, CD4+ helper and CD8+ T-cells are more susceptible to glucose availability compared with Tregs. Moreover, highly glycolytic tumor cells promote an acidic TME by secreting lactate, a byproduct of glycolysis.

Lactic acid inhibits the survival and effector function of T-cells [33]. Lactic metabolites also inhibit CD4+ and CD8+ T-cell mobility [34]. Similarly, knockdown of the lactate-producing enzyme lactate dehydrogenase A promoted CD8+ T-cell infiltration in the TME while decreasing the number of Tregs in mice tumors. In addition, neutralizing the TME increased CD8+ T-cell effector functions and infiltration and reinforced the response to immunotherapy in mouse tumor models [35–37]. Therefore, increased tumor glycolysis can modulate the immune profile of the TME by restricting the availability of glucose to immune cells and secreting immunosuppressive lactic acid. Consistently, this study confirmed the inverse correlation between HK2 tumor expression and the ratio of CD8+ T-cells to Tregs in human lung cancer and colon cancer tissues.

The evaluation of Treg infiltration as a prognostic marker has shown conflicting results, which might be due to higher Treg infiltration in an inflamed TME rich in CD8+ T-cells and other immune cells [29]. In this study, increased Treg infiltration was associated with a favorable prognosis, which is a paradoxical finding given their immunosuppressive (i.e., pro-tumoral) effects (Supplementary Figure S3). This might be because Treg infiltration was positively correlated with CD8+ T-cell infiltration. Therefore, some researchers have evaluated the prognostic significance of Treg infiltration compared with CD8+ T-cell infiltration (e.g., the ratio of Tregs to CD8+ T-cells or the ratio of CD8+ T-cells to Tregs), rather than Treg infiltration alone [38–40].

In this study, a high number of CD8+ TILs or a high ratio of CD8+ T-cells to Tregs was associated with a favorable prognosis. In contrast, a low ratio of CD8+ T-cells to Tregs was consistently associated with a poor prognosis in patients with early- and advanced-stage lung SqCC. Although the statistical significance was marginal, a high CD8+ T-cell to Treg ratio was associated with better prognosis in lung cancer patients treated with immunotherapy. These findings reflect the biological roles of CD8+ T-cells and Tregs in anti-tumor immunity

and emphasize the value of ratio of CD8+ T-cell to Treg ratio as a prognostic biomarker as well as a potential predictive biomarker. However, the implications of the CD8+ T-cell to Treg ratio as a predictive biomarker for immunotherapy need further study.

Tumor cells rely on increased metabolism, especially glycolysis for their survival, growth, and proliferation. This metabolic reprogramming also increases the aggressiveness of the tumor indirectly by modulating the TME into a pro-tumorigenic milieu. Therefore, attempts have been made to target tumor metabolism to enhance treatment responses and patient outcomes [41, 42]. HK2 is an interesting target for modulating metabolism. Several HK2 inhibitors have been evaluated in vitro and in mouse models, but no successful agents that clinically suppress tumor HK2 have been discovered due to their low specificity to tumor cells [43]. Even though there are more hurdles to overcome for developing clinically applicable HK2 inhibitors, there could be promising combination agents to enhance immunotherapy or chemotherapy responses by releasing immune cells from metabolic competition and directly decreasing tumor cell survival.

In summary, this study demonstrated that tumor HK2 expression/glycolysis can influence the immune profile of the TME, and an increase in tumor HK2 expression/glycolysis might be linked to a decreased CD8+ T-cell to Treg ratio, implying a poor prognosis. Although we did not investigate a detailed causal relationship between glycolysis and the TME, the above findings suggest that tumor metabolism may influence the tumor immune microenvironment into becoming pro- or anti-tumorigenic.

Conclusion

An increase in tumor HK2 expression/glycolysis may contribute to shaping the immunosuppressive/pro-tumorigenic tumor immune microenvironment by modulating/decreasing the ratio of tumor-infiltrating CD8+ T-cells to regulatory T-cells in human cancers. Thus, tumor HK2 expression/glycolysis should be further investigated as potential biomarkers or therapeutic targets in the era of cancer immunotherapy.

Abbreviations

ADC: Adenocarcinoma; AJCC: American Joint Committee on Cancer; CR: Complete response; CRC: Colorectal carcinoma; FAO: Fatty acid oxidation; HK2: Hexokinase-2; HNPCC: Hereditary nonpolyposis colorectal cancer; MD: Moderately differentiated; MDSC: Myeloid-derived suppressor cell; MSI-L or H: Microsatellite instability-low or high; MSS: Microsatellite stable; NA: Not assessable; OS: Overall survival; OXPHOS: Oxidative phosphorylation; PD: Poorly differentiated or Progressive disease; PFS: Progression free survival; PR: Partial response; PY: Pack-year; SD: Stable disease; SqCC: Squamous cell carcinoma; TAM: Tumor-associated macrophage; TIL: Tumor-infiltrating lymphocyte; TME: Tumor microenvironment; Treg: Regulatory T-cell; WD: Well differentiated.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12885-022-10239-6>.

Additional file 1: Supplementary Data S1. Clinicopathological features and immunoprofiles of patients with lung adenocarcinoma subjected to flow cytometry. **Supplementary Data S2.** Clinicopathological features and immune profiles of patients with lung adenocarcinoma subjected to immunohistochemistry. **Supplementary Data S3.** Clinicopathological features and immune profiles of patients with lung squamous cell carcinoma subjected to immunohistochemistry. **Supplementary Data S4.** Clinicopathological features and immune profiles of patients with colon adenocarcinoma subjected to immunohistochemistry. **Supplementary Data S5.** Clinicopathological features and immune profiles of non-small cell lung cancer patients with PD-1/PD-L1 blockade subjected to immunohistochemistry.

Additional file 2: Supplementary Table S1. Clinicopathological features of patients with lung adenocarcinoma subjected to flow cytometry for comprehensive immunoprofiling. **Supplementary Table S2.** Clinicopathological features of patients with lung adenocarcinoma subjected to immunohistochemistry. **Supplementary Table S3.** Clinicopathological features of patients with lung squamous cell carcinoma subjected to immunohistochemistry. **Supplementary Table S4.** Clinicopathological features of patients with colon adenocarcinoma subjected to immunohistochemistry. **Supplementary Table S5.** Clinicopathological features of non-small cell lung cancer patients with PD-1/PD-L1 blockade subjected to immunohistochemistry. **Supplementary Table S6.** HK2 expression according to clinicopathologic parameters in lung adenocarcinoma patients. **Supplementary Table S7.** HK2 expression according to clinicopathologic parameters in lung squamous cell carcinoma patients. **Supplementary Table S8.** HK2 expression according to clinicopathologic parameters in colorectal cancer patients.

Additional file 3: Supplementary Figure S1. Gating strategies for lymphoid and myeloid cells in lung adenocarcinoma. **Supplementary Figure S2.** Correlation between CD8+ T-cell infiltration and Treg infiltration in human cancer tissues. **Supplementary Figure S3.** Survival analysis of patients with lung SqCC according to the number of Tregs.

Additional file 4. Supplementary Methods.

Acknowledgements

The authors thank Medical Illustration & Design, part of the Medical Research Support Services of Yonsei University College of Medicine, for all artistic support related to this work.

Authors' contributions

Designed and supervised the study: DHC, YKJ. Performed the experiments and analyzed the data: SK, JK, SSG, JY. Performed statistical analysis: SK. Performed acquisition of clinical data: MK, BK, YTK, JK. Wrote the paper: SK, YKJ. All authors read and approved the final manuscript.

Funding

This work was supported by the Basic Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science and ICT (MSIT) (grant No: 2020R1A4A1017515).

Availability of data and materials

The datasets analyzed during the current study are available in Additional file 1.

Declarations

Ethics approval and consent to participate

This study was performed in accordance with the World Medical Association Declaration of Helsinki and was approved by the institutional review board (IRB) of Seoul National University Hospital (SNUH) (H-1404-100-572 and H-1905-115-1035). Informed consent for participation in the IHC study only cohorts was waived by the IRB of Seoul National University Hospital on the

basis that this study was a retrospective study using archived material, and did not increase risk to the patients. Fresh tumor samples from 41 lung ADC patients were used for flow cytometry analysis. All patients in this cohort agreed to the sample collection and data analysis. Written informed consent was provided by all participants for this cohort.

Consent for publication

Not applicable.

Competing interests

The authors have declared no conflicts of interest.

Author details

¹Department of Pathology, Seoul National University College of Medicine, 103 Daehak-ro, Jongno-gu, 03080 Seoul, Republic of Korea. ²Department of Pathology, Yonsei University College of Medicine, Seoul, Republic of Korea. ³Department of Biomedical Sciences, Seoul National University College of Medicine, Seoul, Republic of Korea. ⁴Department of Internal Medicine, Seoul National University College of Medicine, Seoul, Republic of Korea. ⁵Department of Thoracic Surgery, Seoul National University Hospital, Seoul National University College of Medicine, Seoul, Republic of Korea. ⁶Seoul National University Cancer Research Institute, Seoul, Republic of Korea. ⁷Department of Pathology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Republic of Korea.

Received: 14 February 2022 Accepted: 26 October 2022

Published online: 01 November 2022

References

- Topalian SL. Targeting Immune Checkpoints in Cancer Therapy. *JAMA*. 2017;318(17):1647–8.
- Reck M, Rodríguez-Abreu D, Robinson AG, Hui R, Csőszi T, Fülöp A, Gottfried M, Peled N, Tafreshi A, Cuffe S, et al. Pembrolizumab versus Chemotherapy for PD-L1–Positive Non–Small-Cell Lung Cancer. *N Engl J Med*. 2016;375(19):1823–33.
- Bellmunt J, de Wit R, Vaughn DJ, Fradet Y, Lee J-L, Fong L, Vogelzang NJ, Climent MA, Petrylak DP, Choueiri TK, et al. Pembrolizumab as Second-Line Therapy for Advanced Urothelial Carcinoma. *N Engl J Med*. 2017;376(11):1015–26.
- Sharma P, Hu-Lieskovan S, Wargo JA, Ribas A. Primary, Adaptive, and Acquired Resistance to Cancer Immunotherapy. *Cell*. 2017;168(4):707–23.
- Davis AA, Patel VG. The role of PD-L1 expression as a predictive biomarker: an analysis of all US Food and Drug Administration (FDA) approvals of immune checkpoint inhibitors. *J Immunother Cancer*. 2019;7(1):278.
- Lagos GG, Izar B, Rizvi NA. Beyond Tumor PD-L1: Emerging Genomic Biomarkers for Checkpoint Inhibitor Immunotherapy. *Am Soc Clin Oncol Educ Book*. 2020;e40:e47–57.
- Lei Y, Li X, Huang Q, Zheng X, Liu M. Progress and Challenges of Predictive Biomarkers for Immune Checkpoint Blockade. *Front Oncol*. 2021;11(609). <https://doi.org/10.3389/fonc.2021.617335>.
- Tang T, Huang X, Zhang G, Hong Z, Bai X, Liang T. Advantages of targeting the tumor immune microenvironment over blocking immune checkpoint in cancer immunotherapy. *Signal Transduct Target Therapy*. 2021;6(1):72.
- Nguyen KB, Spranger S. Modulation of the immune microenvironment by tumor-intrinsic oncogenic signaling. *J Cell Biol*. 2019;219(1). <https://doi.org/10.1083/jcb.201908224>.
- Kumagai S, Togashi Y, Sakai C, Kawazoe A, Kawazu M, Ueno T, Sato E, Kuwata T, Kinoshita T, Yamamoto M, et al. An Oncogenic Alteration Creates a Microenvironment that Promotes Tumor Progression by Conferring a Metabolic Advantage to Regulatory T Cells. *Immunity*. 2020;53(1):187–203.e188.
- Peng W, Chen JQ, Liu C, Malu S, Creasy C, Tetzlaff MT, Xu C, McKenzie JA, Zhang C, Liang X, et al. Loss of PTEN Promotes Resistance to T Cell-Mediated Immunotherapy. *Cancer Discov*. 2016;6(2):202–16.
- Koyama S, Akbay EA, Li YY, Aref AR, Skoulidis F, Herter-Sprie GS, Buczkowski KA, Liu Y, Awad MM, Denning WL, et al. STK11/LKB1 Deficiency Promotes Neutrophil Recruitment and Proinflammatory Cytokine

- Production to Suppress T-cell Activity in the Lung Tumor Microenvironment. *Cancer Res.* 2016;76(5):999–1008.
13. Marco M, Chen C-T, Choi S-H, Pelossof R, Shia J, Garcia-Aguilar J. A KRAS mutation is associated with an immunosuppressive tumor microenvironment in mismatch-repair proficient colorectal cancer. *J Clin Oncol.* 2019;37(4_suppl):609–9.
 14. Ischenko I, D'Amico S, Rao M, Li J, Hayman MJ, Powers S, Petrenko O, Reich NC. KRAS drives immune evasion in a genetic model of pancreatic cancer. *Nat Commun.* 2021;12(1):1482.
 15. Chang CH, Qiu J, O'Sullivan D, Buck MD, Noguchi T, Curtis JD, Chen Q, Gindin M, Gubin MM, van der Windt GJ, et al. Metabolic Competition in the Tumor Microenvironment Is a Driver of Cancer Progression. *Cell.* 2015;162(6):1229–41.
 16. Kim S, Jang JY, Koh J, Kwon D, Kim YA, Paeng JC, Ock CY, Keam B, Kim M, Kim TM, et al. Programmed cell death ligand-1-mediated enhancement of hexokinase 2 expression is inversely related to T-cell effector gene expression in non-small-cell lung cancer. *J Exp Clin Cancer Res.* 2019;38(1):462.
 17. Warburg O. On the Origin of Cancer Cells. *Science.* 1956;123(3191):309–19.
 18. Guerra L, Bonetti L, Brenner D. Metabolic Modulation of Immunity: A New Concept in Cancer Immunotherapy. *Cell Rep.* 2020;32(1):107848.
 19. Leone RD, Powell JD. Metabolism of immune cells in cancer. *Nat Rev Cancer.* 2020;20(9):516–31.
 20. Buszko M, Shevach EM. Control of regulatory T cell homeostasis. *Curr Opin Immunol.* 2020;67:18–26.
 21. Weinberg SE, Singer BD, Steinert EM, Martinez CA, Mehta MM, Martínez-Reyes I, Gao P, Helmin KA, Abdala-Valencia H, Sena LA, et al. Mitochondrial complex III is essential for suppressive function of regulatory T cells. *Nature.* 2019;565(7740):495–9.
 22. Delgoffe GM, Kole TP, Zheng Y, Zarek PE, Matthews KL, Xiao B, Worley PF, Kozma SC, Powell JD. The mTOR Kinase Differentially Regulates Effector and Regulatory T Cell Lineage Commitment. *Immunity.* 2009;30(6):832–44.
 23. Wang H, Franco F, Ho P-C. Metabolic Regulation of Tregs in Cancer: Opportunities for Immunotherapy. *Trends in Cancer.* 2017;3(8):583–92.
 24. Viola A, Munari F, Sánchez-Rodríguez R, Scolaro T, Castegna A. The Metabolic Signature of Macrophage Responses. *Front Immunol.* 2019;10(1462). <https://doi.org/10.3389/fimmu.2019.01462>.
 25. Veglia F, Sanseviero E, Gabrilovich DI. Myeloid-derived suppressor cells in the era of increasing myeloid cell diversity. *Nat Rev Immunol.* 2021;21(8):485–98.
 26. Amin MBESBAJCoC. *AJCC cancer staging manual.* 2017.
 27. Edge SB. *AJCC Cancer Staging Manual.* New York: Springer; 2011.
 28. Koh J, Kim HY, Lee Y, Park IK, Kang CH, Kim YT, Kim JE, Choi M, Lee WW, Jeon YK, et al. IL23-Producing Human Lung Cancer Cells Promote Tumor Growth via Conversion of Innate Lymphoid Cell 1 (ILC1) into ILC3. *Clin Cancer Res.* 2019;25(13):4026–37.
 29. Saleh R, Elkord E. FoxP3 + T regulatory cells in cancer: Prognostic biomarkers and therapeutic targets. *Cancer Lett.* 2020;490:174–85.
 30. Spranger S, Spaepen RM, Zha Y, Williams J, Meng Y, Ha TT, Gajewski TF. Upregulation of PD-L1, IDO, and T(regs) in the melanoma tumor microenvironment is driven by CD8(+) T cells. *Sci Transl Med.* 2013;5(200):200ra116.
 31. Macintyre AN, Gerriets VA, Nichols AG, Michalek RD, Rudolph MC, Deoliveira D, Anderson SM, Abel ED, Chen BJ, Hale LP, et al. The glucose transporter Glut1 is selectively essential for CD4 T cell activation and effector function. *Cell Metab.* 2014;20(1):61–72.
 32. Michalek RD, Gerriets VA, Jacobs SR, Macintyre AN, MacIver NJ, Mason EF, Sullivan SA, Nichols AG, Rathmell JC. Cutting edge: distinct glycolytic and lipid oxidative metabolic programs are essential for effector and regulatory CD4 + T cell subsets. *J Immunol.* 2011;186(6):3299–303.
 33. Brand A, Singer K, Koehl GE, Koltz M, Schoenhammer G, Thiel A, Matos C, Bruss C, Klobuch S, Peter K, et al. LDHA-Associated Lactic Acid Production Blunts Tumor Immunosurveillance by T and NK Cells. *Cell Metab.* 2016;24(5):657–71.
 34. Haas R, Smith J, Rocher-Ros V, Nadkarni S, Montero-Melendez T, D'Acquisto F, Bland EJ, Bombardieri M, Pitzalis C, Perretti M, et al. Lactate Regulates Metabolic and Pro-inflammatory Circuits in Control of T Cell Migration and Effector Functions. *PLoS Biol.* 2015;13(7):e1002202.
 35. Calcinotto A, Filipazzi P, Grioni M, Iero M, De Milito A, Ricupito A, Cova A, Canese R, Jachetti E, Rossetti M, et al. Modulation of microenvironment acidity reverses anergy in human and murine tumor-infiltrating T lymphocytes. *Cancer Res.* 2012;72(11):2746–56.
 36. Pilon-Thomas S, Kodumudi KN, El-Kenawi AE, Russell S, Weber AM, Luddy K, Damaghi M, Wojtkowiak JW, Mulé JJ, Ibrahim-Hashim A, et al. Neutralization of Tumor Acidity Improves Antitumor Responses to Immunotherapy. *Cancer Res.* 2016;76(6):1381–90.
 37. Zhang YX, Zhao YY, Shen J, Sun X, Liu Y, Liu H, Wang Y, Wang J. Nanoenabled Modulation of Acidic Tumor Microenvironment Reverses Anergy of Infiltrating T Cells and Potentiates Anti-PD-1 Therapy. *Nano Lett.* 2019;19(5):2774–83.
 38. Sato E, Olson SH, Ahn J, Bundy B, Nishikawa H, Qian F, Jungbluth AA, Frosina D, Gnjatic S, Ambrosone C, et al. Intraepithelial CD8 + tumor-infiltrating lymphocytes and a high CD8+/regulatory T cell ratio are associated with favorable prognosis in ovarian cancer. *Proc Natl Acad Sci USA.* 2005;102(51):18538–43.
 39. Takada K, Kashiwagi S, Goto W, Asano Y, Takahashi K, Takashima T, Tomita S, Ohsawa M, Hirakawa K, Ohira M. Use of the tumor-infiltrating CD8 to FOXP3 lymphocyte ratio in predicting treatment responses to combination therapy with pertuzumab, trastuzumab, and docetaxel for advanced HER2-positive breast cancer. *J Translational Med.* 2018;16(1):86.
 40. Shang B, Liu Y, Jiang S-J, Liu Y. Prognostic value of tumor-infiltrating FoxP3 + regulatory T cells in cancers: a systematic review and meta-analysis. *Sci Rep.* 2015;5(1):15179.
 41. Koudhi S, Ben Ayed F, Benamar Elgaaied A. Targeting Tumor Metabolism: A New Challenge to Improve Immunotherapy. *Front Immunol.* 2018;9(353). <https://doi.org/10.3389/fimmu.2018.00353>.
 42. Farhadi P, Yarani R, Dokaneheifard S, Mansouri K. The emerging role of targeting cancer metabolism for cancer therapy. *Tumor Biology.* 2020;42(10):1010428320965284.
 43. Ciscato F, Ferrone L, Masgras I, Laquatra C, Rasola A. Hexokinase 2 in Cancer: A Prima Donna Playing Multiple Characters. *Int J Mol Sci.* 2021;22(9):4716.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

