

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

## International Journal of Surgery Case Reports

journal homepage: [www.casereports.com](http://www.casereports.com)

# Intrahepatic portal vein thrombosis due to postoperative biliary obstruction successfully treated by a partial thrombectomy combined with thrombolytic drug therapy

Naoya Imamura<sup>a</sup>, Atsushi Nanashima<sup>a,\*</sup>, Yuki Tsuchimochi<sup>a</sup>, Takeomi Hamada<sup>a</sup>, Koichi Yano<sup>a</sup>, Masahide Hiyoshi<sup>a</sup>, Yoshiro Fujii<sup>a</sup>, Kunihide Nakamura<sup>b</sup>

<sup>a</sup> Division of Hepato-Biliary-Pancreas Surgery, Department of Surgery, University of Miyazaki Faculty of Medicine, 5200 Kihara Kiyotake, Miyazaki, 889-1692, Japan

<sup>b</sup> Division of Cardiovascular Surgery, Department of Surgery, University of Miyazaki Faculty of Medicine, 5200 Kihara Kiyotake, Miyazaki, 889-1692, Japan

## ARTICLE INFO

## Article history:

Received 9 August 2017

Accepted 21 November 2017

Available online 27 November 2017

## Keywords:

Pancreatic head cancer

Stricture of biliary anastomosis

Organized portal thrombosis

Thrombolytic therapy

Surgery

Case report

## ABSTRACT

**INTRODUCTION:** This case report aims to inform pancreatic surgeons about our perioperative management of intrahepatic portal vein thrombosis caused by an obstruction of hepaticojejunostomy (HJ) after pancreaticoduodenectomy (PD).

**CASE PRESENTATION:** A 65-year-old woman was diagnosed with pancreas head carcinoma involving the superior mesenteric vein (SMV). Pancreaticoduodenectomy combined with SMV resection was followed by HJ. Twisting or narrowing was not evident during anastomosis. Total bilirubin values progressively increased to 13 mg/dL on day 5. At that time, we suspected anastomotic occlusion and found complete portal thrombosis of the left liver. Therefore, emergency re-anastomosis of the HJ was followed by thrombectomy, which was not completely successful and did not completely recover initial portal flow. Thrombolytic drugs improved obstructive jaundice, eradicated the organized thrombosis and recovered the portal flow by day 30. The post-operative course was uneventful.

**DISCUSSION:** A thrombosis immediately formed in the portal vein due to biliary obstruction of an anastomotic site. We speculated that biliary dilation and related inflammation caused a relative increase in arterial flow and decreased portal flow at the localized part of the umbilical portion. Although early surgical thrombectomy was attempted soon after the primary operation, the organized thrombosis persisted. However, thrombolytic therapy eradicated the thrombosis.

**CONCLUSION:** Careful anastomosis of HJ during PD was necessary to avoid postoperative biliary stricture. This type of complication affects intrahepatic blood flow, particularly via the portal vein. Although immediate re-anastomosis or thrombectomy is applied, organized thrombosis cannot always be surgically removed.

© 2017 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

The only curative treatment option for pancreatic carcinoma is R0 resection [1]. However, this often requires combined vascular resections of the portal vein to avoid exposing the tumor surface at the dissection plane [2]. Portal flow during combined vascular resection can become transiently blocked within 30 min of vascular reconstruction. A 30-min block without a portal bypass is probably safe, but postoperative portal vein thrombosis (PVT) can arise after invasive procedures as hepatectomy or pancreaticoduodenectomy (PD) [3–5]. The mechanism of PVT might involve various factors such as the intraoperative occlusion of portal flow, a kinked


portal trunk, stenosis of a vascular anastomosis and surrounding inflammation [3–8]. We have often experienced portal stenosis after postoperative intraabdominal infection or inflammation. Thus, we speculate that postoperative surrounding inflammation is one cause of postoperative PVT. The present patient underwent PD with a portal vein anastomosis due to combined resection of the portal and superior mesenteric veins. These procedures were immediately followed by postoperative PVT and obstructive jaundice due to stenosis of a hepatico-jejunostomy (HJ).

## 2. Case presentation

A 65-year-old woman was admitted to our institution with a tumor of the pancreatic head. Endoscopic fine needle aspiration confirmed pancreas head ductal adenocarcinoma.


\* Corresponding author.

E-mail address: [a.nanashima@med.miyazaki-u.ac.jp](mailto:a.nanashima@med.miyazaki-u.ac.jp) (A. Nanashima).



**Fig 1.** Enhanced computed tomography findings.

(a) Black arrow, pancreas head carcinoma has invaded SMV. (b) White arrow, resection and anastomosis of involved SMV combined with pancreaticoduodenectomy.



**Fig. 2.** Postoperative enhanced computed tomography findings at day 5 after PD.

(a) Black arrow, anastomotic stricture has caused dilated intrahepatic duct; white arrow accompanying complete thrombosis of left liver. (b) White arrow, removal of a part of the portal vein thrombosis.


The main PA involved the superior mesenteric vein (SMV; Fig. 1a) and PD combined with SMV and vascular anastomosis was scheduled. Distant tumor metastasis was not evident and thus the scheduled operation proceeded followed by direct anastomosis of the portal trunk and the SMV (Fig. 1b). R0 resection was achieved without tumor exposure at the dissection plane. Intraoperative Doppler ultrasonography confirmed sufficient flow in the intrahepatic portal vein (PV) and artery. The size of the orifice of the hepatic duct stump was about 10 mm and the HJ was achieved by end-to-side anastomosis using 5-0 absorbable surgical thread. Twisting or narrowing of the anastomotic site was not evident at that time.

Hyperbilirubinemia was confirmed on day 1 after PD and total bilirubin values progressively increased to 13 mg/dL on day 5 after PD. The entire intrahepatic bile duct was dilated and we suspected an occluded anastomotic site. Follow-up ultrasonography showed decreased left PV flow and enhanced computed tomography at day 5 showed complete portal thrombosis of the left liver (Fig. 2a). Emergency surgery for the biliary stricture and PV thrombosis immediately proceeded on the same day. Macroscopic assessment of the intraoperative findings did not show a constricted HJ, but the sutures were removed and the HJ was re-anastomosed with careful placement of an intraductal tube biliary

stent. We then attempted thrombectomy via the umbilical portal vein or the main portal trunk under cut-down, but little thrombus was removed (Fig. 2b). We considered that the intra-portal thrombosis might be organized, and if so, it would be very difficult to mechanically remove during surgery. The thrombectomy was not sufficient enough to allow good portal flow. We therefore administered 15,000 units of intravenous heparin for 10 days followed by oral warfarin to dissolve the PV thrombosis. This approach dramatically improved obstructive jaundice and obliterated the organized thrombosis by day 30 (Fig. 3). Thus, the patient was discharged on day 31 without any further complications. The post-operative course followed up at an out-patient clinic was uneventful.

### 3. Discussion

The cause and mechanism of PVT after PD in the present patient could not be defined, but PV reconstruction can result in PVT by transient occlusion [3–5]. However, Doppler ultrasonography confirmed intrahepatic portal flow. The wide lymphadenectomy around the portal vein might also have caused PVT via mechanical injury caused by holding or lifting the portal vein. Considering these operative procedures, the subsequent events might have been avoided. The PV anastomosis required no more than 20 min



**Fig. 3.** Postoperative enhanced computed tomography findings at day 30 after PD. Biliary obstruction has improved. Impacted portal thrombus has disappeared and portal flow in left liver has recovered after thrombolytic drug therapy.

and vascular stenosis at the anastomotic site was not consistently evident during the surgical procedures. The most likely cause of PVT was postoperative obstructive jaundice due to an obstruction of HJ anastomosis. A biliary stricture was not evident during the procedure and the hepatic duct was about 10 mm in length. The hepatic duct was thin, but twisting of the anastomosis was not visible at either the first or the second operation. A tube stent was not deployed as the clinical usefulness of such stents or drainage has not yet been established [8]. Jaundice and the intrahepatic biliary dilation progressed and subsequent PVT in the left liver were remarkable by day 5 after PD in our patient. Not only can cholangitis can cause PVT [9–11], but PVT can also cause cholangitis or biliary stenosis [12,13]. We suppose that cholangitis induces arterial flow in the Glissonian pedicle, which might relatively reduce portal flow. Adjacent biliary inflammation might influence portal stenosis or reduced flow because of a thin venous wall and low venous flow.

Acute PVT after surgery is a lethal complication and emergency thrombectomy is necessary when an organ is injured or a vital abnormality is identified [11,14].

Anti-coagulant or thrombolytic drugs are usually administered to stable patients [15–17]. However, the PVT was completely organized and mechanical thrombectomy was not feasible by any reasonable route. The acute stage of PVT might be soft and floating and a balloon catheter can usually achieve thrombectomy. Thus, we eventually abandoned this strategy. Liver function and jaundice gradually recovered after re-anastomosis of the HJ. Image analysis showed that the intrahepatic PVT persisted after the jaundice improved. Intravenous heparin followed by oral warfarin obviously recovered the occluded portal flow. Although intraoperative thrombectomy was not accomplished, thrombolytic or anticoagulant drugs were useful to achieve recanalization. This work has been reported in line with the SCARE criteria [18].

#### 4. Conclusion

Postoperative complications comprising biliary occlusion and organized PVT in the liver of a patient with pancreatic head cancer were treated by PD with portal vein reconstruction due to combined portal vein resection. Obstructive jaundice at the anastomotic site in the HJ was treated by re-anastomosis, but surgical thrombectomy failed. Thrombolytic and anti-coagulant drugs administered late after re-operation recovered the obstructive jaundice and eliminated the PVT.

#### Conflict of interest

Naoya Imamura, Atsushi Nanashima, Yuki Tsuchimochi, Takeomi Hamada, Kouichi Yano, Masahide Hiyoshi, Yoshiro Fujii, and Kunihide Nakamura, declare that they have no conflict of interest.

#### Funding

Naoya Imamura, Atsushi Nanashima, Yuki Tsuchimochi, Takeomi Hamada, Kouichi Yano, Masahide Hiyoshi, Yoshiro Fujii, and Kunihide Nakamura, declare that they have no sources of funding.

#### Ethical approval

We have received ethics approval at our institute without reference number in this case.

#### Consent

Written consent obtained from patients.

#### Author contribution

Naoya Imamura; Attending physician, writing paper.  
 Atsushi Nanashima; Editing, contributor and supervisor. Chairman and Director of the department and a main operator.  
 Yuki Tsuchimochi; Attending physician.  
 Takeomi Hamada; Attending physician.  
 Kouichi Yano; Attending physician.  
 Masahide Hiyoshi; Attending physician.  
 Yoshiro Fujii; Attending physician.  
 Kunihide Nakamura; Attending physician. cardiovascular surgeon and Chairman of a different division of the Department of Surgery, University of Miyazaki, Japan.

#### Guarantor

The guarantor is Atsushi Nanashima.  
 Professor Kunihide Nakamura is a cardiovascular surgeon and Chairman of a different division of the Department of Surgery, University of Miyazaki, Japan.

## References

- [1] J.M. Alamo, L.M. Marín, G. Suarez, C. Bernal, J. Serrano, L. Barrera, M.A. Gómez, J. Muntané, F.J. Padillo, Improving outcomes in pancreatic cancer: key points in perioperative management, *World J. Gastroenterol.* 20 (2014) 14237–14245.
- [2] T.C. Chua, A. Saxena, Extended pancreaticoduodenectomy with vascular resection for pancreatic cancer: a systematic review, *J. Gastrointest. Surg.* 14 (2010) 1442–1452.
- [3] M.J. Kang, J.Y. Juang, Y.R. Chang, W. Jung, S.W. Kim, Portal vein patency after pancreaticoduodenectomy for periampullary cancer, *BJS* 102 (2015) 77–84.
- [4] I. Watanabe, Y. Ito, E. Akimoto, Y. Sekine, Y. Haruyama, K. Amemiya, S. Miyano, T. Kosaka, M. Machida, T. Kitabatake, K. Kojima, Postoperative portal vein thrombosis and gastric hemorrhage associated with late-onset hemorrhage from the common hepatic artery after pancreaticoduodenectomy, *Korean J. Hepatobiliary Pancreat. Surg.* 20 (2016) 44–47.
- [5] N.J. Zyromski, T.J. Howard, Acute superior mesenteric-portal vein thrombosis after pancreaticoduodenectomy: treatment by operative thrombectomy, *Surgery* 143 (2008), 566–556.
- [6] M. Kikuchi, W. Nishizaki, K. Tsuruya, I. Hamada, T. Higashi, K. Sakuma, H. Shiozawa, J. Aoki, R. Nagashima, J. Koizumi, Y. Arase, K. Shiraishi, M. Matsushima, T. Mine, Acute portal vein thrombosis due to chronic relapsing pancreatitis: a fistula between a pancreatic pseudocyst and the splenic vein, *Clin. J. Gastroenterol.* 7 (2014) 52–57.
- [7] A.H. Stockland, D.L. Willingham, R. Paz-Fumagalli, H.P. Grewal, J.M. McKinney, C.B. Hughes, E.M. Walser, Pancreas transplant venous thrombosis: Role of endovascular interventions for graft salvage, *Cardiovasc. Intervent. Radiol.* 32 (2009) 279–283.
- [8] N.O. Glebova, C.W. Hicks, K.M. Piazza, C.J. Abularrage, A.M. Cameron, R.D. Schlick, C.L. Wolfgang, J.M. Black III, Technical risk factors for portal vein reconstruction thrombosis in pancreatic resection, *J. Vasc. Surg.* 62 (2015) 424–433.
- [9] L.R. Warren, M.D. Chandrasegaram, D.J. Madigan, P.M. Dolan, E.L. Neo, C.S. Worthley, Falciform ligament abscess from left sided portal pyaemia following malignant obstructive cholangitis, *World J. Surg. Oncol.* 10 (2012) 278.
- [10] C. Cuffari, E. Seidman, J. DuBois, P. Brochu, F. Alvarez, *Eur. J. Pediatr.* 156 (1997) 186–189.
- [11] H.C. Oh, D.W. Seo, S.H. Kim, Portal vein thrombosis after EUS-guided pancreatic cyst ablation, *Dig. Dis. Sci.* 57 (2012) 1965–1967.
- [12] R. Nyman, H. Al-Suhaibani, I. Kagevi, Portal vein thrombosis mimicking tumour and causing obstructive jaundice. A case report, *Acta Radiol.* 37 (1996) 685–687.
- [13] C.H. Lin, G.H. Lo, K.H. Lai, K.T. Mok, H.B. Pan, W.B. Teng, Obstructive jaundice caused by calcified portal venous thrombosis: a case report, *Chin. Med. J.* 58 (1996) 50–53.
- [14] P. Handa, M. Crowther, J. Douketis, Portal vein thrombosis: a clinician-oriented and practical review, *Clin. Appl. Thromb./Hemost.* 20 (2014) 498–506.
- [15] A.M. Sharma, D. Zhu, Z. Henry, Portal vein thrombosis: when to treat and how? *Vasc. Med.* 21 (2016) 61–69.
- [16] M. Schultheiß, D. Bettinger, R. Thimme, Nonsurgical therapeutic options in portal vein thrombosis, *Viszeralmedizin* 30 (2014) 388–392.
- [17] J.L. Kaplan, S.L. Weintraub, J.P. Hunt, A. Gonzalez, J. Lopera, A. Brazzini, Treatment of superior mesenteric and portal vein thrombosis with direct thrombolytic infusion via an operatively placed mesenteric catheter, *Am. Surg.* 70 (2004) 600–604.
- [18] R.A. Agha, A.J. Fowler, A. Saetta, I. Barai, S. Rajmohan, D.P. Orgill, for the SCARE Group, The SCARE statement: consensus-based surgical case report guidelines, *Int. J. Surg.* 34 (2016) 180–186.

## Open Access

This article is published Open Access at [sciencedirect.com](https://www.sciencedirect.com). It is distributed under the [IJSCR Supplemental terms and conditions](#), which permits unrestricted non commercial use, distribution, and reproduction in any medium, provided the original authors and source are credited.