

# Bone morphogenetic protein 4 (BMP-4) and epidermal growth factor (EGF) inhibit metalloproteinase-9 (MMP-9) expression in cancer cells

Nathalie Bibens Lulan<sup>1</sup> and Yves St-Pierre <sup>1</sup>

<sup>1</sup> INRS-Institut National de la Recherche Scientifique, INRS-Institut-Armand-Frappier, Boul. des Prairies, Laval, Québec, Canada

Correspondence to: Yves St-Pierre, email: yves.st-pierre@iaf.inrs.ca

Keywords: Gene expression, matrix metalloproteinase-9, bone morphogenetic protein-4, EGF, gremlin

Received: February 02, 2015

Accepted: March 16, 2015

Published: March 23, 2015

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## ABSTRACT

**Matrix metalloproteinase-9 (MMP-9) plays a central role in the progression of the cancer. While a large number of studies have contributed to our understanding of the molecular mechanisms responsible for upregulating MMP-9 gene expression in normal and cancer cells, our knowledge on the signals that suppress MMP-9 expression is much more limited. Here, we report that EGF and BMP-4 cooperate to inhibit MMP-9 expression in cancer cells. Treatment with EGF reduces the expression of MMP-9 at both mRNA while augmenting BMP-4 expression. Interestingly, recombinant BMP-4 suppressed constitutive and PMA-induced MMP-9 expression in both fibrosarcoma and breast cancer cells. Addition of gremlin a natural inhibitor of BMP-4, inhibited the suppression of MMP-9 by EGF. The suppression of MMP-9 by BMP-4 likely occurs at the transcriptional level since BMP-4 suppressed MMP-9 mRNA expression and activation of a reporter vector encoding the human MMP-9 promoter. The suppressive effect of BMP-4 occurs via Smad1/5/8 and is specific since BMP-4 did not inhibit MMP-2 while BMP-2 was ineffective in suppressing MMP-9. Taken together, these results are consistent with a new paradigm for the role of EGF and BMPs in controlling MMP gene expression in cancer cells.**

## INTRODUCTION

Tumor cell proliferation, invasion and metastasis are mediated, at least in part, through the degradation of the extracellular matrix by metalloproteinases (MMPs) locally produced by tumor and stromal cells. A point in case is MMP-9, which plays a significant role in cell invasion and metastasis in several cancers [1-4]. Consequently, a large number of studies have focused their attention on identifying and characterizing the molecular mechanisms that are responsible for increased expression of MMP-9 in cancer. In contrast, our knowledge on the molecular mechanisms that suppress MMP-9 remains largely unexplored and largely focused on epigenetic mechanisms, including DNA methylation [5-7]. Using an experimental mouse lymphoma model, where MMP-9 plays a central role, our group has recently shown that the EGF/EGR1 (epidermal growth factor/ early growth response 1) pathway was involved in the repression of MMP-9

expression by stromal cells [8]. More specifically, we found that expression of MMP-9 in stromal was repressed by the activation of the *EGR1* gene induced by EGF secreted by tumor cells and it correlates with decreased MMP-9. How EGF/EGR-1 suppresses MMP-9, however, remains unknown.


Bone morphogenetic proteins (BMPs) were initially identified by Marshall Urist in 1960 [9]. Today, more than 30 members the BMP family have been reported. BMPs are extracellular proteins that bind membrane anchored serine/threonine receptors and induce a signal through intracellular R-Smads (1/5/8) signalization. Activated R-Smads form a heteromeric complex with the classic Smad4 which translocates to the nucleus and activates the transcription of BMP target genes [10]. Until recently, BMPs have been well known for their role in bone formation and as being powerful developmental regulators. The functions of BMPs, however, seems to go well beyond their role in bone formation. They also

control a broad range of biological activities. During embryonic development, for example, BMP-4 is involved in gastrulation, mesoderm formation, hematopoiesis and development of several organs and tissues [11-13]. In fact, their role as critical signaling molecules that regulate cell fate decision, cell differentiation, cell survival and vasculogenesis, motility and cell adhesion are now well recognized. Not surprisingly, these properties have attracted the attention of an increasing number of investigators in the field of cancer. Abnormally high levels of BMPs have been reported in many cancer and have been associated with a poor prognosis, consistent with their ability promote dissemination, invasiveness and migration [14, 15]. Their role in cancer progression, however, remains ambiguous since several studies have shown that BMPs have anti-tumor functions [16, 17]. For example, while we know that BMPs can increase MMP expression in some cancer cells, such as gastric or prostate cancer cells [18, 19], others have shown that BMP-4 treatment of C3HT101/2 stem cells blocks MMP-3 and MMP-13 expression [20]. Such an ability to suppress MMPs has also been reported for BMP-4 and -6 [16, 21]. These results suggest the existence of a functional relationship between BMPs and MMPs. Here, we have investigated whether BMPs could be involved in the suppression of MMP-9 by EGF/EGR-1.


## RESULTS

### MMP-9 and BMP-4 expression in HT1080 cells following EGF/EGR-1 activation

Using an *in vivo* mouse model, we have previously shown that EGF suppresses MMP-9 gene expression [8]. Using the human HT1080 cells, an *in vitro* cell model commonly used to study the molecular mechanisms regulating human MMP-9, we have confirmed that EGF can suppress MMP-9 in a dose-dependent manner at the mRNA level (Fig. 1A). Similar results were obtained at the protein level as shown by Western blot and zymography (Fig. 1B-C). Such inhibition of MMP-9 by EGF was also observed in human breast cancer cell lines (Supplementary Fig.S1). Because BMP-4 has recently been shown to inhibit choroidal neovascularization by inhibiting VEGF and MMP-9 expression [21], we next investigated whether BMP-4 could be involved in the suppression of MMP-9 expression by EGF/EGR-1. We indeed found that suppression of MMP-9 by EGF in HT1080 cells correlated with increased *BMP-4* gene expression (Fig. 2A). Such increased in BMP-4 expression was also observed in stable transfectants of HT1080 cells expressing constitutive levels of EGR-1, a transcription factor known to be activated EGF (Fig. 2B). We thus tested whether treatment of HT1080 cells with recombinant BMP-4 (rBMP-4) could



**Figure 1: Suppression of MMP-9 in HT1080 cells following treatment with EGF.** (A) MMP-9 mRNA expression in absence or presence of recombinant EGF. Levels of transcripts were measured 16 h after adding EGF. GAPDH was used as loading and specificity control. Suppression of MMP-9 expression by increasing doses of EGF was confirmed at the protein level using (B) Western-blot analysis and (C) gelatin zymography. Data are representative of at least two independent experiments.



**Figure 2: BMP-4 expression in HT1080 cells by EGF/EGR1.** (A) BMP-4 mRNA expression in absence or presence of recombinant EGF. Levels of transcripts were measured 16 h after adding EGF. GAPDH was used as loading and specificity control. (B), BMP-4 and MMP-9 expression in HT1080 cells following stable expression of an expression vector encoding human EGR1. GAPDH was used as loading and specificity control. Data are representative of at least two independent experiments.

downregulate MMP-9 expression in HT1080 cells. Our results showed that rBMP-4 suppressed the expression of MMP-9 by HT1080 cells at both mRNA and protein levels (Fig. 3 A-B). The ability of BMP-4 to inhibit MMP-9 was specific since rBMP-4 did not modulate MMP-2. Moreover, no detectable decrease of MMP-9 was observed following treatment with recombinant BMP-2 (Supplementary Fig. S2). We also confirmed that rBMP-4 was biologically active, as shown by its ability to induce phosphorylation of Smad1/5/8 (Fig. 3C). The use of a luciferase reporter containing the human MMP-9 promoter further indicated that BMP-4 is likely to suppress MMP-9 expression at the transcriptional level (Fig. 3D). Ectopic expression of flagged-BMP-4 in HT1080 cells was also effective in suppressing MMP-9, further supporting the specificity of the inhibition (Fig. 4A-C). The ability of BMP-4 to inhibit MMP-9 expression was not restricted to HT1080 cells. We observed a similar decrease of MMP-9 by BMP-4 at both mRNA and proteins levels using the human breast epithelial MDA-MB-231, MDA-MB-468 and SK-BR3 cells, which express MMP-9 following stimulation with PMA (Fig. 5).


### Restoration of MMP-9 gene expression with gremlin

To further confirm that BMP-4 inhibits the expression of MMP-9, we used gremlin, a natural

antagonist of BMP-4 which is highly expressed in cancer-associated stromal cells [22]. Our results showed that treatment of HT1080 cells with increasing concentrations of recombinant gremlin induced a detectable increase in MMP-9 expression in HT1080 cells at both mRNA and protein levels (Fig. 6A-B). Gremlin also blocked the EGF-mediated MMP-9 suppression (Fig. 6C). It also restored PMA-induced MMP-9 expression in cells treated with BMP-4. We also observed a significant increase in MMP-9 transcription and protein in MDA-MB-231 during treatment with gremlin (Supplementary Fig. S2). The effectiveness of gremlin to block BMP-4-induced signals was further confirmed by monitoring the phosphorylation of Smad1/5/8 (Fig. 6D).

### BMP-4 over-expression inhibited HT1080 cellular invasion

Because elevated level of MMP-9 is well known to increase the migratory properties of tumor cells *in vivo* and *in vitro* [1, 2, 4], we next tested whether BMP-4 was able to suppress the invasion of HT1080 cells. For this purpose, we compared the *in vitro* transmigration of HT1080 cells through Matrigel in absence and presence of recombinant BMP-4. Our results showed that BMP-4 did indeed reduce the invasiveness of HT1080 cells as compared to control cells (Fig. 7).



**Figure 3: MMP-9 is decreased following BMP-4 stimulation in HT1080 cells.** (A) MMP-9 mRNA expression 24 and 48h following stimulation with recombinant BMP-4 (200 ng/ml). GAPDH was used as loading and specificity control. In (B), a zymogram (top gel) showing reduced MMP-9 secretion in supernatants of HT1080 cells treated for 16h with recombinant BMP-4 (200ng/ml). The lower panel shows the MMP-9 mRNA level of the treated cells. (C) Western blot analysis showing expression of MMP-9 and phosphorylation of Smad1/5 after treatment with human recombinant BMP-4. (D) Luciferase activity of in HT1080 cells transfected with a luciferase reporter vector containing the MMP-9-promoter following treatment with human recombinant BMP-4. Statistical analyses were carried out using Student's t test for unpaired samples. (\* =  $p \leq 0,05$ ; \*\* =  $p \leq 0,005$ ).

## Smad6 affect the down-regulation of MMP-9 induced by BMP-4


BMPs are well-known for their ability to induce signaling through the canonical pathway of Smad1/5/8, co-Smad (Smad4) and inhibitory Smad6 and 7. To determine whether this molecular pathway was involved in MMP-9 suppression by BMP-4, we carried a transient transfection of HT1080 cells using an expression vector encoding Smad6, which dimerizes with Smad1/5/8, thereby blocking its translocation to the nucleus [23]. We first confirmed that *de novo* expression of Smad6 was effective in blocking the phosphorylation of Smad1/5/8 while reducing the ability of BMP-4 to suppress MMP-9 (Fig. 8).

## DISCUSSION


A large number of studies have contributed to our understanding of the molecular mechanisms responsible for upregulating MMP-9 gene expression in normal and cancer cells. In contrast, our knowledge on the repressive signals that suppress MMP-9 expression remains fragmented. In a previous study, we had found that EGF suppressed the expression of MMP-9 *in vivo* and in

HT1080 cells [8]. Here we have extended this work by showing that: 1) EGF-mediated suppression of MMP-9 was also observed in other cancer cells, most notably in human breast cancer cells; 2) stimulation with EGF also induced expression of BMP-4; this effect was observed in both HT1080 fibrosarcoma cells and MDA-MB-231, MDA-MB-468, SK-BR3 mammary epithelial cells; 3) the suppression of MMP-9 by BMP-4 likely occurs at the transcriptional level since BMP-4 suppresses MMP-9 mRNA expression and suppresses the activation of a reporter vector encoding the human MMP-9 promoter; BMP-4-mediated suppression of MMP-9 likely involves the Smad1/5/8 pathway. Moreover, transfection of a vector encoding Smad6 suggests that the suppressive effect of BMP-4 possibly involves the Smad1/5/8 pathway. Finally, we found that the effect of BMP-4 on MMP-9 was specific since BMP-4 did not inhibit MMP-2 while BMP-2 was ineffective in suppressing MMP-9.

BMPs are often associated with a poor prognosis in patients with different types of cancers. This seems to be largely due to the fact that BMPs target genes encoding osteoblast proteins and osteoblast-specific transcription factors [24]. For example, in prostate cancer, BMP-6 promotes osteoblastic activity of prostate cancer cells and confer them with a more invasive phenotype [25, 26]. Similarly, BMP-7 promotes expression of VEGF which




**Figure 4: *De novo* expression of BMP-4 reduces MMP-9 gene expression.** (A) MMP-9 mRNA expression in mock-transfected HT1080 cells or HT1080 cells transfected with an expression vector encoding a flagged human BMP-4 (pCMV-BMP-4-Flag). GAPDH was used as loading and specificity control. The lower panel represents the control Western blot gels showing *de novo* expression of flagged BMP-4 in transfected cells. (B) Zymography (left panel) showing MMP-9 in the supernatant of transfected cells as in (A). (C) Quantitative analyses of MMP-9 expression by imaging densitometry is shown on the right histogram, which represents the means of independent experiments shown in (A). Data are representative of at least three independent experiments. (\* =  $p \leq 0,05$ )




**Figure 5: MMP-9 expression is suppressed by BMP-4 in MDA-MB-231, MDA-MB-468 and SKBR3 cells.** MMP-9 mRNA expression in PMA-stimulated cells (20ng/ml) treated with or without recombinant BMP-4 (50 or 100 ng/ml) in MDA-MB-231 cells (A), SK-BR3 (B) and MDA-MB-468 cells (C). GAPDH was used as loading and specificity control. Data are representative of at least three independent experiments.

contributes to the osteoblastic metastasis lesions [27]. A number of studies have shown, however, that BMPs may have a dual role in cancer and may in fact hamper tumor progression. For example, the bone morphogenetic protein pathway is active in human colon adenomas but inactivated in colorectal cancer [28]. A mutation of the receptor BMPRI1A (bone morphogenetic protein receptor type IA) also promotes colon cancer [29]. Other studies have further shown that BMPs may also exert an anti-tumorigenic effect in breast cancer [16, 17]. In patients with familial adenomatous polyposis, BMP-2 can also block cell growth while promoting apoptosis of mature epithelial cells [30]. Such suppressive effect for BMPs seems to be particularly true for BMP-4. A recent clinical study has shown that patients high levels of BMP-4 have a better chance for survival when compared to patients with high-grade glioma who have a lower expression level of BMP-4 [31]. Also, BMP-4 negatively regulates tumor-


initiating cells [32]. Whether such dual role in cancer is due to the ability to regulate MMPs is an interesting possibility since MMPs have also been reported to have a dual role in cancer. Our results herein and those from other groups showing that BMPs can suppress MMP gene



**Figure 6: MMP-9 is increased following treatment of HT1080 cells with gremlin.** (A) MMP-9 mRNA expression in cells treated with increasing concentrations of gremlin. GAPDH was used as loading and specificity control. (B) Zymogram showing MMP-9 in the supernatant of HT1080 cells treated with recombinant BMP-4 in absence or presence of gremlin. (C) Kinetic analysis showing MMP-9 expression and phosphorylation of Smad1/5 in cells treated treatment with human recombinant BMP-4 with or without gremlin. Data are representative of at least three independent experiments. (D) MMP-9 mRNA expression in absence or presence of recombinant EGF and gremlin in HT1080 cells.



**Figure 7: BMP-4 inhibits *in vitro* HT1080 cell invasion.** (A) HT1080 cell migration across Matrigel with or without recombinant BMP-4. (B) Migrated HT1080 cells were quantified. Data are average  $\pm$  SEM; n = 10 unit areas. Data are representative of at least three independent experiments. (\*\*\*)  $p \leq 0,001$ .



**Figure 8: Smad6 reduces the ability of BMP-4 to suppress MMP-9 expression.** (A) MMP-9 mRNA expression in mock-transfected HT1080 cells or HT1080 cells transfected with an expression vector encoding a flagged Smad6 (pCS2-Smad6-Flag). MMP-9 expression was measured 16h after transfection. The lower panel represents the control Western blot gels showing *de novo* expression of Smad6 in transfected cells. GAPDH and actin were used as loading and specificity controls for RT-PCR and Western blot analyses respectively. In (B), a zymogram showing MMP-9 in the supernatant of HT1080 transfected cells. Data are representative of at least three independent experiments.

expression supports this possibility. For example, Otto et al., have shown that BMP-4 blocks expression of MMP-3 and MMP-13 in C3HT101/2 stem cells [20]. BMP-6 has also been shown to inhibit MMP-9 expression in breast cancer cells [16]. Such suppression of MMP-9 by BMPs has also been reported during choroïdal neovascularization [21]. Our results showing that EGF induced BMP-4 further suggest the existence of a new functional relationship between both proteins that may be critical in the control of the invasive behavior of cancer cells. Taken together, these results are consistent with a new paradigm for the role of BMPs in controlling MMP gene expression in cancer cells. Future investigations will nevertheless be needed to solidly establish such a model and how the canonical Smad pathway is involved. While our preliminary studies with Smad6 are interesting in this regard, it is important to note that BMPs can activate the non-canonical MAPK (p38, MEK/ERK), Tak1/Tab1 and PI3K pathways [33, 34]. Interestingly, these non-canonical signal pathways are widely recognized to activate various transcription factors, including Nuclear factor-kappaB (NF-κB) [35], a transcription factor well-known for its ability to modulate MMP-9 expression [36]. Future investigations will thus be needed to identify which of these pathways are involved.

Overall, our study supports the idea that targeting EGF/BMPs could be a valuable alternative to inhibit MMP-dependent tumor progression, at least in some types of cancer. Cautions need to be taken, however, since MMP activity is not limited to extracellular matrix degradation and extends to proteolysis of various proteins bound to the cell membrane or even proteins secreted by cells. The repertoire of MMP substrates, or degradome, does indeed vary depending on the type and the different grades of tumors and in many cases, local increase of MMPs may have different impact on tumor progression [37]. Clearly, prudence will be needed to fully exploit the functional relationship between BMPs and MMPs for the treatment of cancer. Lessons from the failure of MMP inhibitors in clinical trials will be valuable to this regard.

## MATERIALS AND METHODS

### Cell lines and reagents

The human MDA-MB-231, MDA-MB-468, SKBR3, MCF7 and HT1080 cell lines were obtained from the American Tissue Culture Collection (ATCC, Manassas, VA). The MDA-MB-231 MDA-MB-468, MCF7 and HT1080 cells were maintained in culture in Dulbecco's modified Eagle complete medium (DMEM) [supplemented with 10% (v/v) FCS, 2 mmol/L L-glutamine and 10 mmol/L Hepes buffer]. The SKBR3 cell was maintained in culture in McCoy's medium [supplemented with 10% (v/v) FCS, 2 mmol/L L-glutamine and 10 mmol/L Hepes

buffer]. Recombinant human EGF was purchased from Prospec Technogene (Ness Ziona, Israel). PMA (Phorbol-12-myristate-13-acetate) was purchased from Sigma-Aldrich (St. Louis, MI). Recombinant Human BMPs and Gremlin were obtained from Peprotech (Rocky Hill, NJ, USA). All other reagents were purchased from Sigma-Aldrich, unless otherwise indicated.

### RNA extraction and semi quantitative RT-PCR

Total RNA was isolated using Trizol reagent according to the manufacturer's instructions (Invitrogen, lifetechnologies, Burlington, ON, Canada). After reverse transcription, cDNAs were amplified using the following conditions: 94°C for 1 min, followed by 30-35 cycles of the following: 94°C for 1 min, 58-64°C for 1 min (depending on the primers), and 72°C for 1 min, followed by a final extension step at 72°C for 10 min. PCR was performed in a thermal cycler (MJ Research, Watertown, MA). PCR assays using equal amounts of RNAs that were reverse-transcribed and amplified by PCR for 25 to 40 cycles with gene-specific primers (Table S1) confirmed that the amplification was in the linear range for each gene. As an internal control, amplification of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) mRNA was carried out by RT-PCR using specific primers. Amplified products were analyzed by electrophoresis on 1 % agarose gels using SYBR Safe DNA gel (Invitrogen) staining and UV illumination.

### Vectors, transfection and luciferase assays

The vector encoding Smad6 (pCS2-Flag-Smad6) was obtained from Addgene (Cambridge, MA). The vector encoding BMP-4 (pCMV-Flag-BMP4) was obtained from Sino Biological Inc. (Benjing, China). The plasmid encoding the luciferase reporter vector containing a fragment encompassing the essential consensus sequences for the transcriptional activity of the *mmp-9* promoter (pGL3-MMP-9) has been described [36]. For transfection, cells were plated at equal density 24h before transfection. Cells were then washed twice and transfected with 2 μg of DNA using DNAfectin 2100 according to manufacturer's protocol (ABM, Richmond, BC, Canada). After transfection, cells were incubated in complete medium at 37°C in 5% CO<sub>2</sub> for 4 h. The culture medium was then changed to complete DNEM medium for 20 h. For zymography only, the culture medium was changed for serum free medium for 16 h. Luciferase activity was measured using the Luciferase Assay System protocol (Promega, Madison, WI, USA) and a luminometer (Lumat LB 9507, Berthold). The transfection efficiency was monitored by co-transfection with the pCMV/βGal plasmid encoding β-galactosidase (Promega, Madison, WI, USA). The β-galactosidase activity was detected

by a colorimetric enzyme assay using o-nitrophenyl- $\beta$ -galactopyranoside as a substrate. The ratio of luciferase to  $\beta$ -galactoside activity in each served as a measure of normalized luciferase activity.

### Gelatin Zymography

Zymography was performed in polyacrylamide gels that had been cast in the presence of gelatin as previously described [36]. Briefly, samples were suspended in loading buffer and, without prior denaturation, were run on a 7.5% SDS-polyacrylamide gel containing 0.5 mg/ml of gelatin. After electrophoresis, gels were washed to remove SDS and incubated for 18 h at 37°C in a denaturing buffer. Gels were subsequently stained with Coomassie brilliant blue G-250 and destained in 30% (v/v) methanol/ 10% acetic acid to detect gelatinase secretion. The proteolytic activity of MMP-9 was identified as a clear band on a blue background.

### Matrigel invasion assay

Invasion assay was carried out using Matrigel-coated invasion chambers (BD Biosciences). Briefly, cells were resuspended in culture medium without FBS at the concentration of  $6 \times 10^4$  cells/ml and 500  $\mu$ l of the cell suspension was seeded on the upper chamber. The lower chamber was filled with 700  $\mu$ l of the culture medium without cells and 10% FBS was added as a chemoattractant. Cells were incubated with or without rhBMP4 for 18 h in a humidified tissue culture incubator, at 37°C, 5% CO<sub>2</sub> atmosphere. After removing non-invasive cells with a cotton swab. Invasive cells adhering to membrane of the upper chamber were fixed with methanol 100% for 1 min and then stained with a Borax 1% toluidine 1% solution for 1 min. Number of cells on the membrane was counted under a light microscope at 40X magnification.

### ACKNOWLEDGEMENTS

This research was supported by a grant from the Canadian Institutes for Health Research of Canada (CIHR).

### CONFLICT OF INTEREST

The authors of this manuscript have no conflicts of interest to declare.

### REFERENCES

1. Bernhard EJ, Gruber SB, Muschel RJ. Direct evidence linking expression of matrix metalloproteinase 9 (92-kDa

- gelatinase/collagenase) to the metastatic phenotype in transformed rat embryo cells. *Proc Natl Acad Sci U S A* 1994; 91:4293-4297.
2. Egeblad M, Werb Z. New functions for the matrix metalloproteinases in cancer progression. *Nat Rev Cancer*. 2002; 2:161-174.
3. Kessenbrock K, Plaks V, Werb Z. Matrix metalloproteinases: regulators of the tumor microenvironment. *Cell*. 2010; 141:52-67.
4. Lopez-Otin C, Matrisian LM. Emerging roles of proteases in tumour suppression. *Nat Rev Cancer*. 2007; 7:800-808.
5. Chicoine E, Esteve PO, Robledo O, Van Themsche C, Potworowski EF, St-Pierre Y. Evidence for the role of promoter methylation in the regulation of MMP-9 gene expression. *Biochem Biophys Res Commun*. 2002; 297:765-772.
6. Couillard J, Demers M, Lavoie G, St-Pierre Y. The role of DNA hypomethylation in the control of stromelysin gene expression. *Biochem Biophys Res Commun*. 2006; 342:1233-1239.
7. Sato N, Maehara N, Su GH, Goggins M. Effects of 5-aza-2'-deoxycytidine on matrix metalloproteinase expression and pancreatic cancer cell invasiveness. *J Natl Cancer Inst*. 2003; 95:327-330.
8. Bouchard F, Belanger SD, Biron-Pain K, St-Pierre Y. EGR-1 activation by EGF inhibits MMP-9 expression and lymphoma growth. *Blood*. 2010; 116:759-766.
9. Nogami H, Urist MR. The classic : a morphogenetic matrix for differentiation of cartilage in tissue culture. *Clin Orthop Relat Res*. 2009; 467:3063-3067.
10. Ten Dijke P, Korchynskyi O, Valdimarsdottir G, Goumans MJ. Controlling cell fate by bone morphogenetic protein receptors. *Mol Cell Endocrinol*. 2003; 211:105-113.
11. Cho KW, Kim JY, Song SJ, Farrell E, Eblaghie MC, Kim HJ, Tickle C, Jung HS. Molecular interactions between Tbx3 and Bmp4 and a model for dorsoventral positioning of mammary gland development. *Proc Natl Acad Sci U S A* 2006; 103:16788-16793.
12. Pizette S, Abate-Shen C, Niswander L. BMP controls proximodistal outgrowth, via induction of the apical ectodermal ridge, and dorsoventral patterning in the vertebrate limb. *Development*. 2001; 128:4463-4474.
13. Winnier G, Blessing M, Labosky PA, Hogan BL. Bone morphogenetic protein-4 is required for mesoderm formation and patterning in the mouse. *Genes Dev*. 1995; 9:2105-2116.
14. Kejner AE, Burch MB, Sweeny L, Rosenthal EL. Bone morphogenetic protein 6 expression in oral cavity squamous cell cancer is associated with bone invasion. *Laryngoscope*. 2013; 123:3061-3065.
15. Owens P, Polikowsky H, Pickup MW, Gorska AE, Jovanovic B, Shaw AK, Novitskiy SV, Hong CC, Moses HL. Bone morphogenetic proteins stimulate mammary fibroblasts to promote mammary carcinoma cell invasion.

16. Wang C, Hu F, Guo S, Mi D, Shen W, Zhang J, Qiao Y, Zhu T, Yang S. BMP-6 inhibits MMP-9 expression by regulating heme oxygenase-1 in MCF-7 breast cancer cells. *J. Cancer Res Clin Oncol.* 2011; 137:985-995.
17. Buijs JT, van der Horst G, van den Hoogen C, Cheung H, de Rooij B, Kroon J, Petersen M, van Overveld PG, Pelger RC, van der Pluijm G. The BMP2/7 heterodimer inhibits the human breast cancer stem cell subpopulation and bone metastases formation. *Oncogene.* 2012; 31:2164-2174.
18. Darby S, Cross SS, Brown NJ, Hamdy FC, Robson CN. BMP-6 over-expression in prostate cancer is associated with increased Id-1 protein and a more invasive phenotype. *J Pathol.* 2008; 214:394-404.
19. Kang MH, Oh SC, Lee HJ, Kang HN, Kim JL, Kim JS, Yoo YA. Metastatic function of BMP-2 in gastric cancer cells: the role of PI3K/AKT, MAPK, the NF-kappaB pathway, and MMP-9 expression. *Exp Cell Res.* 2011; 317:1746-1762.
20. Otto TC, Bowers RR, Lane MD. BMP-4 treatment of C3H10T1/2 stem cells blocks expression of MMP-3 and MMP-13. *Biochem Biophys Res Commun.* 2007; 353:1097-1104.
21. Xu J, Zhu D, Sonoda S, He S, Spee C, Ryan SJ, Hinton DR. Over-expression of BMP4 inhibits experimental choroidal neovascularization by modulating VEGF and MMP-9. *Angiogenesis.* 2012; 15:213-227.
22. Sneddon JB, Zhen HH, Montgomery K, van de Rijn M, Tward AD, West R, Gladstone H, Chang HY, Morganroth GS, Oro AE, Brown PO. Bone morphogenetic protein antagonist gremlin 1 is widely expressed by cancer-associated stromal cells and can promote tumor cell proliferation. *Proc Natl Acad Sci U S A.* 2006; 103:14842-14847.
23. Hata A, Lagna G, Massague J, Hemmati-Brivanlou A. Smad6 inhibits BMP/Smad1 signaling by specifically competing with the Smad4 tumor suppressor. *Genes Dev.* 1998; 12:186-197.
24. Reddi AH, Reddi A. Bone morphogenetic proteins (BMPs): from morphogens to metabologens. *Cytokine Growth Factor Rev.* 2009; 20:341-342.
25. Dai JL, Keller J, Zhang J, Lu Y, Yao Z, Keller ET. Bone morphogenetic protein-6 promotes osteoblastic prostate cancer bone metastases through a dual mechanism. *Cancer Res.* 2005; 65:8274-8285.
26. Darby S, Cross SS, Brown NJ, Hamdy FC, Robson CN. BMP-6 over-expression in prostate cancer is associated with increased Id-1 protein and a more invasive phenotype. *J Pathol.* 2008; 214:394-404.
27. Dai J, Kitagawa Y, Zhang J, Yao Z, Mizokami A, Cheng S, Nör J, McCauley LK, Taichman RS, Keller ET. Vascular endothelial growth factor contributes to the prostate cancer-induced osteoblast differentiation mediated by bone morphogenetic protein. *Cancer Res.* 2004; 64:994-999.
28. Kodach LL, Bleuming SA, Musler AR, Peppelenbosch MP, Hommes DW, van den Brink GR, van Noesel CJ, Offerhaus GJ, Hardwick JC. The bone morphogenetic protein pathway is active in human colon adenomas and inactivated in colorectal cancer. *Cancer.* 2008; 112:300-306.
29. Kodach LL, Wiercinska E, de Miranda NF, Bleuming SA, Musler AR, Peppelenbosch MP, Dekker E, van den Brink GR, van Noesel CJ, Morreau H, Hommes DW, Ten Dijke P, Offerhaus GJ, Hardwick JC et al. The bone morphogenetic protein pathway is inactivated in the majority of sporadic colorectal cancers. *Gastroenterology.* 2008; 134:1332-1341.
30. Hardwick JC, Van Den Brink GR, Bleuming SA, Ballester I, Van Den Brande JM, Keller JJ, Offerhaus GJ, Van Deventer SJ, Peppelenbosch MP. Bone morphogenetic protein 2 is expressed by, and acts upon, mature epithelial cells in the colon. *Gastroenterology.* 2004; 126:111-121.
31. Ivanova T, Zouridis H, Wu Y, Cheng LL, Tan IB, Gopalakrishnan V, Ooi CH, Lee J, Qin L, Wu J, Lee M, Rha SY, Huang D, Liem N, Yeoh KG, Yong WP, Teh BT, Tan P. Integrated epigenomics identifies BMP4 as a modulator of cisplatin sensitivity in gastric cancer. *Gut.* 2013; 62:22-33.
32. Piccirillo SG, Reynolds BA, Zanetti N, Lamorte G, Binda E, Broggi G, Brem H, Olivi A, Dimeco F, Vescovi AL. Bone morphogenetic proteins inhibit the tumorigenic potential of human brain tumour-initiating cells. *Nature.* 2006; 444:761-765.
33. Barneda-Zahonero B, Miñano-Molina A, Badiola N, Fadó R, Xifró X, Saura CA, Rodríguez-Alvarez J. Bone morphogenetic protein-6 promotes cerebellar granule neurons survival by activation of the MEK/ERK/CREB pathway. *Mol Biol Cell.* 2009; 20:5051-5063.
34. Zhou Q, Heinke J, Vargas A, Winnik S, Krauss T, Bode C, Patterson C, Moser M. ERK signaling is a central regulator for BMP-4 dependent capillary sprouting. *Cardiovasc Res.* 2007; 76:390-399.
35. Zhang YE. Non-Smad pathways in TGF-beta signaling. *Cell Res.* 2009; 19:128-139.
36. Estève PO, Chicoine E, Robledo O, Aoudjit F, Descoteaux A, Potworowski EF, St-Pierre Y. Protein kinase C-zeta regulates transcription of the matrix metalloproteinase-9 gene induced by IL-1 and TNF-alpha in glioma cells via NF-kappa B. *J Biol Chem.* 2002; 277:35150-35155.
37. Cauwe B, Martens E, Proost P, Opdenakker G. Multidimensional degradomics identifies systemic autoantigens and intracellular matrix proteins as novel gelatinase B/MMP-9 substrates. *Integr Biol.* 2009; 1:404-426.