

ORAL PRESENTATION

Open Access

# Toxoplasmosis: a rare cause of IRIS in HIV infected patients. Case series

Ruxandra Moroti<sup>1,2\*</sup>, Daniela Munteanu<sup>1</sup>, Mihaela Rădulescu<sup>1,2</sup>, Adriana Hristea<sup>1,2</sup>, Iulia Niculescu<sup>1,2</sup>, Raluca Mihăilescu<sup>1</sup>, Roxana Petre<sup>1,3</sup>, Raluca Hrișcă<sup>1,3</sup>, Raluca Jipa<sup>1,3</sup>, Ana Maria Petrescu<sup>1</sup>, Maria Nica<sup>4</sup>, Mihai Lazăr<sup>1,2</sup>, Anca-Ruxandra Negru<sup>1</sup>, Irina Lăpădat<sup>1</sup>, Angelica Teniță<sup>1</sup>, Victoria Aramă<sup>1,2</sup>

From The 9th Edition of the Scientific Days of the National Institute for Infectious Diseases Prof Dr Matei Bals Bucharest, Romania. 23-25 October 2013

## Background

Cerebral toxoplasmosis is one of the main 3 intracerebral opportunistic infections in HIV positive patients, along with cryptococcosis and tuberculosis. In comparison to these last 2 entities, toxoplasmosis does not provoke or very rarely provokes reconstitution syndromes.

## Methods

We analyzed a case series of 3 patients with cerebral toxoplasmosis admitted in the Adults III Department of the National Institute for Infectious Diseases "Prof. Dr. Matei Bals" in 2012-2013.

## Results

Three patients, one male and 2 women, aged 55 years old, respectively 41 and 42 year-old, all 3 diagnosed concomitantly with HIV infection (as very late presenters) and cerebral toxoplasmosis, with a CD4 count of 6, 6 and 7/cmm respectively, viral loads (VL) of 254,000, 57,000 and 156,000 copies/mL respectively, and CSF viral load below the plasmatic VL in all 3 cases. We recorded minimal abnormalities of CSF analysis regarding the number of cells and biochemical exams; all had positive PCR for *Toxoplasma gondii* in the CSF and positive serology (IgG). All 3 had intracerebral lesions (abscesses) and all were biopsied at the neurosurgery department for diagnostic purpose before knowing their HIV-positive status. They received high doses of oral trimethoprim/sulfamethoxazole (T/S) for toxoplasmosis and antiretroviral therapy in the first 2 weeks after the diagnosis. They repeated cerebral imagery (MRI) after 3 weeks of T/S and had no regression of the

size of lesions (although with the decreasing of perilesional edema) and new lesions, in two cases without having corresponding symptoms; in all 3 cases the CD4 count increased in the first month more than 100%. The search for another cause for the augmentation of their brain lesions was negative. Maintaining the same medication, the next imagery exams showed improvement in 2 out of 3 cases, in which the outcome was favorable with almost complete neurological recovery. In the remaining case the evolution was unfavorable (death).

## Conclusions

In our 3 cases we presumed a paradoxical toxoplasmosis IRIS, with little or no clinical deterioration strictly linked with imagery exams depreciation in 2 out of 3 cases but with a fatal evolution in one case. Even rarely reported, the toxoplasmosis IRIS could be taken into account in some situations.

## Authors' details

<sup>1</sup>National Institute for Infectious Diseases "Prof. Dr. Matei Bals", Bucharest, Romania. <sup>2</sup>Carol Davila University of Medicine and Pharmacy, Bucharest, Romania. <sup>3</sup>Central University Emergency Military Hospital Dr Carol Davila, Bucharest, Romania. <sup>4</sup>Clinical Hospital of Infectious and Tropical Diseases "Dr. Victor Babeș", Bucharest, Romania.

Published: 16 December 2013

doi:10.1186/1471-2334-13-S1-O7

Cite this article as: Moroti et al.: Toxoplasmosis: a rare cause of IRIS in HIV infected patients. Case series. *BMC Infectious Diseases* 2013 13(Suppl 1):O7.

\* Correspondence: ruxandra\_moroti@yahoo.com

<sup>1</sup>National Institute for Infectious Diseases "Prof. Dr. Matei Bals", Bucharest, Romania

Full list of author information is available at the end of the article