



EMPIRICAL/THEORETICAL STUDY

Expanding the scope and relevance of health interventions: Moving beyond clinical trials and behavior change models

KHARY K. RIGG, PhD¹, HILARY H. COOK, PhD² & JOHN W. MURPHY, PhD³

¹Department of Mental Health Law & Policy, Louis de la Parte Florida Mental Health Institute, University of South Florida, Tampa, FL, USA, ²Social Sciences Department, Miami-Dade College, Miami, FL, USA, and ³Department of Sociology, University of Miami, Coral Gables, FL, USA

Abstract

An overemphasis on clinical trials and behavior change models has narrowed the knowledge base that can be used to design interventions. The overarching point is that the process of overanalyzing variables is impeding the process of gaining insight into the everyday experiences that shape how people define health and seek treatment. This claim is especially important to health decision-making and behavior change because subtle interpretations often influence the decisions that people make. This manuscript provides a critique of traditional approaches to developing health interventions, and theoretically justifies what and why changes are warranted. The limited scope of these models is also discussed, and an argument is made to adopt a strategy that includes the perceptions of people as necessary for understanding health and health-related decision-making. Three practical strategies are suggested to be used with the more standard approaches to assessing the effectiveness and relevance of health interventions.

Key words: *health interventions, clinical trials, behavior change models, photovoice, ethnography, narratives, theory, public health*

(Accepted: 27 June 2014; Published: 21 July 2014)

Epidemiology is the study of population health. This work is most commonly done by empirical assessments of populations, in order to identify risk factors and calculate the odds that persons with certain characteristics will contract a disease. Epidemiologists and public health practitioners use the concept of risk to link various sociodemographic factors to the incidence or prevalence of a problem. Risk factors are used, in part, because they provide fast and simple answers to guide public health policy.

This strategy has been referred to as “black box” epidemiology (Susser & Susser, 1996). What is suggested by this phrase is that very little insight is gained into how persons view or will likely respond to certain social or environmental conditions by the use of this model. The so-called objective features of a community may indicate that undesirable conditions are present (e.g., poverty, violence), but how persons interpret these factors is equally important for dealing with health and illness. For this reason, Krieger (2005) recommends that epidemiologists

should focus on data that are more “embodied” in communities.

However, epidemiologists and others who work in the area of public health (e.g., social epidemiologists, community psychologists) have tried to be holistic and attuned to community sentiment. For example, one of the major organizing concepts within modern public health is the epidemiological triad (Page, Cole, & Timmreck, 1995). This triad includes an agent, host, and environment as important considerations. This conceptualization of what epidemiology encompasses demonstrates a high level of flexibility and holism, but most studies are often relatively shallow. And although this strategy may include contextual variables, how these factors are interpreted is not necessarily the focus of attention. Most often, several convenient contextual variables are highlighted but merely treated as supplementary factors.

Especially noteworthy is that public health is still heavily burdened by the influence of a biomedical model because of the historical context of the

Correspondence: Khary Rigg, Department of Mental Health Law & Policy, Louis de la Parte Florida Mental Health Institute, University of South Florida, 13301 Bruce B. Downs Blvd., 33612 Tampa, FL, USA. E-mail: rigg@usf.edu

© 2014 K. K. Rigg et al. This is an Open Access article distributed under the terms of the Creative Commons CC-BY 4.0 License (<http://creativecommons.org/licenses/by/4.0/>), allowing third parties to copy and redistribute the material in any medium or format and to remix, transform, and build upon the material for any purpose, even commercially, provided the original work is properly cited and states its license.

Citation: Int J Qualitative Stud Health Well-being 2014, 9: 24743 - <http://dx.doi.org/10.3402/qhw.v9.24743>

(page number not for citation purpose)

discipline. At its inception as a discipline in the 19th century, epidemiology sought to address the spread of infectious diseases. As part of this legacy, today's standard epidemiological methods are not focused on theoretical premises, but on improving methodological and statistical designs (Morabia, 2007; Saracci, 2007). Generally, the emphasis of public health research is on refining clinical trial approximations through the use of increasingly complex mathematical models of health problems. But critics claim that this methodology is inappropriate for a field that is supposed to be as much about humans in their environmental context as pathogens and disease (Agar, 2003; Krieger, 2012).

Because quantitative indicators seem straightforward and manageable as a starting point, policy makers often rely on such studies to inform interventions. Although this strategy can be useful, especially for rapid assessment and emergency response, these studies may not capture the reality of the persons who will be affected by such programs. As a result, many development projects have the potential to be irrelevant and, in some cases, harmful. Other writers (Biehl, 2007) have shared similar concerns about the dehumanizing potential that health interventions can have when persons' everyday experiences and perspectives are not seriously considered.

Advances in public health research

Like all disciplines, public health has gone through different stages of development (Tulchinsky & Varivakova, 2009). The first era of public health was characterized by the sanitation revolution and general environmental and hygienic advances. The second stage focused on individual health behaviors, and is the target of the critique undertaken. The third era of public health is what has been called for since the 1970s, but it is yet to be fully realized. The third era is described by the World Health Organization's "Health for All" goals, and abandons the emphasis on individual level behaviors in exchange for understanding health more comprehensively and aiming to improve quality of life as a human right (Kickbusch, 2003).

One of the major institutional documents that changed the way that public health is approached originated in Canada in 1974 and is called the Lalonde Report (Lalonde, 1981). This document set off what is considered the third era of public health and challenged the purely biomedical view of health. This document was the beginning of a redefinition of health promotion that shifts the focus from individual interventions and supports a holistic approach to health.

In the mid-1980s, an international discussion began that embraces this new public health. Essentially, several important documents were published by the World Health Organization that situated health outcomes at the center of development policy, redefined the goal of health policies to include social and economic well-being, and called for a serious reorientation away from individual risk factors toward addressing context and meaning in the social environment (Kickbusch, 2003). Simultaneous discussion in academia stressed the need to modernize the field of public health that led to an attempt to promote a model that better connects the needs of people and communities, along with broadening of the disciplinary base to include other fields that address more ecologically the health of populations. Although much research in the discipline remains stuck in the second wave, the individual and behaviorally focused era of public health, there are some writers that have accepted the challenge to move public health forward. Agar (2003), for example, outlines how the epidemiological triad of host-agent-environment paradigm that runs through epidemiology can be greatly enhanced by broadening this scheme to consider the "person-in-context."

Traditional epidemiology and epistemology

A major philosophical influence at the time epidemiology emerged, and still operative today in most sciences, was René Descartes (1985) who argued that the way to truth is to separate the subjective (mind) from the objective (body). This Cartesian dualism is the epistemology that sustains the scientific method and led to the use of the experimental design as the gold standard for clinical medicine (Zaner, 1988). As a consequence of this dualism, the study of human illness became viewed as a natural (objective) science, whereby subjectivity (human experience) was reduced in importance (Winkelman, 2009). Accordingly, the biomedical model began to permeate the field of epidemiology. The success of the biomedical model is based on imagery that was heavily influenced both by Louis Pasteur's work in germ theory in the mid-1800s and bacteriology that gained momentum with Robert Koch's breakthroughs in the later part of that century. Nonetheless, dualism lent credibility to this emergent type of causal analysis.

During Koch's time of great scientific advancements, physicians viewed bacteriology as the way of improving health and believed that focusing on bacteria was the correct path to cure disease (Gradmann, 2009). This belief is still dominant today, and bacteriology's reductionism and isolationism permeates both medical practice and the field of

epidemiology, in the form of searching for unique causal agents.

Hence, epidemiology and medicine has the methodological goal of implementing experimental designs, but mostly through the use of sophisticated statistical models. The question remains whether this method is the best way to understand the intersection of pathogens and the human context. In other words, researchers are supposed to study disease in the human context, and not merely the biological transmission of pathogens. In this regard, experiments create a very strict situation with minimal variety. This process severely limits what is considered relevant to studies in health. Nonetheless, the assumption is that general laws can be found through these restricted cases, and moreover assumes that these laws are similar to those found in physics (Cassell, 1991; Reiser, 1981). This elevation of experimental design, especially in relation to something as tentative as human decision-making relating to health, may be too reductionist to generate meaningful insights.

This traditional outlook assumes that people inhabit a “real world” made of objects with properties that exist outside of interpretation (Murphy & Min Choi, 1997). Variables such as socioeconomic status or poor mental health are assumed to impinge on individual behavior in predictable ways. This viewpoint supports today’s dominant perspective on society and generally reinforces the logic and use of biomedicine to study disease (Engel, 1977). Nonetheless, the social and behavioral sciences may not be acquiring knowledge that is relevant to actual human beings. In the health field in particular, applying the “gold standard” randomized control trial, without including other approaches, decouples the medical from the human and may result in complete irrelevance to everyday life and the implementation of health interventions (Hesse-Biber, 2012).

Contemporary perspectives, such as phenomenology (Husserl, 2012; Merleau-Ponty, 1996) and social constructionism (Berger & Luckmann, 1966; Lock & Strong, 2010), contend that attempting to distill objective data from lived experience is a wrongheaded approach to the study of human behavior. In addition, this narrow focus on specific definitions of rationality limits the questions, theories, and methodologies that can be used by researchers (Susser, 2004). Indeed, numerous writers argue that the study of human beings and their social worlds should not mimic the natural sciences. Instead of separating the world into objective facts and subjective perceptions, all knowledge is understood to involve a knower (Schutz, 1967). In other words, data are always entrenched in some perspective. In this regard, the classic epidemiological triad

that seeks to understand health in terms of agent-environment-host demonstrates that the goal should be to try to place disease in an interpretive human context (Page et al., 1995).

Critical overview

Critics argue that the social and behavioral sciences are more complicated than the natural sciences. Specifically important is that constructs in the social sciences are “second degree,” or models of the concepts that are used by people as they function in their everyday worlds (Schutz, 1953). Because of the theoretical orientations that are dominant in public health, these abstractions are thought to be more valid than the schemas persons use to make sense of their lives. However, this method may obscure how they think about their health or any other issue. Valid concepts, in other words, may be more closely tied to the strategies persons use to interpret and manage events (Berger & Luckmann, 1966).

A development within public health called the “new public health” (Hills, 2000) is an attempt to refocus these efforts, and include community input in evaluations of health and the planning of interventions (Frenk, 1993; Murphy & Rigg, 2014), because the traditional methodologies overlook how people actively construct the realities in which they live. The point is that people do not necessarily think of themselves as composites of risk factors, and quite often decisions and events that are not overtly about health influence how they think about their well-being. In a recent study in the Dominican Republic (Cook, 2013), for example, a direct question about personal health prompted persons to discuss economic and environmental conditions. Most important about this insight is that these persons expect that any effective intervention must extend beyond personal or even interpersonal solutions.

What constitutes a reasonable intervention, therefore, depends on the type of “local reasoning” that is typically not captured by standardized instruments. In this sense, the imagery used in risk factor modeling does not coincide with how people view themselves and their health (Raphael, 2003). As a result, the traditional approach to assessing health has limited the scope of health assessments to variables that can be neatly circumscribed by checklists and introduced into predictive models (Smith, 1998).

For most researchers, only knowledge that is separated from judgments is truthful, and the way to attain this knowledge is by equating reliable and valid data with technical operations (Murphy, 1992). Nonetheless, by seeking to neatly operationalize the factors that persons consider when making decisions

regarding their health, researchers can easily lose sight of the context or the web of meaning surrounding health-relevant phenomena. Although there is a place for these types of studies in public health and social science, the first step to any meaningful understanding of health phenomena should not be the obfuscation of the human element.

To be able to understand a phenomenon and create valid constructs that truly represent a group of people, researchers must be comfortable with getting close to the data. Having data that do not fit easily into categorical variables should be valued in research, especially when first investigating a health issue or health behavior in a previously under-studied socio-environmental context (Quimby, 2006). In order to attain an informed understanding of how people make health-relevant decisions, researchers must move beyond the simple identification of risk factors and grasp how people interpret or construct the meanings of things, events, and relationships. Specifically, health planners should be expected to engage a community in genuine dialog, in order to grasp the various “hidden” issues that might be operating (Murphy & Rigg, 2014; Wallerstein & Duran, 2010).

Promising strategies underutilized in public health

We suggest three strategies to help accomplish this goal. First, narratives (Agnew, 2006; Charmaz, 1999; Rigg & Murphy, 2013b) are a useful, but underutilized approach that reflects the idea that a person’s reality is situational. Illness narratives (Williams, 1984), in particular, provide an innovative and useful way of approaching the subjective aspects of the illness experience (Ezzy, 2000). The norms of a patient’s local context, which narratives can help identify, should ideally be used to guide an intervention. The need and course of intervention, in other words, should be driven by the various narratives and storylines in a patient’s life and not preconceived disease frameworks (Frank, 1998; Rigg & Murphy, 2013a). Narrative approaches are sensitive to the situations of people (Kleinman & Kleinman, 1996), and can bring into view relevant details of persons’ lives that in most cases are of critical importance to the success of health interventions and programs (Charon, 2011).

Narrative medicine, for example, has been receiving some attention as a new model for clinical practice (Charon, 2001). This approach to medicine recognizes the value of patients’ narratives in practice, research, and education. Proponents of narrative medicine have argued that most medical schools train physicians to treat health problems as simply a medical issue, without taking into account the

specific psychological and social history of the patient (Remen, 2002). Narrative medicine places patients’ stories at the center of medical practice and education to encourage a more patient-centered approach to care. Not so much a new specialty as a new frame for clinical work and research, narrative medicine can validate the experience of the patient and encourage physician self-reflection, leading to improved health outcomes (Greenhalgh & Hurwitz, 1999).

Second, photovoice is a research method well-suited to capturing and conveying the point of view of participants. Succinctly defined as a process by which people can identify, represent, and enhance their community through a specific photographic technique (Wang, Cash, & Powers, 2000), photovoice entails providing study participants with cameras, allowing them to record, discuss, and communicate to others their realities as seen through their eyes (Wang & Burris, 1997). The production of a photograph and the photographer’s description of the photo provide immediate data and the foundation for building shared knowledge (Newman, 2010). Photovoice is consistent with the core tenets of community-based participatory research (e.g., empowerment, participation) and has been used successfully to address a variety of serious health issues including HIV (Rhodes & Hergenrather, 2007) and chronic pain (Baker & Wang, 2006).

Photovoice methodology has the potential to strengthen the quality and validity of research by using the perspectives of participants to generate new understanding about health-related experiences and decision-making. In this way, the identification and documentation of problems (i.e., barriers to health care utilization) are made by the participants themselves. It is important to note that the use of photovoice also provides an opportunity for vulnerable populations (i.e., homeless) to share their experiences with the research community, service providers, and policymakers; groups that these populations typically have little access to (Lopez, Eng, Robinson, & Wang, 2005). Photovoice methodology sets out to capture and convey the point of view of the participant or patient and allows their experiences to reveal themselves in their own terms (Booth & Booth, 2003). In a recent project (True, 2014), photovoice was used successfully to engage military veterans in communicating their experiences regarding challenges to getting their health care needs met. The data generated by this project not only informed new policy decisions by the Veterans Health Administration but were also used to create a photo exhibit that showed community members how the aftermath of war can affect the health of returning soldiers.

Third, ethnography has the unique capacity to help capture the cultural nuances and context that are often missing from traditional decision-making models. Ethnography is well-suited to resituate and rethink how practitioners conceptualize the health behaviors of persons because of its focus on capturing the social meanings and the ordinary activities of persons in naturally occurring settings (Biehl, 2007). Through the use of participant observation, field notes, and in-depth interviews to capture “everyday life,” ethnographic research reflects the knowledge and the system of meanings in the lives of a particular group. Because of this, ethnography is uniquely qualified to confront and humanize the ways public health problems are framed and interventions carried out (Biehl, 2007).

Although these approaches are not new, they have failed to become mainstream in public health research and are still viewed as ancillary. These approaches have unique potential to bring the experiences of patients into conversations about clinical practice and prevention. Therefore, these and other similar approaches need to become far more normative in public health and epidemiological research.

“Rational” decision-making

Much of the recent crossover from epidemiological to sociological approaches to public health have focused on creating models that attempt to take into account variables that are beyond the traditional scope of epidemiology. Some of these efforts have focused on neighborhood disorder (Browning & Cagney, 2003; Ross & Mirowsky, 2001), cultural traits (Abraido-Lanza, Chao, & Flórez, 2005; Lee, Sobal, & Frongillo, 2000), inequality (Subramanian & Kawachi, 2003), and the resulting health outcomes. Much of this information, however, is attained through structured questionnaires or analysis of medical records. In general, such studies have the goal of developing path models based on rational choice decisions (Pescosolido, 1992).

Rational choice theory has been adopted by many disciplines (Hechter & Kanazawa, 1997; Li, Zhang, & Sarathy, 2010), and has become widely used to explain a range of behaviors. Rational choice theory focuses on the cost–benefit analyses that actors purportedly make when deciding to take certain actions (Boudon, 1998). Most important at this juncture is that rational choice theory underlies many of the behaviorally based models applied to public health problems, and assumes that people make decisions designed to maximize health outcomes. Recognizing this epistemological influence is incredibly important, especially because research shows that the human mind does not necessarily work “rationally” (Boudon,

1998). That is, persons seem to make relevant decisions that vary along a continuum, instead of pursuing an idealized logic.

The public health literature is replete with models that attempt to describe the decision-making of persons with respect to their health. These models are almost all within the rational choice paradigm and attempt to create formulae designed to predict expected behaviors. But sometimes a person simply does not have the means to act, let alone the luxury to make an informed decision. Examples of commonly used health behavior models in this tradition include the theory of reasoned action–planned behavior model (Montaña & Kasprzyk, 2008; Redding, Rossi, Rossi, Velicer, & Prochaska, 2000), the health belief model (Rosenstock, Strecher, & Becker, 1994; Strecher & Rosenstock, 1997), social cognitive theory (Bandura, 1986, 1990), the transtheoretical model (Prochaska & Velicer, 1997; Redding et al., 2000), and the health behavior model (Andersen, 1995).

These models range from simplistic to complex and include varying levels of contextual variables. The problem is that they all rely on the assumption that human thought and behavior follow a set of predictable rules. What actual community members believe and the value systems that pervade a community is often not the focus of attention. Simply put, the community’s world is thus overlooked (Pollner, 2010). Because of this, these types of models, which attempt to map human cognition and decision-making, portray inaccurate representations of how people think and act.

Traditional models in public health approach human decision-making as if people are constantly evaluating objective indices pertaining to health. As researchers work to refine these models—in order to become more holistic—they often seek to identify “missing” variables to understand how to better predict human decisions. But these models assume that the parameters of choosing health are clear, like the rules of a game (Murphy, 1992), and that identical factors influence, although perhaps with different weighting depending on sociodemographic characteristics, the computations of people when they make decisions about their health or seeking treatment.

Consistent with this view of decision-making, behavioral analyses have tended to be equated with risk factor research in public health. Specifically, researchers operationalize variables in order to isolate certain aspects of behavior or the environment (Riffenburgh, 2006). But this approach to modeling health overlooks how persons actively construct the reality in which they live. Comparing studies of condom use based on a rational choice health model

versus a more holistic method provides a useful demonstration of the reductionism in these studies.

As HIV becomes an increasingly feminized epidemic (Wingood, 2003), public health researchers try to understand how men and women make safer sex decisions to use protections such as condoms with sexual partners. Attempts to study this issue have traditionally focused on knowledge about disease transmission and condom use. Medical anthropologists, however, have delved into this topic and found that some of the underlying social determinants are a woman's power to insist on condom use (Dunkle et al., 2004; Pulerwitz, Amaro, DeJong, Gortmaker, & Rudd, 2002); how trust, fidelity, and self-esteem are constructed in relationships (Sobo, 1995; Sterk, Klein, & Elifson, 2004; Syvertsen et al., 2013); whether women who are at high risk have alternative economic options aside from receiving financial assistance from their primary partners (Parker, Easton, & Klein, 2000); and the social meaning for women of being perceived as having a faithful partner (Cook, 2013; Sobó, 1995). These types of in-depth studies that look at the social meaning of condom use can more accurately assess the situation from the perspective of the actors, and therefore lead to more socially sensitive and, thus, efficacious interventions.

Conclusion

As a discipline, public health is struggling to advance the third wave that moves beyond individual level behaviors to critically understand health in the community context. This transition includes more holistic definitions of health and broadens what is thought to fall under the purview of health, including a critique of the dualistic assumptions that underpin public health and health care. Understanding the assumptions of traditional public health is important to appreciate their impact on health-related decision-making.

The connection between theory and public health is important, particularly in view of recent theoretical shifts that define knowledge and alter how decision-making is evaluated (Ferguson, 2006). This manuscript provides a critique of traditional public health approaches and theoretically justifies what and why changes are warranted. The overarching point is that the process of overanalyzing variables is impeding the process of gaining insight into the experiential-based pathways persons follow when defining health and seeking treatment. This claim is especially important to behavior change and health decision-making, because intuitive and subtle interpretations influence the decisions that people make (Krieger, 2005).

Strategies that are grounded in the traditions of, for example, phenomenology or participatory action research, can be used to understand not just knowledge about disease transmission, but how people view their options, barriers, and the likely success of a particular intervention.

Conflict of interest and funding

The authors have not received any funding or benefits from industry or elsewhere to conduct this study.

References

- Abraido-Lanza, A., Chao, M., & Flórez, K. (2005). Do healthy behaviors decline with greater acculturation?: Implications for the Latino mortality paradox. *Social Science and Medicine*, 61(6), 1243–1255.
- Agar, M. (2003). Toward a qualitative epidemiology. *Qualitative Health Research*, 13(7), 974–986.
- Agnew, R. (2006). Storylines as a neglected cause of crime. *Journal of Research in Crime and Delinquency*, 43(2), 119–131.
- Andersen, R. (1995). Revisiting the behavior model and access to medical care: Does it matter? *Journal of Health and Social Behavior*, 36, 1–10.
- Baker, T. A., & Wang, C. C. (2006). Photovoice: Use of a participatory action research method to explore the chronic pain experience in older adults. *Qualitative Health Research*, 16(10), 1405–1413.
- Bandura, A. (1986). *Social foundations of thought and action: A social cognitive theory*. Englewood Cliffs, NJ: Prentice Hall.
- Bandura, A. (1990). Perceived self-efficacy in the exercise of control over AIDS infection. *Evaluation and Program Planning*, 13, 9–17.
- Berger, P., & Luckmann, T. (1966). *The social construction of reality: A treatise in the sociology of knowledge*. Garden City, NY: Anchor Books.
- Biehl, J. G. (2007). *Will to live: AIDS therapies and the politics of survival*. Princeton, New Jersey: Princeton University Press.
- Booth, T., & Booth, W. (2003). In the frame: Photovoice and mothers with learning difficulties. *Disability & Society*, 18(4), 431–442.
- Boudon, R. (1998). Limitations of rational choice theory. *American Journal of Sociology*, 104(3), 817–828.
- Browning, C., & Cagney, K. (2003). Moving beyond poverty: Neighborhood structure, social processes, and health. *Journal of Health and Social Behavior*, 44, 552–571.
- Cassell, E. J. (1991). *The nature of suffering and the goals of medicine*. New York: Oxford University Press.
- Charmaz, K. (1999). Stories of suffering: Subjective tales and research narratives. *Qualitative Health Research*, 9(3), 362–382.
- Charon, R. (2001). Narrative medicine: A model for empathy, reflection, profession, and trust. *JAMA*, 286(15), 1897–1902.
- Charon, R. (2011). The novelization of the body, or, how medicine and stories need one another. *Narrative*, 19(1), 33–50.
- Cook, H. (2013). Buscando el Moro: A constructivist approach to health decision-making in a Dominican Batey. *Open access dissertations*. Paper 1048. http://scholarlyrepository.miami.edu/oa_dissertations.

- Descartes, R. (1985). *The philosophical writings of Descartes* (Vol. 2). Cambridge, England: Cambridge University Press.
- Dunkle, K., Jewkes, R., Brown, H., Gray, G., McIntyre, J., & Harlow, S. (2004). Gender-based violence, relationship power, and risk of HIV infection in women attending antenatal clinics in South Africa. *The Lancet*, *363*(9419), 1415–1421.
- Engel, G. (1977). The need for a new medical model: A challenge for biomedicine. *Science*, *196*(4286), 129–136.
- Ezzy, D. (2000). Illness narratives: Time, hope and HIV. *Social Science & Medicine*, *50*(5), 605–617.
- Ferguson, H. (2006). *Phenomenological society: Experience and insight in modern society*. Thousand Oaks, CA: Sage.
- Frank, A. W. (1998). Just listening: Narrative and deep illness. Families, Systems. *Health*, *16*(3), 197.
- Frenk, J. (1993). The new public health. *Annual Review of Public Health*, *14*, 469–490.
- Gradmann, C. (2009). *Laboratory disease: Robert Koch's medical bacteriology*. Baltimore, MD: The Johns Hopkins University Press.
- Greenhalgh, T., & Hurwitz, B. (1999). Why study narrative? *BMJ*, *318*(7175), 48–50.
- Hills, M. (2000). Human science research in public health: The contribution and assessment of a qualitative approach. *Canadian Journal of Public Health*, *91*(6), I4–I7.
- Hechter, M., & Kanazawa, S. (1997). Sociological rational choice theory. *Annual Review of Sociology*, *23*, 191–214.
- Hesse-Biber, S. (2012). Weaving a multimethodology and mixed methods praxis into randomized control trials to enhance credibility. *Qualitative Inquiry*, *18*(10), 876–889.
- Husserl, E. (2012). *Ideas: General introduction to pure phenomenology*. New York: Routledge.
- Kickbusch, I. (2003). The contribution of the World Health Organization to a new public health and health promotion. *American Journal of Public Health*, *93*(3), 383–388.
- Kleinman, A., & Kleinman, J. (1996). The appeal of experience; the dismay of images: Cultural appropriations of suffering in our times. *Daedalus*, *125*, 1–23.
- Krieger, N. (2005). Embodiment: A conceptual glossary for epidemiology. *Journal of Epidemiology and Community Health*, *59*(5), 350–355.
- Krieger, N. (2012). Methods for the scientific study of discrimination and health: An ecosocial approach. *American Journal of Public Health*, *102*(5), 936–944.
- Lalonde, M. (1981). *A new perspective on the health of Canadians. Minister of Supply and Services Canada*. Retrieved September 15, 2012, from http://www.hc-sc.gc.ca/hcs-sss/alt_formats/hpb-dgps/pdf/pubs/1974-lalonde/lalonde-eng.pdf
- Lee, S. K., Sobal, J., & Frongillo, E. (2000). Acculturation and health in Korean Americans. *Social Science and Medicine*, *51*(2), 159–173.
- Li, H., Zhang, J., & Sarathy, R. (2010). Understanding compliance with internet use policy from the perspective of rational choice theory. *Decision Support Systems*, *48*(4), 635–645.
- Lock, A., & Strong, T. (2010). *Social constructionism: Sources and stirrings in theory and practice*. Cambridge, UK: Cambridge University Press.
- Lopez, E., Eng, E., Robinson, N., & Wang, C. (2005). Photovoice as a community-based participatory research method. In B. Israel, E. Eng, A. Schulz, & E. Parker (Eds.), *Methods in community-based participatory research for health* (pp. 326–348). San Francisco, CA: Jossey-Bass.
- Morabia, A. (2007). Epidemiological methods and concepts in the nineteenth century and their influences on the twentieth century. In W. Holland, J. Olsen, & C. V. Florey (Eds.), *The development of modern epidemiology: Personal reports from those who were there* (pp. 17–29). New York: Oxford University Press.
- Montaño, D., & Kasprzyk, D. (2008). Theory of reasoned action, theory of planned behavior, and the integrated behavioral model. In K. Glanz, B. Rimer, & K. Viswanath (Eds.), *Health behavior and health education: Theory, research, and practice (chapter 4)* (pp. 67–95). San Francisco, CA: John Wiley & Sons.
- Merleau-Ponty, M. (1996). *Phenomenology of perception*. New York: Motilal Banarsidass.
- Murphy, J. (1992). Reason, bounded rationality and the lebenswelt: Socially sensitive decision making. *American Journal of Economics and Sociology*, *51*(3), 293–304.
- Murphy, J., & Min Choi, J. (1997). *Postmodernism, unraveling racism, and democratic institutions*. Westport, CT: Praeger.
- Murphy, J. W., & Rigg, K. K. (2014). Clarifying the philosophy behind the Community Mental Health Act and community-based interventions. *Journal of Community Psychology*, *42*(3), 285–298.
- Newman, S. D. (2010). Evidence-based advocacy: Using photovoice to identify barriers and facilitators to community participation after spinal cord injury. *Rehabilitation Nursing*, *35*(2), 47–59.
- Page, R., Cole, G., & Timmreck, T. (1995). *Basic epidemiological methods and biostatistics, a practical guidebook*. Sudbury, MA: Jones and Bartlett.
- Parker, R., Easton, D., & Klein, C. (2000). Structural barriers and facilitators in HIV prevention: A review of international research. *AIDS*, *14*(Special Issue), 22–32.
- Pescosolido, B. (1992). Beyond rational choice: The social dynamics of how people seek help. *American Journal of Sociology*, *97*(4), 1096–1138.
- Pollner, M. (2010). *Mundane reason: Reality in everyday and sociological discourse*. Cambridge, England: Cambridge University Press.
- Prochaska, J., & Velicer, W. (1997). The transtheoretical model of health behavior change. *American Journal of Health Promotion*, *12*(1), 38–48.
- Pulerwitz, J., Amaro, H., DeJong, W., Gortmaker, S., & Rudd, R. (2002). Relationship power, condom use, and HIV risk among women in the USA. *AIDS Care*, *14*(6), 789–800.
- Quimby, E. (2006). Ethnography's role in assisting mental health research and clinical practice. *Journal of Clinical Psychology*, *62*(7), 859–879.
- Raphael, D. (2003). Barriers to addressing the societal determinants of health: Public health units and poverty in Ontario Canada. *Health Promotion Management*, *18*(4), 397–405.
- Reiser, S. J. (1981). *Medicine and the reign of technology*. Cambridge, England: Cambridge University Press.
- Redding, C., Rossi, J., Rossi, S., Velicer, W., & Prochaska, J. (2000). Health behavior models. *The International Electronic Journal of Health Education*, *3*(Special Issue), 180–193.
- Remen, R. N. (2002). *Kitchen table wisdom: Stories that heal*. New York: Riverhead Books, Pan Australia.
- Rhodes, S. D., & Hergenrather, K. C. (2007). Recently arrived immigrant Latino men identify community approaches to promote HIV prevention. *American Journal of Public Health*, *97*(6), 984–985.
- Riffenburgh, R. (2006). *Statistics in medicine* (2nd ed.). San Diego, CA: Elsevier Academic Press.
- Rigg, K., & Murphy, J. (2013a). Storylines as a neglected tool for mental health service providers and researchers. *International Journal of Mental Health & Addiction*, *11*(4), 431–440.
- Rigg, K., & Murphy, J. (2013b). Understanding the etiology of prescription opioid abuse: Implications for prevention and treatment. *Qualitative Health Research*, *23*(7), 963–975.

- Rosenstock, I., Strecher, V., & Becker, M. (1994). The health belief model and HIV risk behavior change. In R. DiClemente & J. Peterson (Eds.), *Preventing AIDS: Theories and methods of behavioral interventions (Chapter 2)* (pp. 5–24). New York, NY: Plenum Press.
- Ross, C., & Mirowsky, J. (2001). Neighborhood disadvantage, disorder, and health. *Journal of Health and Social Behavior*, *42*, 258–276.
- Saracci, R. (2007). Epidemiological concepts Pre-1950 and their relation to work in the second half of the century. In W. Holland, J. Olsen, & C. de V. Florey (Eds.), *The development of modern epidemiology: Personal reports from those who were there* (pp. 32–38). New York: Oxford University Press.
- Schutz, A. (1953). Common-sense and scientific interpretation of human action. *International Phenomenological Society*, *14*(1), 1–38.
- Schutz, A. (1967). *The Phenomenology of the social world*. Evanston, IL: Northwestern University Press.
- Smith, M. (1998). Community-based epidemiology: Community involvement in defining social risk. In J. Pardeck, C. Longino, & J. Murphy (Eds.), *Reason and rationality in health and human services delivery* (pp. 51–65). New York: Haworth Press.
- Sobo, E. (1995). Finance, romance, social support, and condom use among impoverished inner-city women. *Human Organization*, *54*(2), 115–128.
- Sterk, C., Klein, H., & Elifson, K. (2004). Predictors of condom-related attitudes among at-risk women. *Journal of Women's Health*, *13*(6), 676–688.
- Strecher, V., & Rosenstock, I. (1997). The health belief model. In A. Baum, S. Newman, J. Weinman, R. West, & C. McManus (Eds.), *Cambridge handbook of psychology, health and medicine* (pp. 113–117). Cambridge, UK: Cambridge University Press.
- Subramanian, S., & Kawachi, I. (2003). Income inequality and health: What have we learned so far? *Epidemiologic Reviews*, *26*, 78–91.
- Susser, E. (2004). Eco-epidemiology: Thinking outside the black box. *Epidemiology*, *15*(5), 519–520.
- Susser, M., & Susser, E. (1996). Choosing a future for epidemiology: I. Eras and paradigms. *American Journal of Public Health*, *86*(5), 668–673.
- Syvertsen, J. L., Robertson, A. M., Palinkas, L. A., Rangel, M. G., Martinez, G., & Strathdee, S. A. (2013). 'Where sex ends and emotions begin': Love and HIV risk among female sex workers and their intimate, non-commercial partners along the Mexico-US border. *Culture, Health & Sexuality*, *15*(5), 540–554.
- True, J. G. (2014, March). *From Research to Advocacy (and Back Again): Experiences from a Photovoice Project with Iraq and Afghanistan War Veterans*. Paper presented at the 74th Annual Meeting of the Society for Applied Anthropology, Albuquerque, NM.
- Tulchinsky, T., & Varivakova, E. (2009). *The new public health* (2nd ed.). San Diego, CA: Elsevier Academic Press.
- Wallerstein, N., & Duran, B. (2010). Community-based participatory research contributions to intervention research: The intersection of science and practice to improve health equity. *American Journal of Public Health*, *100*, S40–S46.
- Wang, C. C., & Burris, M. (1997). Photovoice: Concept, methodology, and use for participatory needs assessment. *Health Education & Behavior*, *24*, 369–387.
- Wang, C. C., Cash, J. L., & Powers, L. S. (2000). Who knows the streets as well as the homeless? Promoting personal and community action through photovoice. *Health Promotion Practice*, *1*, 81–89.
- Williams, G. (1984). The genesis of chronic illness: Narrative re-construction. *Sociology of Health & Illness*, *6*(2), 175–200.
- Wingood, G. M. (2003). Feminization of the HIV epidemic in the United States: Major research findings and future research needs. *Journal of Urban Health*, *80*(3), iii67–iii76.
- Winkelman, M. (2009). *Culture and health: Applying medical anthropology*. San Francisco, CA: John Wiley and Sons.
- Zaner, R. M. (1988). *Ethics and the clinical encounter*. Englewood Cliffs, NJ: Prentice Hall.