

***Fbxw7* Deletion Accelerates *Kras*^{G12D}-Driven Pancreatic Tumorigenesis via Yap Accumulation^{1,2,3}**



Qiang Zhang^{*}, Yaqing Zhang[†], Joshua D. Parsels^{*}, Ines Lohse^{*}, Theodore S. Lawrence^{*}, Marina Pasca di Magliano[†], Yi Sun^{*,‡,§} and Meredith A. Morgan^{*}

^{*}Department of Radiation Oncology, University of Michigan Medical School, Ann Arbor, MI 48109, USA; [†]Department of Surgery, University of Michigan Medical School, Ann Arbor, MI 48109, USA; [‡]Institute of Translational Medicine, Zhejiang University School of Medicine, Hangzhou, Zhejiang, PR China; [§]Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, Zhejiang University, Hangzhou, China

Abstract

Pancreatic cancers driven by *KRAS* mutations require additional mutations for tumor progression. The tumor suppressor FBXW7 is altered in pancreatic cancers, but its contribution to pancreatic tumorigenesis is unknown. To determine potential cooperation between *Kras* mutation and *Fbxw7* inactivation in pancreatic tumorigenesis, we generated P48-Cre;LSL-*Kras*^{G12D};*Fbxw7*^{fl/fl} (KFC^{fl/fl}) compound mice. We found that KFC^{fl/fl} mice displayed accelerated tumorigenesis: all mice succumbed to pancreatic ductal adenocarcinoma (PDA) by 40 days of age, with PDA onset occurring by 2 weeks of age. PDA in KFC^{fl/fl} mice was preceded by earlier onset of acinar-to-ductal metaplasia (ADM) and pancreatic intraepithelial neoplasia (PanIN) lesions, and associated with chromosomal instability and the accumulation of Fbxw7 substrates Yes-associated protein (Yap), c-Myc, and Notch. Using KFC^{fl/fl} and FBXW7-deficient human pancreatic cancer cells, we found that Yap silencing attenuated growth promotion by Fbxw7 deletion. Our data demonstrate that Fbxw7 is a potent suppressor of *Kras*^{G12D}-induced pancreatic tumorigenesis due, at least in part, to negative regulation of Yap.

Neoplasia (2016) 18, 666–673

Introduction

Pancreatic cancer has an overall 5-year survival rate of 8% and is currently the fourth leading cause of cancer-related death in the United States. Nearly all human pancreatic cancers are characterized by mutations in *KRAS* followed in order of mutational frequency by *P16*, *P53*, and *DPC4* [1]. Genetically engineered *Kras* mutant mouse models of pancreatic cancer recapitulate the human disease process beginning with acinar-to-ductal metaplasia (ADM), followed by pancreatic intraepithelial neoplasia (PanIN) formation, and ultimately pancreatic ductal adenocarcinoma (PDA) [2]. The introduction of other mutations such as *p53*, *p16*, or *Dpc4* further accelerates the development of PDA. Using an inducible *Kras* mutant (G12D) mouse model, it was shown that *Kras* mutation is involved not only in initiation but also in the maintenance of PDA, as inactivation of mutant *Kras* in established tumors leads to tumor regression [3]. Furthermore, tumor regrowth following inactivation of mutant *Kras* involves expression of Yap which is also required for initiation of PDA in *Kras* mutant mice [4,5].

FBXW7, a component of the Skp1-Cullin1-Fbox E3 ubiquitin ligase complex, is a characterized tumor suppressor. For instance, conditional *Fbxw7* deletion cooperates with *APC* mutation or *P53*

Address all correspondence to: Meredith A. Morgan, Department of Radiation Oncology, University of Michigan Medical School, Room 4326B Medical Sciences I, Ann Arbor, MI, 48109-5637.

E-mail: mmccrack@med.umich.edu

¹One-sentence summary: This study for the first time identifies FBXW7 as a suppressor of mutant *Kras*-driven tumorigenesis.

²Grant support: This work was funded by National Institutes of Health grants R01CA163895 (M.A.M.), P50CA130810 (T.S.L.), R01CA118762 and R01CA171277 (Y.S.), and R01CA151588 (M.P.d.M.) and the American Cancer Society (M.P.d.M.).

³Disclosure of potential conflicts of interest: none.

Received 18 August 2016; Accepted 24 August 2016

© 2016 The Authors. Published by Elsevier Inc. on behalf of Neoplasia Press, Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1476-5586

<http://dx.doi.org/10.1016/j.neo.2016.08.009>

deletion to accelerate intestinal tumorigenesis in a haploinsufficient manner [6,7]. Although *FBXW7* is most frequently mutated in colorectal, uterine, and bladder cancers, it is also a significantly mutated gene in pancreatic cancers [8,9]. Furthermore, in a Sleeping Beauty transposon insertional mutagenesis model of *Kras*^{G12D}-induced pancreatic tumorigenesis, *Fbxw7* mutation cooperated with *Kras*^{G12D} to accelerate PDA formation with a high frequency (24%), suggesting that *Fbxw7* may be an important tumor suppressor in *Kras*-driven pancreatic cancers [10]. In addition to mutation, there is also emerging evidence to suggest that *FBXW7* is altered by other mechanisms in pancreatic cancers including by reduced gene expression and heightened protein degradation which correlate with reduced survival [11].

Although the tumor suppressor functions of *FBXW7* have largely been attributed to its proteolytic regulation of oncogenic substrates such as Cyclin E, c-MYC, Notch, and c-JUN, *FBXW7* regulates over 30 substrates and likely unknown substrates as well [12,13]. Loss of *FBXW7* promotes genomic instability in part by increased Cyclin E expression [14] which may contribute to tumorigenesis. We have recently reported that *FBXW7* regulates nonhomologous end-joining (NHEJ) via K63-linked polyubiquitination of XRCC4, promoting the interaction of XRCC4 with DNA double-strand breaks and other core NHEJ proteins [15]. This direct involvement of *FBXW7* in NHEJ may also contribute to the genomic instability and tumorigenesis in *FBXW7*-deficient models.

Because pancreatic cancer is characterized most frequently by *KRAS* mutation (~100%) [1] as well as mutations or reduced expression of *FBXW7* in a subset of human pancreatic cancers [8,9,11], in this study, we investigated whether *FBXW7* acts as a tumor suppressor in *Kras*^{G12D}-driven pancreatic tumorigenesis. Using a pancreas conditional P48-Cre;LSL-*Kras*^{G12D}; *Fbxw7*^{fl/fl} mouse model, we found that homozygous deletion of *Fbxw7* causes a dramatic acceleration of *Kras*^{G12D}-driven pancreatic tumorigenesis. Based on this finding, we went on to investigate the contribution of *Fbxw7* substrates, such as Yap, to the mechanisms of tumorigenesis in both mouse and human pancreatic cancer cells.

Materials and Methods

Mouse Strains

The conditional LSL-*Fbxw7*^{fllox/fllox} mouse was a gift from Dr. Iannis Aifantis (New York University) [16]. Conditional LSL-*Fbxw7*^{fllox/fllox}; LSL-*Kras*^{G12D/+}; and P48-Cre strains were interbred to obtain LSL-*Kras*^{G12D/+}; LSL-*Fbxw7*^{fllox/fllox}; P48-Cre triple mutant animals (KFC). The mice were housed in specific pathogen-free facilities, and all the studies were conducted in compliance with the Institutional Committee on Use and Care of Animals guidelines. Genotyping was carried out by tail clipping from mice 2 weeks after birth. Tail specimens were incubated for 24 hours at 55°C in tail-lysis buffer containing proteinase K. NaCl was added, and cellular debris were pelleted by centrifugation. DNA was precipitated by the addition of isopropanol and washed with 70% ethanol. DNA pellets dissolved in water were used for polymerase chain reaction (PCR) analysis.

Cell Lines

Panc-1 cells were obtained from the American Type Culture Collection. Cells were cultured in Dulbecco's modified Eagle medium supplemented with 10% FBS, 2 mM glutamine, and antibiotics. The KFC cell line was derived from the tumor of a 3-week-old KFC mouse. Briefly, mouse pancreas was harvested and

finely minced and incubated with a Collagenase V/HBSS (with Ca and Mg) solution at 37°C under constant shaking for 1.5 hours. Collagenase digestion of pancreas tissue was neutralized by RPMI-1640 medium containing 10% FBS. Digested tumor cells were filtered through a 40- μ m filter and then cultured in RPMI-1640 medium containing 10% FBS, 2 mM glutamine, and antibiotics.

Acinar Cell Culture

The 3D culture of pancreatic acinar cells was prepared by digesting pancreata from 1-week-old mice with Collagenase P followed by culture in Matrigel as previously described [17]. Briefly, pancreata from KFC, KC, and control mice were cut into small pieces and digested with 2 mg/ml of Collagenase P (Roche Diagnostics) in HBSS for 15 minutes at 37°C. Cells were then washed three times with HBSS with 5% FBS and filtered through 100- μ m nylon meshes. After centrifugation, the cell suspension was mixed 1:1 with Matrigel and plated onto the collagen layer. The acinar cell/Matrigel mix was allowed to solidify for 1 hour at 37°C before adding medium. The formation of ductlike structures was observed at days 1, 2, and 3.

Histopathologic Analysis

Histopathologic analysis was conducted by a pathologist (Y.Z.) on all identified hematoxylin and eosin (H&E)-stained slides. Pancreata sections were evaluated for ADM, PanIN1, PanIN2, PanIN3, and PDA lesions based on a previously reported classification system [2]. Pancreata were diagnosed according to the most severe phenotype observed, and data were expressed as the percentage of animals with each phenotype.

Immunohistochemistry and Immunoblotting

Immunohistochemistry was conducted as previously described [3]. Images were acquired with an Olympus BX-51 microscope, Olympus DP71 digital camera, and DP Controller software. Antibodies used for immunohistochemistry include those recognizing YAP (Cell Signaling, #4912, 1:500), c-MYC (Abcam, ab32072, 1:200), Notch (Cell Signaling, #3608, 1:250), Ki67 (BD Bioscience, Cat. 550609, 1:500), CK19 (Abcam, ab87000, 1:500), and Amylase (Sigma-Aldrich, A8273, 1:500).

Immunoblotting was conducted as previously described [15]. *FBXW7* was detected by *FBXW7* immunoprecipitation followed by immunoblotting. Antibodies used for immunoblotting in this study include those recognizing *FBXW7* (Bethyl, A301-720A, 1:1000, overnight, 4°C), YAP (Cell Signaling, #4912, 1:3000, overnight, 4°C), c-MYC (Abcam, ab32072, 1:2000, overnight, 4°C), Notch (Cell Signaling, #3608, 1:2000, overnight, 4°C), c-JUN (Cell Signaling, #9165, 1:1000, overnight, 4°C), MCL-1 (Santa Cruz, sc-819, 1:1000, overnight, 4°C), Cyclin E (Santa Cruz, sc-198, 1:1000, overnight, 4°C), mTOR (Cell Signaling, #2972, 1:1000, overnight, 4°C), and actin (Sigma Aldrich, Clone AC-40, 1:5000, overnight, 4°C).

Lentivirus-Based shRNA Infection and siRNA Transfection

Cells were infected with lentivirus (control 5'-GCAAGCT GACCCTGAAGTTCAT-3'; *FBXW7*: 5'-ACAGGACAGTGTTTA CAAA-3') in the presence of 8 μ g/ml of polybrene. Silencing efficiency was detected 72 hours after infection by immunoblot. For Yap RNAi experiments, cells were transfected with Oligofectamine (Invitrogen) transfection reagent. The two independent siYap oligonucleotide sequences used were 5'-CCACCAAGCUAGAUAAAGA-3' (siYap-2) and 5'-GCACCUAUCACUCUCGAGA-3' (siYap-4). Nonspecific siRNA was used as a control.

Colony Formation Assay

Cells were infected with shRNA (FBXW7 or control) and then transfected with Yap siRNA. After 3 days, cells were seeded in triplicate in 60-mm dishes and allowed to form colonies for 9 to 14 days. KFC pancreatic tumor cells were transfected with two independent siYap oligonucleotides. Two days posttransfection, KFC cells were seeded to form colonies for 14 days. Colonies were fixed with 100% methanol, stained with methylene blue in methanol, and then scored.

Statistical Analysis

Survival curves were calculated according to the Kaplan-Meier method, and statistical differences were analyzed by the log-rank and Gehan-Breslow-Wilcoxon tests using GraphPad Prism. A two-sided, unpaired Student's *t* test was used for other statistical analyses. *P* values of < .05 were considered statistically significant.

Results

Acceleration of *Kras*^{G12D}-Pancreatic Tumorigenesis by *Fbxw7* Deletion

To investigate the involvement of FBXW7 in mutant *KRAS*-driven pancreatic tumorigenesis, we crossed P48-Cre;LSL-*Kras*^{G12D} mice [2] with *Fbxw7*^{fl/fl} mice [16] to generate mice with pancreas specific activation of *Kras*^{G12D} and heterozygous or homozygous deletion of *Fbxw7* (designated KFC^{fl/+} or KFC^{fl/fl}, respectively; Figure 1A). We first confirmed deletion of *Fbxw7* by examining *Fbxw7* protein levels in pancreata from 1-week-old mice and found *Fbxw7* protein to be substantially reduced in pancreata from KFC^{fl/fl} mice compared with those from KC control mice or KFC^{fl/+} mice (Figure 1B). To begin to establish a phenotype for these mice, survival was monitored in KC, KFC^{fl/+}, KFC^{fl/fl} as well as in FC^{fl/fl} mice. We observed a profound effect of homozygous *Fbxw7* deletion on survival of KFC^{fl/fl} mice (Figure 1C) with a median survival time of 28.5 days. Most KFC^{fl/fl} mice displayed early symptoms of morbidity and were euthanized; 100% of KFC^{fl/fl} mice succumbed to PDA by 40 days compared with only 7% of KFC^{fl/+} and none of the KC or FC mice during the observation period. Gross examination of KFC^{fl/fl} mice revealed smaller body size, enlarged pancreata, ascites, and duodenum adhesions in most mice with pancreas main duct obstruction and liver metastases occurring in a subset of mice (Supplementary Figure 1, A–E). Histological examination of pancreas sections from 1-month-old KFC^{fl/fl} mice revealed moderately to poorly differentiated PDA accompanied by PanIN3 lesions (Figure 1D). In contrast, pancreata from KC or FC^{fl/fl} mice appeared largely normal, although early-stage PanIN lesions were occasionally observed in KFC^{fl/+} pancreata (Supplementary Figure 1F). To further explore the effects of heterozygous *Fbxw7* deletion, we also examined pancreata from 6-month-old mice. Whereas KC mice showed some PanIN lesions, KFC^{fl/+} contained more PanIN lesions, ranging from PanIN1A through PanIN3 in both genotypes, but no PDA (Figure 1D). Pancreata from FC^{fl/fl} mice were overall normal with the exception of occasional enlarged ductal cells, consistent with a prior study [18] (Supplementary Figure 1G).

Tumorigenesis in *Kras*^{G12D}; *Fbxw7*-Deleted Pancreata Is Preceded by ADM and PanIN and Associated with Chromosomal Instability

We then sought to characterize the progression of precursor lesions leading to PDA in KFC^{fl/fl} mice. To this end, pancreata from 1-, 2-, and 3-week-old KC, KFC^{fl/+}, and KFC^{fl/fl} mice were examined

histologically and scored for the presence of ADM, PanINs, and PDA. In KC and KFC^{fl/+} mice at 1 to 2 weeks of age, pancreata were composed mainly of acinar cells with no observable ADM or PanIN lesions (Figure 2, A and B). By 3 weeks of age, pancreata from KC and KFC^{fl/+} mice began to develop ADM with progression toward PanIN2 in the KFC^{fl/+} mice. In striking contrast, KFC^{fl/fl} pancreata contained PanIN1 and PanIN2 lesions as well as stroma accumulation at 1 week, which advanced to a combination of PanIN3 and PDA at 2 weeks. By 3 weeks of age, 100% of KFC^{fl/fl} pancreata contained PDA. To confirm the acceleration of ADM in KFC^{fl/fl} pancreata, we used 3D cultures of acinar clusters and observed the formation of ductlike structures by as early as 1 day in KFC^{fl/fl} acini (Figure 2, C and D). Further characterization of KFC^{fl/fl} pancreata showed increased CK19, decreased amylase, and increased Alcian Blue staining, confirming ADM and progression toward PanIN lesions, respectively (Supplementary Figure 2).

Genomic deletion or mutation of *FBXW7* has been associated with genomic instability in various cancers [6,14,19,20]. Given the high degree of chromosomal instability (CIN) in human PDA, we examined KFC^{fl/fl} pancreatic tumors for signs of CIN and found abnormal mitotic figures (Supplementary Figure 3A). Furthermore, using a cell line derived from KFC^{fl/fl} pancreatic tumors (Supplementary Figure 3B), we found increased numbers of micronuclei, which are commonly observed in cells with CIN (Supplementary Figure 3C). Consistent with this finding, flow cytometry also revealed that the KFC tumor cell line, as well as the established KPC cell line (with *p53* deletion), is aneuploid (Supplementary Figure 3D). These data suggest that CIN is associated with the accelerated pancreatic tumorigenesis observed in KFC^{fl/fl} mice.

Accumulation of *Fbxw7* Substrates in *Fbxw7*-Deleted Pancreata

To begin to determine the mechanisms leading to accelerated pancreatic tumorigenesis in the KFC^{fl/fl} mice, pancreata from 1-week-old mice were analyzed for expression of oncogenic substrates of *Fbxw7* [12,13]. We hypothesized that *Fbxw7* substrates involved in *Kras*^{G12D}-driven tumorigenesis would be accumulated in response to homozygous *Fbxw7* deletion. Of the substrates investigated, we found that the levels of c-Myc, Yap, and Notch, but not c-Jun, Cyclin E, and Mcl-1, were elevated in KFC^{fl/fl} pancreata as well as in the KFC^{fl/+} pancreata, although to a lesser extent (Figure 3A). Evaluation of these proteins immunohistochemically confirmed their increased expression within aberrant ductal structures in KFC^{fl/fl} pancreata (Figure 3B). In addition, we also assessed the proliferative marker Ki67 and found significantly higher levels of Ki67 in KFC^{fl/fl} pancreata (Figure 3B, Supplementary Figure 4A). These changes in KFC^{fl/fl} pancreata appeared to be independent of *Kras* activity, as downstream effectors of *Kras* (pErk1/2 and pAkt) were not elevated in KFC^{fl/fl} pancreata (Supplementary Figure 4B). Taken together, these data demonstrate an association between increased c-Myc, Yap, and Notch expression and *Kras*^{G12D}-induced tumorigenesis in *Fbxw7* homozygous deleted mice.

Rescue of Growth Promotion Caused by *Fbxw7* Deletion by Depletion of Yap

Based on recent studies implicating Yap in *Kras*^{G12D}-induced pancreatic tumorigenesis [4,5], we hypothesized that *Fbxw7* may promote tumorigenesis through a Yap-dependent mechanism. To test this hypothesis, we first evaluated the ability of *Fbxw7* to regulate Yap protein levels in a cell line established from KFC^{fl/fl} pancreatic




Figure 1. Deletion of mouse *Fbxw7* accelerates *Kras*^{G12D}-driven pancreatic tumorigenesis. (A) Genetic makeup of the KFC model. (B) Western blot analysis of Fbxw7 protein levels in KC, KFC^{fl/+}, and KFC^{fl/fl} pancreata specimens from 1-week-old mice. (C) The survival of KC, FC, KFC^{fl/+}, and KFC^{fl/fl} mice is expressed using the Kaplan-Meier method, and statistically significant differences from KC mice are shown ($P < .0001$ ****). (D) H&E staining of pancreata at 1 and 6 months. Scale bar, 100 μ m. Each image is representative of at least three independent animals.

tumors. Consistent with the low levels of Yap observed in pancreata from KC mice (relative to KFC), exogenously expressed FBXW7 reduced Yap, c-Myc, and Notch protein levels in primary KFC-derived cells (Figure 4A) without substantial changes in other Fbxw7 substrates (Cyclin E, Mcl-1, mTOR; Supplementary Figure 5A). We then depleted Yap from KFC cells to determine its contribution to the increased growth of KFC cells and found that Yap depletion caused a significant decrease in the growth of KFC cells which was similar to the effect observed by exogenous FBXW7 expression (Figure 4B). Interestingly, Yap depletion was also associated with reduced c-Myc protein levels (Figure 4A). To further define the role of YAP in KFC^{fl/fl} tumorigenesis, we utilized human pancreatic cancer cells

that were deleted of FBXW7 by shRNA and then further silenced for YAP by siRNA. FBXW7 shRNA-treated Panc-1 and MiaPaCa-2 cells demonstrated increased levels of YAP and c-MYC levels without change in other FBXW7 substrates (Figure 4C; Supplementary Figure 5, B and C). Silencing of YAP by either of two independent siRNAs caused a reduction in c-MYC protein levels and, more importantly, significantly inhibited the growth of FBXW7-deficient human pancreatic cancer cells (Figure 4D, Supplementary Figure 5D). These results are consistent with the accelerated tumorigenesis and increased Yap levels observed in KFC^{fl/fl} pancreata. Taken together, these data demonstrate that Yap accumulation is causally related, at least in part, to the accelerated growth observed in Fbxw7-deficient, *Kras* mutant pancreatic cancer cells.




Figure 2. Pancreatic tumor progression in KFC^{fl/fl} mice. (A) H&E staining of KC, KFC^{fl/+}, and KFC^{fl/fl} pancreata at 1, 2, and 3 weeks. Scale bar, 100 μ m. (B) Pathologic analysis of the lesions of KC, KFC^{fl/+}, and KFC^{fl/fl} pancreata at 1, 2, and 3 weeks. Data represent at least three independent mice for each genotype and time point. (C) Deletion of *Fbxw7* accelerates ADM. Transmitted light images of control, KC, and KFC^{fl/fl} pancreatic cell clusters in 3D culture from day 0 to 3 are shown. Scale bar, 100 μ m. (D) Quantification of ductlike structures. Statistically significant differences were determined by a two-sided, unpaired Student's *t* test and are indicated ($P < .01^{**}$).

Discussion

In this study, we show for the first time that Fbxw7 is a suppressor of *Kras* mutation-driven pancreatic tumorigenesis. The dramatically accelerated tumorigenesis observed in the KFC^{fl/fl} mice was associated with accumulation of oncogenic Fbxw7 substrates such as c-Myc, Notch, and Yap. Importantly, we found that Yap expression was a driving mechanism of the accelerated tumorigenesis in response to

Fbxw7 deletion given that Yap depletion attenuated growth in both KFC primary cells as well as FBXW7-deficient human pancreatic cancer cells. These findings suggest that alterations in FBXW7 may be an important contributor to human pancreatic tumorigenesis. Furthermore, our data suggest that Fbxw7 is a relatively potent suppressor of pancreatic tumorigenesis given the development of tumors by less than 1 month in response to *Fbxw7* deletion relative to




Figure 3. Analysis of Fbxw7 substrates in pancreata. (A) Western blot analysis of the indicated proteins in lysates obtained from pancreata of 1-week-old mice for each of the indicated genotypes. (B) Immunohistochemical assessment of Yap, c-Myc, Notch, and Ki67 in KC, KFC^{fl/+}, and KFC^{fl/fl} pancreata at 1 week. Scale bar, 100 μ m.

latency periods of 2 to 9 months in other *Kras*^{G12D} models (e.g., *p16*^{Ink4a}, *P53*, and *Dpc4*) [21–23].

Although *FBXW7* is most frequently mutated in colorectal, uterine, and bladder cancers, it is also a significantly mutated gene in pancreatic cancers [8,9]. *FBXW7* mutations are generally heterozygous, missense, loss-of-function mutations occurring in the

WD domain that function in a dominant negative manner [13,24]. There are however emerging evidence also to suggest that *FBXW7* is altered by other mechanisms in pancreatic cancers including by reduced gene expression and heightened protein degradation [11]. It is important for future studies to define the spectrum of genetic and posttranslational alterations of *FBXW7* in both primary and




Figure 4. Depletion of Yap rescues growth in Fbxw7-deficient mouse and human pancreatic cancer cells. (A) Western blot analysis of c-Myc, Yap, and Notch proteins in KFC primary tumor cells with depletion of Yap. FLAG-FBXW7 overexpression in KFC tumor cells was used as a control. (B) The growth of KFC primary cells was measured with or without Yap depletion by colony forming assays. (C) Western blot detection of YAP, c-MYC, Notch, and FBXW7 levels in Panc-1 cells with or without FBXW7 and/or YAP depletion by colony formation assays. (D) The growth of Panc-1 cells with or without FBXW7 and/or YAP depletion was assessed by colony formation assays. Images are representative of at least two independent experiments (A, C). Data are expressed as the mean \pm SE of three independent experiments, and statistical significance is indicated ($P < .05^*$, $< 0.01^{**}$, or $< 0.001^{***}$) (B, D).

metastatic human pancreatic cancers. In addition, given the frequency of *KRAS* and *FBXW7* mutations in colorectal cancers, studies to investigate their effects on intestinal tumorigenesis are warranted.

Prior studies have shown that Fbxw7 is a haploinsufficient tumor suppressor [7,25]. In mice with *P53* deletion, heterozygous *Fbxw7* deletion resulted in the development of multiple tumor types [25]. Similarly, intestinal tumorigenesis in APC mice was accelerated by heterozygous deletion of *Fbxw7* [7]. In the current study, we found that homozygous deletion of *Fbxw7* produced a dramatic acceleration of pancreatic tumorigenesis that was not observed under heterozygous conditions. However, consistent with a haploinsufficient function of Fbxw7, KFC^{fl/fl} mice did have an increased frequency of PanIN lesions relative to KC mice (Figure 2B), leading to the development of PDA in a subset of these mice (7%). Taken together, these results are consistent with a haploinsufficient effect of Fbxw7 on PanIN development but clearly demonstrate that the most profound effects of Fbxw7 on *Kras*^{G12D}-induced tumorigenesis require homozygous deletion.

Our findings implicate Yap accumulation as being causally related, at least in part, to the accelerated growth observed in Fbxw7-deficient,

Kras mutant pancreatic cancer cells. Furthermore, it is also possible that YAP accumulation is associated with the observed genomic instability, as other studies have shown that YAP can promote genomic instability [26]. Interestingly, we also found c-Myc, another established Fbxw7 substrate, consistently elevated in KFC^{fl/fl} pancreata and FBXW7-deficient human pancreatic cancer cells. Although it is conceivable that direct regulation of c-Myc by Fbxw7 could be a mechanism of tumorigenesis, Yap depletion led to reduced expression of c-Myc (Figure 4, A and C, and Supplementary Figure 5C). This finding, together with a recent report demonstrating YAP-dependent c-MYC expression [27], suggests that c-Myc may function downstream of Yap in *Kras*^{G12D}-induced pancreatic tumorigenesis. Future studies are required to fully define the mechanisms of Fbxw7-mediated regulation of Yap and c-Myc in pancreatic tumorigenesis.

In conclusion, this study demonstrates that Fbxw7 suppresses *Kras*^{G12D}-induced pancreatic tumorigenesis via a Yap-dependent mechanism. These findings suggest that YAP and the Hippo pathway should be explored as a therapeutic strategy in pancreatic cancers with

FBXW7 alterations. Furthermore, based on the genomic instability associated with FBXW7 deletion [14,15], investigation of therapeutic strategies exploiting DNA damage repair pathways is warranted for cancers with FBXW7 alterations.

Acknowledgements

We thank Dr. Ingrid Bergin for pathology assistance.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.neo.2016.08.009>.

References

- [1] Rozenblum E, Schutte M, Goggins M, Hahn SA, Panzer S, Zahurak M, Goodman SN, Sohn TA, Hruban RH, and Yeo CJ, et al (1997). Tumor-suppressive pathways in pancreatic carcinoma. *Cancer Res* **57**, 1731–1734.
- [2] Hingorani SR, Petricoin EF, Maitra A, Rajapakse V, King C, Jacobetz MA, Ross S, Conrads TP, Veenstra TD, and Hitt BA, et al (2003). Preinvasive and invasive ductal pancreatic cancer and its early detection in the mouse. *Cancer Cell* **4**, 437–450.
- [3] Collins MA, Bednar F, Zhang Y, Brisset JC, Galban S, Galban CJ, Rakshit S, Flannagan KS, Adsay NV, and Pasca di Magliano M (2012). Oncogenic Kras is required for both the initiation and maintenance of pancreatic cancer in mice. *J Clin Invest* **122**, 639–653.
- [4] Kapoor A, Yao W, Ying H, Hua S, Liewen A, Wang Q, Zhong Y, Wu CJ, Sadanandam A, and Hu B, et al (2014). Yap1 activation enables bypass of oncogenic Kras addiction in pancreatic cancer. *Cell* **158**, 185–197.
- [5] Zhang W, Nandakumar N, Shi Y, Manzano M, Smith A, Graham G, Gupta S, Vietsch EE, Laughlin SZ, and Wadhwa M, et al (2014). Downstream of mutant KRAS, the transcription regulator YAP is essential for neoplastic progression to pancreatic ductal adenocarcinoma. *Sci Signal* **7**, ra42.
- [6] Grim JE, Knoblaugh SE, Guthrie KA, Hagar A, Swanger J, Hespelt J, Delrow JJ, Small T, Grady WM, and Nakayama KI, et al (2012). Fbw7 and p53 cooperatively suppress advanced and chromosomally unstable intestinal cancer. *Mol Cell Biol* **32**, 2160–2167.
- [7] Sancho R, Jandke A, Davis H, Diefenbacher ME, Tomlinson I, and Behrens A (2010). F-box and WD repeat domain-containing 7 regulates intestinal cell lineage commitment and is a haploinsufficient tumor suppressor. *Gastroenterology* **139**, 929–941.
- [8] Calhoun ES, Jones JB, Ashfaq R, Adsay V, Baker SJ, Valentine V, Hempen PM, Hilgers W, Yeo CJ, and Hruban RH, et al (2003). BRAF and FBXW7 (CDC4, FBW7, AGO, SEL10) mutations in distinct subsets of pancreatic cancer: potential therapeutic targets. *Am J Pathol* **163**, 1255–1260.
- [9] Bailey P, Chang DK, Nones K, Johns AL, Patch AM, Gingras MC, Miller DK, Christ AN, Bruxner TJ, and Quinn MC, et al (2016). Genomic analyses identify molecular subtypes of pancreatic cancer. *Nature* **531**, 47–52.
- [10] Perez-Mancera PA, Rust AG, van der Weyden L, Kristiansen G, Li A, Sarver AL, Silverstein KA, Grutzmann R, Aust D, and Rummele P, et al (2012). The deubiquitinase USP9X suppresses pancreatic ductal adenocarcinoma. *Nature* **486**, 266–270.
- [11] Ji S, Qin Y, Shi S, Liu X, Hu H, Zhou H, Gao J, Zhang B, Xu W, and Liu J, et al (2015). ERK kinase phosphorylates and destabilizes the tumor suppressor FBW7 in pancreatic cancer. *Cell Res* **25**, 561–573.
- [12] Tu K, Yang W, Li C, Zheng X, Lu Z, Guo C, Yao Y, and Liu Q (2014). Fbxw7 is an independent prognostic marker and induces apoptosis and growth arrest by regulating YAP abundance in hepatocellular carcinoma. *Mol Cancer* **13**, 110.
- [13] Davis RJ, Welcker M, and Clurman BE (2014). Tumor suppression by the Fbw7 ubiquitin ligase: mechanisms and opportunities. *Cancer Cell* **26**, 455–464.
- [14] Rajagopalan H, Jallepalli PV, Rago C, Velculescu VE, Kinzler KW, Vogelstein B, and Lengauer C (2004). Inactivation of hCDC4 can cause chromosomal instability. *Nature* **428**, 77–81.
- [15] Zhang Q, Karnak D, Tan M, Lawrence TS, Morgan MA, and Sun Y (2016). FBXW7 Facilitates Nonhomologous End-Joining via K63-Linked Polyubiquitylation of XRCC4. *Mol Cell* **61**, 419–433.
- [16] Thompson BJ, Jankovic V, Gao J, Buonamici S, Vest A, Lee JM, Zavadil J, Nimer SD, and Aifantis I (2008). Control of hematopoietic stem cell quiescence by the E3 ubiquitin ligase Fbw7. *J Exp Med* **205**, 1395–1408.
- [17] Zhang Y, Morris JP, Yan W, Schofield HK, Gurney A, Simeone DM, Millar SE, Hoey T, and Hebrok M (2013). Pasca di Magliano M. Canonical wnt signaling is required for pancreatic carcinogenesis. *Cancer Res* **73**, 4909–4922.
- [18] Sancho R, Gruber R, Gu G, and Behrens A (2014). Loss of Fbw7 reprograms adult pancreatic ductal cells into alpha, delta, and beta cells. *Cell Stem Cell* **15**, 139–153.
- [19] Rajagopalan H and Lengauer C (2004). hCDC4 and genetic instability in cancer. *Cell Cycle* **3**, 693–694.
- [20] Siu KT, Xu Y, Swartz KL, Bhattacharyya M, Gurbuxani S, Hua Y, and Minella AC (2014). Chromosome instability underlies hematopoietic stem cell dysfunction and lymphoid neoplasia associated with impaired Fbw7-mediated cyclin E regulation. *Mol Cell Biol* **34**, 3244–3258.
- [21] Aguirre AJ, Bardeesy N, Sinha M, Lopez L, Tuveson DA, Horner J, Redston MS, and DePinho RA (2003). Activated Kras and Ink4a/Arf deficiency cooperate to produce metastatic pancreatic ductal adenocarcinoma. *Genes Dev* **17**, 3112–3126.
- [22] Hingorani SR, Wang L, Multani AS, Combs C, Deramandt TB, Hruban RH, Rustgi AK, Chang S, and Tuveson DA (2005). Trp53R172H and KrasG12D cooperate to promote chromosomal instability and widely metastatic pancreatic ductal adenocarcinoma in mice. *Cancer Cell* **7**, 469–483.
- [23] Bardeesy N, Cheng KH, Berger JH, Chu GC, Pahler J, Olson P, Hezel AF, Horner J, Lauwers GY, and Hanahan D, et al (2006). Smad4 is dispensable for normal pancreas development yet critical in progression and tumor biology of pancreas cancer. *Genes Dev* **20**, 3130–3146.
- [24] Akhondji S, Sun D, von der Lehr N, Apostolidou S, Klotz K, Maljukova A, Cepeda D, Fiegl H, Dafou D, and Marth C, et al (2007). FBXW7/hCDC4 is a general tumor suppressor in human cancer. *Cancer Res* **67**, 9006–9012.
- [25] Mao JH, Perez-Losada J, Wu D, Delrosario R, Tsunematsu R, Nakayama KI, Brown K, Bryson S, and Balmain A (2004). Fbxw7/Cdc4 is a p53-dependent, haploinsufficient tumour suppressor gene. *Nature* **432**, 775–779.
- [26] Fernandez LA, Squatrito M, Northcott P, Awan A, Holland EC, Taylor MD, Nahle Z, and Kenney AM (2012). Oncogenic YAP promotes radioresistance and genomic instability in medulloblastoma through IGF2-mediated Akt activation. *Oncogene* **31**, 1923–1937.
- [27] Xiao W, Wang J, Ou C, Zhang Y, Ma L, Weng W, Pan Q, and Sun F (2013). Mutual interaction between YAP and c-Myc is critical for carcinogenesis in liver cancer. *Biochem Biophys Res Commun* **439**, 167–172.