

Short report

Open Access

Relationship between tonic inhibitory currents and phasic inhibitory activity in the spinal cord lamina II region of adult mice

Toyofumi Ataka^{1,2} and Jianguo G Gu^{*1}

Address: ¹Department of Oral and Maxillofacial Surgery, McKnight Brain Institute and College of Dentistry, University of Florida, Gainesville, Florida 32610, USA and ²Division of Anesthesiology, Niigata University Graduate School of Medical and Dental Sciences, Niigata 951-8510, Japan

Email: Toyofumi Ataka - ata-p@med.niigata-u.ac.jp; Jianguo G Gu* - jgu@dental.ufl.edu

* Corresponding author

Published: 27 November 2006

Received: 01 November 2006

Molecular Pain 2006, **2**:36 doi:10.1186/1744-8069-2-36

Accepted: 27 November 2006

This article is available from: <http://www.molecularpain.com/content/2/1/36>

© 2006 Ataka and Gu; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Phasic and tonic inhibitions are two types of inhibitory activities involved in inhibitory processing in the CNS. In the spinal cord dorsal horn, phasic inhibition is mediated by both GABAergic and glycinergic inhibitory postsynaptic currents. In contrast to phasic inhibitory currents, using patch-clamp recording technique on spinal cord slices prepared from adult mice we revealed that tonic inhibitory currents were mediated by GABA_A receptors but not by glycine receptors in dorsal horn lamina II region. We found that there was a linear relationship ($r = 0.85$) between the amplitude of tonic inhibitory currents and the frequency of GABAergic inhibitory postsynaptic currents. Analysis of charge transfer showed that the charges carried by tonic inhibitory currents were about 6 times of charges carried by phasic inhibitory currents. The prominent charge transfer by tonic inhibitory currents and their synaptic activity dependency suggest a significant role of tonic inhibition in sensory processing.

Background

GABA (γ -Aminobutyric acid) and glycine are two principle inhibitory neurotransmitters in the spinal cord dorsal horn. They are either released separately or co-released from presynaptic terminals of inhibitory neurons. Upon the binding to GABA_A receptors and glycine receptors at postsynaptic membrane, they elicit inhibitory postsynaptic currents (IPSCs). IPSCs provide phasic inhibition in neuronal network and are important for information processing. In addition to its action at synaptic sites, recent studies in several brain regions of matured animals have indicated that low concentrations of ambient GABA can activate high affinity GABA_A receptors that are expressed at extrasynaptic sites to elicit a sustained inhibitory current [1-5]. A term 'tonic inhibitory currents' has been used to describe this sustained inhibitory current [6]. Functionally, tonic GABAergic inhibition has been shown

to control neuronal excitability in the brain [7-9]. Tonic inhibitory currents have been identified in rat cerebellar granule cells [1], granule cells of the dentate gyrus [2], thalamocortical relay neurons of the ventral basal complex [3], layer V pyramidal neurons in the somatosensory cortex [4], inhibitory interneurons in the CA1 region of the hippocampus [5]. However, not all CNS neurons that were examined displayed tonic inhibitory currents under normal conditions. For example tonic inhibitory currents were normally not observed in hippocampal pyramidal cells in brain slices from adult animals [[5,10], but see [11]].

The lamina II of the spinal dorsal horn (substantia gelatinosa) plays an important role in processing nociceptive input from fine myelinated A δ – and unmyelinated C-primary afferents from the periphery [12]. In this area, inhib-

itory neurons produce feedback and feed-forward inhibition to control nociceptive input, and a reduction of inhibitory activity in lamina II can result in central sensitization [13,14]. Previous studies on inhibitory controls in the spinal cord dorsal horn have been mainly focused on inhibitory postsynaptic currents, i.e. phasic inhibition. Little is known about whether tonic inhibitory currents are present in this region and if so, whether tonic inhibitory currents are mediated by GABA receptors and/or glycine receptors.

Materials and methods

The methods for preparing thick adult mouse spinal cord slices, as well as blind whole-cell patch-clamp recording techniques, have been described previously [15]. In brief, transverse spinal cord slices (500–600 μm in thickness) were prepared from L5 spinal cords of adult mice (Harlan, IN, USA) aged between 6 and 9 weeks. In each experiment, a spinal cord slice was transferred to a recording chamber (volume of 0.5 ml). The slice was supported at the bottom by a nylon mesh in the recording chamber. The slice was superfused with Krebs solution at flow rate of 10 ml/min. The Krebs solution contained (in mM): NaCl 117, KCl 3.6, CaCl_2 2.5, MgCl_2 1.2, NaH_2PO_4 1.2, NaHCO_3 25, and glucose 11. The solution was equilibrated with 95% O_2 and 5% CO_2 , maintained at room temperature (22°C), and the pH of the Krebs solution was 7.35. Under a dissecting microscope with 40 \times magnification, lamina regions were identified based on morphological features. The lamina II was clearly discernible as a relatively translucent band across the superficial dorsal horn. Under visual guidance, the patch electrode was inserted vertically into the lamina II.

Whole-cell patch-clamp recordings were made from lamina II neurons with electrodes (5–10 M Ω) filled with an internal solution containing (in mM): K-gluconate 120, KCl 20, MgCl_2 2, Na_2ATP 2, NaGTP 0.5, HEPES 20, EGTA 0.5, and pH 7.2 adjusted with NaOH. Signals were amplified and filtered at 2 kHz and sampled at 5 kHz (Axopatch 200B). Spontaneous inhibitory postsynaptic currents (sIPSCs) were recorded with cells being held at 0 mV. Each recording was performed on a cell in a fresh slice without prior application of any agonist or antagonist. Isolation of GABAergic sIPACs was accomplished by including 2 μM strychnine in the bath solution. Tonic inhibitory currents were revealed following the application of 20 μM bicuculline for a period of 3 min. All compounds tested were applied through the bath solution at a flow rate of 10 ml/min. Analysis of sIPSCs, including threshold setting and peak identification criteria, were performed according to a method previously described [15]. Decay time constant (τ) of sIPSCs was analyzed using Clampfit 9 (Axon Instruments, Inc., Sunnyvale, CA, USA). sIPSC frequency, amplitude, and average charge transfer (Q_{sIPSCs} , integrated


area under sIPSCs) were analyzed using Mini Analysis Program (Synaptosoft, Inc., Decatur, GA, USA). Charge transfer (Q_{PC}) associated with sIPSCs in a given time (t) was calculated using the equation $Q_{\text{PC}} = f \times Q_{\text{sIPSCs}} \times t$, where f is the frequencies (Hz) of sIPSCs, Q_{sIPSCs} is the average charge transfer per sIPSC during a 60-s baseline recording, and t is the duration (60 s), respectively. The charge transfer associated with tonic currents was calculated according to the equation: $Q_{\text{TC}} = I_{\text{TC}} \times t$, where Q_{TC} is the charge transfer produced by tonic currents, I_{TC} is the current amplitude at steady-state, and t is time (60 s). Unless otherwise indicated, data represent Mean \pm SEM, Student's t -tests were used for statistical comparison, and significance was considered at the $P < 0.05$ level.

Results

Under our voltage-clamp condition with cells being held at 0 mV, spontaneous inhibitory postsynaptic currents (sIPSCs) recorded from lamina II neurons were outward currents, and spontaneous excitatory postsynaptic currents (sEPSCs) were not detectable because the holding potential of 0 mV is at the reversal potential for sEPSCs. Phasic inhibitory activities (or sIPSCs) showed three types, rapid, slow, and mixed types, based on the kinetics of their decay phases. When these different types of sIPSCs are integrated together, it yields an sIPSC that best fits into a two-exponential equation (Figure 1A). Following the application of 2 μM strychnine to block glycine receptors, rapid type of sIPSCs disappeared but slow sIPSCs remained. When bath solution contained 20 μM bicuculline but not strychnine, only rapid type of sIPSCs could be observed (Figure 1A). All sIPSCs could be completely blocked in the presence of both bicuculline (2 μM) and strychnine (20 μM , not shown).

Application of 20 μM bicuculline not only inhibited slow types of sIPSCs, but also caused a shift of baseline holding current (Figure 1B). The shift of baseline hold currents represents the presence of tonic inhibitory currents mediated by GABA $_A$ receptors [6]. The amplitudes of the tonic inhibitory currents revealed by bicuculline were 8.4 ± 0.7 pA ($n = 30$, Figure 1D). After washing off bicuculline, hold currents returned to the baseline levels.

Glycinergic inhibitory postsynaptic currents contributed to phasic inhibitory activity in lamina II region, and application of glycine receptor antagonist strychnine (2 μM) blocked glycinergic sIPSCs. However, there was no significant change of baseline holding currents following the application 2 μM strychnine (Figure 1C). The baseline holding currents were -0.3 ± 0.5 pA ($n = 6$) in lamina II, not significantly different from the baseline noise level (Figure 1D). These results suggested that glycine receptors did not significantly account for the tonic inhibitory currents in dorsal horn lamina II neurons of adult mice.

**Figure 1**

Revealing tonic inhibitory currents in lamina II neurons of adult mice. A). Traces on the left side show sIPSCs recorded in normal bath solution (control), following the applications of either 2 μM strychnine (Stry) or 20 μM bicuculline (Bic). Traces on the right side are the average of 100 sIPSCs. τ is time constant for averaged trace. B). Sample trace (top panel) shows the shift of baseline holding current following the application of 20 μM bicuculline. The lower traces show at an expanded scale the baselines before and during bicuculline application as well after wash off bicuculline. C). Sample trace shows that application of strychnine (2 μM) inhibited some sIPSCs but did not affect baseline holding current. D). Bar graph shows pooled results of baseline shift following the application of either 20 μM bicuculline ($n = 30$) or 2 μM strychnine ($n = 6$). Bicuculline or strychnine was applied for 3 min in each experiment.

Amplitude of tonic inhibitory currents showed large variations among different recordings. One possible cause of this variation might be due to the differences of resting membrane potentials of these neurons. Therefore, we measured resting membrane potentials of each recorded neurons and plotted amplitudes of tonic inhibitory currents against resting membrane potentials of each recording. No relationship was found between tonic inhibitory current amplitudes and resting membrane potentials ($r = 0.27$, $n = 30$, Figure 2A). We determined whether amplitude of tonic inhibitory currents were in proportional to inhibitory synaptic inputs by plotting tonic inhibitory current amplitude against sIPSC frequency. We also did not find a strong association between the amplitudes of tonic inhibitory currents and the frequency of total sIPSCs ($r = 0.5$, $n = 30$, Figure 2B).

Total sIPSCs in the spinal cord dorsal horn included both glycinergic and GABAergic inhibitory synaptic activity. We isolated GABA_A receptor-mediated sIPSCs from total sIPSCs by including 2 μ M strychnine in bath solution. Under this condition, glycine receptor-mediated sIPSCs were completely abolished. We then applied 20 μ M bicuculline to reveal tonic currents, and we found that neurons receiving higher frequency of GABAergic synaptic inputs usually had larger amplitudes of tonic inhibitory currents (Figure 3A), and neurons receiving lower frequency of GABAergic synaptic inputs usually had smaller amplitudes of tonic inhibitory currents (Figure 3B). There was a good linear relationship between the amplitude of GABAergic tonic inhibitory currents and the frequency of GABAergic IPSCs ($r = 0.8515$, $n = 14$, Figure 3C).

Charge transfer through GABA_A receptors is a measure of inhibition for both phasic and tonic inhibitory currents. We determined charge transfer carried by phasic inhibitory currents and by tonic inhibitory currents in a period of 60 sec recording (Figure 4A). Charge transfer was 1.5 ± 0.31 pC ($n = 44$) for the total phasic currents mediated by both GABAergic and glycinergic inhibitory postsynaptic currents (Figure 4B). On the other hand, charge transfer was 8.7 ± 0.89 pC ($n = 44$) for tonic inhibitory currents mediated by GABA_A receptors, and was about 6 times of the total charge transfer mediated by both GABAergic and glycinergic inhibitory postsynaptic currents (Figure 4B).

Discussion

Using spinal cord slice preparations from adult mice, the present study shows that tonic inhibitory currents are present in lamina II of the dorsal horn and was solely mediated by GABA_A receptors, that the extent of tonic inhibition is proportional to GABAergic inhibitory synaptic activity, and that tonic currents transfer charges substantially higher than phasic currents. The results provide




Figure 2
Lack of correlation between tonic inhibitory currents and resting membrane potentials as well as between tonic inhibitory currents and total sIPSC activity. A). Graph shows a plot of tonic inhibitory currents against membrane potentials for each recorded neuron ($n = 30$). B). Graph shows a plot of tonic inhibitory current against sIPSC frequency ($n = 30$). Tonic inhibitory currents were revealed by the applications of 20 μ M bicuculline. Both resting membrane potentials and sIPSCs were measured before the application of bicuculline.

new information about inhibitory activities in a nociceptive processing region.

We have observed a large variation of the size of tonic inhibitory currents. It has been suggested that ambient GABA concentrations around the extrasynaptic domains of neurons is a factor that determines the size of tonic inhibitory currents. The concentrations of ambient GABA




Figure 3
Correlation between tonic inhibitory currents and GABAergic inhibitory synaptic activity. A). Sample trace shows bicuculline-induced baseline shift in a lamina II neuron with high GABAergic inhibitory synaptic activity. B). Sample trace shows bicuculline-induced baseline shift in another lamina II neuron with low GABAergic inhibitory synaptic activity. C). Plot of GABAergic tonic inhibitory currents against frequency of GABAergic sIPSCs ($n = 14$). Linear regress coefficient (r) = 0.8515. All experiments were performed in the presence of 2 μ M strychnine.

vary from tens of nanomolar to a few micromolar based on *in vivo* microdialysis studies in the brain [16-19]. One possible cause of the variation may be regional and temporal differences in GABAergic neuron activity. We have shown that GABAergic sIPSC frequencies recorded in lamina II neurons have a big variation, and GABAergic sIPSC frequency and tonic current size are correlated. It is very likely that neurons with higher GABAergic sIPSC frequency have higher concentrations of ambient GABA around them due to more frequent spillover of GABA from inhibitory synapses. Therefore, these neurons have larger-sized tonic inhibitory currents mediated by extrasynaptic GABA_A receptors. This is consistent with previous




Figure 4
Charge transfers mediated by phasic inhibitory currents and by tonic inhibitory currents. A). Diagram illustrates the measurement of phasic inhibitory current charge transfer (Q_{PC}) and tonic inhibitory current charge transfer (Q_{TC}). B). Bar Graph shows the comparison between Q_{PC} and Q_{TC} ($n = 32$). Tonic currents were revealed by the application of 20 μ M bicuculline. Strychnine was not included in bath solution.

studies in cerebellar granule cells, which shows that ambient GABA concentration is maintained by action potential-dependent vesicular release and is responsible for tonic GABA_A receptor activation [1,7,20-23].

Although phasic inhibition is mediated by both GABA_A and glycine receptors in spinal cord lamina II of adult mice, we did not observe any contribution of glycine receptors to tonic inhibitory currents. The lack of glycine receptor-mediated tonic inhibitory currents could be mainly due to the lack of high affinity glycine receptor isoforms. Of different glycine receptor isoforms ($\alpha 1\beta$, $\alpha 2\beta$, $\alpha 3\beta$) identified in the CNS, their EC₅₀ values for glycine were normally above 50 μ M [24]. In contrast, extrasynaptic GABA receptors that contribute to tonic currents were found to be high affinity isoforms (e.g. $\alpha 6\beta\gamma\delta$, $\alpha 4\beta\gamma\delta$ and $\alpha 5\beta\gamma\gamma_2$) with EC₅₀ in nanomolar range [6].

The large charge transfer carried by GABAergic tonic inhibitory currents shown in this study indicates a persistent increase in the input conductance of lamina II dorsal horn neurons. The increase of input conductance in neurons can decrease the size and duration of the excitatory postsynaptic potentials (EPSPs), and make neurons less likely to generate action potentials. Indeed, tonic currents have been shown to be a critical determinant that controls neuron excitability in cerebellar granule cells [7-9]. In spinal cord lamina II, decreases of neuron excitability by GABAergic tonic inhibitory current may be an important

mechanism to control nociceptive inputs and to prevent central hyper-sensitization.

Competing interests

The author(s) declare that they have no competing interests.

References

- Kaneda M, Farrant M, Cull-Candy SG: **Whole-cell and single-channel currents activated by GABA and glycine in granule cells of the rat cerebellum.** *J Physiol (Lond)* 1995, **485**:419-435.
- Nusser Z, Mody I: **Selective modulation of tonic and phasic inhibitions in dentate gyrus granule cells.** *J Neurophysiol* 2002, **87**:2624-28.
- Porcello DM, Huntsman MM, Mihalek RM, Homanics GE, Huguenard JR: **Intact synaptic GABAergic inhibition and altered neurosteroid modulation of thalamic relay neurons in mice lacking δ subunit.** *J Neurophysiol* 2003, **89**:1378-86.
- Yamada J, Yamamoto S, Ueno S, Furukawa T, Fukuda A: **GABAA receptor-mediated tonic inhibition in rat somatosensory cortex.** *FENS Forum Abstr* 2004, **2**: A083.027.
- Semyanov A, Walker MC, Kullmann DM: **GABA uptake regulates cortical excitability via cell type-specific tonic inhibition.** *Nat Neurosci* 2003, **6**:484-90.
- Farrant M, Nusser Z: **Variations on an inhibitory theme: phasic and tonic activation of GABA_A receptors.** *Nat Rev Neurosci* 2005, **6**:215-29.
- Brickley S, Cull-Candy S, Farrant M: **Development of a tonic form of synaptic inhibition in rat cerebellar granule cells resulting from persistent activation of GABAA receptors.** *J Physiol (Lond)* 1996, **497**:753-759.
- Hamann M, Rossi DJ, Attwell D: **Tonic and spillover inhibition of granule cells control information flow through cerebellar cortex.** *Neuron* 2002, **33**:625-33.
- Chadderton P, Margrie TW, Haussler M: **Integration of quanta in cerebellar granule cells during sensory processing.** *Nature* 2004, **428**:856-60.
- Caraiscos VB, Elliott EM, You-Ten KE, Cheng YV, Belelli D, Newell JG, Jackson MF, Lambert JJ, Rosahl TW, Wafford KA, MacDonald JF, Orser BA: **Tonic inhibition in mouse hippocampal CA1 pyramidal neurons is mediated by α 5 subunit-containing γ -aminobutyric acid type A receptors.** *Proc Natl Acad Sci USA* 2004, **101**:3662-7.
- Bai D, Zhu G, Pennefather P, Jackson MF, MacDonald JF, Orser BA: **Distinct functional and pharmacological properties of tonic and quantal inhibitory postsynaptic currents mediated by γ -aminobutyric acid_A receptors in hippocampal neurons.** *Mol Pharmacol* 2001, **59**:814-824.
- Yoshimura M, Jessell T: **Amino acid-mediated EPSPs at primary afferent synapses with substantia gelatinosa neurones in the rat spinal cord.** *J Physiol* 1990, **430**:315-35.
- Moore KA, Kohno T, Karchewski LA, Scholz J, Baba H, Woolf CJ: **Partial peripheral nerve injury promotes a selective loss of GABAergic inhibition in the superficial dorsal horn of the spinal cord.** *J Neurosci* 2002, **22**:6724-31.
- Baba H, Ji RR, Kohno T, Moore KA, Ataka T, Wakai A, Okamoto M, Woolf CJ: **Removal of GABAergic inhibition facilitates polysynaptic A fiber-mediated excitatory transmission to the superficial spinal dorsal horn.** *Mol Cell Neurosci* 2003, **24**:818-830.
- Nakatsuka T, Chen M, Takeda D, King C, Ling J, Xing H, Ataka T, Vierck C, Yezierski R, Gu JG: **Substance P-driven feed-forward inhibitory activity in the mammalian spinal cord.** *Mol Pain* 2005, **1**:20.
- Lerma J, Herranz AS, Herreras O, Abaira V, Martin del Rio R: **In vivo determination of extracellular concentration of amino acids in the rat hippocampus. A method based on brain dialysis and computerized analysis.** *Brain Res* 1986, **384**:145-55.
- Tossman U, Jonsson G, Ungerstedt U: **Regional distribution and extracellular levels of amino acids in rat central nervous system.** *Acta Physiol Scand* 1986, **127**:533-45.
- Kennedy RT, Thompson JE, Vickroy TW: **In vivo monitoring of amino acids by direct sampling of brain extracellular fluid at ultralow flow rates and capillary electrophoresis.** *J Neurosci Methods* 2002, **114**:39-49.
- Xi ZX, Ramamoorthy S, Shen H, Lake R, Samuvel DJ, Kalivas PW: **GABA transmission in the nucleus accumbens is altered after withdrawal from repeated cocaine.** *J Neurosci* 2003, **23**:3498-505.
- Tia S, Wang JF, Kotchabhakdi N, Vicini S: **Developmental changes of inhibitory synaptic currents in cerebellar granule neurons: role of GABAA receptor α 6 subunit.** *J Neurosci* 1996, **16**:3630-40.
- Wall MJ, Usowicz MM: **Development of action potential-dependent and independent spontaneous GABAA receptor-mediated currents in granule cells of postnatal rat cerebellum.** *Eur J Neurosci* 1997, **9**:533-48.
- Brickley SG, Cull-Candy SG, Farrant M: **Vesicular release of GABA contributes to both phasic and tonic inhibition of granule cells in the cerebellum of mature mice.** *J Physiol* 2003:547. P, C30.
- Carta M, Mameli M, Valenzuela CF: **Alcohol enhances GABAergic transmission to cerebellar granule cells via an increase in Golgi cell excitability.** *J Neurosci* 2004, **24**:3746-51.
- Aguayo LG, van Zundert B, Tapia JC, Carrasco MA, Alvarez FJ: **Changes on the properties of glycine receptors during neuronal development.** *Brain Res Brain Res Rev* 2004, **47**:33-5.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

