

ORIGINAL ARTICLE

Adseverin mediates RANKL-induced osteoclastogenesis by regulating NFATc1

Min-Kyoung Song, Zang Hee Lee and Hong-Hee Kim

Adseverin is a Ca^{2+} -dependent actin filament-severing protein that has been reported to regulate exocytosis via rearrangements of the actin cytoskeleton in secretory cells. However, the role of adseverin in bone cells has not yet been well characterized. Here, we investigated the role of adseverin in osteoclastogenesis using primary osteoclast precursor cells. Adseverin expression was upregulated during RANKL (receptor activator of nuclear factor- κ B ligand)-induced osteoclast differentiation. Moreover, genetic silencing of adseverin decreased the number of osteoclasts generated by RANKL. Adseverin knockdown also suppressed the RANKL-mediated induction of nuclear factor of activated T-cell c1 (NFATc1), which is a key transcription factor in osteoclastogenesis. In addition, adseverin knockdown impaired bone resorption and the secretion of bone-degrading enzymes from osteoclasts. These effects were accompanied by decreased NFATc1 expression and the activation of nuclear factor- κ B. Collectively, our results indicate that adseverin has a crucial role in osteoclastogenesis by regulating NFATc1.

Experimental & Molecular Medicine (2015) 47, e199; doi:10.1038/emm.2015.94; published online 18 December 2015

INTRODUCTION

Bone is a complex tissue that is constantly remodeled and maintained by cycling between bone formation and resorption throughout life. This dynamic phenomenon is mediated by two types of cells, osteoblasts and osteoclasts. Osteoblasts form the bone matrix by depositing organic and inorganic components, whereas osteoclasts resorb bone by secreting degrading enzymes and protons. Osteoclasts are multinuclear cells that differentiate from the monocyte/macrophage lineage of hematopoietic cells.¹ Two main factors are needed for osteoclast differentiation, including macrophage-colony-stimulating factor (M-CSF) and the receptor activator of nuclear factor- κ B ligand (RANKL). M-CSF is important for proliferation of the monocyte/macrophage lineage and osteoclast survival,² whereas RANKL induces differentiation during osteoclastogenesis.³ When RANKL binds to its receptor (RANK), TNF receptor-associated factor 6 (TRAF6) is recruited, thereby initiating the nuclear factor- κ B (NF- κ B) and mitogen-activated protein kinase (MAPK) signaling pathways. The calcium signaling pathway is also activated by RANKL in response to additional stimulation of surface immune receptors. These pathways all culminate in the activation of nuclear factor of activated T-cell (NFAT) c1 (NFATc1), the master transcription factor of osteoclastogenesis. Taken together with other accessory transcription factors, NFATc1 drives the expression of a number of

osteoclast-specific genes.⁴ Upon expression of osteoclast-specific genes including tartrate-resistant acid phosphatase (TRAP), *cathepsin K* and dendritic cell-specific transmembrane protein (*DC-STAMP*), osteoclasts become mature and functionally activated. Mature osteoclasts secrete bone-degrading enzymes into the resorption lacunae by exocytosis. By sealing off the resorption lacunae with an actin belt, osteoclasts induce the effective concentration of enzymes and protons in the resorbing space.

Adseverin, also called *scinderin*, is a member of the gelsolin superfamily of actin-binding proteins. Although adseverin has the highest degree of homology with gelsolin (~60% at the amino-acid level), the expression of adseverin is more restricted.⁵ Adseverin was first discovered in chromaffin cells of the adrenal medulla, and it is now believed to be expressed in all secretory cells. In secretory cells, adseverin mediates actin cytoskeletal remodeling during exocytosis.^{6–8} Similar to gelsolin, adseverin contains six gelsolin-like (G) domains, three actin binding sites and two calcium binding sites with different affinities.^{7,9} Phosphatidylinositol-4,5-bisphosphate has been shown to inhibit the interaction between actin and adseverin, most likely through competition between actin and phosphatidylinositol-4,5-bisphosphate.^{10,11} Upon stimulation, adseverin interacts with F-actin in the presence of Ca^{2+} and decreases the viscosity of actin gels by severing actin filaments.⁷

Immunofluorescence experiments have revealed that adseverin is mainly localized in the subplasmalemmal region, near a mesh of actin filaments, in chromaffin cells.⁷ Although the role of adseverin in chromaffin cells has been well characterized, the role of adseverin in other secretory cells, including osteoclasts, is relatively unknown. One report found that gelsolin^{-/-} mice exhibited increased bone thickness, high bone mass and defects in podosome assembly and bone resorption,¹² suggesting that additional gelsolin superfamily proteins such as adseverin may be involved in bone remodeling.

In the present study, we found that adseverin expression is upregulated during the differentiation of primary bone marrow-derived macrophages (BMMs) into osteoclasts. Using small interfering RNA (siRNA)-mediated gene silencing, we found that adseverin promotes osteoclast differentiation by upregulating NFATc1 and activating NF- κ B signaling, especially p65 phosphorylation and nuclear translocation.

MATERIALS AND METHODS

Reagents

Recombinant human M-CSF and human soluble RANKL were purchased from PeproTech (Rocky Hill, NJ, USA). Antibodies against ERK, JNK, p38, Akt, p65, I κ B and PARP were obtained from Cell Signaling Technology (Cambridge, MA, USA), as were phospho-specific antibodies for ERK (Thr202/Tyr204), JNK (Thr182/Tyr185), p38 (Thr180/Tyr182), Akt (Ser473), p65 (Ser536) and I κ B (Ser32). Monoclonal antibodies against β -actin (AC-74) and secondary antibodies were obtained from Sigma-Aldrich (St Louis, MO, USA). Antibodies against adseverin (N17), NFATc1 (7A6), c-Fos (H125), lamin B (M-20) and α -tubulin (TU-02) were purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). The siRNA targeting adseverin and the negative control siRNA were also obtained from Santa Cruz Biotechnology. All other reagents were obtained from Sigma-Aldrich.

Culture of osteoclasts

To obtain mouse bone marrow-derived osteoclasts, 5-week-old female ICR mice were killed and their bone marrow was flushed from their tibiae and femora. After eliminating erythrocytes using hypotonic buffer, cells were incubated overnight in α -modified Eagle's medium (α -MEM) supplemented with 10% fetal bovine serum in culture dishes. Adherent cells were discarded, and nonadherent cells were further incubated with 30 ng ml⁻¹ M-CSF in Petri dishes. After 3 days, adherent BMMs were collected by scraping and used as osteoclast precursor cells. BMMs were differentiated into osteoclasts by culturing them with 30 ng ml⁻¹ M-CSF and 120 ng ml⁻¹ RANKL.

TRAP staining

Cells were fixed in 3.7% formaldehyde and permeabilized using 0.1% Triton X-100. Leukocyte acid phosphatase (TRAP) staining was performed in the dark for 5–15 min using a commercially available kit (Sigma-Aldrich) according to the manufacturer's instructions.

siRNA-mediated knockdown

BMMs were transfected with 30 nM adseverin-specific siRNAs (Santa Cruz Biotechnology). Briefly, siRNAs and HiPerFect (Qiagen, Hilden, Germany) or Lipofectamine 2000 (Invitrogen, Carlsbad, CA, USA) transfection reagent were mixed with serum- and antibiotic-free

α -MEM. The siRNA mixtures were incubated for 20 min at room temperature and then added to the cells. The cells were incubated for 6–14 h, and then the culture medium was replaced with α -MEM supplemented with serum.

Reverse transcription-polymerase chain reaction

Total RNA was isolated from cells using TRIzol reagent (Invitrogen) according to the manufacturer's instructions. Next, 3 μ g RNA was used for cDNA synthesis using Superscript II reverse transcriptase (Invitrogen). The PCR reaction was performed in a thermal cycler (Bio-Rad, Hercules, CA, USA), and the thermocycling conditions were as follows: an initial denaturation step at 95 °C for 5 min, followed by 27 cycles of denaturation at 95 °C for 30 s, annealing at 57 °C for 30 s and extension at 72 °C for 30 s. The following primers were used for amplification: adseverin, 5'-GAGGAACAGACCCAGCAAAT-3' and 5'-GCCACAGAGGCCTGTATCTT-3'; and HPRT, 5'-CCTAAGATG AGCGCAAGTTGAA-3' and 5'-CCACAGGGACTAGAACACCTGC TAA-3'. Real-time PCR was performed using an ABI7300 real-time system (Applied Biosystems, Warrington, UK) and a KAPA SYBR FAST qPCR Kit (Kapa Biosystems, Wilmington, MA, USA). Thermocycling conditions were as follows: an initial activation step at 95 °C for 3 min, followed by 40 cycles of denaturation at 95 °C for 3 s and amplification at 60 °C for 33 s. All reactions were run in triplicate, and the housekeeping gene *HPRT* was used for normalization. Gene expression levels were determined as fold changes using the cycle threshold comparison method. The following primer sets were used: adseverin, 5'-TCCAGAGCAGAGAGCTCAA-3' and 5'-TTCGCA GTCAGATCATTGGT-3'; NFATc1, 5'-CCAGTATACCAGCTCTG CCA-3' and 5'-GTGGGAAGTCAGAAGTGGGT-3'; TRAP, 5'-CGA CCATTGTTAGCCACATACG-3' and 5'-TCGTCCTGAAGATACTG CAGGT-3'; cathepsin K, 5'-ATATGTGGGCCACCATGAAAGTT-3' and 5'-TCGTTCCCCACAGGAATCTCT-3'; DC-STAMP, 5'-GGGTG CTGTTTGCCGCTG-3' and 5'-CGACTCCTTGGGTTCCCTTGCT-3'; v-ATPase (*Atp6v0d2*), 5'-GGGAGACCCTCTTCCCCACC-3' and 5'-CCACCGACAGCGTCAAACAAA-3'; and HPRT, 5'-CCTAAG ATGAGCGCAAGTTGAA-3' and 5'-CCACAGGGACTAGAACACC TGCTAA-3'.

Immunocytochemistry

BMMs were cultured on 15-mm glass coverslips in 24-well plates and then fixed with 3.7% formaldehyde. After permeabilization with 0.1% Triton X-100, nonspecific binding sites were blocked using phosphate-buffered saline containing 1% bovine serum albumin for 1 h and 30 min. Cells were then incubated with anti-adseverin antibodies overnight at 4 °C. Cells were then washed and stained using rhodamine-phalloidin and fluorescein isothiocyanate-conjugated secondary antibodies for 2 h. After washing the cells with phosphate-buffered saline, they were mounted in the presence of 4',6-diamidino-2-phenylindole (DAPI). Images were obtained using a ZEISS LSM700 confocal microscope (Carl Zeiss MicroImaging GmbH, Goettingen, Germany).

Transwell migration assay

BMMs were seeded in Petri dishes and transfected with adseverin and control siRNAs. After culturing the cells with 30 ng ml⁻¹ M-CSF and 120 ng ml⁻¹ RANKL for 2 days, the cells were scraped and transferred to the upper chambers of transwell plates (Corning, Corning, NY, USA). The lower chambers contained 240 ng ml⁻¹ RANKL as an attractant. After 16 h of incubation in a CO₂ incubator at 37 °C, cells were fixed and stained with crystal violet.

Resorption assay

BMMs were seeded on calcium phosphate-coated 48-well plates and cultured with 30 ng ml^{-1} M-CSF and 120 ng ml^{-1} RANKL. After 7 days, the plates were rinsed with distilled water and stained with von Kossa reagents. Photographs were taken under a light microscope, and the resorption area was quantified using the ImageJ software (NIH, Bethesda, MD, USA).

MMP assay

BMMs were cultured with 30 ng ml^{-1} M-CSF and 120 ng ml^{-1} RANKL for 2 days, after which the medium was replaced with serum-free α -MEM. After 12 h, the conditioned medium was collected, and matrix metalloproteinase (MMP) activity was measured using an MMP Activity Assay Kit (Abcam, Cambridge, UK) according to the manufacturer's instructions.

Western blotting

Cells were lysed using RIPA buffer (120 mM Tris-HCl (pH 7.5), 150 mM NaCl, 1 mM Na_2EDTA , 1 mM EGTA, 0.5% NP-40, 2.5 mM sodium pyrophosphate, 1 mM β -glycerophosphate, 1 mM Na_3VO_4 , 1 mM NaF and a protease inhibitor cocktail; Roche, Mannheim, Germany). The protein concentrations of the cell lysates were measured using a DC Protein Assay Kit (Bio-Rad), and equal amounts of protein were loaded onto 8 or 10% sodium dodecyl sulfate-polyacrylamide gels for electrophoresis. After transfer onto nitrocellulose membranes (Amersham Pharmacia, Uppsala, Sweden), nonspecific binding sites on the membranes were blocked with 5% non-fat skim milk for 1 h, and the membranes were then incubated with primary antibodies overnight at 4°C . The membranes were then washed several times with TBST (Tris-buffered saline with Tween-20) and incubated with horseradish peroxidase-conjugated secondary antibodies in 2% skim milk for 1 h. Immunoreactive bands were detected using ECL reagents in the dark. Nuclear and cytoplasmic protein fractions were prepared using NE-PER nuclear and

cytoplasmic extraction reagents (Pierce, Rockford, IL, USA) according to the manufacturer's protocols.

Statistical analysis

Statistical differences between results were tested using Student's *t*-tests. *P*-values < 0.05 were regarded as significant.

RESULTS

Adseverin expression is upregulated during RANKL-induced osteoclastogenesis

In our preliminary studies, we used microarrays to identify the transcriptional changes that accompany osteoclast differentiation. We found that adseverin expression was upregulated 12-fold at day 3 after RANKL treatment (data not shown). To validate this result, BMMs were cultured with M-CSF and RANKL, and the level of adseverin mRNA was measured using quantitative real-time PCR. Adseverin expression was found to be increased up to 50-fold at day 3 (Figure 1a). Upregulation of TRAP, an osteoclastogenesis marker gene, was also detected, confirming that effective osteoclast differentiation occurred in the primary precursor cells. Moreover, RANKL treatment increased the level of adseverin mRNA in a dose-dependent manner at concentrations up to 120 ng ml^{-1} (Figure 1b). The level of adseverin protein was also increased, peaking at day 3 post-RANKL treatment (Figure 1c).

Adseverin depletion impairs osteoclast differentiation and actin ring formation

To investigate whether adseverin has a role in osteoclastogenesis, adseverin expression was silenced in BMMs using adseverin-specific siRNAs (Figure 2a). As expected, culturing control cells with M-CSF and RANKL resulted in a




Figure 1 The expression of adseverin during receptor activator of nuclear factor- κ B ligand (RANKL)-induced osteoclastogenesis. (a) Bone marrow-derived macrophages (BMMs) were cultured with 30 ng ml^{-1} macrophage-colony-stimulating factor (M-CSF) and 120 ng ml^{-1} RANKL for 0, 1, 2 or 3 days. Total RNA was isolated and analyzed using quantitative real-time PCR. The mRNA level of adseverin was normalized to the level of HPRT (hypoxanthine-guanine phosphoribosyltransferase). The mRNA level of tartrate-resistant acid phosphatase (TRAP), a marker of osteoclast differentiation, was also analyzed. (b) BMMs were stimulated with the indicated doses of RANKL in the presence of 30 ng ml^{-1} M-CSF. Cells were cultured for 3 days and then analyzed using real-time PCR. (c) Western blot analysis was performed using antibodies against adseverin and β -actin. Protein lysates were prepared from BMMs cultured for 0, 1, 2, 3 or 5 days with 30 ng ml^{-1} M-CSF and 120 ng ml^{-1} RANKL. The data are presented as the means \pm s.d. $**P < 0.005$.




Figure 2 Adseverin depletion impairs osteoclast differentiation and actin ring formation. **(a)** Bone marrow-derived macrophages (BMMs) were transfected with 30 nm of control or adseverin-specific small interfering RNA (siRNA) and cultured for 2 days with 120 ng ml^{-1} receptor activator of nuclear factor- κ B ligand (RANKL) and 30 ng ml^{-1} macrophage-colony-stimulating factor (M-CSF). Adseverin mRNA expression was then analyzed using reverse transcription-polymerase chain reaction (RT-PCR). **(b)** siRNA-transfected BMMs were cultured with M-CSF and RANKL for 4 days. Cells were fixed and then stained for tartrate-resistant acid phosphatase (TRAP). Images were captured using a light microscope (magnification $\times 100$), and TRAP-positive multinucleated cells (MNCs) were counted. **(c)** siRNA-transfected BMMs were cultured in 96-well plates. CCK assays were performed on days 0, 1, 2 and 3 post-RANKL treatment. **(d)** BMMs were cultured with 30 ng ml^{-1} M-CSF and 120 ng ml^{-1} RANKL for 4 days. The actin ring was labeled with rhodamine-phalloidin (red), and cells were also immunostained using anti-adseverin-FITC (fluorescein isothiocyanate) antibodies. Nuclei were counterstained with 4',6-diamidino-2-phenylindole (DAPI). **(e)** As in **d**, fixed cells were stained with DAPI and rhodamine-phalloidin and then counted for the number of nuclei per osteoclast. **(f)** BMMs were cultured for 2 days with M-CSF and RANKL, and migration assays were performed for 16 h using transwell plates. The data are presented as the means \pm s.d. * $P < 0.05$; ** $P < 0.005$.

considerable number of multinucleated cells that were positive for the osteoclast marker TRAP at day 4. However, only a few multinucleated cells were observed in the adseverin-knockdown cells; moreover, most of the multinucleated cells that were formed were smaller and harbored fewer nuclei than were observed in the control cells (Figure 2b). To examine whether adseverin controls osteoclast differentiation by regulating the proliferation of osteoclast precursors, a cell proliferation assay was performed. However, no significant difference was observed between adseverin-knockdown cells and control cells (Figure 2c). Although adseverin has been reported to exert an anticancer effect by inhibiting cell proliferation,¹³ adseverin does not appear to affect the proliferation of osteoclast precursors.

Next, we assessed actin ring formation. The actin ring is a typical morphological feature and a functionally important

cytoskeletal structure in mature osteoclasts. In control cells, actin rings clearly formed around the plasma membranes, and adseverin was primarily located in the subplasmalemmal region near the actin ring (Figure 2d). However, the actin belt was rarely observed in the adseverin-knockdown cells, and most of these cells had only one or two nuclei (Figure 2d). Because cell fusion is a process that is characteristic of osteoclastogenesis, we analyzed the fusion index by counting the number of nuclei per osteoclast. As shown in Figure 2e, adseverin knockdown led to a lower fusion index compared with the index observed in the control knockdown cells. Finally, we examined whether cell migration ability was affected by adseverin knockdown. Adseverin-knockdown cells showed a higher capacity to migrate compared with control cells (Figure 2f), suggesting that adseverin-knockdown cells might have less time to be in contact with neighboring cells for fusion. Taken together, these

results indicate that adseverin positively regulates osteoclast differentiation, fusion and actin ring formation without affecting proliferation.

Adseverin-knockdown osteoclasts display defects in resorption and enzyme secretion

Given the morphological defects that were observed in adseverin-knockdown osteoclasts, we next examined osteoclast function in these cells. To resorb bone, osteoclasts secrete enzymes such as TRAP, cathepsin K and MMPs. We predicted that resorption activity would also be impaired in the adseverin-knockdown osteoclasts because of their clearly defective multinucleation and actin ring formation. To test this hypothesis, BMMs were cultured on calcium phosphate-coated plates, which mimic the inorganic part of the bone. At day 7, the remaining calcium phosphate was stained using von Kossa reagents. This experiment revealed that adseverin-knockdown osteoclasts were defective in calcium phosphate resorption. Specifically, the knockdown cells exhibited ~80% less resorption compared with control cells (Figure 3a). Next, we examined enzyme secretion from osteoclasts using bone particles to mimic the resorption environment.¹⁴ The levels of TRAP and cathepsin K in the conditioned medium collected from control cells increased over time; however, conditioned medium from adseverin-knockdown osteoclasts showed lower levels of these enzymes (Figure 3b). Moreover, the amount of activated MMPs in the conditioned medium of adseverin-knockdown cells was ~63% lower compared with

the amount in the medium of control cells (Figure 3c). This finding suggests that adseverin regulates osteoclast function by modulating the secretion of bone-degrading enzymes.

NFATc1 expression and nuclear translocation are regulated by adseverin

NFATc1 is a key transcription factor during osteoclastogenesis, and its expression is strongly induced by RANKL.¹⁵ Furthermore, NFATc1 expression is autoamplified, and NFATc1 also mediates the expression of many osteoclast-specific genes.¹⁶ To investigate whether adseverin regulates NFATc1 expression during RANKL-induced osteoclastogenesis, we determined the expression level of NFATc1 in adseverin-knockdown BMMs. After RANKL stimulation, the protein level of NFATc1 was significantly lower in the knockdown cells compared with that in control cells (Figure 4a). c-Fos, which is a transcription factor required for the induction of NFATc1, regulates the transcription of osteoclast-specific genes through NFATc1.^{4,15} c-Fos was also decreased at day 1 (Figure 4a). Adseverin is therefore likely to regulate NFATc1 expression at least, in part, by modulating of c-Fos expression. Because NFATc1 autoregulates its own expression, a decrease in NFATc1 expression is expected to be accompanied by a decrease in NFATc1 nuclear translocation. Therefore, we assessed the nuclear translocation of NFATc1 following adseverin knockdown. As expected, both nuclear and cytoplasmic NFATc1 were decreased in adseverin-knockdown cells




Figure 3 Adseverin-knockdown osteoclasts display defects in resorption and enzyme secretion. **(a)** Bone marrow-derived macrophages (BMMs) were seeded on calcium phosphate-coated 48-well plates. Cells were transfected with 30 nm adseverin small interfering RNA (siRNA) and then cultured with macrophage-colony-stimulating factor (M-CSF) and receptor activator of nuclear factor- κ B ligand (RANKL) for 7 days. After the cells were washed, von Kossa staining was performed, and the resorption areas were analyzed. **(b)** After siRNA transfection, BMMs were cultured with M-CSF and RANKL for 2 days. The cells were further incubated in serum-free α -modified Eagle's medium (α -MEM) with bone particles obtained by grinding dentin. Supernatants were collected after 1, 6 and 12 h of incubation and analyzed using western blot analysis with anti-tartrate-resistant acid phosphatase (TRAP) and anti-cathepsin K antibodies. **(c)** As in **b**, supernatants were collected from cells cultured without bone particles for 12 h. Total matrix metalloproteinase (MMP) activity was measured using a commercial MMP Assay Kit. The data are presented as the means \pm s.d. ** $P < 0.005$.




Figure 4 Adseverin regulates nuclear factor of activated T cell c1 (NFATc1) expression and nuclear translocation. **(a)** Protein lysates were prepared from macrophage-colony-stimulating factor (M-CSF)/receptor activator of nuclear factor- κ B ligand (RANKL)-treated bone marrow-derived macrophages (BMMs) after small interfering RNA (siRNA) transfection. Western blot analysis was performed using antibodies for c-Fos, NFATc1 and adseverin. **(b)** After siRNA transfection, BMMs were cultured for 2 days in the presence of M-CSF and RANKL. Nuclear and cytoplasmic fractions were prepared and analyzed using western blot analysis. Lamin B and α -tubulin were used as controls for nuclear and cytoplasmic proteins, respectively. **(c)** BMMs were transfected with adseverin siRNA and then cultured with M-CSF and RANKL for 3 days. The expression levels of adseverin, NFATc1, tartrate-resistant acid phosphatase (TRAP), cathepsin K, dendritic cell-specific transmembrane protein (DC-STAMP) and v-ATPase were analyzed using real-time PCR. The data are presented as the means \pm s.d. ****** $P < 0.005$.

(Figure 4b). Moreover, the expression of osteoclast-specific genes that are directly induced by NFATc1, such as TRAP, cathepsin K, DC-STAMP and v-ATPase, were also markedly decreased during osteoclast differentiation in adseverin-deficient cells compared with control cells (Figure 4c). These results indicate that adseverin regulates osteoclast differentiation by modulating the expression of c-Fos and NFATc1, and, consequently, the expression of NFATc1-driven genes.

Adseverin regulates osteoclastogenesis via the NF- κ B signaling pathway and not the MAPK signaling pathway

To investigate the mechanisms by which adseverin regulates NFATc1 expression during RANKL-induced osteoclastogenesis, we examined the effects of adseverin knockdown on the MAPK signaling pathway. MAPKs have been reported to be activated by RANKL and to upregulate NFATc1 expression through AP-1. After elevating adseverin expression by culturing cells

with RANKL for 2 days, the activation of MAPK signaling proteins was assessed using western blot analysis with phospho-specific MAPK antibodies. RANKL-mediated activation of ERK, JNK and p38 was observed in both the adseverin-knockdown cells and the control cells at similar levels (Figure 5a). Next, Akt and NF- κ B activation were determined. There was no difference in RANKL-induced Akt phosphorylation between the adseverin-knockdown and the control cells (Figure 5b). The canonical NF- κ B signaling pathway, which involves the phosphorylation and degradation of I κ B and the subsequent nuclear translocation of NF- κ B, was activated by RANKL. There was no clear distinction between the adseverin-knockdown and control cells (Figure 5c). However, RANKL-induced phosphorylation of p65 NF- κ B was significantly reduced (Figure 5c). The nuclear translocation of p65 was also decreased in the adseverin-knockdown cells compared with the control cells (Figure 5d). NF- κ B is a key signaling molecule that




Figure 5 Adseverin regulates osteoclastogenesis via the nuclear factor- κ B (NF- κ B) signaling pathway but not the mitogen-activated protein kinase (MAPK) signaling pathway. (**a–c**) After small interfering RNA (siRNA) transfection, bone marrow-derived macrophages (BMMs) were cultured with 30 ng ml^{-1} macrophage-colony-stimulating factor (M-CSF) and 120 ng ml^{-1} receptor activator of nuclear factor- κ B ligand (RANKL) for 2 days. Cells were serum-starved for 4 h in α -modified Eagle's medium (α -MEM) and then stimulated with 600 ng ml^{-1} RANKL. Stimulated cells were harvested after 0, 5, 15 and 30 min, and protein lysates were then prepared. Western blot analysis was performed using antibodies against the phosphoforms or total forms of the indicated proteins. (**d**) Cells were stimulated with RANKL as in **a–c** for 0, 30 and 60 min. Nuclear and cytoplasmic fraction proteins were prepared and subjected to western blot analysis.

is activated by RANKL that is known to regulate NFATc1 expression in osteoclastogenesis.^{4,16} Therefore, these data demonstrate that adseverin has a role in RANKL-induced osteoclastogenesis by regulating the noncanonical NF- κ B signaling pathway but not the MAPK signaling pathway.

DISCUSSION

Ca^{2+} signaling is one of the main intracellular events that is involved in osteoclast differentiation. Adseverin is a Ca^{2+} -regulated protein that has been shown to have Ca^{2+} -dependent actin-severing activity. Based on these observations, we hypothesized that adseverin has a role in osteoclast differentiation. We found that adseverin expression was increased during osteoclast differentiation (Figure 1). Furthermore, our results provide evidence showing that adseverin controls osteoclast differentiation by regulating NFATc1 expression and NF- κ B signaling (Figures 4 and 5).

NF- κ B has been demonstrated to have a crucial role in osteoclastogenesis. The p50 and p65 NF- κ B subunits are recruited to the NFATc1 promoter in response to RANKL stimulation.¹⁶ In addition, dehydroxymethylepoxyquinomicin, an NF- κ B inhibitor, has been shown to attenuate RANKL-induced osteoclastogenesis by downregulating NFATc1.¹⁷ Moreover, NF- κ B1 (p50) and NF- κ B2 (p52) double knockout mice display a severe osteopetrotic phenotype.¹⁸ Therefore, NF- κ B has a crucial role in osteoclastogenesis by mediating NFATc1 expression. We observed that p65 phosphorylation was reduced following adseverin knockdown (Figure 5c). Because p65 phosphorylation enhances the transactivation potential of NF- κ B,^{19,20} it is likely that adseverin regulates

NF- κ B transcription activity during osteoclastogenesis. In addition, we found that the nuclear translocation of p65 was regulated by adseverin (Figure 5d). However, there was no difference indicating its involvement in I κ B degradation (Figure 5c), which excludes the possibility of adseverin-mediated regulation of the canonical NF- κ B pathway. Sasaki *et al.*²¹ reported a noncanonical NF- κ B pathway in which the nuclear translocation of phosphorylated p65 was independent of regulation by I κ B. Taken together, these data indicate that adseverin regulates p65 phosphorylation and nuclear translocation in a noncanonical manner during RANKL-induced osteoclast differentiation.

Adseverin is known to be expressed in most secretory cells and is believed to have a role in actin cytoskeletal rearrangements during exocytosis. In osteoclasts, numerous vesicles containing lysosomal enzymes are transported to the cell surface, and their contents are released by exocytosis.²² These released enzymes degrade the organic components of the bone matrix. Consistent with this role of adseverin in exocytosis, we observed that adseverin knockdown reduced the secretion of bone-degrading enzymes by osteoclasts (Figure 3). This reduction in secretion is likely due to the effect of adseverin knockdown on actin cytoskeleton dynamics, in addition to the downregulation of NFATc1 levels, which reduces the expression of these enzymes.

Adseverin and gelsolin share a common function, the severing of actin filaments, and they also have high homology. Consistent with a report showing that gelsolin-knockout mice have high bone mass and display defects in osteoclast podosome assembly,¹² we found that adseverin-knockdown

osteoclasts possessed numerous defects in their differentiation and function (Figures 2 and 3). Although adseverin and gelsolin have some common features, they also have a number of different properties. Importantly, adseverin and gelsolin have different Ca^{2+} -binding affinities.⁷ This finding may explain why adseverin is more specialized for exocytosis and why it is expressed in fewer tissues than gelsolin. Therefore, it is possible that gelsolin and adseverin regulate bone metabolism via different mechanisms. Gelsolin is known to regulate bone mass by affecting osteoclast podosome assembly, whereas adseverin may also regulate osteoclasts by acting as a signal transducer during the early stage of differentiation. In support of this hypothesis, we observed that NFATc1 expression was decreased at one day post-RANKL treatment (Figure 4).

Recently, actin as demonstrated to be present in both the nucleus and the cytoplasm. Furthermore, the functions of several nuclear actin-binding proteins have been identified. Many members of the gelsolin superfamily, such as flightless I, supervillin and gelsolin, which are present in the nucleus, are coactivators of nuclear receptors. Supervillin promotes the transactivation of the androgen receptor,^{23,24} whereas gelsolin facilitates the nuclear translocation of the androgen receptor by acting as a coregulator.²⁵ In addition, mounting evidence supports a role for gelsolin as an upstream factor in many signaling cascades. For example, gelsolin cleaves caspase-3, the products of which are effectors of apoptosis.²⁶ Moreover, gelsolin has been shown to regulate collagen phagocytosis via Rac-dependent control of the cytoskeleton.²⁷ Although the potential roles of adseverin as a signal transducer or transcriptional regulator have not yet been clearly defined, some reports have concluded that adseverin affects signal-transduction pathways either directly or indirectly via cytoskeletal rearrangements and that adseverin regulates the differentiation of chondrocytes and megakaryoblastic leukemia cells.^{28,29} These reports, along with data from other gelsolins, raise the possibility that adseverin may have additional functions beyond its well-known role in regulating actin polymerization in the cytoplasm.

The data presented here demonstrate that adseverin has a role in osteoclastogenesis. Specifically, adseverin regulates NFATc1 expression via NF- κ B signaling, which is necessary for effective osteoclast differentiation and resorption. This study is the first to demonstrate a role for adseverin beyond its role in controlling the actin cytoskeleton. Our data thus identify a novel role for adseverin in signal transduction and osteoclastogenesis.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

This work was supported by a grant from the National Research Foundation of Korea (NRF-2014R1A2A1A10050406) to H-HK.

- Hayashi S, Yamane T, Miyamoto A, Hemmi H, Tagaya H, Tanio Y *et al*. Commitment and differentiation of stem cells to the osteoclast lineage. *Biochem Cell Biol* 1998; **76**: 911–922.
- Valledor AF, Borrás FE, Culléll-Young M, Celada A. Transcription factors that regulate monocyte/macrophage differentiation. *J Leukoc Biol* 1998; **63**: 405–417.
- Yasuda H, Shima N, Nakagawa N, Yamaguchi K, Kinoshita M, Mochizuki S *et al*. Osteoclast differentiation factor is a ligand for osteoprotegerin/osteoclastogenesis-inhibitory factor and is identical to TRANCE/RANKL. *Proc Natl Acad Sci USA* 1998; **95**: 3597–3602.
- Kim JH, Kim N. Regulation of NFATc1 in osteoclast differentiation. *J Bone Metab* 2014; **21**: 233–241.
- Kwiatkowski DJ. Functions of gelsolin: motility, signaling, apoptosis, cancer. *Curr Opin Cell Biol* 1999; **11**: 103–108.
- Silacci P, Mazzolai L, Gauci C, Stergiopoulos N, Yin HL, Hayoz D. Gelsolin superfamily proteins: key regulators of cellular functions. *Cell Mol Life Sci* 2004; **61**: 2614–2623.
- Rodriguez Del Castillo A, Lemaire S, Tchakarof L, Jeyapragasan M, Doucet JP, Vitale ML *et al*. Chromaffin cell scinderin, a novel calcium-dependent actin filament-severing protein. *EMBO J* 1990; **9**: 43–52.
- Tchakarof L, Vitale ML, Jeyapragasan M, Rodriguez Del Castillo A, Trifaro JM. Expression of scinderin, an actin filament-severing protein, in different tissues. *FEBS Lett* 1990; **268**: 209–212.
- Chumnarnsilpa S, Lee WL, Nag S, Kannan B, Larsson M, Burtnick LD *et al*. The crystal structure of the C-terminus of adseverin reveals the actin-binding interface. *Proc Natl Acad Sci USA* 2009; **106**: 13719–13724.
- Maekawa S, Sakai H. Inhibition of actin regulatory activity of the 74-kDa protein from bovine adrenal medulla (adseverin) by some phospholipids. *J Biol Chem* 1990; **265**: 10940–10942.
- Zhang L, Marcu MG, Nau-Staudt K, Trifaro JM. Recombinant scinderin enhances exocytosis, an effect blocked by two scinderin-derived actin-binding peptides and PIP2. *Neuron* 1996; **17**: 287–296.
- Chellaiyah M, Kizer N, Silva M, Alvarez U, Kwiatkowski D, Hruska KA. Gelsolin deficiency blocks podosome assembly and produces increased bone mass and strength. *J Cell Biol* 2000; **148**: 665–678.
- Wang D, Sun SQ, Yu YH, Wu WZ, Yang SL, Tan JM. Suppression of SCIN inhibits human prostate cancer cell proliferation and induces G0/G1 phase arrest. *Int J Oncol* 2014; **44**: 161–166.
- Cremasco V, Decker CE, Stumpo D, Blackshear PJ, Nakayama KI, Nakayama K *et al*. Protein kinase C-delta deficiency perturbs bone homeostasis by selective uncoupling of cathepsin K secretion and ruffled border formation in osteoclasts. *J Bone Miner Res* 2012; **27**: 2452–2463.
- Takayanagi H, Kim S, Koga T, Nishina H, Isshiki M, Yoshida H *et al*. Induction and activation of the transcription factor NFATc1 (NFAT2) integrate RANKL signaling in terminal differentiation of osteoclasts. *Dev Cell* 2002; **3**: 889–901.
- Asagiri M, Sato K, Usami T, Ochi S, Nishina H, Yoshida H *et al*. Autoamplification of NFATc1 expression determines its essential role in bone homeostasis. *J Exp Med* 2005; **202**: 1261–1269.
- Takatsuna H, Asagiri M, Kubota T, Oka K, Osada T, Sugiyama C *et al*. Inhibition of RANKL-induced osteoclastogenesis by (-)-DHMEQ, a novel NF- κ B inhibitor, through downregulation of NFATc1. *J Bone Miner Res* 2005; **20**: 653–662.
- Itsova V, Caamano J, Loy J, Yang Y, Lewin A, Bravo R. Osteopetrosis in mice lacking NF- κ B1 and NF- κ B2. *Nat Med* 1997; **3**: 1285–1289.
- Sakurai H, Chiba H, Miyoshi H, Sugita T, Toriumi W. I κ B kinases phosphorylate NF- κ B p65 subunit on serine 536 in the transactivation domain. *J Biol Chem* 1999; **274**: 30353–30356.
- Doyle SL, Jefferies CA, O'Neill LA. Bruton's tyrosine kinase is involved in p65-mediated transactivation and phosphorylation of p65 on serine 536 during NF- κ B activation by lipopolysaccharide. *J Biol Chem* 2005; **280**: 23496–23501.
- Sasaki CY, Barberi TJ, Ghosh P, Longo DL. Phosphorylation of RelA/p65 on serine 536 defines an I κ B-independent NF- κ B pathway. *J Biol Chem* 2005; **280**: 34538–34547.
- Vaananen HK, Zhao H, Mulari M, Halleen JM. The cell biology of osteoclast function. *J Cell Sci* 2000; **113**(Pt 3): 377–381.
- Sampson ER, Yeh SY, Chang HC, Tsai MY, Wang X, Ting HJ *et al*. Identification and characterization of androgen receptor associated coregulators in prostate cancer cells. *J Biol Regul Homeost Agents* 2001; **15**: 123–129.

- 24 Ting HJ, Yeh S, Nishimura K, Chang C. Supervillin associates with androgen receptor and modulates its transcriptional activity. *Proc Natl Acad Sci USA* 2002; **99**: 661–666.
- 25 Nishimura K, Ting HJ, Harada Y, Tokizane T, Nonomura N, Kang HY *et al*. Modulation of androgen receptor transactivation by gelsolin: a newly identified androgen receptor coregulator. *Cancer Res* 2003; **63**: 4888–4894.
- 26 Geng YJ, Azuma T, Tang JX, Hartwig JH, Muszynski M, Wu Q *et al*. Caspase-3-induced gelsolin fragmentation contributes to actin cytoskeletal collapse, nucleolysis, and apoptosis of vascular smooth muscle cells exposed to proinflammatory cytokines. *Eur J Cell Biol* 1998; **77**: 294–302.
- 27 Arora PD, Glogauer M, Kapus A, Kwiatkowski DJ, McCulloch CA. Gelsolin mediates collagen phagocytosis through a rac-dependent step. *Mol Biol Cell* 2004; **15**: 588–599.
- 28 Nurminsky D, Magee C, Faverman L, Nurminskaya M. Regulation of chondrocyte differentiation by actin-severing protein adseverin. *Dev Biol* 2007; **302**: 427–437.
- 29 Zunino R, Li Q, Rose SD, Romero-Benitez MM, Lejen T, Brandan NC *et al*. Expression of scinderin in megakaryoblastic leukemia cells induces differentiation, maturation, and apoptosis with release of plateletlike particles and inhibits proliferation and tumorigenesis. *Blood* 2001; **98**: 2210–2219.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>