RESEARCH

BMC Public Health



Socio-demographic disparities in basic undertwo immunization coverage: insights from the 2016 Malawi demographic and health survey



Ololade Julius Baruwa^{1*}, Monica Ewomazino Akokuwebe², Oluwafemi John Adeleye³ and Babatunde Makinde Gbadebo³

Abstract

Background Childhood immunization is a vital component of public health, preventing the spread of infectious diseases and reducing child mortality. This study examines variations in basic immunization coverage across districts and explores socio-demographic disparities in immunization coverage among children aged 12–23 months in Malawi.

Methods The study employed a cross-sectional design, utilizing data from the 2016 MDHS, a nationally representative survey. The analysis included 3,248 children aged 12–23 months. Socio-demographic variables, including the child's sex, maternal age, marital status (currently married), education, place of residence, region, wealth status, and employment status, were analysed using multivariate logistic regression models and a choropleth map to assess variations in basic immunization coverage across all 28 districts in Malawi.

Results The results showed that only 77.1% of children aged 12–23 months received basic immunization. The findings also highlighted significant variations in immunization coverage across different socio-demographic groups and among the 28 districts of Malawi. The highest coverage rates, ranging from 84.9 to 90.7%, were observed in Mwanza and Chiradzulu districts, while the lowest rates, between 65.3% and 68.0%, were found in Ntchisi, Blantyre, and Machinga districts. Multivariable analysis further indicated that children whose mothers were from a high household wealth index (OR = 1.45, 95% CI = 1.15–1.82), residing in rural areas (OR = 1.55, 95% CI = 1.20–2.01), currently married (OR = 1.33, 95% CI = 1.10–1.61), and with secondary or higher education (OR = 1.57, 95% CI = 1.11–2.21) were more likely to receive basic immunization.

Conclusion The study highlights low coverage of basic immunization in some districts as well as socio-demographic disparities in under-two immunization coverage in Malawi, necessitating tailored interventions such as educational campaigns and region-specific strategies.

Keywords Immunization, Vaccination, Malawi, Socio-demographic, Demographic and health survey

*Correspondence: Ololade Julius Baruwa baruwaololade@gmail.com ¹Centre for Social Sciences Research, University of Cape Town, Cape Town, Rondebosch, South Africa

²SAMRC Developmental Pathways for Health Research Unit, Department of Paediatrics, School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa ³Department of Epidemiology and Medical Statistics, Faculty of Public Health, College of Medicine, University of Ibadan, Ibadan, Nigeria

© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

Introduction

Immunization for children under two years old (0–23 months) has significantly improved childhood survival and health outcomes, preventing millions of deaths globally [1, 2]. It serves as a vital intervention for disease prevention and an effective strategy for controlling infectious diseases among newborns and older infants [3, 4]. However, over the past decade, childhood vaccination coverage has stagnated, with over 14 million children worldwide either unvaccinated or under-vaccinated [2–4]. In response, the World Health Organization (WHO) developed the Immunization Agenda 2030 (IA2030) as a global strategy to enhance vaccine coverage and ensure that no one is left behind.

Vaccination coverage among newborns and older infants is unevenly distributed across settings and populations [1, 2]. Measuring socio-demographic inequalities is critical for identifying and addressing gaps in vaccine coverage that affect specific sub-populations. While childhood immunization coverage has increased globally and regionally since 1980, disparities persist, with high-income countries achieving substantially higher and more equitable coverage rates than low-income counterparts [5–7]. Reducing these inequalities is crucial for achieving the United Nations Sustainable Development Goals (SDGs).

Studies show that 60% of the world's unvaccinated children reside in low- or middle-income countries, including Nigeria, India, Pakistan, Indonesia, Ethiopia, the Philippines, the Democratic Republic of Congo, Brazil, Angola, and Vietnam [4, 8]. Socioeconomic status, urban versus rural residence, maternal education, and other demographic characteristics significantly influence vaccination coverage, particularly in low- and middle-income countries [9, 10]. Marginalized communities remain under-vaccinated and disproportionately unvaccinated [10]. Early childhood disparities in immunization can have long-term effects on health outcomes, making it imperative to address these gaps.

In Malawi, the Expanded Programme on Immunization (EPI) was established in 1979, introducing vaccines such as Bacillus Calmette-Guérin (BCG) for tuberculosis, diphtheria, tetanus, and pertussis (DTP), oral polio vaccine (OPV), and measles-containing vaccine (MCV) [11, 12]. Despite these efforts, only 73% of children aged 12–23 months were fully vaccinated in 2016, falling short of the Malawian government's goal of over 80% coverage [13]. However, recent data on whether significant improvements have been made remain limited.

Several studies have cited poor access to immunization services, the need to reach vulnerable pregnant and nursing women in rural communities, and inadequate supervision of routine immunization activities as key barriers to vaccine uptake [14–17]. Additionally, the COVID-19

pandemic further disrupted routine immunization and sensitization campaigns, hindering progress toward IA2030 goals [18–20]. A sustained and comprehensive intervention is urgently needed to ensure equitable, effective, and resilient immunization services in Malawi.

To enhance vaccination uptake and implement effective intervention strategies, it is crucial to identify both enablers and barriers to immunization coverage. Numerous studies have highlighted individual, household, and community-level determinants, including maternal age, education, household income, prenatal care attendance, skilled birth attendance, place of residence, geographical setting, awareness of immunization benefits, proximity to immunization clinics, cultural factors, and parental awareness of children's vaccinations [21–25]. However, most studies have focused on either full or incomplete immunization, and no study has comprehensively examined variations in basic immunization coverage across Malawi's districts or socio-demographic disparities in immunization among children aged 12–23 months.

This study aims to analyze socio-demographic disparities in basic immunization coverage among children aged 12–23 months in Malawi. Specifically, it investigates variations in coverage across districts and explores disparities based on socio-demographic factors. The findings will provide valuable insights for policymakers and health professionals, supporting the development of targeted interventions to improve childhood vaccination uptake and overall immunization coverage in Malawi.

Materials and methods

Study setting

Malawi is a low-income country with a population of approximately 20.4 million people, nearly 85% of whom reside in rural areas [4, 7]. Childhood vaccination is administered through local health facilities, either at fixed clinics or mobile outreach programs conducted by Health Surveillance Assistants (HSAs), who provide coverage for every village in Malawi [2, 11]. Vaccination services are free of charge. However, Malawi faces a significant shortage of healthcare personnel, often resulting in one health worker attending to a large number of infants seeking vaccination. This shortage can lead to inadequate documentation and follow-up, particularly in rural areas, where vaccines may not be administered on time or at all. Additionally, major stock-outs of Bacillus Calmette-Guérin (BCG) and diphtheria, pertussis, and tetanus (DPT) vaccines were reported at the central level in 2007 [26, 27]. Despite rapid urbanization, Malawi remains largely rural, with nearly 85% of the population still living in rural areas [28]. In Malawi, urban areas are classified as the four major metropolitan cities - Lilongwe, Blantyre, Mzuzu, and Zomba-along

with secondary cities, including townships and district centers. All other areas in the country are designated as rural.

Data source

The data utilized in this research was sourced from the children (KR file) dataset of the 2015-16 Malawi Demographic and Health Survey (MDHS) [27]. The MDHS 2015/16 was implemented by the National Statistics Office (NSO) and the Community Health Science Unit (CHSU). The government of Malawi, the National AIDS Commission (NAC), the United Nations Population Fund (UNFPA), the United Nations Children's Fund (UNICEF), and the United Kingdom's Department for International Development (DFID), the Centre for Disease Control and Prevention (CDC), and the United States Agency for International Development (USAID) provided the funds. Technical assistance on the survey was provided by ICF Macro through the MEASURE DHS program, a USAIDfunded project. Using a two-stage cluster sampling technique based on enumeration areas defined in Malawi's 2008 census, the survey gathered data on 17,395 children under the age of five. After thorough data wrangling and cleaning, the final analysis for this study was conducted on a subset of 3,248 children aged 12-23 months. Detailed information on sample size estimation and sampling strategies is accessible in the complete MDHS published reports at https://www.dhsprogram.com/.

Study outcome

The outcome variable in this study is basic child immunization. A child is considered basic immunization if they have received all the recommended vaccinations, according to the vaccination policy framework in Malawi [27]. Specifically, basic immunization includes one dose of BCG, three doses of DPT-HepB-Hib, three doses of oral polio vaccine (excluding the birth dose), two doses of rotavirus vaccine, three doses of pneumococcal vaccine, and one dose of measles vaccine [27]. Children who received all the recommended vaccinations were coded as "1" and categorized as "yes," while those who missed at least one recommended vaccine were coded as "0" and categorized as "no."

Explanatory variables

In this study, socio-demographic factors were considered as the explanatory variables. These factors were considered because they had a statistically significant association with basic child immunization in previous studies [11, 12, 14, 29]. The explanatory variables include sex of the child (male, female), maternal age [15–34], maternal education (no education, primary, and secondary or higher), mother is currently married (no and yes), place of residence (urban and rural), region (northern, central and southern), maternal wealth status (poor, middle and rich), and maternal employment status (working and not working).

Statistical analyses

This analysis includes all relevant factors to control for potential confounding, with variable selection guided by the literature for their theoretical and policy relevance. This approach ensures both rigorous control and alignment with existing research. To explore the variations in basic immunization coverage by districts and to investigate the socio-demographic disparities in basic immunization coverage among 12-to-23-month-old children in Malawi, we followed four analytic steps. In the first step, we used descriptive statistical methods to describe the sample characteristics of respondents (frequency and percentages) (Table 1). The second step involved conducting a bivariate analysis, examining the cross-tabulation of all explanatory variables against the outcome variable using the Chi-square test (Table 1).

In the third step, a spatial mapping analysis was conducted using ArcGIS Pro 2.8 software to determine district-specific proportions of basic immunization coverage in Malawi (Fig. 1). The spatial distribution of immunization prevalence was visualized through coloured density polygons, which depicted coverage by district. Spatial and spatiotemporal mapping methodologies are commonly used in public health research for their effectiveness in illustrating the distribution of health indicators, facility locations, and geographical accessibility. These methods also provide valuable insights into underserved areas and regions lacking sufficient healthcare interventions [26, 27, 30, 31]. For this analysis, the graduated symbol properties in ArcGIS were configured using a CSV file that contained the proportions of basic immunization coverage across Malawi's districts. This data was integrated with the attribute table of the base map (shapefile), which included the x and y coordinates of each district's polygon grid within Malawi's geographical framework. The resulting analysis produced graduated colour maps representing basic immunization coverage for each district. Additionally, a coloured legend indicating varying proportions of immunization coverage was generated to enhance interpretability.

The fourth step involved using binary logistic regression to examine socio-demographic disparities in basic immunization coverage among children aged 12–23 months (Table 2). The analysis was performed in two stages (models). The first model assessed the independent relationship of each socio-demographic factor with immunization coverage, calculating unadjusted odds ratios (uOR) for each variable. The second model incorporated all the socio-demographic factors simultaneously to assess their combined influence on immunization

	Total sample	No Basic Immunization	Basic Immunization	Chi-square (χ²) ρ-value
Variables	N = 3248	22.9 (743)	77.1 (2505)	
Sex of Child	% (n)	% (n)	% (n)	
Male	49.9 (1622)	24.2 (393)	75.8 (1229)	0.067
Female	50.1 (1626)	21.5 (350)	78.5 (1276)	
Maternal age				
15–19	10.4 (337)	27.3 (92)	72.7 (245)	0.115
20-24	31.9 (1036)	23.5 (243)	76.5 (793)	
25–29	22.7 (738)	21.0 (155)	79.0 (583)	
30-34	18.3 (595)	20.7 (123)	79.3 (472)	
35 and above	16.7 (542)	24.0 (130)	76.0 (412)	
Education				
No education	10.9 (355)	25.9 (92)	74.1 (263)	< 0.001*
Primary	68.2 (2216)	24.1 (534)	75.9 (1682)	
Secondary/Higher	20.8 (677)	17.3 (117)	82.7 (560)	
Wealth Status				
Poor	45.9 (1491)	25.2 (375)	74.9 (1116)	< 0.001*
Middle	19.8 (643)	24.1 (155)	75.9 (488)	
Rich	34.3 (1114)	19.1 (213)	80.9 (901)	
Place of Residence				
Urban	15.6 (508)	24.8 (126)	75.2 (382)	0.260
Rural	84.4 (2740)	22.5 (617)	77.5 (2123)	
Region				
Northern region	18.6 (603)	20.4 (123)	79.6 (480)	0.274
Central region	35.3 (1147)	23.5 (270)	76.5 (877)	
Southern region	46.1 (1498)	23.4 (350)	76.6 (1148)	
Currently Married				
No	22.6 (733)	27.2 (199)	72.9 (534)	< 0.001*
Yes	77.4 (2515)	21.6 (544)	78.4 (1971)	
Employment Status				
Not Working	35.1 (1141)	24.4 (278)	75.6 (863)	0.137
Working	64.9 (2107)	22.1 (465)	77.9 (1642)	

Table 1 Study participants' characteristics and outcome and its Chi-Square association

Source: Authors' compilation; * - Significant at 5%

coverage (adjusted odds ratios [aOR]). The binary logistic regression model was applied to investigate the predictive relationship between the dichotomous dependent variable, using the value 0 or 1 (i.e. failure or success), and a set of explanatory variables. Therefore, the binary logistic regression model is expressed as [32, 33]:

$$Log (\rho / 1 - \rho) = a + \Sigma \beta i Xi + \mathfrak{e},$$

where ρ is the event of occurrence (outcome variable), Xi is the explanatory variables, β i is the size of the coefficient of explanatory variables, and \mathfrak{e} is the margin of error. For model assessment, the log-likelihood ratio (LLR) was employed and sample weight was applied to address related concerns of over-and under-sampling. However, the SVY command was used to account for the complex survey design and generalizability of the findings. The statistical significance for all tests was set at $\rho < 0.05$. The data were analysed with STATA version 15.4 for MacOs, and three steps were pursued to analyze the data. All variables were weighted to guarantee the representativeness of the survey data and account for the complex sampling strategy used in the survey.

Results

Socio-demographic characteristics of the study population Table 1 presents the characteristics of the study population. Findings revealed that 77.1% of children aged 12–23 months had received the basic immunization. About 50% (49.9%) of the children are male, and 31.9% are born to mothers who are between the ages of 20–24 years. The table further shows that 68.2% of the mothers have primary education, 45.9% are from poor households, 84.4% live in rural areas, 46.1% are from the Southern region, 46.1% are from the Southern region, 77.4% are currently married, and 64.9% are employed (Table 1).



Fig. 1 Choropleth map showing distribution of basic immunization coverage across districts in Malawi

Prevalence and bivariate associations of basic immunization coverage among children aged 12–23 months by socio-demographic factors

Table 1 also presents the bivariate analysis. The bivariate analysis showed that education, wealth status, and marriage status have significant relationships with immunization coverage among children aged 12–23 months. Basic immunization coverage was higher among female children (78.5%) than male children (75.8%). Basic immunization coverage was also more prevalent among children born to mothers aged 30–34 years (79.3%) compared to those born to mothers aged 15–19 years (72.7%). The findings revealed that coverage of basic immunization was more pronounced among children whose mothers had secondary or higher education (82.7%) compared to those whose mothers had no formal education (74.1%).

The prevalence of basic immunization coverage was higher (80.9%) among children whose mothers are from rich households compared to those of poor and middle-wealth status (74.9% and 75.9%, respectively). Further, the prevalence of basic immunization was 77.5% among children whose mothers resided in rural areas, 79.6% among children whose mothers lived in the northern region, 78.4% among children whose mothers were currently

married, and 77.9% among children whose mothers were working (Table 1).

Choropleth map showing the basic immunization coverage among children aged 12–23 months across districts in Malawi

The distribution of basic immunization coverage across the 28 districts in Malawi is illustrated in Fig. 1 below. *High coverage* (\geq 90%) was observed in Mwanza and Chiradzulu districts, with proportions ranging from 84.9 to 90.7%. *Moderate coverage* (80-89%) was found in the districts of Karonga, Rumphi, Chitipa, Mchinji, Dowa, Salima, Thyolo, Neno, and Phalombe, with coverage ranging from 78.1 to 84.8%. *Low coverage* (<80%) was recorded in Ntchisi, Blantyre, and Machinga districts, with proportions ranging from 65.3 to 68.0% (Fig. 1). This reclassification follows the WHO recommendations on immunization coverage, which categorizes high coverage as \geq 90%, moderate coverage as 80-89%, and low coverage as <80% [35].

Table 2 Logistic regression model showing the association
between socio-demographic factors and under-two (12-23
months) basic immunization coverage in Malawi

X • • • •	00 (050) (01)	
Variables	uOR (95% C.I)	aOR (95% C.I)
Sex of Child		
Male	1	1
Female	1.17 (0.99–1.37)	1.15 (0.98–1.36)
Maternal age		
15–19	1	1
20–24	1.23 (0.93–1.62)	1.09 (0.82–1.45)
25–29	1.41 (1.05–1.9)**	1.21 (0.89–1.65)
30–34	1.44 (1.06–1.97)**	1.27 (0.93–1.75)
35 and above	1.19 (0.87–1.62)	1.09 (0.78–1.51)
Education		
No education	1	1
Primary	1.1 (0.85–1.42)	1.08 (0.83-1.42)
Secondary/Higher	1.67 (1.23–2.28)***	1.57 (1.11–2.21)**
Wealth Status		
Poor	1	1
Middle	1.06 (0.85–1.31)	1.02 (0.82–1.27)
Rich	1.42 (1.18–1.72)***	1.45 (1.15–1.82)***
Place of Residence		
Urban	1	1
Rural	1.14 (0.91–1.42)	1.55 (1.2–2.01)***
Region		
Northern region	1	1
Central region	0.83 (0.65–1.06)	0.93 (0.72–1.19)
Southern region	0.84 (0.67-1.06)	0.96 (0.76–1.22)
Currently Married		
No	1	1
Yes	1.35 (1.12–1.63)***	1.33 (1.10–1.61)***
Employment Status		
Not Working	1	1
Working	1.14 (0.96–1.35)	1.13 (0.95–1.35)

* p<0.1, ** p<0.05, *** p<0.01; uOR – Unadjusted Odds Ratio; aOR – Adjusted Odds Ratio; C.I – Confidence Interval

Multivariable analysis of factors associated with basic immunization coverage among children aged 12–23 months

Table 2 presents the odds ratio and confidence interval of the relationship between socio-demographic factors and basic immunization coverage among children aged 12–23 months in Malawi. The multivariable analysis showed that mothers' educational level, household wealth status, and marital status (currently married) are protective of basic immunization coverage in the unadjusted and adjusted analyses. Maternal age showed a significant association with basic immunization coverage in the unadjusted analysis but not in the adjusted, while place of residence showed a significant association in the adjusted analysis but in the unadjusted. The multivariate analysis showed that the odds of basic immunization coverage among children aged 12–23 months increase with the level of the mother's education. For instance, the adjusted logistic regression analysis showed that the likelihood of basic immunization coverage is 1.57 times higher among children whose mothers had secondary or higher education (aOR = 1.57; 95%CI = 1.11-2.21) compared to those with no formal education. Likewise, the odds of basic immunization coverage were 1.45 times higher among children born to mothers from rich households (aOR = 1.45; 95%CI = 1.15 - 1.82) compared to their counterparts from poor households (Table 2). The likelihood of basic immunization coverage was 1.55 times higher among children whose mothers reside in a rural area (aOR = 1.55; 95%CI = 1.20-2.01) compared to those whose mothers reside in an urban area. The odds of basic immunization coverage were 1.33 times higher among children whose mothers were currently married (aOR = 1.33; 95%CI = 1.10-1.61) compared to those whose mothers were currently unmarried (Table 2).

Discussion

This study explores socio-demographic disparities in basic immunization coverage among children aged 12-23 months in Malawi. Our findings indicate that only 77.1% of these children had received basic immunization, closely aligning with a previous study that reported 76% coverage in the same age group [28]. This coverage rate falls short of the World Health Organization's (WHO) recommended 90% immunization coverage benchmark, highlighting a significant gap in achieving optimal vaccination rates for this vulnerable population. The low immunization rates pose a potential public health risk, underscoring the need for targeted interventions and awareness campaigns. Healthcare policymakers and practitioners must collaborate to develop strategies that enhance immunization coverage for infants and toddlers, thereby safeguarding their health and contributing to overall community well-being.

Our study reveals significant disparities in basic immunization coverage across Malawi's 28 districts. Mwanza and Chiradzulu recorded the highest immunization rates, while Ntchisi, Blantyre, and Machinga exhibited the lowest. These variations highlight the importance of exploring district-specific factors that influence immunization uptake. The high coverage observed in Mwanza and Chiradzulu may be attributed to several factors, including proximity to healthcare facilities, which plays a crucial role in vaccine accessibility. Residing closer to vaccination services has been linked to increased immunization rates, whereas longer travel distances often hinder access and reduce coverage. Conversely, districts such as Ntchisi, Blantyre, and Machinga, which reported lower immunization rates, may face challenges such as limited access to healthcare services, lower socioeconomic conditions, and inadequate health education [36]. Addressing these barriers is essential for improving immunization coverage.

To enhance immunization rates in underperforming districts, targeted interventions are required. These may include strengthening healthcare infrastructure to minimize travel distances and addressing socioeconomic barriers that restrict healthcare access. By drawing lessons from districts with high coverage and implementing localized strategies, Malawi can work toward achieving equitable immunization coverage across all regions.

Our multivariable analysis found a significant association between maternal age and basic immunization coverage. Specifically, the likelihood of completing basic immunization schedules increased with maternal age, a finding that aligns with previous studies [28, 34]. Older mothers may have socio-economic advantages and better health literacy, enabling them to engage more effectively with healthcare systems. These advantages include a greater understanding of immunization's importance, increased autonomy in decision-making, and improved access to healthcare facilities.

Consistent with previous studies [28, 34], our findings indicate that children of educated mothers are more likely than those of non-educated mothers to receive the basic under-two immunization. The possible explanation is that maternal education enhance health literacy, allowing mothers to recognize the importance of immunizing their children [14, 37]. Educated mothers are also more capable of navigating healthcare systems, ensuring they attend immunization clinics on time and adhere to vaccination schedules. Furthermore, through exposure to health education programmes and media, educated mothers may possess greater knowledge about preventive healthcare practices, such as immunization. A positive attitude toward vaccination and a proactive approach to making sure their children receive their vaccinations on time can be attributed to this increased awareness [38].

An interesting finding of this study is that children born to mothers in rural settings have a higher likelihood of receiving basic under-two immunization compared to those in urban areas. This finding contradicts previous studies [37, 40]. However, this observed pattern could be the result of several factors. First, due to the possible lack of access to centralized healthcare facilities, communitybased healthcare interventions, such as immunization programmes, may be given more importance in rural areas. Hence, community networks that promote collective participation in preventive health practices and facilitate the dissemination of health information may be beneficial to mothers living in rural areas. Second, people living in rural areas may have a greater understanding of the value of immunization as a preventative measure against common diseases that are common in these areas, as these populations frequently face unique health challenges [39–41]. Third, rural communities' geographic dispersion may also lead to focused outreach initiatives that bring healthcare services, such as immunization, closer to families. However, it is important to consider that Malawi's population is predominantly rural, which may influence this association.

The multivariable analysis indicating that children born to mothers from rich households were more likely to receive basic under-two immunization aligns with a previous study [40, 41]. Higher socioeconomic status, as reflected in the rich household wealth index, often provides families with the financial capacity to cover transportation costs, purchase health insurance, and manage other expenses related to healthcare services, including immunization. Additionally, economic stability in richer households may contribute to better health literacy, heightened awareness of the importance of immunization, and a stronger ability to prioritize healthcare needs. Rich households are more likely to have access to information, resources, and opportunities that enable them to utilize health services more effectively, ensuring their children receive timely vaccinations.

Finally, the finding that the odds of basic immunization coverage were higher among children whose mothers were currently married compared to those whose mothers were unmarried suggests that marital status plays a critical role in healthcare-seeking behaviour and child health outcomes, including immunization. Married mothers may benefit from greater social and emotional support from their spouses, family members, or the broader community [24, 41]. This support can make it easier to prioritize healthcare needs like immunization, especially in low-resource settings where logistical and financial barriers exist. Spouses may assist with household responsibilities, childcare, or transport to healthcare facilities, thereby reducing the burden on mothers and enabling them to adhere to vaccination schedules.

Strengths and limitations

Our study highlights socio-demographic disparities in basic immunization coverage among children aged 12-23 months in Malawi, utilizing nationally representative data from the 2016 Demographic and Health Survey (DHS). While our findings provide valuable insights into factors influencing immunization coverage, it is essential to acknowledge the study's limitations. First, recall bias may affect the accuracy of immunization data, as mothers might not always accurately recall their children's vaccination history. Second, the cross-sectional nature of the data limits our ability to establish causal relationships, meaning observed associations should be interpreted with caution. Additionally, while the DHS provides robust data, it may not capture all potential determinants of immunization coverage, such as healthcare accessibility and cultural influences on immunization. Despite these limitations, our study contributes to the existing

body of knowledge by identifying key socio-demographic disparities in immunization coverage. These findings can inform future research and targeted interventions aimed at reducing inequities and improving vaccination rates among young children in Malawi.

Conclusions

This study highlights the need to address socio-demographic disparities in basic immunization coverage among children aged 12–23 months in Malawi, where coverage remains below the WHO's recommended 90% target. The findings suggest that maternal age, education, wealth status, marital status, and rural residence are associated with basic immunization, underscoring the importance of targeted interventions to improve healthcare access and awareness. Notably, children from rural areas, wealthier households, and those with educated or married mothers had higher basic immunization coverage. However, variations in coverage across districts and socio-economic groups highlight inequities that require policy intervention.

Efforts should focus on expanding access in underserved districts and urban settings and supporting mothers through socio-economic initiatives. Addressing systemic barriers to healthcare can help ensure more equitable immunization opportunities for all children, regardless of geographic or socio-economic background. Collaborative approaches involving policymakers, healthcare providers, and community stakeholders will be key to closing the immunization gap and advancing global health goals.

Acknowledgement

This work is based on research supported by the UKRI GCRF Accelerating Achievement for Africa's Adolescents (Accelerate) Hub (ES/S008101/1); Oak Foundation [OFIL-20-057 and R46194/AA001]; Wellspring Philanthropic Fund (Grant no. 16204). Also, Monica Ewomazino Akokuwebe, PhD acknowledged the support by the DSI-NRF Centre of Excellence (CoE) in Human Development at the University of the Witwatersrand, Johannesburg, South Africa (Postdoctoral Research Fellowship Grant Reference/UID Number: 91489). The content is solely the responsibility of the authors and does not reflect the views of the UKRI GCRF Accelerating Achievement for Africa's Adolescents (Accelerate) Hub and the DSI-NRF CoE in Human Development. The authors also acknowledge Dr. Mwiza Gideon Singini for providing contextual insights on Malawi setting.

Author contributions

All authors (Dr Ololade Julius Baruwa, Dr Monica Ewomazino Akokuwebe, Mr Oluwafemi John Adeleye, Dr Babatunde Makinde Gbadebo) were involved in the conceptualization, analysis, and discussion of the manuscript.

Funding

This research was not funded by any organization or institution.

Data availability

The study used secondary data from dhsprogram.org, which is publicly available at https://dhsprogram.com/data/available-datasets.cfm.

Declarations

Ethical approval

This study used data that was obtained from the 2015-16 Malawi Demographic and Health Survey (MDHS). Informed consent was obtained from each respondent before the interviews. We obtained approval to use the data from the DHS repository (https://dhsprogram.com/data/available-datase ts.cfm accessed on 14 March 2023).

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Received: 18 August 2024 / Accepted: 28 February 2025 Published online: 05 March 2025

References

- Colomé-Hidalgo M, Donado Campos J, Gil de Miguel A. Monitoring inequality changes in full immunization coverage in infants in Latin America and the Caribbean. Rev Panam Salud Publica. 2020;44:e56. https://doi.org/10.26633/R PSP.2020.56.
- Ali HA, Hartner AM, Echeverria-Londono S, Roth J, Li X, Abbas K, Portnoy A, Vynnycky E, Woodruff K, Ferguson NM, Toor J, Gaythorpe Katy AM. Vaccine equity in low and middle income countries: a systematic review and metaanalysis. Int J Equity Health. 2022;21:82. https://doi.org/10.1186/s12939-022-0 1678-5.
- Lyons C, Nambiar D, Johns NE, Allorant A, Bergen N, Hosseinpoor AR. Inequality in childhood immunization coverage: A scoping review of data sources, analyses, and reporting methods. Vaccines. 2024;12(8):850. https://doi.org/10. 3390/vaccines12080850.
- Dimitrova A, Carrasco-Escobar G, Richardson R, Benmarhnia T. Essential childhood immunization in 43 low- and middle-income countries: analysis of Spatial trends and socioeconomic inequalities in vaccine coverage. PLoS Med. 2022;20(1). https://doi.org/10.1371/journal.pmed.1004166.
- Arat A, Burström B, Östberg V, et al. Social inequities in vaccination coverage among infants and pre-school children in Europe and Australia – a systematic review. BMC Public Health. 2019;19:290. https://doi.org/10.1186/s12889-01 9-6597-4.
- Sharma S, Maheshwari S, Jaiswal AK, Mehra S. Income-based inequality in full immunization coverage of children aged 12–23 months in Eastern India: A decomposition analysis. Clin Epidemiol Global Health. 2021;11:100738.
- Eagan RL, Larson HJ. Recent trends in vaccine coverage and confidence: A cause for concern. Hum Vaccines Immunotherapeutics. 2023;19(2). https://do i.org/10.1080/21645515.2023.2237374.
- 8. Lai X, et al. Estimating global and regional between-country inequality in routine childhood vaccine coverage in 195 countries and territories from 2019 to 2021: a longitudinal study. eClinicalMedicine. 2023;60:102042.
- Shiferie F, Gebremedhin S, Andargie G, Tsegaye DA, Alemayehu WA, Mekuria LA, Wondie T, Fenta TG. Vaccination dropout and wealth related inequality among children aged 12–35 months in remote and underserved settings of Ethiopia: A cross-sectional evaluation survey. Front Pead. 2023;11:1280746. ht tps://doi.org/10.3389/fped.2023.1280746.
- D Frenkel L. The global burden of vaccine-preventable infectious diseases in children less than 5 years of age: implications for COVID-19 vaccination. How can we do better? Allergy Asthma Proc. 2021;42(5):378–85. https://doi.org/10. 2500/aap.2021.42.210065.
- Mmanga K, Mwenyenkulu TE, Nkoka O, Ntenda PAM. Tracking immunization coverage, dropout and equity gaps among children ages 12–23 months in Malawi - bottleneck analysis of the Malawi demographic and health survey. Int Health. 2022;14(3):250–9. https://doi.org/10.1093/inthealth/ihab038.
- Lubanga AF, Bwanali AN, Munthali L, Mphepo M, Chumbi GD, Kangoma M, Khuluza C. Malawi vaccination drive: an integrated immunization campaign against typhoid, measles, Rubella, and polio; health benefits and potential challenges. Hum Vaccines Immunotherapeutics. 2023;19(2). https://doi.org/1 0.1080/21645515.2023.2233397.
- UNICEF. Immunization and nutrition supplies budget process mapping. UNICEF Malawi Resour Mobilization Strategy 2021–2023. 1–78. https://www.

unicef.org/esa/media/10521/file/UNICEF-Malawi-Immunization-Nutrition-Su pplies-Budget-Process-Mapping-2021.pdf.

- Bar-Zeev N, Kapanda L, King C, Beard J, Phiri T, Mvula H, Crampin AC, Mwansambo C, Costello A, Parashar U, Tate JE, Verani JR, Whitney CG, Heyderman RS, Cunliffe NA, French N, Consortium V. Methods and challenges in measuring the impact of National Pneumococcal and rotavirus vaccine introduction on morbidity and mortality in Malawi. Vaccine. 2015;33(23):2637–45. https://doi. org/10.1016/j.vaccine.2015.04.053.
- Hu Y, Chen Y, Liang H, Wang Y. An overview of coverage of BCG vaccination and its determinants based on data from the coverage survey in Zhejiang Province. Int J Environ Res Public Health. 2018;15(6):1155. https://doi.org/10.3 390/ijerph15061155.
- Evans B, Keiser O, Kaiser L, Jombart T. Analysis of global routine immunisation coverage shows disruption and stagnation during the first two-years of the COVID-19 pandemic with tentative recovery in 2022. Vaccine: X. 2023;15:100383. https://doi.org/10.1016/j.jvacx.2023.100383.
- Charoenwat B, Suwannaying K, Paibool W, Laoaroon N, Sutra S, Thepsuthammarat K, Sirirattanakul S. The impact of rotavirus vaccination on acute diarrhea in Thai children under 5 years of age in the first year of universal implementation of rotavirus vaccines in the National immunization program (NIP) in Thailand: a 6-year analysis. BMC Public Health. 2023;23:2109. https://doi.org/10.1186/s12889-023-16958-0.
- United Nations International Children's Emergency Fund (UNICEF). (2023). Levels and trends of child mortality in West and Central Africa. Accessed on October 20th, 2024 from https://www.unicef.org/wca/media/9501/file/hild% 20Mortality%20in%20WCAR_Technical%20brief_finale_20230929.pdf
- Lawn JE, Bhutta ZA, Ezeaka C, Saugstad O. Ending preventable neonatal deaths: multicountry evidence to inform accelerated progress to the sustainable development goal by 2030. Neonatology. 2023;120(4):491–9. https://doi. org/10.1159/000530496.
- Watson OJ, Barnsley G, Toor J, Hogan AB, Winskill P, Ghani AC. Global impact of the first year of COVID-19 vaccination: a mathematical modelling study. Lancet Infect Dis. 2022;22(9):1293–302. https://doi.org/10.1016/S1473-3099(2 2)00320-6.
- Touray E, Barrow A, Kinteh B, Badjie M, Nget M, Touray J, Kinteh S, Jatta LSS, P.S., Ceesay L. Childhood vaccination uptake and associated factors among children 12–23 months in rural settings of the Gambia: a community-based cross-sectional study. BMC Public Health. 2021;21:1740. https://doi.org/10.11 86/s12889-021-11810-9.
- Grand-Guillaume-Perrenoud JA, Origlia P, Cignacco E. Barriers and facilitators of maternal healthcare utilisation in the perinatal period among women with social disadvantage: A theory-guided systematic review. Midwifery. 2022;105:103237. https://doi.org/10.1016/j.midw.2021.103237.
- Akokuwebe ME, Idemudia ES. Factors associated with choice of home birth as place of delivery among women of reproductive age in South Africa. Afr J Reprod Health. 2023;27:22–40.
- Addo IY, Acquah E, Nyarko SH, Boateng K, E. N., Dickson KS. Factors associated with unskilled birth attendance among women in sub-Saharan Africa: A multivariate-geospatial analysis of demographic and health surveys. PLoS ONE. 2023;18(2):e0280992. https://doi.org/10.1371/journal.pone.0280992.
- Osuafor GN, Akokuwebe ME, Idemudia ES. Male involvement in family planning decisions in Malawi and Tanzania: what are the determinants?? Int J Environ Res Public Health. 2023;20(6):5053. https://doi.org/10.3390/ijerph200 65053.
- 26. Food and Agriculture Organization of the United Nations (FAO). (2011). Gender inequalities in rural employment in Malawi: an overview. Malawi country profile. Prepared by the Gender, Equity and Rural Employment Division of FAO. p. viii.
- Malawi Demographic and Health Survey (MDHS). (2015-16). 2015-16 Malawi Demographic and Health Survey. National Statistical Office (NSO) [Malawi] and ICF. 2017. Zomba, Malawi, and Rockville, Maryland, USA. NSO and ICF.
- 28. Mmanga K, Mwenyenkulu TE, Nkoka O, Ntenda PA. Tracking immunization coverage, dropout and equity gaps among children ages 12–23 months in

Malawi–bottleneck analysis of the Malawi demographic and health survey. Int Health. 2022;14(3):250–9.

- King C, Bar- Zeev N, Phiri T, Beard J, Mvula H, Crampin A, Heinsbroek E, Hungerford D, Lewycka S, Verani J, Whitney C, Costello A, Mwansambo C, Cunliffe N, Heyderman R, French N, for the VacSurv Consortium. Population impact and effectiveness of sequential 13-valent Pneumococcal conjugate and monovalent rotavirus vaccine introduction on infant mortality: prospective birth cohort studies from Malawi. BMJ Global Health. 2020;5:e002669. https:// doi.org/10.1136/bmjgh-2020-002669.
- Bobo FT, Asante A, Woldie M, Dawson A, Hayen A. Child vaccination in sub-Saharan Africa: increasing coverage addresses inequalities. Vaccine. 2022;40(1):141–50. https://doi.org/10.1016/j.vaccine.2021.11.005.
- Ameyaw EK, Kareem YO, Ahinkorah BO, et al. Decomposing the rural-urban gap in factors associated with childhood immunisation in sub-Saharan Africa: evidence from surveys in 23 countries. BMJ Global Health. 2021;6:e003773.
- 32. Johns NE, Hosseinpoor AR, Chisema M, Danovaro-Holliday MC, Kirkby K, Schlotheuber A, Shibeshi M, Sodha SV, Zimba B. Original research: association between childhood immunisation coverage and proximity to health facilities in rural settings: A cross-sectional analysis of service provision assessment 2013–2014 facility data and demographic and health survey 2015–2016 individual data in Malawi. BMJ Open. 2022;12(7). https://doi.org/10.1136/bmj open-2022-061346.
- Ratta AK, Meshram PD. Study of determinants of immunization status among under-five children attending OPD in a rural health center of the tertiary health care Institute. J Family Med Prim Care. 2020;9(6):2724–9. https://doi.or g/10.4103/jfmpc_jfmpc_387_20.
- Local Burden of Disease Vaccine Coverage Collaborators. Mapping routine measles vaccination in low- and middle-income countries. Nature. 2021;589:415–9. https://doi.org/10.1038/s41586-020-03043-4.
- 35. World Health Organization (WHO). (2021). Immunization coverage. https://w www.ho.int/news-room/fact-sheets/detail/immunization-coverage
- 36. Johns, N. E., Hosseinpoor, A. R., Chisema, M., Danovaro-Holliday, M. C., Kirkby, K.,Schlotheuber, A.,... Zimba, B. (2022). Association between childhood immunisation coverage and proximity to health facilities in rural settings: a cross-sectional analysis of Service Provision Assessment 2013–2014 facility data and demographic and Health Survey 2015–2016 individual data in Malawi. BMJ open, 12(7), e061346.
- Hardhantyo M, Chuang YC. Urban-rural differences in factors associated with incomplete basic immunization among children in Indonesia: A nationwide multilevel study. Pediatr Neonatology. 2021;62(1):80–9.
- 38. Ekezie W, Awwad S, Krauchenberg A, Karara N, Dembiński Ł, Grossman Z, Del Torso S, Dornbusch HJ, Neves A, Copley S, Mazur A, Hadjipanayis A, Grechukha Y, Nohynek H, Damnjanović K, Lazić M, Papaevangelou V, Lapii F, Stein-Zamir C, Rath B, For The ImmuHubs Consortium. Access to vaccination among disadvantaged, isolated and difficult-to-reach communities in the WHO European region: A systematic review. Vaccines (Basel). 2022;10(7):1038.
- Bangura JB, Xiao S, Qiu D, Ouyang F, Chen L. Barriers to childhood immunization in sub-Saharan Africa: A systematic review. BMC Public Health. 2020;20:1–15.
- Lakew Y, Bekele A, Biadgilign S. Factors influencing full immunization coverage among 12–23 months of age children in Ethiopia: evidence from the National demographic and health survey in 2011. BMC Public Health. 2015;15:1–8.
- Mekonnen AG, Bayleyegn AD, Ayele ET. Immunization coverage of 12–23 months old children and its associated factors in Minjar-Shenkora district, Ethiopia: a community-based study. BMC Pediatr. 2019;19:1–8.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.