



OPEN ACCESS

Case report

Giant gluteal lipoma surgical management

Alexis Litchinko, Floryn Cherbanyk,^{ORCID} Markus Menth, Bernhard Egger

Surgery, HFR Fribourg Hopital Cantonal, Fribourg, Switzerland

Correspondence to

Dr Alexis Litchinko, alexis.litchinko@gmail.com

Accepted 14 July 2019

SUMMARY

Surgical treatment of lipomas is typically only considered when they are painful or unsightly. We present the case of a massive hip lipoma; with this extreme case, we show that the global prolongation of life expectancy can lead to other indications of removal.

BACKGROUND

Lipomas are benign and common soft-tissue tumours with many different clinical presentations. Surgical management can sometimes be tricky, and surgeons must keep in mind the potential risk of malignancy with large tumours.

CASE PRESENTATION

We describe the case of a 90-year-old patient with a giant gluteal lipoma having evolved over more than 20 years. The patient was well aware of this slow-growing painless mass on his right hip but refused any treatment. He was addressed to our surgical consultation by his family doctor for lameness and unbalanced walking. (figure 1A, B).

INVESTIGATIONS

Even though a benign lipoma was clearly suspected, a CT scan was performed. However, the lesion was so large that it could not be scanned entirely (figure 2). The tumour-like lipoma seemed to be in contact with the sacrum, the tuber ischiadicum and the greater trochanter. Furthermore, there was a suspicion of an infiltration of the flexor's muscles of the right leg. Vascularisation of the lipoma was mainly provided by an enlarged superior gluteal artery.




Figure 1 (A) Posterior view of the patient's lipoma. (B) Lateral view of the patient's lipoma.




Figure 2 CT scan of the right hip lipoma.

TREATMENT

After discussing the pros and cons of an intervention with the patient and his family, the decision was taken to remove the mass surgically. The patient was installed in a dorsal decubitus position (figure 3A). The operation resulted in the removal of a 30×60 cm 20 kg lipoma (figure 3B).

OUTCOME AND FOLLOW-UP

The pathology report confirmed that the tumour was benign. Skin was preserved in order to allow a functional reconstruction of the gluteal region. Wound management was simple, and follow-up was unremarkable. Additionally, the patient received intensive physiotherapy for walk recovery and after 5 weeks he could achieve completely independent walking.

DISCUSSION

What makes this case interesting is the fact that even a benign pathology such as a lipoma progressed into a situation requiring delicate surgical management, both due to the technical aspect of the operation itself, and because of the patient's advanced age.

With no difference in prevalence across genders and a 40–60 years old average age,¹ lipomas are the most frequent benign soft-tissue tumours. Whether isolated or multiple, congenital or post-traumatic,² lipomas are frequent in our daily routine; some studies mention the genetic implication in more than 50% of them.³ Even with minimal risk of malignant evolution, the first step of imaging should be ultrasonography,⁴ sometimes associated with MRI for large or deep tumours.⁵ Various treatments can be suggested for mechanical or aesthetic concerns. Surgery is the most appropriate treatment, associated with other techniques⁶ (suction, soft-tissue reconstruction, flap and so on).



© BMJ Publishing Group Limited 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Litchinko A, Cherbanyk F, Menth M, et al. *BMJ Case Rep* 2019;**12**:e229842. doi:10.1136/bcr-2019-229842




Figure 3 (A) Dorsal decubitus surgical position. (B) Lipoma total dissection.

Various pathologies imitating lipomas, such as lipomatosis or lipodystrophy, should be distinguished from it.⁷ The WHO gives a standard classification for all types of lipomas.⁸

Giant lipomas are a specific entity as their precise definition is only based on the general aspect of the tumour, with a specific size and weight cut-off given by Sanchez *et al.*⁹ In order for a lipoma to be called 'giant', the lesion must be at least 10 cm in one dimension or weigh a minimum of 1 kg. Based on recent literature, the case we report here is the second largest sub-cutaneous lipoma ever described (table 1).

Learning points

- ▶ Giant lipomas are above all aesthetics concerns but in advanced form a functional problem.
- ▶ Benign tumours with low risk of malignant evolution.
- ▶ Surgery as if aesthetic or functional disorders.
- ▶ The transformation of a large lipoma (>10 cm) to a liposarcoma is rare.

Contributors AL wrote the article; MM, FC and BE verified the analytical methods and supervised the findings of this work. All authors discussed the results and contributed to the final manuscript.

Table 1 Largest giant lipomas in French/English-language literature (>10 cm and >1000 g)

Reference	Year	Localisation	Pathology	Size (cm)/weight (kg)
Current case	2017	Gluteal	Unknown	30×60/20
Emegoakor <i>et al</i> ¹⁰	2017	Lower limb	Unknown	22×17/??
Mascarenhas <i>et al</i> ¹¹	2017	Gluteal	Liposarcoma	17/??
Guler <i>et al</i> ^{12, 13}	2016	Back	Unknown	38×22×21/3.575
Grimaldi <i>et al</i> ¹⁴	2015	Back	Lipoma	36×40×24/5.75
Dabloun <i>et al</i> ¹⁵	2015	Back	Unknown	25×25×18/??
Silistrelil <i>et al</i> ¹⁶	2004	Back	Unknown	??/12.350
Martin <i>et al</i> ¹⁷	1928	Neck	Unknown	??/12.5
Brandler <i>et al</i> ¹⁸	1894	Back	Unknown	??/22.7

?? represents that others authors did not provide a number for this specific information.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- 1 Inampudi P, Jacobson JA, Fessell DP, *et al.* Soft-tissue lipomas: accuracy of sonography in diagnosis with pathologic correlation. *Radiology* 2004;233:763–7.
- 2 Copcu E, Sivrioglu NS. Posttraumatic lipoma: analysis of 10 cases and explanation of possible mechanisms. *Dermatol Surg* 2003;29:215–20.
- 3 Turc-Carel C, Limon J, Dal Cin P, *et al.* Cytogenetic studies of adipose tissue tumors. II. Recurrent reciprocal translocation t(12;16)(q13;p11) in myxoid liposarcomas. *Cancer Genet Cytogenet* 1986;23:291–9.
- 4 Kuwano Y, Ishizaki K, Watanabe R, *et al.* Efficacy of diagnostic ultrasonography of lipomas, epidermal cysts, and ganglions. *Arch Dermatol* 2009;145:761–4.
- 5 Fuchs A, Henrot P, Walter F, *et al.* [Lipomatous tumors of soft tissues in the extremities and the waist in adults]. *J Radiol* 2002;83:1035–57.
- 6 Al-basti HA, El-Khatib HA. The use of suction-assisted surgical extraction of moderate and large lipomas: long-term follow-up. *Aesthetic Plast Surg* 2002;26:114–7.
- 7 Delgado L, Ingen-Housz-Oro S. Maladies du tissu adipeux: lipomes, lipomatoses, lipodystrophies. *EMC - Dermatologie* 2011;6:1–9.
- 8 Vilanova JC. WHO Classification of Soft Tissue Tumors. *Imaging of soft tissue tumors*: Springer International Publishing, 2017:187–96.
- 9 Sanchez MR, Golomb FM, Moy JA, *et al.* Giant lipoma: case report and review of the literature. *J Am Acad Dermatol* 1993;28(2 Pt 1):266–8.
- 10 Emegoakor CD, Echezona CN, Onwukamuche ME, *et al.* A report of two cases. *Niger J Gen Pract* 2017;15:46–9.
- 11 Mascarenhas MRM, Mutti LA, Paiva JMG, *et al.* Giant atypical lipoma. *An Bras Dermatol* 2017;92:546–9.
- 12 Righi A, Pantalone O, Tagliaferri G. Giant lipoma of the thigh: a case report. *J Ultrasound* 2012;15:124–6.
- 13 Guler O, Mutlu S, Mahiroglu M. Giant lipoma of the back affecting quality of life. *Ann Med Surg* 2015;4:279–82.
- 14 Grimaldi L, Cuomo R, Castagna A, *et al.* Giant lipoma of the back. *Indian J Plast Surg* 2015;48:220–1.
- 15 Dabloun S, Khechimi M, Jenzeri A, *et al.* Lipome géant du dos : à propos d'un cas. *Annales de Dermatologie et de Vénérologie* 2015;142:5353.
- 16 Silistrelil OK, Durmuş EU, Ulusal BG, *et al.* What should be the treatment modality in giant cutaneous lipomas? Review of the literature and report of 4 cases. *Br J Plast Surg* 2005;58:394–8.
- 17 Martin HS. Massive lipoma of the subcutaneous tissue of the back. *J Am Med Assoc* 1928;90:2013.
- 18 Brandler TI. Large fibrolipoma. *Br Med J* 1894;1:574.

Copyright 2019 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow