



Cite this: *RSC Adv.*, 2025, 15, 10634

Transition-metal-free regioselective synthesis of spiro-oxazolidines through [3 + 2] annulation reactions of azadienes with haloalcohols†

Amol T. Savekar,^{ID ‡} Sonali M. Vitnor,[‡] Vishal B. Karande and Suresh B. Waghmode^{ID *}

The transition-metal-free regioselective [3 + 2] annulation of azadienes with haloalcohols for the preparation of highly functionalized spiro-oxazolidines is experimentally simple and proceeds under mild conditions. The metal-free protocols have more significance than the metal-catalyzed ones when the toxicity associated with the metal catalyst is considered. This transformation features a broad substrate scope, good yields, and excellent regioselectivity. Moreover, large-scale synthesis and representative transformations of spiro-oxazolidines were carried out to provide additional evidence on the practicality of this approach.

Received 27th February 2025

Accepted 14th March 2025

DOI: 10.1039/d5ra01423a

rsc.li/rsc-advances

Introduction

Spirocyclic skeletons are privileged structural moieties in many biologically active natural products owing to their inherent three-dimensional architecture.¹ Among the useful classes of oxygen- and nitrogen-based heterocycles, oxazolidine is a significant motif present in various natural products and pharmaceuticals (as shown in Fig. 1), and it is used as a chiral auxiliary and chiral ligand in diverse asymmetric transformations.^{2,3} In particular, spiro-oxazolidines, with heteroatom-substituted quaternary stereocenters, are valuable building blocks to construct medicinally relevant compounds.⁴

Since the existence of heteroatoms can bring new synthetic and biological values, the focus has recently been shifted to utilizing azadienes (1 and 4) to prepare heterocycles containing

multi-heteroatoms.⁵ Azadiene contains exocyclic alkylidene and an imine unit, which acts as a four-atom synthon in various Michael additions and subsequent cascade reactions, delivering diverse cyclic compounds through [4 + *n*] cycloaddition reactions due to the driving force for aromatization.⁶ Recently, Zhao's group reported [4 + 3] annulation through 1,4 addition of α -bromohydroxamates to azadienes to access benzofuran-fused seven-membered heterocycles (Scheme 1a).⁷ In contrast to well-developed [4 + *n*] cyclizations through 1,4-addition, which are applicable only for the synthesis of aromatized heterocycles, these azadienes can also serve as two-atom synthons to undergo [2 + *n*] annulation to synthesize spirocyclic motifs.⁸ Very recently, Liu's group developed a methodology for the synthesis of spirocyclopentanone through [3 + 2] cycloaddition of cyclopropanes with azadienes (Scheme 1b).⁹ Except a few reports, the synthesis of spiro-oxazolidine scaffolds has not been developed yet.¹⁰ The 1,2-addition reactions are comparatively less developed; Trost's group have recently demonstrated Pd-catalyzed [3 + 2] spiroannulation from azadienes and vinyl epoxides (Scheme 1c).¹¹ These reported methods involved noble metal-





Fig. 1 Representative examples of bioactive oxazolidine scaffolds.

Department of Chemistry, Savitribai Phule Pune University (Formerly University of Pune), Ganeshkhind, Pune-411007, India. E-mail: suresh.waghmode@unipune.ac.in; suresh.waghmode@gmail.com

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d5ra01423a>

‡ These authors contributed equally to this work.



Scheme 1 Annulation reactions of azadienes with various reaction partners.



catalyzed protocol, high cost, multistep cascade reactions, drastic reaction conditions, and limited substrate scope, adding to their drawbacks. The quest to explore alternatives to transition metal catalysts is mainly attributed to the toxicity inherent in such systems, especially when it comes to synthesizing heterocycles of biological relevance. Hence, protocols that proceed under transition-metal-free conditions are always desirable among the scientific community.¹²

However, to our knowledge, the regioselective transition-metal-free [3 + 2] spiroannulation of azadienes has not been studied yet. Therefore, it is highly desirable to synthesize spiro-oxazolidines under transition-metal-free and ambient reaction conditions. Herein, we describe the development of an approach for the regioselective synthesis of spiro-oxazolidines under ambient reaction conditions.

Recently, haloalcohols (Br, Cl, and I) and their homologs were explored and developed as Michael donors for [3 + 2]/[4 + 2] annulation reactions with various Michael acceptors for the synthesis of an important class of heterocyclic compounds.¹³ Based on this background and as a continuation of our interest in bioactive-fused polycyclic structure synthesis,¹⁴ we carried out [3 + 2] annulation of azadienes with haloalcohols under milder reaction conditions to obtain the corresponding spiro-


oxazolidines, which are constituents of various natural products and pharmaceuticals (Scheme 1d).

Results and discussion


The investigation started by evaluating the reaction of benzofuran-derived azadienes **1a** (1.0 equiv.), and 2-bromoethanol **2a** (1.5 equiv.) with, 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) (2.5 equiv.) as a base in MeCN at room temperature (*i.e.*, 25 °C) for 3 h, and the desired (*Z*)-2-benzylidene-3'-tosyl-2*H*-spiro[benzofuran-3,2'-oxazolidine] product **3a** was generated in 28% yield (Table 1; entry 1). After extensive screening of various bases such as K₃PO₄, K₂CO₃, Et₃N, NaH, Cs₂CO₃, and *t*-BuOK, surprisingly it was found that Cs₂CO₃ provided the highest 81% yield in 3 h (entries 2–7). Similarly, organic solvents were also studied, namely 1,4-dioxane, DMF, DMSO, THF, dichloromethane (DCM), toluene, ethyl acetate (EtOAc), and acetone; among the studied solvents, acetone (83%, yield) proved to be the best (entries 8–15). Furthermore, variation in the stoichiometry of 2-bromoethanol did not show promising results (entries 16 and 17). A lower yield was observed when the base loading decreased or increased (entries; 18 and 19).

After optimizing the reaction conditions, we began exploring the scope of the [3 + 2] annulation reaction (Scheme 2). When various *N*-protecting groups such as *N*-SO₂Ph, *N*-SO₂PhBr, *N*-Ns, *N*-SO₂PhCF₃, and *N*-Ms containing azadienes were used as substrates, the corresponding target products **3b–3f** were obtained in good yields (63–76%). We observed that spirocyclization was facilitated by 2-chloroethanol and 2-iodoethanol provided the product with lower yields (20–31%). Various substitutions at *ortho*, *meta*, and *para* positions on the phenyl of azadienes provided products in good to high yields (**3g–3n**). The


Table 1 Optimization of reaction conditions for the synthesis of spiro-oxazolidine derivatives^a

					
Sr. no.	2a (equiv.)	Base	Solvent	Time (h)	Yield ^b (%)
1	1.5	DBU	MeCN	3	28
2	1.5	K ₃ PO ₄	MeCN	6	78
3	1.5	K ₂ CO ₃	MeCN	12	52
4	1.5	Et ₃ N	MeCN	12	Trace
5	1.5	NaH	MeCN	12	40
6	1.5	Cs ₂ CO ₃	MeCN	3	81
7	1.5	<i>t</i> -BuOK	MeCN	12	40
8	1.5	Cs ₂ CO ₃	1,4-Dioxane	12	10
9	1.5	Cs ₂ CO ₃	DMF	6	72
10	1.5	Cs ₂ CO ₃	DMSO	12	Trace
11	1.5	Cs ₂ CO ₃	THF	12	5
12	1.5	Cs ₂ CO ₃	DCM	12	nr
13	1.5	Cs ₂ CO ₃	Toluene	12	nr
14	1.5	Cs ₂ CO ₃	EtOAc	2	30
15	1.5	Cs ₂ CO ₃	Acetone	2	83
16	1	Cs ₂ CO ₃	Acetone	2	75
17	2	Cs ₂ CO ₃	Acetone	2	83
18 ^c	1.5	Cs ₂ CO ₃	Acetone	2	73
19 ^d	1.5	Cs ₂ CO ₃	Acetone	2	81


^a All reactions were performed with **1a** (0.27 mmol, 1.0 equiv.), **2a** (0.40 mmol, 1.5 equiv.), and base (0.67 mmol, 2.5 equiv.) in solvent (2.0 mL) at room temperature (rt) under N₂. ^b Isolated yields. ^c Cs₂CO₃ (0.53 mmol, 2.0 equiv.). ^d Cs₂CO₃ (0.80 mmol, 3.0 equiv.), nr = no reaction.



Scheme 2 Substrate scope for the synthesis of spiro-oxazolidine derivatives.^{a,b} ^aAll reactions were performed with **1** (0.27 mmol, 1.0 equiv.), **2** (0.40 mmol, 1.5 equiv.), and Cs₂CO₃ (0.67 mmol, 2.5 equiv.) in 2.0 mL acetone at room temperature for 2 h. ^bIsolated yields. ^c2-Chloroethanol and *t*-BuOK. ^d2-Iodoethanol and *t*-BuOK.




Scheme 4 Synthesis of spiro-oxazolidines derivatives from isatin-derived ketimines.




Scheme 5 Plausible reaction mechanism for the synthesis of spiro-oxazolidines.

To demonstrate the synthetic potential of this transformation, we investigated the scaled-up preparation of **3a** and **5a** under the standard reaction conditions (Scheme 6a and b). The reaction of 1.0 g of **1a** and **4a** proceeded smoothly, to deliver products **3a** and **5a** with yields of 75% and 58% in 3 h, respectively. Furthermore, to investigate the potential utility and enrich the spirocyclic compound's molecular complexity, we carried out the derivatization of the product **3a** (Scheme 7). When *m*-CPBA was used for the epoxidation of **3a** in DCM, epoxide **8** was obtained with excellent yields and



Scheme 6 Gram-scale synthesis.



Scheme 7 Synthetic transformation of product **3a**.

diastereoselectivity (91%, >20:1 dr) in 12 h (Scheme 7a). Various epoxides are valuable building blocks in chemical synthesis and such structural motifs are present in biologically active molecules.¹⁵ Hydrogenation of **3a** readily generated product **9** in good yields with moderate diastereoselectivity (Scheme 7b).

Conclusions

In summary, we have developed a highly regioselective and chemoselective [3 + 2] spiroannulative transformation of azadienes. The reaction was carried out in one pot without transition metals and under mild reaction conditions. This method demonstrated good substrate generality, and was successfully applied to large-scale synthesis. Product derivations further confirmed the applicability of this method. However, this feeding sequence-controlled divergent reaction was limited to a special type of substrate.

Data availability

The data supporting this study are available in the published article and its ESI.[†]

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

ATS thanks UGC New Delhi for the research fellowship (JUNE-18-144311), and SMV thanks NF-OBC (NBCFDC Ref. No. 231610106961). SBW and VBK thank SERB-SURE project SUR/2022/000608 for financial support. They also thank CIF SPPU, Pune, for the spectral analysis facility.

References

- (a) C. M. Marson, *Chem. Soc. Rev.*, 2011, **40**, 5514–5533; (b) Y. Zheng, C. M. Tice and S. B. Singh, *Bioorg. Med. Chem. Lett.*, 2014, **24**, 3673–3682; (c) R.-Y. Q. Li-Ming Zhou and G.-F. Yang, *Expert Opin. Drug Discovery*, 2020, **15**, 603–625; (d) K. Hiesinger, D. Dar'in, E. Proschak and M. Krasavin, *J. Med. Chem.*, 2021, **64**, 150–183; (e) L. Yu, A. Dai, W. Zhang, A. Liao, S. Guo and J. Wu, *J. Agric. Food Chem.*, 2022, **70**, 10693–10707.
- (a) T. Fukuyama, L. Li, A. A. Laird and R. K. Frank, *J. Am. Chem. Soc.*, 1987, **109**, 1587–1589; (b) K. Ii, S. Ichikawa, B. Al-Dabbagh, A. Bouhss and A. Matsuda, *J. Med. Chem.*, 2010, **53**, 3793–3813; (c) A. Kamal, P. Swapna, R. Shetti, A. B. Shaik, M. P. Narasimha Rao and S. Gupta, *Eur. J. Med. Chem.*, 2013, **62**, 661–669.
- (a) C. Agami and F. Couty, *Eur. J. Org. Chem.*, 2004, 677–685; (b) C. Wolf and H. Xu, *Chem. Commun.*, 2011, **47**, 3339–3350.
- (a) J. D. Scott and R. M. Williams, *J. Am. Chem. Soc.*, 2002, **124**, 2951–2956; (b) U. Rix, J. Zheng, L. L. Remsing Rix, L. Greenwell, K. Yang and J. Rohr, *J. Am. Chem. Soc.*, 2004,

- 126, 4496–4497; (c) H. U. Kaniskan and P. Garner, *J. Am. Chem. Soc.*, 2007, **129**, 15460–15461; (d) T. T. Talele, *J. Med. Chem.*, 2020, **63**, 13291–13315.
- 5 (a) Q. Deng and X. Meng, *Chem.-Asian J.*, 2020, **15**, 2838–2853; (b) S. K. Saha, A. Bera, S. Singh and N. K. Rana, *Eur. J. Org. Chem.*, 2023, **26**, e202201470; (c) Y. Song, G. Zhang, Y. Zhang, H. Zhang, X. Meng and L. Zhang, *Org. Chem. Front.*, 2022, **9**, 4598–4605; (d) B. Shi, M. Xiao, J.-P. Zhao, Z. Zhang, W. J. Xiao and L.-Q. Lu, *J. Am. Chem. Soc.*, 2024, **146**, 26622–26629; (e) Y.-Z. Liu, C. Zheng and S.-L. You, *ACS Catal.*, 2024, **14**, 15743–15750.
- 6 (a) J. Chen and Y. Huang, *Org. Lett.*, 2017, **19**, 5609–5612; (b) L.-C. Yang, Z.-Q. Rong, Y.-N. Wang, Z. Y. Tan, M. Wang and Y. Zhao, *Angew. Chem., Int. Ed.*, 2017, **56**, 2927–2931; (c) C.-S. Wang, T.-Z. Li, Y.-C. Cheng, J. Zhou, G.-J. Mei and F. Shi, *J. Org. Chem.*, 2019, **84**, 3214–3222; (d) V. Laina-Martin, J. Humbrias-Martin, R. Mas-Balleste, J. A. F. Salas and J. Aleman, *ACS Catal.*, 2021, **11**, 12133–12145; (e) G.-H. Chen, Y. Ye, D.-X. Zhang, H.-J. Li, N. Zhang, G. J. Liang, D. Zhang, J. Zhou and H. Zhou, *Org. Chem. Front.*, 2023, **10**, 4698–4702; (f) Y. Wu, Z. Wang, Y. Shan, Y. Ma, T. Li, C. Yuan, H. Guo and B. Mao, *Chem. Sci.*, 2024, **15**, 9703–9708.
- 7 Q. Y. Fang and L. M. Zhao, *Chem. Commun.*, 2020, **56**, 14079–14082.
- 8 (a) Q. Y. Fang, M. H. Yi, X. X. Wu and L.-M. Zhao, *Org. Lett.*, 2020, **22**, 5266–5270; (b) K. Liu, J. Yang and X. Li, *Org. Lett.*, 2021, **23**, 826–831; (c) L. Tu, L. Gao, Q. Wang, Z. Cao, R. Huang, Y. Zheng and J. Liu, *J. Org. Chem.*, 2022, **87**, 3389–3401; (d) L. Yang, C. Pi, Y. Wu and X. Cui, *Org. Lett.*, 2022, **24**, 7502–7506; (e) R. Khuntia, S. K. Mahapatra, L. Roy and S. C. Pan, *Chem. Sci.*, 2023, **14**, 10768–10776; (f) J. Y. Su, J. Zhang, Z.-Y. Xin, H. Li, H. Zheng and W. P. Deng, *Org. Chem. Front.*, 2024, **11**, 4539–4545; (g) J. Huang, A. Yu, L. Zhang and X. Meng, *Org. Lett.*, 2024, **26**, 8945–8950.
- 9 S. Li, Z.-H. Dong, S.-Y. Dan, M.-J. Zheng, T. Long, J. Zhan, Q. Zhou, W.-D. Chu and Q. Z. Liu, *Org. Chem. Front.*, 2024, **11**, 2905–2910.
- 10 (a) T. Rajasekaran, G. Karthik, B. Sridhar and B. V Subba Reddy, *Org. Lett.*, 2013, **15**, 1512–1515; (b) C. Parida, B. Mondal, A. Ghosh and S. C. Pan, *J. Org. Chem.*, 2021, **86**, 13082–13091; (c) M. G. Ordonez, L. Martin, A. Maestro and J. M. Andrés, *Org. Biomol. Chem.*, 2023, **21**, 2361–2369; (d) S. Jiang, K. Li, J. Yan, K. Shi, C. Zhao, L. Yang and G. Zhong, *J. Org. Chem.*, 2017, **82**, 9779–9785.
- 11 (a) B. M. Trost and Z. Zuo, *Angew. Chem., Int. Ed.*, 2021, **60**, 5806–5810; (b) Z.-H. Dong, S. Li, T. Long, J. Zhan, C. Ruan, X. Yan, W.-D. Chu, K. Yuan and Q.-Z. Liu, *Org. Lett.*, 2024, **26**, 3235–3240.
- 12 (a) V. P. Mehta and B. Punji, *RSC Adv.*, 2013, **3**, 11957–11986; (b) C.-L. Sun and Z.-J. Shi, *Chem. Rev.*, 2014, **114**, 9219–9280; (c) A. Ibrar, I. Khan, N. Abbas, U. Farooq and A. Khan, *RSC Adv.*, 2016, **6**, 93016–93047; (d) B. Borah and L. R. Chowhan, *RSC Adv.*, 2021, **11**, 37325–37353; (e) K. T. Ashitha, A. Krishna, D. Basavaraja and S. B. Somappa, *Org. Chem. Front.*, 2022, **9**, 5306–5357; (f) M. Ramanathan and Z. Moussa, *Org. Chem. Front.*, 2025, **12**, 256–327.
- 13 (a) M. K. Ghorai, D. Shukla and K. Das, *J. Org. Chem.*, 2009, **74**, 7013–7022; (b) R. A. Gaikwad, A. T. Savekar and S. B. Waghmode, *J. Org. Chem.*, 2023, **88**, 9987–10001; (c) A. T. Savekar, R. A. Gaikwad and S. B. Waghmode, *J. Org. Chem.*, 2024, **89**, 9389–9404; (d) K. K. Gond and M. R. Maddani, *New J. Chem.*, 2024, **48**, 8667–8671.
- 14 (a) R. A. Gaikwad, S. B. Kamble and S. B. Waghmode, *Chem.-Asian J.*, 2022, **17**, e202200931; (b) A. T. Savekar, V. B. Karande, D. G. Hingane and S. B. Waghmode, *Org. Biomol. Chem.*, 2024, **22**, 8945–8950.
- 15 (a) M. González-Pérez, R. Gómez-Bombarelli, J. Arenas-Valgañón, M. T. Pérez-Prior, M. P. García-Santos, E. Calle and J. Casado, *Chem. Res. Toxicol.*, 2012, **25**, 2755–2762; (b) J. Zhang, C. Wang, L. Ji and W. Liu, *Chem. Res. Toxicol.*, 2016, **29**, 841–850; (c) S. Meninno and A. Lattanzi, *ACS Org. Inorg. Au*, 2022, **2**, 289–305; (d) G.-W. Yang, R. Xie, Y.-Y. Zhang, C.-K. Xu and G.-P. Wu, *Chem. Rev.*, 2024, **124**, 12305–12380.