

Case Report

Biloma Secondary to Percutaneous Liver Biopsy Case Report

Marcos Aranda , Jacqueline Mulhall, Alexander Friedman, and Joel Brockmeyer

Dwight D Eisenhower Army Medical Center, Fort Gordon, Georgia

Correspondence should be addressed to Marcos Aranda; marcos.c.aranda.mil@mail.mil

Received 30 December 2019; Accepted 13 February 2020; Published 4 March 2020

Academic Editor: Tahsin Colak

Copyright © 2020 Marcos Aranda et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Biloma and biliary leak after percutaneous liver biopsy (PLB) are rare. Previous cases are largely in the setting of transplant, oncology, and cirrhotic patients. Patients can be asymptomatic, peritoneal, or present with obstructive symptoms, including bilirubinemia. A 55-year-old male referred for transaminitis attributed to nonalcoholic fatty liver disease (NAFLD) underwent an ultrasound- (US-) guided PLB. He returned the same day with abdominal pain, normal vitals, a nontender abdomen, and a leukocytosis. He was found to have a subcapsular fluid collection attributed to a hematoma. He underwent observation and was discharged. He presented 4 days later with fever, tachycardia, leukocytosis, and bilirubinemia. CT demonstrated growth of the subcapsular fluid collection. Percutaneous drainage revealed bilious fluid. He was transferred for endoscopic retrograde cholangiopancreatography (ERCP). A right biliary branch was stented, and he was discharged the following day with antibiotics. US should be utilized for percutaneous biopsies to avoid biliary complications. Typical presentations of biliary complications include abdominal pain and biliary obstruction. The differential diagnosis for perihepatic and subcapsular fluid collections after PLB should include bile. ERCP should be offered for the treatment of larger or symptomatic collections.

1. Introduction

Percutaneous liver biopsy (PLB) is frequently indicated in the workup of many common conditions including nonalcoholic fatty liver disease (NAFLD), and the procedure is completed frequently with few complications [1]. Common complications include pain, superficial bleeding, hematomas, hemobilia, and intraperitoneal hemorrhage [2, 3]. Less common complications include transient bacteremia, biloma, biliary leak, injury to additional organs, such as the right lung or colon, and death [4, 5]. Boyum et al. found an overall complication rate of 0.7% in over 6000 biopsies of which 0.5% were complications related to bleeding [2]. Seeff et al. found only two episodes of biliary complications in over 2500 patients, and both were related to direct injury of the gallbladder [3].

Paymani et al. reported 4 cases of bile leak in transplant patients, one of which was intraparenchymal and three were transhepatic, and they all reported obstructive symptoms [6]. Don et al. reported one case out of 99 PLB in pediatric transplant patients [7]. Avener et al. reported one case in 1979 of an asymptomatic leak diagnosed incidentally after biliary obstruction secondary to lymphoma [8]. Ruben et al. reported one case of bile peritonitis in the setting of uncom-

pensated cirrhosis [9]. Ahluwalia and LaBrecque described the only other published case of a symptomatic biloma in a nontransplant patient without cirrhosis, which resulted in mass effect with gastric outlet obstruction [10].

2. Case Presentation

A 55-year-old male was referred to a gastroenterologist for a persistent mild transaminitis of an AST in the 40s and an ALT in the 80s. He had a history of hyperlipidemia, obesity with a BMI of 38, and sleep apnea. He took simvastatin 80 mg daily with no antiplatelets or anticoagulants. He consumed about 3 to 4 alcoholic beverages per week, and he had no family history of cirrhosis. Viral serologies were negative, and iron and copper studies were normal. Serologies for autoimmune hepatitis, primary biliary cirrhosis (PBC), and primary sclerosing cholangitis (PSC) had been normal.

He was referred to a radiologist for an ultrasound- (US-) guided biopsy. His INR was 1.1, and his platelets were 236,000 cells/ μ l. A 17 G needle was used for anesthesia and a 14 G needle for sampling. After the procedure, he had a collagen thrombin slurry embolization of the tract. Later that day, he presented with right upper quadrant pain, transient




FIGURE 1: 2 cm nonenhancing subcapsular fluid collection coronal.




FIGURE 2: 2 cm nonenhancing subcapsular fluid collection axial.

tachycardia to the low 100s, and tenderness in the right upper quadrant without rebound. Studies revealed a leukocytosis of $13,900 \text{ cells}/\mu\text{l}$, a total bilirubin of 1.2 mg/dl , and a CT of the abdomen and pelvis with intravenous (IV) contrast revealed a 2 cm subcapsular fluid collection which was attributed to a hematoma (Figures 1 and 2). He was admitted for 24 hours of observation with improved pain control and was discharged.

He returned 4 days later with worsening pain, shortness of breath, and loss of appetite. He was febrile to 102.3 F and tachycardic to the 110s and was tender without peritoneal signs in the right upper quadrant. He had a persistent leukocytosis, and his total bilirubin increased to 3.3 mg/dl . A CT with IV contrast was repeated with enlargement to 12 cm of his subcapsular fluid collection (Figures 3 and 4).

The next day, he underwent percutaneous drainage with return of 20 m of green bilious fluid, and a pigtail catheter was left in place after the diagnosis of biloma was confirmed. The same day he was transferred to a facility with ERCP (endoscopic retrograde cholangiopancreatography) capabilities which demonstrated a bile leak from a small, distal duct of the right lobe which was stented with a 5 cm, 10 F plastic stent of which no images were recorded. He was discharged the following day on a week of amoxicillin/clavulanate.

At a clinic follow-up a week later, his pain had improved, his vitals had normalized, and his drain output had diminished. The drain was removed, and he was arranged a follow-up 6 weeks later for stent removal.

3. Discussion

While the initial presentation after biopsy may have represented subcapsular bleeding rather than bile, the bile seen on percutaneous drainage and bile leak on ERCP more strongly suggest an early and clear association between the biopsy and the bile leak. Our case varies from the current literature in that our patient had no history of cirrhosis, malignancy, or transplant. While our patient was symptomatic, he did not have peritoneal signs or signs of obstruction, which is relatively uncommon.



FIGURE 3: 12 cm subcapsular fluid collection coronal.




FIGURE 4: 12 cm subcapsular fluid collection axial.

Both gastrointestinal and radiologic societies have developed guidelines regarding the prevention of bleeding, but there are few specifics to the prevention of biliary complications [1, 11, 12]. General guidelines to prevent biliary

complications include utilizing US, which can assist in avoiding large bile ducts or the gallbladder [1]. At least a 16 G, 2 cm needle should be utilized to obtain a sufficient pathologic specimen, and small needles should not be used to avoid biliary complications [1]. Once recognized, percutaneous drainage can be performed to diagnose and treat the complication [13–15]. Large or persistent fluid collections require ERCP to diagnose and prevent ongoing leak [13–15]. If a collection is infected, adequate drainage should be assured and antibiotics should be initiated [13, 15].

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

The views expressed in this publication are those of the authors and do not necessarily reflect the official policy or position of the Department of the Army, the Department of Defense, nor the US Government. The identification of any commercial product in this publication does not imply Federal, Department of Defense, or US Army endorsement.

References

- [1] D. C. Rockey, S. H. Caldwell, Z. D. Goodman, R. C. Nelson, A. D. Smith, and American Association for the Study of Liver Diseases, “Liver biopsy,” *Hepatology*, vol. 49, no. 3, pp. 1017–1044, 2009.
- [2] J. H. Boyum, T. D. Atwell, G. D. Schmit et al., “Incidence and Risk Factors for Adverse Events Related to Image-Guided Liver Biopsy,” *Mayo Clinic Proceedings*, vol. 91, no. 3, pp. 329–335, 2016.
- [3] L. B. Seeff, G. T. Everson, T. R. Morgan et al., “Complication Rate of Percutaneous Liver Biopsies Among Persons With Advanced Chronic Liver Disease in the HALT-C Trial,” *Clinical Gastroenterology and Hepatology*, vol. 8, no. 10, pp. 877–883, 2010.
- [4] F. Piccinino, E. Sagnelli, G. Pasquale et al., “Complications following percutaneous liver biopsy: A multicentre retrospective study on 68 276 biopsies,” *Journal of Hepatology*, vol. 2, no. 2, pp. 165–173, 1986.
- [5] D. H. Van Thiel, J. S. Gavalier, H. Wright, and A. Tzakis, “Liver biopsy. Its safety and complications as seen at a liver transplant center,” *Transplantation*, vol. 55, no. 5, pp. 1087–1089, 1993.
- [6] M. Paymani, A. B. Zajko, and W. L. Campbell, “Bile leakage as a complication of liver biopsy in liver transplants,” *Abdominal Imaging*, vol. 18, no. 3, pp. 258–260, 1993.
- [7] S. Don, K. K. Kopecky, M. D. Pescovitz, and R. S. Filo, “Ultrasound-guided pediatric liver transplant biopsy using a spring-propelled cutting needle (biopsy gun),” *Pediatric Radiology*, vol. 24, no. 1, pp. 21–24, 1994.
- [8] D. L. Avenier and M. M. Berenson, “Asymptomatic bilious ascites following percutaneous liver biopsy,” *Archives of Internal Medicine*, vol. 139, no. 2, pp. 245–246, 1979.
- [9] R. A. Ruben and S. Chopra, “Bile peritonitis after liver biopsy: nonsurgical management of a patient with an acute abdomen: a case report with review of the literature,” *American Journal of Gastroenterology*, vol. 82, no. 3, pp. 265–268, 1987.
- [10] J. P. Ahluwalia and D. R. LaBrecque, “A large biloma causing gastric outlet obstruction after a percutaneous liver biopsy,” *Journal of Clinical Gastroenterology*, vol. 38, no. 6, pp. 535–539, 2004.
- [11] S. Gupta, M. J. Wallace, J. F. Cardella et al., “Quality improvement guidelines for percutaneous needle biopsy,” *Journal of Vascular and Interventional Radiology*, vol. 21, no. 7, pp. 969–975, 2010.
- [12] I. J. Patel, J. C. Davidson, B. Nikolic et al., “Consensus guidelines for periprocedural management of coagulation status and hemostasis risk in percutaneous image-guided interventions,” *Journal of Vascular and Interventional Radiology*, vol. 23, no. 6, pp. 727–736, 2012.
- [13] P. G. Jackson and S. R. Evans, “Chapter 54: Biliary system,” in *Sabiston Textbook of Surgery*, C. M. Townsend, D. Beauchamp, and M. Evers, Eds., pp. 1482–1519, Elsevier, Philadelphia, 20th edition, 2017.
- [14] S. B. Solomon, J. J. Mezhir, H. Schoellenast, and P. J. Allen, “Chapter 27: Postoperative complications requiring intervention: diagnosis and management,” in *Blumgart’s Surgery of the Liver, Biliary Tract and Pancreas*, W. R. Jarnigan, Ed., pp. 459–474, Elsevier, Philadelphia, 6th edition, 2017.
- [15] D. M. Notrica, “Chapter 16: Abdominal and renal trauma,” in *Holcomb and Ashcraft’s Pediatric Surgery*, G. W. Holcomb, J. P. Murphy, and S. D. St. Peter, Eds., pp. 236–253, Elsevier, China, 7th edition, 2019.