

Cost Analysis in Shoulder Surgery

A Systematic Review

Thomas Tischer,^{*†} Prof, MD, MBA, Robert Lenz,[†] MD, Jochen Breinlinger-O'Reilly,[‡] Prof, and Christoph Lutter,[†] MD, MHBA

Investigation performed at the Department of Orthopaedic Surgery, University Medicine Rostock, Rostock, Germany

Background: Cost analysis studies in medicine were uncommon in the past, but with the rising importance of financial considerations, it has become increasingly important to use available resources most efficiently.

Purpose: To analyze the current state of cost-effectiveness analyses in shoulder surgery.

Study Design: Systematic review; Level of evidence, 4.

Methods: A systematic review of the current literature was performed following PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines. All full economic analyses published since January 1, 2010 and including the terms “cost analysis” and “shoulder” were checked for usability. The methodological quality of the studies was assessed using the Oxford Centre for Evidence-Based Medicine levels of evidence and established health economic criteria (Quality of Health Economic Studies [QHES] instrument).

Results: A total of 34 studies fulfilled the inclusion criteria. Compared with older studies, recent studies were of better quality: one level 1 study and eight level 2 studies were included. The mean QHES score was 87 of 100. The thematic focus of most studies ($n = 13$) was rotator cuff tears, with the main findings as follows: (1) magnetic resonance imaging is a cost-effective imaging strategy, (2) primary (arthroscopic) rotator cuff repair (RCR) with conversion to reverse total shoulder arthroplasty in case of failure is the most cost-effective strategy, (3) the platelet-rich plasma augmentation of RCR seems not to be cost-effective, and (4) the cost-effectiveness of double-row RCR remains unclear. Other studies included shoulder instability ($n = 3$), glenohumeral osteoarthritis ($n = 3$), proximal humeral fractures ($n = 4$), subacromial impingement ($n = 4$), and other shoulder conditions ($n = 7$).

Conclusion: Compared with prior studies, the quality of recently available studies has improved significantly. Current studies could help decision makers to appropriately and adequately allocate resources. The optimal use of financial resources will be of increasing importance to improve medical care for patients. However, further studies are still necessary.

Keywords: economic evaluation; cost-effectiveness; rotator cuff; instability

*Address correspondence to Thomas Tischer, Prof, MD, MBA, Department of Orthopaedic Surgery, University Medicine Rostock, Doberaner Strasse 142, 18057 Rostock, Germany (email: thomas.tischer@med.uni-rostock.de).

[†]Department of Orthopaedic Surgery, University Medicine Rostock, Rostock, Germany.

[‡]Berlin School of Economics and Law, Berlin, Germany.

Final revision submitted December 3, 2019; accepted January 21, 2020.

One or more of the authors has declared the following potential conflict of interest or source of funding: T.T. is a paid consultant for Arthrex and Bauerfeind. AOSSM checks author disclosures against the Open Payments Database (OPD). AOSSM has not conducted an independent investigation on the OPD and disclaims any liability or responsibility relating thereto.

The Orthopaedic Journal of Sports Medicine, 8(5), 2325967120917121

DOI: 10.1177/2325967120917121

© The Author(s) 2020

For a long time, economic evaluations including cost analysis studies were uncommon in medicine, especially in orthopaedic surgery. Because of the increasing importance of financial factors, the efficient use of available resources has become more important. In 2017, the United States spent 17.2% of its gross domestic product in the health care sector, much more than any other country (eg, Germany: 11.3%; Japan: 10.7%) and much more than in 2000, when only 12.5% of the United States gross domestic product was spent in the health care sector.⁴⁴

The purpose of economic evaluations is to analyze alternative courses of action (eg, different treatments) in terms of both their costs and consequences (eg, patient outcomes).¹⁵ The aim should be to increase value for the patients, with “value” defined as the health outcome per dollar spent.⁴³ The outcome is not only the achieved health status but also the process of regeneration and the achievement of sustainable health. If a more expensive treatment

TABLE 1
Overview and Explanations of Abbreviations, Terms, and Concepts^a

Cost-effective threshold	Value that a society is willing to pay for health. Used as a rough guide to help determine whether a particular investment constitutes a reasonable value. Varies from country to country. Referencing a threshold of US\$50,000/QALY has, in practice, implied adding new favorable interventions but without displacing any unfavorable interventions. An increase of the threshold from \$25,000 to \$100,000 results in more money needed in the health care system.
Discounting	Economic evaluation practice of weighting future gains and losses in health care less heavily than those in the present.
EQ-5D	Standardized instrument for measuring generic health status.
ICER	Outline of the incremental cost associated with 1 additional unit of the measure of effect. $ICER = \Delta \text{cost} / \Delta \text{health gain}$ (eg, life years, shoulder function).
Markov model	Used to show how a hypothetical cohort of patients moves between different health states over time. The model simulation ends when all patients in the cohort are “dead” to compare long-term health and cost outcomes.
Monte Carlo simulation	Method for the calculation of cost-effectiveness ratios and uncertainties involved in the analysis. Includes the mean value and variance of cost-effectiveness ratio, probability distribution function, skewness, and cumulative frequency distribution.
Probabilistic sensitivity analysis	Demonstrates the parameter uncertainty in a decision problem. Involves sampling parameters from their respective distributions (rather than simply using mean/median values).
QHES	Validated questionnaire to assess the quality of economic studies with 16 criteria (maximum score = 100). Values of 80-100 points are generally attributed to high-quality studies, and studies <50 points are considered not worthy of publication.
QALY	Routinely used as a summary measure of health outcomes for economic evaluations, which incorporates the impact on both the quantity and the quality of life. $QALY = T_1Q_1 - T_0Q_0$ (with “T” being survival year, “Q” being health status, “1” being perfect health status, and “0” being death).
Sensitivity analysis	Cost-effectiveness analysis is one of the main tools of economic evaluations. Every cost-effectiveness analysis is based on a number of assumptions, some of which may not be accurate, introducing uncertainty. Sensitivity analysis formalizes ways to measure and evaluate this uncertainty.
SF-6D	Evaluation tool of health status used in economics as a variable in the QALY calculation to determine the cost-effectiveness of a treatment.

^aEQ-5D, EuroQol 5 Dimensions; ICER, incremental cost-effectiveness ratio; QALY, quality-adjusted life year; QHES, Quality of Health Economic Studies; SF-6D, Short Form-6 Dimensions.

method leads to faster regeneration and earlier return to work, then this method can still be cost-effective.

Cost analysis can be performed using different methods, with cost-effectiveness analysis, cost-utility analysis, and cost-benefit analysis being most common.^{4,15} As a measure of outcome, the use of quality-adjusted life years (QALYs) has been widely established.^{4,50} QALYs facilitate the comparison of different treatment options, and incremental cost analysis can be performed (incremental cost-effectiveness ratio [ICER]). QALYs are calculated using generic quality of health measurement tools such as the EuroQol 5 Dimensions index (EQ-5D) or Short Form-6 Dimensions (SF-6D).^{5,56} However, better measurement tools for the “outcome” have not yet been established, and suitable instruments that are more specific than the EQ-5D/SF-6D but still allow comparisons between different diseases are still under development.^{40,49} Investigators should study and report evidence-based results from validated, disease-specific, and patient-reported outcome measures as well as health-related quality of life measures.

By defining the threshold of what a QALY (1 year in perfect health, usually US\$50,000-\$100,000/QALY) may cost, resources can be better distributed among different treatment strategies.^{37,39} For example, it would be more useful to support a therapy that increased quality of life by 2.0 QALYs for the cost of \$50,000 than a similar therapy

that also cost \$50,000 but increased the QALY only by 1.0. For mathematical calculations, different models (eg, Markov, Monte Carlo) are used and various statistical methods applied to compensate for uncertainty (eg, probabilistic sensitivity analysis, 1- or 2-way sensitivity analysis) as well as the future inflation of costs (eg, discounting). See Table 1 for a brief explanation of key concepts.

Multiple cost analysis studies and reviews have been published in sports orthopaedics over the past years, especially focusing on anterior cruciate ligament reconstruction.^{41,53} In shoulder surgery, cost analysis studies have rarely been performed or are of lower quality, as stated in a review from 2012 that included literature before 2010.²⁷ Therefore, the aim of this work was to analyze the current state of cost-effectiveness analyses in shoulder surgery since 2010.

METHODS

This systematic review was performed in accordance with the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines.³⁶ A systematic electronic search of the PubMed database was conducted in November 2019 to identify clinical studies investigating cost analysis in shoulder surgery. The following search terms were used in all fields: (cost analysis AND shoulder).

Studies were limited to articles published after December 2009. In addition to the electronic search, the reference lists of all included articles were manually searched for additional relevant articles.

For the purpose of this systematic review, only clinical studies that specifically investigated full economic analyses (defined by comparing ≥ 2 alternative treatments and including both costs and effects) were included. Additional inclusion criteria were as follows: English-language studies, studies published online, or studies in print in a peer-reviewed journal. The exclusion criteria were as follows: narrative or systematic reviews, meeting abstracts, and proceedings.

The titles and abstracts of all articles were screened for relevance according to the inclusion and exclusion criteria. If no abstract was available, the full-text article was obtained to assess the relevance of the study. The full text was obtained and subsequently reviewed for possible inclusion in the systematic review for all articles that were not excluded during the initial screening process.

Quality Assessment

All included studies were assessed and assigned a level of evidence (level 1-5) according to the Oxford Centre for Evidence-Based Medicine (OCEBM) criteria for economic studies.³² Level 1 and 2 studies include complex computer simulation models (Monte Carlo simulation, Markov model) with inputs derived solely from level 1 or 2 studies.³² Additionally, the Quality of Health Economic Studies (QHES) instrument was used to assess the quality of the included studies as used in other studies in orthopaedic surgery.^{41,43} There were 2 authors (T.T. and C.L.) who independently performed the evaluation, and a consensus was reached by a discussion. The QHES instrument consists of 16 “yes” or “no” questions, with each question weighted between 1 and 9 points and a total score range from 0 to 100. Scores in the range of 80 to 100 are generally considered high quality, whereas scores below 50 are typically not deemed worthy of publication.⁴³

RESULTS

In the scope of this systematic literature review, 406 articles published after the end of 2009 were found that reported on cost analysis and the shoulder (Figure 1). After the exclusion criteria were applied, 34 studies that performed full economic analyses were included (Table 2). The detailed results are displayed in individual tables: 13 studies considered the topic of rotator cuff tears (RCTs) (Table 3), 3 considered shoulder instability (Appendix Table A1), 3 considered glenohumeral osteoarthritis (Appendix Table A2), 4 considered proximal humeral fractures (Appendix Table A3), 4 considered subacromial impingement (Appendix Table A4), and 7 considered other shoulder conditions (Appendix Table A5). The high-quality studies included in this systematic review show cost-effective therapeutic strategies.




Figure 1. Flowchart of the search strategy in accordance with PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines.

Quality Analysis

The quality of the studies included in this systematic review was higher compared with studies before 2010. In total, there was one level 1 study, eight level 2 studies, nine level 3 studies, nine level 4 studies, and seven level 5 studies. The eight level 2 studies used complex Markov models and appropriate sensitivity analyses (Monte Carlo simulation, probabilistic sensitivity analysis). Levels of evidence and QHES scores are shown in Table 2. Before 2010, sensitivity analyses and discounting were rarely used in the shoulder literature, whereas these methods have now become common in the included studies.²⁷ Table 2 shows the country in which the economic analysis was performed and which models and sensitivity analyses were used. Another important criterion for a high-quality study is whether the aim of the study is clear, specific, and measurable.⁸ This criterion was fulfilled by all the included studies. The quality of the data on which the models were based was the most difficult aspect to evaluate. In all except 1 presented study,⁵⁷ expert opinion was included in various amounts, as opposed to data. Therefore, only 1 level 1 study according to the OCEBM criteria could be included in this review. The mean QHES score showed that the overall study quality/methodology was good, with a mean score of 87 (range, 38-100) (Table 2). Figure 2 presents information regarding how each QHES criterion was met by the 34 studies.

Cost Analysis in RCTs

In total, there were 13 studies focusing on RCTs. Only 1 study fulfilled the criteria of a level of evidence 2 study, while 7 studies were level 3 (Table 3). The main findings were the following: (1) Although additional factors must be considered, such as available resources and workflow,

TABLE 2
Characteristics of Included Studies^a

Author (Year)	Journal (Country)	Primary Outcome	Data Source	Study Perspective	QALY Threshold	Methods	Sensitivity Analysis	Time Frame (Discounting)	Level of Evidence ³²	QHES Score
Rotator cuff tear (n = 13)										
Nicholson ³⁸ (2019)	<i>Bone Joint J</i> (UK)	ICER	n = 92	NHS	£20,000		Yes	No	4	87
Castagna ⁷ (2019)	<i>Int Orthop</i> (Italy)	ICER	Literature review and experts	Health care	\$50,000	Decision analytic model	1-/2-way	2 y (no information)	4	73
Gyftopoulos ¹⁹ (2017)	<i>J Shoulder Elbow Surg</i> (USA)	ICER	Literature review and experts	Health care	\$100,000	Decision analytic model	1-/2-way	2 y (3%/y)	3	99
Dornan ¹³ (2017)	<i>Arthroscopy</i> (USA)	ICER	Literature review and experts	Health care	\$50,000	Markov decision model	1-/2-way	Lifetime (3%/y)	3	95
Huang ²³ (2017)	<i>J Bone Joint Surg Am</i> (Canada)	ICER	Literature review and experts	Health care	\$50,000	Decision analytic model	1-way, PSA	2 y (5%/y)	4	93
Kang ²⁶ (2017)	<i>Orthopedics</i> (USA)	ICER	Literature review and experts	Health care	\$50,000	Markov decision model	1-/2-way	Lifetime (3%/y)	3	99
Makhni ³⁰ (2016)	<i>Arthroscopy</i> (USA)	ICER	Literature review and experts	Health care	\$50,000 and \$100,000	Decision analytic model	1-/2-way	Lifetime (—)	3	99
Samuelson ⁵⁴ (2016)	<i>Arthroscopy</i> (USA)	ICER	Literature review and experts	Health care	\$50,000	Markov decision model	1-way	10 y (3%/y)	3	99
Carr ⁶ (2015)	<i>Health Technol Assess</i> (UK)	ICER/OSS score	n = 273	Health care	£20,000-£30,000	Trial	Adjusting for covariates	2 y (—)	4	92
Mather ³³ (2013)	<i>J Bone Joint Surg Am</i> (USA)	ICER	Literature review and experts	Society	\$50,000	Markov decision model	1-/2-/3-way, PSA	Lifetime (3%/y)	2	99
Renfree ⁵¹ (2013)	<i>J Shoulder Elbow Surg</i> (USA)	Cost/QALY	n = 30	Health care	\$50,000	Trial	No	2 y (3%/y)	5	38
Coe ⁹ (2012)	<i>J Shoulder Elbow Surg</i> (USA)	ICER	Literature review and experts	Society and payer	\$100,000	Markov decision model	1-way	Lifetime (3%/y)	3	96
Genuario ¹⁶ (2012)	<i>J Bone Joint Surg Am</i> (USA)	ICER	Literature review	Society	\$100,000	Decision analytic model	1-/2-way	Lifetime (3%/y)	3	99
Shoulder instability (n = 3)										
Min ³⁵ (2018)	<i>J Shoulder Elbow Surg</i> (USA)	ICER	Literature review/trial	Health care	<\$57,300	Markov decision model	Multivariate, PSA	Lifetime (no information)	2	85
Makhni ²⁹ (2016)	<i>Arthroscopy</i> (USA)	ICER	Literature review and experts	Health care	\$50,000 and \$100,000	Decision analytic model	1-/2-way	Lifetime (no information)	4	84
Crall ¹¹ (2012)	<i>Arthroscopy</i> (USA)	Cost/QALY	Literature review and experts	Payer	\$25,000	Markov decision model	Microsimulation, PSA	15 y (5%/y)	2	100
Osteoarthritis (n = 3)										
Bhat ³ (2016)	<i>Clin Orthop Relat Res</i> (USA)	ICER	Literature review and experts	Health care	\$50,000	Markov decision model	1-way, PSA	Lifetime (no information)	2	92

(continued)

Table 2 (continued)

Author (Year)	Journal (Country)	Primary Outcome	Data Source	Study Perspective	QALY Threshold	Methods	Sensitivity Analysis	Time Frame (Discounting)	Level of Evidence ³²	QHES Score
Bachman ² (2016)	<i>World J Orthop</i> (USA)	Cost/QALY	n = 15 vs study with 224 THA	Health care	\$30,000-\$50,000	Markov decision model	No	Lifetime (3%/y)	5	71
Mather ³⁴ (2010)	<i>J Shoulder Elbow Surg</i> (USA)	ICER	Literature review and experts	Society	\$50,000	Markov decision model	Multivariate	Lifetime (3%/y)	2	99
Proximal humeral fracture (n = 4)										
Osterhoff ⁴⁵ (2017)	<i>Value Health</i> (Canada)	ICER	Literature review and experts	Health care (single payer)	Can\$50,000	Decision tree and Markov modeling	1-/2-way, PSA	Lifetime (5%/y)	2	99
Nwachukwu ⁴² (2016)	<i>J Shoulder Elbow Surg</i> (USA)	ICER	Literature review and experts	US payers and hospitals	\$100,000	Markov decision model	Deterministic sensitivity analysis, PSA	Lifetime (3%/y)	2	99
Corbacho ¹⁰ (2016)/ Handoll ²⁰ (2015)/ Handoll ²¹ (2017) ^b	<i>Bone Joint J (UK)/ Health Technol Assess (UK)/ Bone Joint J (UK)</i>	Cost/QALY	n = 250	NHS	£20,000-£30,000	Trial	Complete case analysis	2 y (3.5%/y)	4	92
Subacromial impingement (n = 4)										
Rombach ⁵² (2019)	<i>Bone Joint J (UK)</i>	QALY/ICER	n = 313	Health care	£20,000	Trial	Yes	12 mo (—)	2	92
Arias-Buria ¹ (2018)	<i>Pain Med</i> (Spain)	QALY/ICER	n = 50	Society	—	Trial	1-way	—	5	89
Marks ⁵¹ (2016)	<i>PLoS One</i> (Australia)	SPADI score	n = 64	Health funder	AUD\$50,000	Trial	1-way	12 wk (—)	5	78
Jowett ²⁵ (2013)	<i>Rheumatology (Oxford)</i> (UK)	ICER	n = 232	Health care	£20,000	Trial	Nonparametric bootstrapping	24 wk (—)	5	87
Other (n = 7)										
Sorensen ⁵⁷ (2020)	<i>J Shoulder Elbow Surg</i> (Denmark)	QALY/ICER	Literature review and experts	Health sector	€34,000	Decision analytic model	1-way	12 mo (—)	1	84
Gyftopoulos ¹⁸ (2018)	<i>AJR Am J Roentgenol</i> (USA)	ICER	Literature review and experts	Health care	\$100,000	Decision analytic model	1-/2-way, PSA	6 mo (no information)	4	81
Paoli ⁴⁶ (2018)	<i>Arthroscopy</i> (USA)	QALY/ICER	Literature review and experts	Society	\$100,000	Decision tree and Markov modeling	1-/2-way, PSA	10 y (—)	3	99
Hatch ²² (2017)	<i>J Shoulder Elbow Surg</i> (USA)	Cost-effectiveness	n = 16	Health care	No	Breakeven analysis	—	—	4	51
Scott ⁵⁵ (2015)	<i>Orthop J Sports Med</i> (USA)	Cost/case because of adverse outcome	Literature review and experts	Society	—	Decision analytic model	Monte Carlo simulation	—	5	75
Dattani ¹² (2013)	<i>Bone Joint J (UK)</i>	Cost/QALY	n = 100 and NHS reference costs	Health care	£30,000	Trial	No	6 mo (—)	5	57
Pearson ⁴⁷ (2010)	<i>J Orthop Trauma</i> (Canada)	ICER	n = 132	Health care	\$50,000	Trial	Yes	25.8 y (3%/y)	3	99

^aICER, incremental cost-effectiveness ratio; NHS, National Health Service; OSS, Oxford Shoulder Score; PSA, probabilistic sensitivity analysis; QALY, quality-adjusted life year; QHES, Quality of Health Economic Studies; SPADI, Shoulder Pain and Disability Index; THA, total hip arthroplasty; —, no information provided.

^bAll 3 studies were related to the PROFHER (PROximal Fracture of the Humerus: Evaluation by Randomisation) trial. The two Handoll et al studies were considered a single study for the purposes of this review.

TABLE 3
Economic Evaluations Concerning RCTs (n = 13)^a

Author (Year)	Purpose	Groups	Results	Conclusions
Nicholson ³⁸ (2019)	Determine cost-effectiveness of arthroscopic RCR over 2-y period and if age adversely affects outcome/cost-effectiveness	(1) Arthroscopic RCR	No significant difference between age <65 or >65 y regarding postoperative shoulder function or EQ-5D score; total mean cost per patient was £3646.94, and mean EQ-5D difference at 1 y was 0.2691, resulting in mean ICER of £13,552.36/QALY; smokers had ICER that was 4 times more expensive	Arthroscopic RCR resulted in excellent patient satisfaction and cost-effectiveness, regardless of age
Castagna ⁷ (2019)	Determine cost-effectiveness of treatment for irreparable RCTs	(1) Subacromial spacer, (2) RCR, (3) TSA, (4) nonoperative treatment	Subacromial spacer cost less and increased effectiveness by 0.06 and 0.10 QALYs, respectively; nonoperative treatment least costly; subacromial spacer gained 0.05 QALYs for additional cost of €522 (ICER, €10,440/QALY)	Subacromial spacer likely to be safe, effective, and cost-effective option for massive irreparable RCTs
Gyftopoulos ¹⁹ (2017)	Determine imaging strategies for full-thickness supraspinatus tears	(1) MRI, (2) ultrasound, (3) ultrasound followed by MRI	Ultrasound least costly strategy (\$1385); MRI most effective (1.332 QALYs; ICER, \$22,756/QALY)	Ultrasound most cost-effective strategy; MRI was preferred strategy based on cost-effectiveness criteria
Dornan ¹³ (2017)	Determine cost-effective treatment for massive RCTs and pseudoparalysis without osteoarthritis	(1) RCR (revised once), (2) RCR (conversion to RTSA in case of failure), (3) primary RTSA	Primary RTSA was cost-effective when utility of RTSA exceeded that of RCR by 0.04 QALYs/y	Primary arthroscopic RCR with conversion to RTSA in case of failure most cost-effective strategy
Huang ²³ (2017)	Determine cost-effectiveness of single row and double row in arthroscopic RCR	(1) Single-row repair, (2) double-row repair	Double-row fixation costlier (\$2134.41 vs \$1654.76, respectively) but more effective than single-row repair (4.073 vs 4.055 QALYs, respectively); ICER of \$26,666.75/QALY for double-row repair	Double-row fixation more cost-effective, especially for larger RCTs (3 cm)
Kang ²⁶ (2017)	Compare different treatments for elderly patients with massive irreparable RCTs	(1) RTSA, (2) HA, (3) arthroscopic decompression/biceps tenodesis, (4) physical therapy	RTSA yielded most QALYs with 7.69, but greater benefits came at higher costs; health utility of RTSA was ≤0.72 (QALY, 7.48), or RTSA probability of no complications was ≤0.83 (QALY of 7.48 at cost of \$23,830)	RTSA considered good value for money compared to other treatments; RTSA most cost-effective treatment in elderly with massive RCTs
Makhni ³⁰ (2016)	Compare cost-effectiveness of arthroscopic RCR to RTSA in patients with symptomatic large/massive RCTs without osteoarthritis	(1) Continued nonoperative treatment and/or observation, (2) primary arthroscopic RCR, (3) primary RTSA	Arthroscopic RCR and RTSA superior to nonoperative care (ICER, \$15,500/QALY and \$37,400/QALY, respectively); arthroscopic RCR dominant over primary RTSA; arthroscopic RCR preferred strategy as long as progression rate to end-stage RCT arthropathy <89%	Arthroscopic RCR, despite high rates of tendon retearing in large/massive RCTs, may be more cost-effective initial treatment strategy compared to RTSA

(continued)

Table 3 (continued)

Author (Year)	Purpose	Groups	Results	Conclusions
Samuelson ⁵⁴ (2016)	Determine if use of PRP products during arthroscopic RCR is cost-effective	(1) With PRP, (2) without PRP	Cost of RCR with and without PRP was \$6775/QALY and \$6612/QALY, respectively; use of PRP to augment RCR not cost-effective; to achieve willingness-to-pay threshold of \$50,000/QALY, addition of PRP would need to be associated with 9.1% reduction in retear rate	Current use of PRP to augment RCR was not cost-effective
Carr ⁶ (2015)	Evaluate clinical effectiveness and cost-effectiveness of arthroscopic and open RCR	(1) Arthroscopic RCR, (2) open RCR	No differences in mean cost; overall treatment cost at 2 y was £2567 ± £176 for arthroscopic surgery and £2699 ± £149 for open surgery; there was difference in total initial procedure-related costs, with arthroscopic repair being costlier; retear rates not different	In patients >50 y with degenerative RCTs, no difference in clinical effectiveness or cost-effectiveness between open repair and arthroscopic repair at 2 y
Mather ³³ (2013)	Examine value of surgical treatment for full-thickness RCTs from societal perspective	(1) Different age groups in RCR	Mean age-weighted total societal savings from RCR compared with nonoperative treatment was \$13,771 over patient's lifetime; savings ranged from \$77,662 (age 30-39 y) to net cost to society of \$11,997 (age 70-79 y); surgical treatment resulted in mean improvement of 0.62 QALYs; estimated lifetime societal savings of \$3.44 billion for ~250,000 RCR procedures performed yearly in US	RCR for full-thickness tears produced net societal cost savings for patients <61 y and greater QALYs for all patients; RCR was cost-effective for all populations
Renfree ⁵¹ (2013)	Prospectively analyze outcomes and costs for primary RTSA	(1) RTSA	Clinical and functional outcomes demonstrated significant improvement; median QALYs improved from 6.56 preoperatively to 7.58 at 2-y follow-up; increase in QALYs calculated from EQ-5D was greater (6.21 to 8.10); mean cost was \$21,536; cost utility at 2 y was \$26,920/QALY by SF-6D and \$16,747/QALY by EQ-5D	EQ-5D and SF-36 (from which the SF-6D is calculated) results demonstrated modestly cost-effective (<\$50,000/QALY) improvement for RCT arthropathy after primary RTSA
Coe ⁹ (2012)	Compare cost-effectiveness of RTSA to HA	(1) RTSA, (2) HA	RTSA could be cost-effective strategy for treatment of RCT arthropathy	RTSA could be cost-effective alternative to HA for RCT arthropathy; cost-effectiveness of RTSA depended on health utility gain, complications, and cost of implant

(continued)

Table 3 (continued)

Author (Year)	Purpose	Groups	Results	Conclusions
Genuario ¹⁶ (2012)	Compare cost-effectiveness of double-row to single-row arthroscopic RCR	(1) Double-row repair, (2) single-row repair	ICER for double-row vs single-row arthroscopic RCR was \$571,500 for RCTs <3 cm and \$460,200 for RCTs of 3 cm; rate of radiographic or symptomatic retears alone had no influence on cost-effectiveness; if increase in cost of double-row repair <\$287 (small or moderate tears) and <\$352 (large or massive tears) compared with single-row repair, double-row repair would represent cost-effective surgical alternative	Double-row RCR not cost-effective for any sized RCTs; variability in costs and probability of retears could have profound effect on results and might create environment in which double-row repair becomes more cost-effective option

^aEQ-5D, EuroQol 5 Dimensions index; HA, hemiarthroplasty; ICER, incremental cost-effectiveness ratio; MRI, magnetic resonance imaging; PRP, platelet-rich plasma; QALY, quality-adjusted life year; RCR, rotator cuff repair; RCT, rotator cuff tear; RTSA, reverse total shoulder arthroplasty; SF-36, 36-Item Short Form Survey; SF-6D, Short Form-6 Dimensions; TSA, total shoulder arthroplasty.




Figure 2. Breakdown of the overall Quality of Health Economic Studies instrument categories for all 34 included studies. The point value for each criterion is shown in parentheses.

magnetic resonance imaging is the first-choice imaging strategy based on cost-effectiveness criteria.¹⁹ (2) Primary (arthroscopic or open) rotator cuff repair (RCR) is the most cost-effective strategy.^{13,30,33} In case of failure, conversion to reverse total shoulder arthroplasty (RTSA) seems to be a cost-effective strategy.¹³ Primarily, RTSA is the most cost-effective treatment option for the elderly (age not specifically defined).²⁶ (3) Platelet-rich plasma as a biological augmentation of RCR seems not to be cost-effective.⁵⁴ (4) The cost-effectiveness of double-row RCR remains unclear. While it does not seem to be cost-effective for any sized RCTs, the variability in costs and probability of retears might have a profound effect on the results and may create an environment in which double-row repair becomes the more cost-effective option.¹⁶ The findings of the five level

4 and 5 studies focusing on the treatment of RCTs can be found in Table 3.

Cost Analysis in Other Abnormalities

Cost analysis for the treatment of shoulder instability showed the cost-effectiveness of both arthroscopic Bankart repair and the open Latarjet procedure in comparison with nonoperative treatment.^{11,35} The Latarjet procedure was more cost-effective than revision arthroscopic Bankart repair.²⁹ Total shoulder arthroplasty was found to be the most cost-effective treatment option for glenohumeral arthritis.^{2,3,34} Regarding proximal humeral fractures, RTSA showed economic advantages over hemiarthroplasty in the elderly (base case of 70 and 72 years) and in complex

fracture situations.^{42,45} A prophylactic administration of local vancomycin powder during arthroplasty was found to be highly cost-effective.²² For the treatment of subacromial impingement, corticosteroid injections plus exercise delivered by a therapist were cost-effective when compared with exercise alone.²⁵ Furthermore, ultrasound-guided injections were found to be the most cost-effective option for the initial steroid-based treatment of adhesive capsulitis.¹⁸ A 2019 level 1 study on the cost-effectiveness of surgical treatment for displaced midshaft clavicular fractures showed that surgical treatment was not cost-effective (considering a threshold of €34,000/QALY).⁵⁷ In patients with professions imposing high loads on the shoulder, however, surgical treatment was described to potentially be more cost-effective compared with nonsurgical treatment.⁵⁷ Paoli et al⁴⁶ compared superior labral from anterior to posterior (SLAP) repair, biceps tenodesis, and nonoperative treatment in their cost-effectiveness study on symptomatic SLAP tears. Primary biceps tenodesis was described to be the most cost-effective treatment option among the 3 evaluated procedures for type II SLAP tears in middle-aged patients.⁴⁶ Detailed findings of the analyzed studies can be found in Appendix Tables A1 to A5.

DISCUSSION

Within the scope of this systematic review, 34 full economic analyses concerning shoulder abnormalities were included that have been published more recently than 2010, expanding the older reviews by Kuye et al²⁷ and Teusink et al.⁵⁸ In contrast to these reviews, only full economic analyses were included. Simple comparisons of costs or cost-minimization studies were disregarded. Most available studies had the perspective of the health care sector. The overall quality of recent studies has significantly improved compared with studies from 2009 and before. The results of cost-effectiveness studies are rarely transferred to daily clinical practice; in most health care systems, medical treatment is customized to the individual patient by doctors, mostly independent of costs. However, exploding costs in health care systems will inevitably lead to an emphasis on more cost-effective therapies in the future.

Representing almost 50% of all included studies, sufficient evidence now exists for the cost-effectiveness of various RCT treatment options. Magnetic resonance imaging was described to be a cost-effective imaging strategy. Primary RCR with conversion to RTSA in case of failure was the most cost-effective strategy. Platelet-rich plasma augmentation of RCR was not cost-effective, and the cost-effectiveness of double-row RCR remained unclear. For displaced proximal humeral fractures (involving the surgical neck), a randomized trial comparing surgical and nonsurgical treatment found no significant between-group differences in quality of life over time.²¹ For most other abnormalities such as shoulder instability, glenohumeral osteoarthritis, or subacromial impingement, fewer high-quality studies have been published so far.

Despite several high-level studies that recently complemented the available body of literature, uncertainty

remains regarding the most cost-effective treatment option for older patients with large or massive RCTs. Makhni et al³⁰ in 2016 analyzed the cost-effectiveness of primary RTSA versus arthroscopic RCR in patients older than 65 years. Although clinical outcomes seemed to be similar in both treatment groups, the authors highlighted the fact that the overall costs were significantly lower in the RCR treatment group, even in cases of postsurgical retears. The authors emphasized this finding by various sensitivity analyses showing that patients with retears would require progression rates to end-stage RCT arthropathy of 89% to justify early RTSA. Similarly, considering a 100% progression rate to RCT arthropathy, the authors calculated a required retear rate of 65% after arthroscopic RCR of RCTs to justify early RTSA. For newer techniques, such as superior capsular reconstruction, no cost-effective analyses are available yet.

Cost analysis on surgical techniques in arthroscopic RCR also revealed varying results. While Genuario et al¹⁶ reported in their analysis on the cost-effectiveness of single-row RCR versus double-row RCR that double-row RCR was not cost-effective for any sized RCTs, Huang et al²³ found double-row RCR to be more cost-effective than single-row RCR in some cases, especially in larger RCTs. This discrepancy most likely results from the varying underlying costs of surgery that were used for calculations, depending on the health care system. While Huang et al based their calculations on surgery costs of \$1654.76 (single-row RCR) and \$2134.41 (double-row RCR) for primary and revision surgery, respectively, Genuario et al referred to surgery costs being 5 to 7 times higher. This difference strongly influences the ICER per QALY and therefore makes a direct comparison of both studies difficult.

Mather et al³³ investigated whether the operative treatment of RCTs is cost-effective from a societal perspective. A complex Markov model was therefore used based on lifelong direct and indirect costs in connection with RCTs. The surgical treatment of RCTs led to cost savings compared with nonoperative treatment depending on the patient's age at the time of treatment (cost savings of \$77,662 in 30- to 39-year-old patients). The model was validated using 1-, 2-, and 3-way sensitivity analyses and Monte Carlo simulations. The response to nonoperative treatment was found to have the biggest impact on the results. If the success rate of nonoperative therapy was increased to 88%, it would be more cost-effective. However, this high success rate is outside of the reported outcome rates for nonoperative therapy. The study by Mather et al³³ emphasizes the complexity of these models, which factors are to be included, and how errors can be minimized using simulations. However, the most important aspect is the data on which the model is based, as highlighted by Castagna et al.⁷ The authors evaluated the subacromial spacer as a new treatment option, but no valid outcome data are available for this treatment method yet.⁷ Therefore, the model used in the study was based on expert opinion and was prone to errors, regardless of the quality level of the study.⁷ The performed sensitivity analysis showed that if the success rate of the subacromial spacer is only reduced by 20%, it would no longer be the

TABLE 4
Different Therapy Options for Various Diseases and Their Economic Evaluation^a

Author (Year)	Treatment	ICER
Huang ²³ (2017)	Double-row vs single-row RCR	\$26,666.75/QALY gained for double-row repair relative to single-row fixation
Makhni ³⁰ (2016)	RTSA vs arthroscopic RCR for symptomatic large and massive rotator cuff tears	\$15,500/QALY (arthroscopic RCR vs nonoperative) and \$37,400/QALY (RTSA vs nonoperative)
Dougherty ¹⁴ (2013)	Arthroscopic RCR	\$13,092/QALY
Dougherty ¹⁴ (2013)	Total knee replacement	\$18,300/QALY
Dougherty ¹⁴ (2013)	Revision total knee replacement	\$31,341/QALY
Lubowitz ²⁸ (2011)	Arthroscopic ACL reconstruction	\$10,326/QALY
Gottlob ¹⁷ (1999)	ACL reconstruction	\$5857/QALY
Huygens ²⁴ (2018)	Transcatheter aortic valve implantation vs standard treatment in inoperable or high-risk operable patients	ICER, €18,421-€120,779
van Asten ⁵⁹ (2018)	Bevacizumab, ranibizumab, and aflibercept for treatment of age-related macular degeneration	ICER, \$278,099/QALY when aflibercept used instead of bevacizumab

^aACL, anterior cruciate ligament; ICER, incremental cost-effectiveness ratio; QALY, quality-adjusted life year; RCR, rotator cuff repair; RTSA, reverse total shoulder arthroplasty.

most cost-effective strategy. This accentuates the immense importance of exact outcome measurements for economic evaluations.^{48,49}

The only study focusing on the prevention of infections after shoulder arthroplasty proved the cost-effectiveness of local vancomycin powder.²² However, these results must be evaluated critically, as only 16 patients were included in a single-center study setup.

Attention should also be paid to the interpretation of the study by Paoli et al,⁴⁶ who focused on the cost-effectiveness of 3 different treatment options for symptomatic SLAP tears in middle-aged patients (mean age, 40 years). While the authors described primary biceps tenodesis to be the most cost-effective treatment option among the 3 evaluated procedures for type II SLAP tears, it needs to be highlighted that biceps tenotomy was not evaluated.⁴⁶ This efficient surgical treatment option, however, might also be cost-effective.⁴⁶

In our study, the QHES score was high (mean, 87), but according to the OCEBM level of evidence, there was only one level 1 study. This crucial difference is based on the fact that the underlying data quality is of utmost importance for OCEBM evidence levels, while the QHES score focuses on study methodology. Thereby, the study quality is rated according to the “best available” source of data. In our opinion, good study quality cannot compensate for poor underlying data, even if they are the “best available.” The example shown for Castagna et al⁷ highlights that fact, and the discrepancy between both tools needs to be considered. In this regard, sensitivity analyses are very valuable.

Compared with other areas in medicine and orthopaedic subspecializations, shoulder surgery was comparable and magnitudes less expensive than, for example, new drugs, in which the ICER could exceed US\$100,000 (Table 4). Therefore, the development and use of suitable quality parameters will be of utmost importance in the future to measure, evaluate, and compare the quality of medical therapies. High-quality studies (level 1) will be required to generate the needed data for economic analyses. An

improved outcome (or more inexpensive therapy with the same outcome) must be proven to justify the increased costs and to compare established therapies, especially for new treatments and new medical products. Surgeons today, more than ever, have to incorporate cost-efficiency into their daily treatment strategies. Nevertheless, decisions must, of course, always be based on patient- and disease-specific factors.

Limitations

This study has several limitations. First, while having tried to assess the available literature as thoroughly as possible by using wide search terms, there is still a chance of having missed available studies. Second, the QHES instrument to assess research methodology represents a potential source of biased scoring. Therefore, two of the authors independently performed the evaluation, and a consensus was reached by a discussion. Third, even though most of the studies are based on good or excellent quality of data, some of the included articles do refer to studies with slightly lower OCEBM levels and QHES scores.

CONCLUSION

Compared with a 2012 review,²⁷ the study quality improved greatly. The current studies might help decision makers to appropriately and adequately allocate resources. However, further studies are still necessary.

REFERENCES

1. Arias-Buria JL, Martin-Saborido C, Cleland J, Koppenhaver SL, Plaza-Manzano G, Fernandez-de-Las-Penas C. Cost-effectiveness evaluation of the inclusion of dry needling into an exercise program for subacromial pain syndrome: evidence from a randomized clinical trial. *Pain Med*. 2018;19(12):2336-2347.

2. Bachman D, Nyland J, Krupp R. Reverse-total shoulder arthroplasty cost-effectiveness: a quality-adjusted life years comparison with total hip arthroplasty. *World J Orthop.* 2016;7(2):123-127.
3. Bhat SB, Lazarus M, Getz C, Williams GR Jr, Namdari S. Economic decision model suggests total shoulder arthroplasty is superior to hemiarthroplasty in young patients with end-stage shoulder arthritis. *Clin Orthop Relat Res.* 2016;474(11):2482-2492.
4. Brazier J, Ratcliffe J, Salomon JA, Tsuchiya A. *Measuring and Valuing Health Benefits for Economic Evaluation.* 2nd ed. New York: Oxford University Press; 2017.
5. Brooks R. EuroQol: the current state of play. *Health Policy.* 1996;37(1):53-72.
6. Carr AJ, Cooper CD, Campbell MK, et al. Clinical effectiveness and cost-effectiveness of open and arthroscopic rotator cuff repair [the UK Rotator Cuff Surgery (UKUFF) randomised trial]. *Health Technol Assess.* 2015;19(80):1-218.
7. Castagna A, Garofalo R, Maman E, Gray AC, Brooks EA. Comparative cost-effectiveness analysis of the subacromial spacer for irreparable and massive rotator cuff tears. *Int Orthop.* 2019;43(2):395-403.
8. Chiou CF, Hay JW, Wallace JF, et al. Development and validation of a grading system for the quality of cost-effectiveness studies. *Med Care.* 2003;41(1):32-44.
9. Coe MP, Greiwe RM, Joshi R, et al. The cost-effectiveness of reverse total shoulder arthroplasty compared with hemiarthroplasty for rotator cuff tear arthroplasty. *J Shoulder Elbow Surg.* 2012;21(10):1278-1288.
10. Corbacho B, Duarte A, Keding A, et al. Cost effectiveness of surgical versus non-surgical treatment of adults with displaced fractures of the proximal humerus: economic evaluation alongside the PROFHER trial. *Bone Joint J.* 2016;98(2):152-159.
11. Crall TS, Bishop JA, Guttman D, Kocher M, Bozic K, Lubowitz JH. Cost-effectiveness analysis of primary arthroscopic stabilization versus nonoperative treatment for first-time anterior glenohumeral dislocations. *Arthroscopy.* 2012;28(12):1755-1765.
12. Dattani R, Ramasamy V, Parker R, Patel VR. Improvement in quality of life after arthroscopic capsular release for contracture of the shoulder. *Bone Joint J.* 2013;95(7):942-946.
13. Dornan GJ, Katthagen JC, Tahal DS, et al. Cost-effectiveness of arthroscopic rotator cuff repair versus reverse total shoulder arthroplasty for the treatment of massive rotator cuff tears in patients with pseudoparalysis and nonarthritic shoulders. *Arthroscopy.* 2017;33(4):716-725.
14. Dougherty CP, Howard T. Cost-effectiveness in orthopedics: providing essential information to both physicians and health care policy makers for appropriate allocation of medical resources. *Sports Med Arthrosc Rev.* 2013;21(3):166-168.
15. Drummond MF, Sculpher MJ, Claxton K, Stoddart GL, Torrance GW. *Methods for the Economic Evaluation of Health Care Programmes.* 4th ed. New York: Oxford University Press; 2015.
16. Genuario JW, Donegan RP, Hamman D, et al. The cost-effectiveness of single-row compared with double-row arthroscopic rotator cuff repair. *J Bone Joint Surg Am.* 2012;94(15):1369-1377.
17. Gottlob CA, Baker CL Jr, Pellissier JM, Colvin L. Cost effectiveness of anterior cruciate ligament reconstruction in young adults. *Clin Orthop Relat Res.* 1999;367:272-282.
18. Gyftopoulos S, Abballe V, Virk MS, Koo J, Gold HT, Subhas N. Comparison between image-guided and landmark-based glenohumeral joint injections for the treatment of adhesive capsulitis: a cost-effectiveness study. *AJR Am J Roentgenol.* 2018;210(6):1279-1287.
19. Gyftopoulos S, Guja KE, Subhas N, Virk MS, Gold HT. Cost-effectiveness of magnetic resonance imaging versus ultrasound for the detection of symptomatic full-thickness supraspinatus tendon tears. *J Shoulder Elbow Surg.* 2017;26(12):2067-2077.
20. Handoll H, Brealey S, Rangan A, et al. The ProFHER (PROximal Fracture of the Humerus: Evaluation by Randomisation) trial: a pragmatic multicentre randomised controlled trial evaluating the clinical effectiveness and cost-effectiveness of surgical compared with non-surgical treatment for proximal fracture of the humerus in adults. *Health Technol Assess.* 2015;19(24):1-280.
21. Handoll HH, Keding A, Corbacho B, Brealey SD, Hewitt C, Rangan A. Five-year follow-up results of the PROFHER trial comparing operative and non-operative treatment of adults with a displaced fracture of the proximal humerus. *Bone Joint J.* 2017;99(3):383-392.
22. Hatch MD, Daniels SD, Glerum KM, Higgins LD. The cost effectiveness of vancomycin for preventing infections after shoulder arthroplasty: a break-even analysis. *J Shoulder Elbow Surg.* 2017;26(3):472-477.
23. Huang AL, Thavorn K, van Katwyk S, MacDonald P, Lapner P. Double-row arthroscopic rotator cuff repair is more cost-effective than single-row repair. *J Bone Joint Surg Am.* 2017;99(20):1730-1736.
24. Huygens SA, Takkenberg JJM, Rutten-van Molken M. Systematic review of model-based economic evaluations of heart valve implantations. *Eur J Health Econ.* 2018;19(2):241-255.
25. Jowett S, Crawshaw DP, Helliwell PS, Hensor EM, Hay EM, Conaghan PG. Cost-effectiveness of exercise therapy after corticosteroid injection for moderate to severe shoulder pain due to subacromial impingement syndrome: a trial-based analysis. *Rheumatology (Oxford).* 2013;52(8):1485-1491.
26. Kang JR, Sin AT, Cheung EV. Treatment of massive irreparable rotator cuff tears: a cost-effectiveness analysis. *Orthopedics.* 2017;40(1):e65-e76.
27. Kuye IO, Jain NB, Warner L, Herndon JH, Warner JJ. Economic evaluations in shoulder pathologies: a systematic review of the literature. *J Shoulder Elbow Surg.* 2012;21(3):367-375.
28. Lubowitz JH, Appleby D. Cost-effectiveness analysis of the most common orthopaedic surgery procedures: knee arthroscopy and knee anterior cruciate ligament reconstruction. *Arthroscopy.* 2011;27(10):1317-1322.
29. Makhni EC, Lamba N, Swart E, et al. Revision arthroscopic repair versus Latarjet procedure in patients with recurrent instability after initial repair attempt: a cost-effectiveness model. *Arthroscopy.* 2016;32(9):1764-1770.
30. Makhni EC, Swart E, Steinhaus ME, et al. Cost-effectiveness of reverse total shoulder arthroplasty versus arthroscopic rotator cuff repair for symptomatic large and massive rotator cuff tears. *Arthroscopy.* 2016;32(9):1771-1780.
31. Marks D, Bisset L, Comans T, et al. Increasing capacity for the treatment of common musculoskeletal problems: a non-inferiority RCT and economic analysis of corticosteroid injection for shoulder pain comparing a physiotherapist and orthopaedic surgeon. *PLoS One.* 2016;11(9):e0162679.
32. Marx RG, Wilson SM, Swiontkowski MF. Updating the assignment of levels of evidence. *J Bone Joint Surg Am.* 2015;97(1):1-2.
33. Mather RC 3rd, Koenig L, Acevedo D, et al. The societal and economic value of rotator cuff repair. *J Bone Joint Surg Am.* 2013;95(22):1993-2000.
34. Mather RC 3rd, Watters TS, Orlando LA, Bolognesi MP, Moorman CT 3rd. Cost effectiveness analysis of hemiarthroplasty and total shoulder arthroplasty. *J Shoulder Elbow Surg.* 2010;19(3):325-334.
35. Min K, Fedorka C, Solberg MJ, Shaha SH, Higgins LD. The cost-effectiveness of the arthroscopic Bankart versus open Latarjet in the treatment of primary shoulder instability. *J Shoulder Elbow Surg.* 2018;27(6 suppl):S2-S9.
36. Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: the PRISMA statement. *J Clin Epidemiol.* 2009;62(10):1006-1012.
37. Neumann PJ, Cohen JT, Weinstein MC. Updating cost-effectiveness: the curious resilience of the \$50,000-per-QALY threshold. *N Engl J Med.* 2014;371(9):796-797.
38. Nicholson JA, Searle HKC, MacDonald D, McBirnie J. Cost-effectiveness and satisfaction following arthroscopic rotator cuff repair: does age matter? *Bone Joint J.* 2019;101(7):860-866.
39. Nwachukwu BU, Bozic KJ. Updating cost effectiveness analyses in orthopedic surgery: resilience of the \$50,000 per QALY threshold. *J Arthroplasty.* 2015;30(7):1118-1120.
40. Nwachukwu BU, Hamid KS, Bozic KJ. Measuring value in orthopaedic surgery. *JBJS Rev.* 2013;1(1):1-9.

41. Nwachukwu BU, Schairer WW, Bernstein JL, Dodwell ER, Marx RG, Allen AA. Cost-effectiveness analyses in orthopaedic sports medicine: a systematic review. *Am J Sports Med.* 2015;43(6):1530-1537.
42. Nwachukwu BU, Schairer WW, McCormick F, Dines DM, Craig EV, Gulotta LV. Arthroplasty for the surgical management of complex proximal humerus fractures in the elderly: a cost-utility analysis. *J Shoulder Elbow Surg.* 2016;25(5):704-713.
43. Ofman JJ, Sullivan SD, Neumann PJ, et al. Examining the value and quality of health economic analyses: implications of utilizing the QHES. *J Manag Care Pharm.* 2003;9(1):53-61.
44. Organization for Economic Co-operation and Development. Health expenditure and financing. https://stats.oecd.org/index.aspx?DataSetCode=HEALTH_STAT. Accessed August 26, 2018.
45. Osterhoff G, O'Hara NN, D'Cruz J, et al. A cost-effectiveness analysis of reverse total shoulder arthroplasty versus hemiarthroplasty for the management of complex proximal humeral fractures in the elderly. *Value Health.* 2017;20(3):404-411.
46. Paoli AR, Gold HT, Mahure SA, et al. Treatment for symptomatic SLAP tears in middle-aged patients comparing repair, biceps tenodesis, and nonoperative approaches: a cost-effectiveness analysis. *Arthroscopy.* 2018;34(7):2019-2029.
47. Pearson AM, Tosteson AN, Koval KJ, et al. Is surgery for displaced, midshaft clavicle fractures in adults cost-effective? Results based on a multicenter randomized, controlled trial. *J Orthop Trauma.* 2010;24(7):426-433.
48. Porter ME. What is value in health care? *N Engl J Med.* 2010;363(26):2477-2481.
49. Porter ME, Larsson S, Lee TH. Standardizing patient outcomes measurement. *N Engl J Med.* 2016;374(6):504-506.
50. Rasanen P, Roine E, Sintonen H, Semberg-Kontinen V, Ryyanen OP, Roine R. Use of quality-adjusted life years for the estimation of effectiveness of health care: a systematic literature review. *Int J Technol Assess Health Care.* 2006;22(2):235-241.
51. Renfree KJ, Hattrup SJ, Chang YH. Cost utility analysis of reverse total shoulder arthroplasty. *J Shoulder Elbow Surg.* 2013;22(12):1656-1661.
52. Rombach I, Merritt N, Shirkey BA, et al. Cost-effectiveness analysis of a placebo-controlled randomized trial evaluating the effectiveness of arthroscopic subacromial decompression in patients with subacromial shoulder pain. *Bone Joint J.* 2019;101(1):55-62.
53. Saltzman BM, Cvetanovich GL, Nwachukwu BU, Mall NA, Bush-Joseph CA, Bach BR Jr. Economic analyses in anterior cruciate ligament reconstruction: a qualitative and systematic review. *Am J Sports Med.* 2016;44(5):1329-1335.
54. Samuelson EM, Odum SM, Fleischli JE. The cost-effectiveness of using platelet-rich plasma during rotator cuff repair: a Markov model analysis. *Arthroscopy.* 2016;32(7):1237-1244.
55. Scott DJ, Sherman S, Dhawan A, Cole BJ, Bach BR Jr, Mather RC 3rd. Quantifying the economic impact of provider volume through adverse events: the case of sports medicine. *Orthop J Sports Med.* 2015;3(3):2325967115574476.
56. Slobogean GP, Noonan VK, O'Brien PJ. The reliability and validity of the Disabilities of Arm, Shoulder, and Hand, EuroQol-5D, Health Utilities Index, and Short Form-6D outcome instruments in patients with proximal humeral fractures. *J Shoulder Elbow Surg.* 2010;19(3):342-348.
57. Sorensen AR, Hammeken LH, Qvist AH, Jensen SL, Ehlers LH. Operative treatment of displaced midshaft clavicular fractures is not cost-effective. *J Shoulder Elbow Surg.* 2020;29(1):27-35.
58. Teusink MJ, Virani NA, Polikandriotis JA, Frankle MA. Cost analysis in shoulder arthroplasty surgery. *Adv Orthop.* 2012;2012:692869.
59. van Asten F, Michels CTJ, Hoyng CB, et al. The cost-effectiveness of bevacizumab, ranibizumab and aflibercept for the treatment of age-related macular degeneration: a cost-effectiveness analysis from a societal perspective. *PLoS One.* 2018;13(5):e0197670.

APPENDIX

TABLE A1
Economic Evaluations Concerning Shoulder Instability (n = 3)^a

Author (Year)	Purpose	Groups	Results	Conclusions
Min ³⁵ (2018)	Analyze cost-effectiveness of arthroscopic Bankart repair and open Latarjet procedure in primary shoulder instability	(1) Arthroscopic Bankart, (2) Latarjet	Overall recurrence rate of 14% after arthroscopic Bankart and 8% after open Latarjet with equal postoperative health utility states; Monte Carlo simulation showed ICER for Bankart of \$4214 and for Latarjet of \$4681 ($P < .001$)	Arthroscopic Bankart and open Latarjet highly cost-effective; Bankart more cost-effective because of lower health utility state after failed Latarjet; Latarjet favored in certain circumstances (ie, critical glenoid bone loss); clinical decisions on case-by-case basis
Makhni ²⁹ (2016)	Examine cost-effectiveness of arthroscopic revision instability repair and Latarjet procedure in patients with recurrent instability after initial arthroscopic instability repair	(1) Nonoperative, (2) revision arthroscopic repair, (3) Latarjet	Latarjet less expensive than revision arthroscopic repair (\$13,672 vs \$15,287, respectively) with improved clinical outcomes (43.78 vs 36.76 QALYs, respectively); arthroscopic repair and Latarjet cost-effective compared with nonoperative treatment (ICER, 3082 and 1141, respectively); sensitivity analyses indicated that with high rates of stability postoperatively, along with improved clinical outcome scores, revision arthroscopic repair becomes increasingly cost-effective	Latarjet for failed instability repair was cost-effective; lower costs and improved clinical outcomes compared with revision arthroscopic instability repair; treatment algorithm must be formed by surgeon's clinical judgment

(continued)

TABLE A1 (continued)

Author (Year)	Purpose	Groups	Results	Conclusions
Crall ¹¹ (2012)	Compare cost-effectiveness of initial observation to surgery for first-time anterior shoulder dislocation	(1) Primary surgery, (2) nonoperative treatment	Primary surgery less costly and more effective for 15-y-old boys, 15-y-old girls, and 25-y-old men; for 25-y-old women and 35-y-old men and women, primary surgery more effective but more costly; however, primary surgery was still very cost-effective (cost/QALY, \$25,000); after 1 recurrence, surgery was less costly and more effective for all scenarios	Primary arthroscopic stabilization was clinically effective and cost-effective for first-time anterior shoulder dislocations; by using willingness-to-pay threshold of \$25,000/QALY, surgery more cost-effective than nonoperative treatment for majority of patients

^aAll values are reported as US dollars. ICER, incremental cost-effectiveness ratio; QALY, quality-adjusted life year.

TABLE A2
Economic Evaluations Concerning Glenohumeral Osteoarthritis (n = 3)^a

Author (Year)	Purpose	Groups	Results	Conclusions
Bhat ³ (2016)	Characterize costs, as expressed by reimbursements for episodes of acute care, and outcomes associated with arthroplasty for osteoarthritis	(1) TSA, (2) HA	Initial HA resulted in average QALY gain of 6.55, and TSA resulted in average QALY gain of 7.96; during lifetime, initial HA led to lifetime revisions of 0.4 per patient, whereas initial TSA led to lifetime revisions of 0.3; during lifetime of 5279 patients, initial HA resulted in \$25,000 per patient-associated direct reimbursements, whereas TSA resulted in \$23,700	On population level, TSA was cost-effective treatment for glenohumeral arthritis in 30- to 50-y-old patients
Bachman ² (2016)	Compare cost-effectiveness of RTSA to THA	(1) RTSA, (2) THA	Cost/QALY was \$3900 for THA and \$11,100 for RTSA; after adjusting model to only include shoulder-specific physical function subscale items, RTSA's QALY improved to 2.8 y, and its cost/QALY decreased to \$8100	Based on industry accepted standards, cost/QALY estimates supported cost-effectiveness of both RTSA and THA; although THA remained "gold standard" in improving quality of life among arthroplasty procedures, cost/QALY estimates supported growing use of RTSA to improve quality of life
Mather ³⁴ (2010)	Compare cost-effectiveness of TSA to HA	(1) TSA, (2) HA	HA had lower number of average QALYs gained at higher average cost to society and was therefore dominated by TSA in glenohumeral osteoarthritis; cost-effectiveness ratio for TSA and HA was \$957/QALY and \$1194/QALY, respectively; sensitivity analysis showed that if utility of TSA is equal to HA, or revision rate is lower than HA, TSA continues to be dominant strategy	TSA with cemented glenoid was cost-effective, with greater utility for patient at lower overall cost to payer; from perspectives of patient and payer, TSA was preferred for certain populations

^aAll values are reported as US dollars. HA, hemiarthroplasty; QALY, quality-adjusted life year; RTSA, reverse total shoulder arthroplasty; THA, total hip arthroplasty; TSA, total shoulder arthroplasty.

TABLE A3
Economic Evaluations Concerning Proximal Humeral Fractures (n = 4)^a

Author (Year)	Purpose	Groups	Results	Conclusions
Osterhoff ⁴⁵ (2017)	Compare cost-effectiveness of RTSA to HA in management of complex proximal humeral fractures	(1) RTSA, (2) HA	Incremental cost/QALY gained for RTSA was Can\$13,679; 1-way sensitivity analysis showed model to be sensitive to RTSA implant and procedural costs; probabilistic sensitivity analysis showed that 92.6% of model simulations favored RTSA	RTSA for treatment of complex proximal humeral fractures in elderly patients was economically preferred; ICER of RTSA was well below standard willingness-to-pay thresholds; cost-effectiveness similar to other highly successful orthopaedic strategies (ie, THA)
Nwachukwu ⁴² (2016)	Assess cost-effectiveness of nonoperative care, HA, and RTSA for complex proximal humeral fractures	(1) Nonoperative, (2) HA, (3) RTSA	From payer perspective, ICER of RTSA was \$8100/QALY; HA was eliminated from payer analysis as cost-ineffective strategy; from hospital perspective, HA was not cost-ineffective, and ICER was \$36,700/QALY, with RTSA providing incremental effectiveness at \$57,400/QALY; RTSA was optimal in 61% and 54% of payer and hospital probabilistic sensitivity analyses, respectively; preferred strategy depended on associated QALY gains, primary RTSA cost, and failure rates for RTSA	RTSA could be cost-effective in surgery of complex proximal humeral fractures; HA also cost-effective depending on perspective (cost-ineffective for payer but cost-effective for hospital); opportunities for increased cost-sharing strategies to alleviate cost burden on hospitals
Corbacho ¹⁰ (2016)/ Handoll ²⁰ (2015)/ Handoll ²¹ (2017) ^b	Compare surgery with no surgery for displaced proximal humeral fractures	(1) Surgery, (2) no surgery	Surgery showed mean greater costs and marginally lower QALYs than nonsurgery; surgery cost £1758 more per patient (95% CI, £1126 to £2389); total QALYs for surgery were smaller than nonsurgery (-0.0101 [95% CI, -0.13 to 0.11]); probability of surgery being cost-effective was <10%, given current NICE willingness-to-pay threshold of £20,000 for additional QALY; results were robust to sensitivity analyses	Current surgical treatment was not cost-effective for majority of displaced fractures of proximal humerus involving surgical neck in NHS

^aAll values are reported as US dollars unless otherwise specified. HA, hemiarthroplasty; ICER, incremental cost-effectiveness ratio; NHS, National Health Service; NICE, National Institute for Health and Care Excellence; QALY, quality-adjusted life year; RTSA, reverse total shoulder arthroplasty; THA, total hip arthroplasty.

^bAll 3 studies were related to the PROFHER (PROximal Fracture of the Humerus: Evaluation by Randomisation) trial. The two Handoll et al studies were considered a single study for the purposes of this review.

TABLE A4
Economic Evaluations Concerning Subacromial Impingement (n = 4)^a

Author (Year)	Purpose	Groups	Results	Conclusions
Rombach ⁵² (2019)	Assess cost-effectiveness of ASD compared with arthroscopic surgery only and with no treatment	(1) ASD, (2) arthroscopic surgery only, (3) no treatment	Statistically significant differences in cumulative QALYs and costs were found at 6 and 12 mo for ASD compared with no treatment; probabilities of ASD being cost-effective compared with no treatment at willingness-to-pay threshold of £20,000/QALY were ~0% at 6 mo and ~50% at 1 y, with probabilities potentially increasing at 2 y	No evidence that ASD was cost-effective during 1-y follow-up period; it could be cost-effective in long term

(continued)

TABLE A4 (continued)

Author (Year)	Purpose	Groups	Results	Conclusions
Arias-Buria ¹ (2018)	Evaluate cost-effectiveness of addition of TrP-DN to exercise program in subacromial pain syndrome	(1) Exercise alone, (2) exercise plus TrP-DN	Those in exercise group made more visits to medical doctors and underwent greater number of other treatments; combination of exercise plus TrP-DN was less costly (mean difference in cost/patient, €517.34); incremental QALYs showed greater benefit for exercise plus TrP-DN; inclusion of TrP-DN into exercise program was more likely to be cost-effective than exercise program alone	From cost-benefit perspective, inclusion of TrP-DN into multimodal management of patients with subacromial pain syndrome should be considered
Marks ³¹ (2016)	Compare clinical effectiveness and cost-effectiveness of corticosteroid injection for shoulder pain by physical therapist or orthopaedic surgeon	(1) Orthopaedic surgeon, (2) physical therapist	64 participants randomized (33 for physical therapist, 31 for orthopaedic surgeon); no significant differences in baseline characteristics between groups; noninferiority of injection by physical therapist demonstrated from total SPADI scores at 6 and 12 wk (upper limit of 95% CI, 13.34 and 7.17 at 6 and 12 wk, respectively); no statistically significant differences between groups on any outcome measures at 6 or 12 wk; from perspective of health funder, physical therapist was less expensive	Corticosteroid injection for shoulder pain, provided by qualified physical therapist, was clinically effective and less expensive; policy makers and service providers should consider implementing this model of care
Jowett ²⁵ (2013)	Analyze cost-effectiveness of subacromial corticosteroid injection with exercise compared with exercise alone in patients with moderate to severe shoulder pain and subacromial impingement syndrome	(1) Subacromial corticosteroid injection combined with exercise, (2) exercise alone	Mean NHS cost/patient (£255 vs £297, respectively) and overall health care cost (£261 vs £318, respectively) were lower in injection plus exercise arm, with no statistical significance; total QALYs gained were very similar in both arms (0.3514 vs 0.3494, respectively), although slightly higher in injection plus exercise arm, indicating that injection plus exercise may be dominant treatment option; at willingness-to-pay threshold of £20,000 per additional QALY gained, there was 61% probability that injection plus exercise was most cost-effective option	Injection plus exercise delivered by therapists was cost-effective, with lower health care costs and less time off work

^aASD, arthroscopic subacromial decompression; NHS, National Health Service; QALY, quality-adjusted life year; SPADI, Shoulder Pain and Disability Index; TrP-DN, trigger point dry needling.

TABLE A5
Remaining Economic Evaluations (n = 7)^a

Author (Year)	Purpose	Groups	Results	Conclusions
Sorensen ⁵⁷ (2020)	Investigate cost utility of plate fixation compared with nonoperative treatment of displaced midshaft clavicular fractures	(1) Plate fixation, (2) nonoperative treatment	Plate fixation was associated with larger QALY gain and higher cost; ICER was estimated to be €182,306/QALY from health sector perspective and €186,158/QALY from societal perspective	Plate fixation was not cost-effective when considering threshold of €34,000/QALY; however, for subgroup of patients with high-loading shoulder professions, plate fixation might be cost-effective

(continued)

TABLE A5 (continued)

Author (Year)	Purpose	Groups	Results	Conclusions
Gyftopoulos ¹⁸ (2018)	Determine cost-effectiveness of landmark-based and imaging-guided intra-articular steroid injections for initial treatment of adhesive capsulitis	(1) Landmark based, (2) imaging guided	Ultrasound-guided injections dominant for base case (least expensive [\$1280] and most effective [0.4096 QALYs]); model sensitive to probabilities of injecting steroid into joint (blind, ultrasound-guided, fluoroscopy-guided) and costs of ultrasound-guided and blind techniques; 2-way sensitivity analyses showed ultrasound-guided injections favored over blind and fluoroscopy-guided injections over range of reasonable probabilities and costs; probabilistic sensitivity analysis showed that ultrasound-guided injections were cost-effective in 44%, 34% for blind injections, and 22% for fluoroscopy-guided injections and over wide range of willingness-to-pay thresholds	Ultrasound-guided injection was most cost-effective option; blind and fluoroscopy-guided injections could also be cost-effective when performed by clinician likely to accurately administer medication into correct location
Paoli ⁴⁶ (2018)	Evaluate cost-effectiveness of nonoperative management, primary SLAP repair, and primary biceps tenodesis for treatment of symptomatic isolated type II SLAP tears	(1) SLAP repair, (2) biceps tenodesis, (3) nonoperative treatment	Primary biceps tenodesis compared with SLAP repair conferred increased effectiveness of 0.06 QALYs, with cost savings of \$1766; compared with nonoperative treatment, both biceps tenodesis and SLAP repair were cost-effective; sensitivity analysis showed that biceps tenodesis was preferred strategy in most simulations (52%); however, for SLAP repair to become cost-effective over biceps tenodesis, its probability of failure would have to be <2.7% or cost of biceps tenodesis would have to be >\$14,644	Compared with primary SLAP repair and nonoperative treatment, primary biceps tenodesis was most cost-effective treatment strategy in middle-aged patients
Hatch ²² (2017)	Assess vancomycin for preventing shoulder replacement infections	(1) Vancomycin, (2) no vancomycin	Efficacy of vancomycin (ARR) evaluated at different unit costs, baseline infection rates, and average costs of treating infection; vancomycin was cost-effective if initial infection rate decreased by 0.04% (ARR); using current costs of vancomycin in literature (range, \$2.50/1000 mg to \$44/1000 mg), vancomycin was cost-effective with ARR range of 0.01% at cost of \$2.50/1000 mg to 0.19% at \$44/1000 mg; baseline infection rates did not influence ARR obtained at any specific cost of vancomycin or cost of treating infection	Breakeven equation to assess efficacy of prophylactic antibiotics during shoulder surgery; prophylactic administration of local vancomycin powder during shoulder arthroplasty was highly cost-effective
Scott ⁵⁵ (2015)	Evaluate if economic savings are realized when procedures are performed by high-volume compared with low-volume providers	(1) High-, (2) medium-, (3) low-volume providers for ACL surgery, RCR, and TSA	Cost/case attributable to adverse outcomes for ACL reconstruction was \$496, \$781, and \$868 for high-, medium-, and low-volume providers; for RCR, it was \$523, \$640, and \$872, respectively; for TSA, it was \$1692, \$1876, and \$2021, respectively; sensitivity analysis showed that 50% increase in number of these 3 procedures performed by high-volume surgeons could save health care system \$23.1 million; if all procedures were performed by high-volume surgeons, it could save \$72 million	Higher provider volumes did convey substantial societal economic benefits; policies to incentivize and facilitate greater portion of procedures being performed by high-volume surgeons might increase efficiency of resource utilization in health care delivery
Dattani ¹² (2013)	Examine shoulder stiffness treated with ACR alone and ACR with ASD	(1) ACR, (2) ACR and ASD	Of 100 patients, 68 underwent ACR alone, and 32 underwent ACR with ASD; ACR showed highly significant improvement in range of movement and functional outcomes (OSS and EQ-5D); mean cost/QALY for ACR and ACR with ASD was £2563 and £3189, respectively	ACR was cost-effective in restoring relatively normal function and health-related quality of life in most patients with contracture of shoulder within 6 mo after surgery; beneficial effects not related to duration of symptoms

(continued)

TABLE A5 (continued)

Author (Year)	Purpose	Groups	Results	Conclusions
Pearson ⁴⁷ (2010)	Determine cost-effectiveness of ORIF of displaced midshaft clavicular fractures	(1) Clavicle ORIF, (2) nonoperative treatment	Base case cost per QALY gained for ORIF was \$65,000; cost-effectiveness improved to \$28,150 per QALY gained when benefit from ORIF was assumed to be permanent, with cost per QALY gained <\$50,000 when functional advantage persisted for ≥ 9.3 y; in other sensitivity analyses, cost per QALY gained for ORIF <\$50,000 when ORIF cost <\$10,465 (base case cost, \$13,668) or long-term utility difference between nonoperative treatment and ORIF was >0.034 (base case difference, 0.014); short-term disutility associated with fracture healing also affected cost-effectiveness, with cost per QALY gained for ORIF <\$50,000 when utility of fracture treated nonoperatively before union was <0.617 (base case utility, 0.706) or when nonoperative treatment increased time to union by 20 wk (base case difference, 12 wk)	Cost-effectiveness of ORIF after acute clavicular fractures depended on durability of functional advantage for ORIF compared with nonoperative treatment; when functional benefits persisted for >9 y, ORIF had favorable value compared with many accepted health interventions

^aAll values are reported as US dollars unless otherwise specified. ACL, anterior cruciate ligament; ACR, arthroscopic capsular release; ARR, absolute risk reduction; ASD, arthroscopic subacromial decompression; EQ-5D, EuroQol 5 Dimensions index; ICER, incremental cost-effectiveness ratio; ORIF, open reduction internal fixation; OSS, Oxford Shoulder Score; QALY, quality-adjusted life year; RCR, rotator cuff repair; SLAP, superior labral from anterior to posterior; TSA, total shoulder arthroplasty.