

The need for neurologists in the care of COVID-19 patients

J. Sellner^{a,b,c} , P. Taba^{d,e} ,
S. Öztürk^f and R. Helbok^g

^aDepartment of Neurology, Landeskrankenhaus Mistelbach-Gänserndorf, Mistelbach, ^bDepartment of Neurology, Christian Doppler Medical Center, Paracelsus Medical University, Salzburg, Austria, ^cDepartment of Neurology, Klinikum rechts der Isar, Technische Universität München, München, Germany, ^dDepartment of Neurology and Neurosurgery, University of Tartu, Tartu, ^eNeurology Clinic, Tartu University Hospital, Tartu, Estonia, ^fDepartment of Neurology, Faculty of Medicine, Selcuk University, Konya, Turkey and ^gNeurological Intensive Care Unit, Department of Neurology, Medical University of Innsbruck, Innsbruck, Austria

Correspondence: J. Sellner, Department of Neurology, Landeskrankenhaus Mistelbach-Gänserndorf, Liechtensteinstr. 67, 2130 Mistelbach, Austria (tel.: +43 2572 9004-11550; fax: +43 2572 9004-49332; e-mail: j.sellner@salk.at).

Keywords: autonomic dysfunction, COVID-19, encephalitis, neurological complications, olfactory bulb, SARS-CoV-2, stroke

doi:10.1111/ene.14257

Received: 3 April 2020
Accepted: 4 April 2020

Coronavirus disease 2019 (COVID-19) is an infectious disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) resulting in severe acute respiratory syndrome in a substantial number of patients. Approximately 5%–10% of patients need intensive care unit (ICU) admission and mechanical ventilation [1]. The primary symptoms of COVID-19 include fever, dry cough and fatigue [2]. However, a recent report from Shandong, China, disclosed that a subset of patients did not suffer from respiratory symptoms but had

neurological signs and symptoms [3]. Moreover, in a retrospective study from Wuhan, China, neurological symptoms were observed in 36.4% of the hospitalized patients with COVID-19 infection and pertained to both the central and peripheral nervous system [4]. Therefore, neurologists need to be part of the multidisciplinary team taking care of the patients. Our call emerges on the one hand from evidence for the neuroinvasiveness of coronaviruses, immunopathology in animal models and observations made in previous SARS epidemics [5]. On the other hand, there is an emerging number of reports of SARS-CoV-2 infection with neurological manifestations and complications which already let us foresee the potential spectrum of disease which we will encounter in the further course of the pandemic.

The lungs are the organs most affected by SARS-CoV-2 because the virus accesses host cells via the enzyme angiotensin-converting enzyme (ACE) 2, which is most abundant in type II alveolar cells. Yet, glial cells and neurons of the central nervous system (CNS) have been reported to express ACE 2, rendering the brain a potential target of the virus [6]. Knowledge about the transneuronal transport of SARS-CoV through the olfactory bulb supports this hypothesis [7]. However, whether the viral invasion of the olfactory bulbs is the neurobiological background for smell and taste disorders reported by infected patients remains to be elucidated [8]. There is a report about a 56-year-old male in China who developed COVID-19 and in whom the virus was detected in the cerebrospinal fluid (CSF) [9]. There are no clinical details outlined beyond the information that this patient recovered and was discharged from hospital. Of note, examination of SARS-CoV-2 in CSF is not a routine examination and may not be consistently available. The concern that patients with neuroinvasive disease and atypical CNS manifestations are likely to be just a matter of time is supported by a case of COVID-19 associated acute necrotizing encephalopathy [10]. The study of the mechanism leading to neuroaxonal injury, which may involve both direct viral damage and bystander

inflammation, is critical for the development of treatment strategies. Whether ACE inhibitors, which are widely used for the treatment of hypertension, suppress the adaptive immune system and the subsequent antiviral response to SARS-CoV-2 is another unsolved question [11].

The neuroinvasive potential of SARS-CoV-2 may play a role in the emergence of respiratory failure in COVID-19 patients. Indeed, coronaviruses were shown to reach the brainstem via a synapse-connected route from the lung and airways [12]. Thus, further characterization of the central cardiac and respiratory dysfunction is key to understanding the underlying mechanisms and identifying patients requiring ICU admission early. The respiratory centre is located in the medulla oblongata and the respiratory rhythm is modulated from various sites of the lower brainstem, including the pons [13]. Therefore, impaired cough and gag reflex may also indicate CNS manifestation. Still, these reflexes are associated with a considerable risk of aerosol transmission and have to be performed with caution. It is unclear if examination of other brainstem reflexes such as corneal reflexes and pupillary reflexes are helpful for early detection of CNS involvement.

Neurologists should also keep in mind the potential risks for para-infectious and post-infectious disorders. Of note, there were cases of acute disseminated encephalomyelitis, vasculopathy and Guillain-Barré syndrome in association with the Middle East Respiratory Syndrome (MERS) CoV [14]. Moreover, there is preliminary evidence for a pro-coagulatory state associated with COVID-19 infection and development of ischaemic stroke [15]. This needs to be considered in addition to the potential risk of cardioembolic stroke due to ACE 2 expression in the heart and subsequent cardiac dysfunction [16]. Moreover, the number of reported cases of COVID-19 patients with intracerebral haemorrhage is increasing [3]. Whether this potential association is related to uncontrolled hypertension by interaction of SARS-CoV-2 with ACE 2 needs to be clarified. Hypertensive encephalopathy may

therefore be an early sign of infection or may develop during the course. As many patients have a high body temperature and electrolyte disturbances, seizures and encephalopathy may develop. Moreover, susceptibility for more severe disease in patients with pre-existing neurological conditions and on immunosuppressive therapy is another concern. Finally, we do not know how many ICU survivors of COVID-19 will develop a post-intensive-care syndrome including critical illness neuropathy and myopathy and suffer from long-term sequelae which again emphasizes the role of neurologists in the care of COVID-19 patients.

We therefore appreciate the concerted efforts by the European Academy of Neurology to better understand neurological manifestations and complications in order to improve the overall outcome of the pandemic and would like to raise awareness for the EANcore COVID-19 activities (<https://www.ean.org/EANcore-Covid-19.4325.0.html>).

Acknowledgements

No financial support.

Competing interests

The authors are elected Chairs of Scientific Panels in the European Academy of Neurology (EAN): JS and PT are heading the EAN Infectious Disease Panel and SÖ and RH the EAN Neurocritical Care Panel. The authors declare that they have no further financial or

non-financial disclosures with respect to this paper.

References

1. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72314 cases from the Chinese Center for Disease Control and Prevention. *JAMA* 2020; **323**: 1239.
2. Huang C, Wang Y, Li X, *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; **395**: 497–506.
3. Wang HY, Li XL, Yan ZR, Sun XP, Han J, Zhang BW. Potential neurological symptoms of COVID-19. *Ther Adv Neurol Dis.* 2020;**13**. <https://doi.org/10.1177/1756286420917830>.
4. Mao L, Wang M, Chen S, *et al.* Neurological manifestations of hospitalized patients with COVID-19 in Wuhan, China: a retrospective case series study. *medRxiv*. 2020doi: 10.1101/2020.02.22.20026500doi.
5. Desforges M, Le Coupanec A, Dubeau P, *et al.* Human coronaviruses and other respiratory viruses: underestimated opportunistic pathogens of the central nervous system? *Viruses* 2019; **12**: 14.
6. Baig AM, Khaleeq A, Ali U, Syeda H. Evidence of the COVID-19 virus targeting the CNS: tissue distribution, host–virus interaction, and proposed neurotropic mechanisms. *ACS Chem Neurosci* 2020; **11**: 995–998.
7. Netland J, Meyerholz DK, Moore S, Cassell M, Perlman S. Severe acute respiratory syndrome coronavirus infection causes neuronal death in the absence of encephalitis in mice transgenic for human ACE2. *J Virol* 2008; **82**: 7264–7275.
8. Giacomelli A, Pezzati L, Conti F, *et al.* Self-reported olfactory and taste disorders in SARS-CoV-2 patients: a cross-sectional study. *Clin Infect Dis* 2020; **71**: 889–890.
9. The Encephalitis SocietyUE. Coronavirus statement. <https://www.encephalitisinfo/blog/coronavirus>. 2020.
10. Poyiadji N, Shahin G, Noujaim D, Stone M, Patel S, Griffith B. COVID-19-associated acute hemorrhagic necrotizing encephalopathy: CT and MRI features. *Radiology* 2020; **201187**.
11. Bernstein KE, Khan Z, Giani JF, Cao DY, Bernstein EA, Shen XZ. Angiotensin-converting enzyme in innate and adaptive immunity. *Nat Rev Nephrol* 2018; **14**: 325–336.
12. Li YC, Bai WZ, Hashikawa T. The neuroinvasive potential of SARS-CoV2 may play a role in the respiratory failure of COVID-19 patients. *J Med Virol* 2020; **92**: 552–555.
13. Duffin J. Functional organization of respiratory neurones: a brief review of current questions and speculations. *Exp Physiol* 2004; **89**: 517–529.
14. Bohmwald K, Galvez NMS, Rios M, Kalergis AM. Neurologic alterations due to respiratory virus infections. *Front Cell Neurosci* 2018; **12**: 386.
15. Hospital Reggio Emilia. Frontline Report: COVID-19 and Impact on Neurology Unit at Reggio Emilia Hospital. <https://www.eanpages.org/2020/03/31/frontline-report-covid-19-impact-on-neurology-unit-at-reggio-emilio-hospital/>. 2020.
16. Madjid M, Safavi-Naeini P, Solomon SD, Vardeny O. Potential effects of coronaviruses on the cardiovascular system. *A Review. JAMA Cardiol* 2020; **5**: 831–840.