

## ARTICLE

# An autoinducible trp-T7 expression system for production of proteins and biochemicals in *Escherichia coli*

Jenny Landberg<sup>1</sup>  | Hemanshu Mundhada<sup>1,2</sup> | Alex Toftgaard Nielsen<sup>1</sup> 

<sup>1</sup>The Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, Kongens Lyngby, Denmark

<sup>2</sup>CysBio ApS, Hørsholm, Denmark

**Correspondence**

Hemanshu Mundhada, CysBio ApS, Agern Allé 1, 2970 Hørsholm, Denmark.  
Email: [hemu@cysbio.com](mailto:hemu@cysbio.com)

**Funding information**

Novo Nordisk Fonden, Grant/Award Numbers: NNF10CC1016517, NNF16CC0020908

**Abstract**

Inducible expression systems can be applied to control the expression of proteins or biochemical pathways in cell factories. However, several of the established systems require the addition of expensive inducers, making them unfeasible for large-scale production. Here, we establish a genome integrated trp-T7 expression system where tryptophan can be used to control the induction of a gene or a metabolic pathway. We show that the initiation of gene expression from low- and high-copy vectors can be tuned by varying the initial concentration of tryptophan or yeast extract, and that expression is tightly regulated and homogenous when compared with the commonly used lac-T7 system. Finally, we apply the trp-T7 expression system for the production of L-serine, where we reach titers of 26 g/L in fed-batch fermentation.

**KEYWORDS**

*Escherichia coli*, inducible promoter, L-serine, tryptophan

## 1 | INTRODUCTION

Cell factories are promising options for the sustainable production of biobased compounds such as proteins and biochemicals (Nielsen & Keasling, 2016). While some products have been commercialized, it is generally difficult to produce those compounds at a price that is competitive with petroleum-derived chemicals (Gustavsson & Lee, 2016; Liu, Zhu, Tan, Zhang, & Ma, 2016). The complex (heterologous) proteins and pathways that are used in cell factories oftentimes compete with native metabolic processes for cellular resources and pathway expression at an early growth phase can drain the cell for energy or result in redox imbalance (Blazeck & Alper, 2013; Brockman & Prather, 2015). Furthermore, if the end-product is a toxic metabolite or protein, production at the beginning of the fermentation results in growth inhibition or poor growth (Saida, Uzan, Odaert, & Bontems, 2006). Expressing the product pathway from a promoter that is induced at a specific timepoint enables a two-phase production process with biomass buildup followed by product accumulation. This approach can increase yield, titer, and productivity and improve process economics (Burg et al., 2016).

There are several inducible promoters and expression systems available for use in *Escherichia coli* (Terpe, 2006). One of the most commonly used systems is the lac-T7 (DE3) system (Studier & Moffatt, 1986). It consists of a T7 RNA polymerase (T7 polymerase) controlled by the lacUV5 promoter and repressed by the lactose operon repressor (LacI). The protein or biochemical pathway is placed under control of the T7 promoter (*pT7*) and expression is induced by the addition of isopropyl  $\beta$ -D-1-thiogalactopyranoside (IPTG) or by growth on lactose (Studier, 2005; Studier & Moffatt, 1986). Gene expression by T7 polymerase enables very high expression levels, which has led to extensive use of the system for protein production (Rosano & Ceccarelli, 2014). Proteins or biochemical pathways are also commonly expressed directly from the lac promoter (*Plac*; Gronenborn, 1976) or from the engineered lac and tryptophan fusion promoters Tac (de Boer, Comstock, & Vasser, 1983) and Trc (Brosius, Erfle, & Storella, 1985), which have lower basal- and higher expression levels compared with *Plac*. All these systems require the addition of IPTG for induction, which is expensive and unfeasible during the production of low- or medium-value compounds (Keasling, 2012). Although IPTG-based

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Biotechnology and Bioengineering* Published by Wiley Periodicals, Inc.

systems can also be induced by switching to lactose-based growth (Studier, 2005), feeding lactose as a sole carbon source to a fed-batch or continuous fermentation is not feasible due to its cost and relatively low solubility in water.

Inducible promoters can be tuned to improve their function and dynamic range. Common approaches to minimize basal level expression and tune the induced expression level include engineering/mutating the repressor, operator and upstream elements, or the  $-10$  and  $-35$  binding sites of the RNA polymerase as well as the spacing in between (Brosius et al., 1985; Chen et al., 2018; Ellefson, Ledbetter, & Ellington, 2018; Guan, Liu, Li, & Zhang, 2013). It has also been shown that novel, synthetic, ligand-responsive promoter sequences can be used to improve expression levels up to 100-fold compared with native sequences (Liu et al., 2019). If the promoter function is unknown, the characterization of important promoter elements can be done by establishing mutant promoter libraries and screening those in a high-throughput manner (Rohllhill, Sandoval, & Papoutsakis, 2017). Using information gained in a promoter library screen, the repressor binding site of the formaldehyde-inducible promoter could be identified and the information could be used to engineer mutated promoter variants with lower basal- and higher induced level expression and a significantly improved dynamic range (Rohllhill et al., 2017). Thermodynamic models are also helpful in the design of inducible promoters, especially to elucidate and predict the interaction between RNA polymerase and DNA (Brewster, Jones, & Phillips, 2012; Chen et al., 2018). In a recent study, Chen et al. (2018) developed a method that can be used to engineer the dynamic range of ligand-inducible promoters. Using a thermodynamic model, they established a promoter library with different  $-10$  and  $-35$  sites and applied the optimal site combinations to build promoters regulated by different combinations of transcription factors, resulting in predictable expression with a variable dynamic range.

Several attempts have been made to find affordable and easy-to-apply (auto)inducible expression systems with low levels of basal expression and high, tunable levels of gene expression. Such systems do for example include those based on sensing of temperature, cell signaling, or depletion/addition of a compound that is naturally occurring in the cultivation medium. For temperature-based induction, the  $p_L$  and  $p_R$  phage promoters can be used to induce gene expression, usually by shifting temperature from below 37 to up to 42°C (Elvin et al., 1990; Kincade & deHaseth, 1991). The system has been coupled to the T7 polymerase for high-level protein production of protein (Chao, Law, Chen, & Hung, 2002). Although an inexpensive method of control, increased cultivation temperatures can induce cellular stress responses and alter protein folding (Valdez-Cruz, Caspeta, Pérez, Ramírez, & Trujillo-Roldán, 2010). In addition, exact and homogenous temperature control can be difficult to achieve in large-scale vessels (Gvazdaitis et al., 1994). For cell signaling-based induction, the native bacterial quorum-sensing system has been applied to establish autoinducible production by coupling gene expression to the concentration of secreted quorum signaling molecules (Gupta, Reizman, Reisch, & Prather, 2017; Kim et al., 2017). This type of system has also been shown to work when coupled to the T7 polymerase for further amplification of protein output

(Zargar, Quan, & Bentley, 2016). While the expression is not always tightly regulated and the system might require tuning for each desired compound, further engineering could provide a system suitable for large-scale fermentation.

The tryptophan promoter (*P<sub>trp</sub>*) is induced as tryptophan is depleted from the medium, or by addition of  $\beta$ -indoleacrylic acid. It has been extensively applied for protein and polypeptide production (Bass & Yansura, 2000) and there are several expression vector systems based on the promoter (Chevalet, Robert, Gueneau, Bonnefoy, & Nguyen, 2000; Rosano & Ceccarelli, 2014). Generally, the native or leaderless (without the *trp* leader sequence *trpL*) promoter and the tryptophan repressor (*trpR*) have been inserted into an expression vector for direct control of a gene of interest (GOI). The promoter is tightly regulated, but does not provide as high expression levels as *lac-T7* and plasmid instability is prevalent (Somerville, 1988).

In this study, we engineered a genome integrated autoinducible expression system by combining the tightly regulated tryptophan promoter with the high-expressing T7 polymerase. We showed that induction of gene expression could be tuned by the initial tryptophan concentration for low- and high-copy vectors, and that expression strength was dependent on the use of the native or leaderless promoter constructs. By cloning the toxin MazF under control of our promoter constructs, we verified that the promoters are tightly regulated and suitable even for toxic protein production. Using flow cytometry, we further demonstrated that population homogeneity was improved compared with the commonly used *lac-T7* system. Finally, we showed that the *trp-T7* expression system can be used for biochemical production by applying it for batch and fed-batch fermentation of L-serine.

## 2 | MATERIALS AND METHODS

### 2.1 | Strains, media, and materials

For cultivation and screening during cloning, 2xYT (16 g/L Bacto tryptone, 10 g/L yeast extract (YE), 5 g/L NaCl) medium and lysogeny broth (LB; 10 g/L tryptone, 5 g/L YE, 10 g/L NaCl) agar plates with appropriate antibiotics were used. Ampicillin, kanamycin, chloramphenicol, and spectinomycin was used with working concentrations of 100, 50, 50, and 50  $\mu$ g/mL, respectively. For cloning and screening of plasmids carrying a tryptophan repressible gene, 0.5 mM tryptophan was added to agar plates and growth medium to repress gene expression. Growth and production experiments were performed in M9 minimal medium with 0.1 mM  $\text{CaCl}_2$ , 2.0 mM  $\text{MgSO}_4$ , 1X trace element solution and 1X M9 salts supplemented with glucose, appropriate antibiotics and other compounds as described below. The stock solution of trace elements was 1,000X and consisted of 15 g/L  $\text{EDTA}(\text{Na}_2)\cdot 2\text{H}_2\text{O}$ , 4.5 g/L  $\text{ZnSO}_4\cdot 7\text{H}_2\text{O}$ , 0.7 g/L  $\text{MnCl}_2\cdot 4\text{H}_2\text{O}$ , 0.3 g/L  $\text{CoCl}_2\cdot 6\text{H}_2\text{O}$ , 0.2 g/L  $\text{CuSO}_4\cdot 2\text{H}_2\text{O}$ , 0.4 g/L  $\text{NaMoO}_4\cdot 2\text{H}_2\text{O}$ , 4.5 g/L  $\text{CaCl}_2\cdot 2\text{H}_2\text{O}$ , 3 g/L  $\text{FeSO}_4\cdot \text{H}_2\text{O}$ , 1 g/L  $\text{H}_3\text{BO}_3$ , and 0.1 g/L KI dissolved in double-distilled water and sterile filtered. The stock solution of M9 salts was 10X and consisted of 6.8 g/L  $\text{Na}_2\text{HPO}_4$  anhydrous, 3 g/L  $\text{KH}_2\text{PO}_4$ , 5 g/L NaCl, and 1 g/L  $\text{NH}_4\text{Cl}$

**TABLE 1** Strains and plasmids used in the study

Strain	Description	Reference/source
<i>E. coli</i> NEB 5-alpha	<i>fhuA2 Δ(argF-lacZ)U169 phoA glnV44 ϕ80 Δ(lacZ)M15 gyrA96 recA1 relA1 endA1 thi-1 hsdR17</i> ; cloning strain	New England Biolabs
<i>E. coli</i> K12 MG1655	F- lambda- <i>ilvG- rfb-50 rph-1</i>	Lab collection
<i>E. coli</i> MG1655 (DE3)	<b>lac-T7</b> : MG1655 λ (DE3)	Mundhada, Schneider, Christensen, and Nielsen (2016)
ALE5	MG1655 $\Delta sdaA\Delta sdaB\Delta tdcG\Delta glyA$ ; serine tolerant strain	Mundhada et al. (2017)
ALE5	MG1655 $\Delta sdaA\Delta sdaB\Delta tdcG\Delta glyA \lambda(DE3)$ ; serine tolerant strain	Mundhada et al. (2017)
JL13	<b>trp-T7</b> : MG1655 attB-186(O): P <sub>trp</sub> ; tryptophan repressor and promoter controlling T7 polymerase expression	This study
JL44	<b>trp*-T7</b> : MG1655 attB-186(O): P <sub>trp</sub> *; tryptophan repressor and truncated promoter controlling T7 polymerase expression	This study
JL45	trp-T7 with pSEVA27-mCherry	This study
JL46	trp*-T7 with pSEVA27-mCherry	This study
JL84	<b>trp-T7-ser</b> : ALE5 attB-186(O): P <sub>trp</sub> ; tryptophan repressor and promoter controlling T7 polymerase expression	This study
JL141	trp-T7 with pCDF-mCherry	This study
JL142	trp*-T7 with pCDF-mCherry	This study
JL151	lac-T7 with pSEVA-mCherry	This study
JL153	lac-T7 with pCDF-mCherry	This study
JL154	trp-T7-ser with pSEVA27-serABC	This study
JL155	trp*-T7-ser with pSEVA27-serABC	This study
JL160	trp-T7 with pSEVA27	This study
JL161	trp-T7 with pSEVA27-mazF	This study
JL162	trp-T7 with pCDF	This study
JL163	trp-T7 with pCDF-mazF	This study
JL164	trp*-T7 with pSEVA27	This study
JL165	trp*-T7 with pSEVA27-mazF	This study
JL166	trp*-T7 with pCDF	This study
JL167	trp*-T7 with pCDF-mazF	This study
Plasmid	Description	Reference/source
pOSIP KO	Cloning-integration plasmid, integrates at phage 186 site; Km <sup>R</sup>	St-Pierre et al. (2013)
pOSIP KO-lacI-T7	Plasmid for integrating T7 polymerase under control of LacI; Km <sup>R</sup>	Mundhada et al. (2017)
pE-FLP	Plasmid for expressing FLP recombinase to excise the integration plasmid backbone; Amp <sup>R</sup>	St-Pierre et al. (2013)
pSEVA27-sl3	<b>pSEVA27</b> : Low-copy number plasmid pSC101 origin (~5 copies per cell); MCS under T7 promoter; Km <sup>R</sup>	Kim, Cavaleiro, Rennig, and Nørholm (2016)
pCDF-1b	<b>pCDF</b> : High-copy number plasmid CloDF13 origin (~20–40 copies per cell); MCS under T7 promoter; Sp <sup>R</sup>	Novagen
pJL3	<b>pOSIP-trp*-T7</b> : pOSIP KO harboring a P <sub>trp</sub> * integration cassette controlling the T7 polymerase; Km <sup>R</sup>	This study
pJL5	<b>pSEVA-mCherry</b> : SEVA27-sl3 plasmid with mCherry expressed from the T7 promoter; Km <sup>R</sup>	This study
pJL9	<b>pOSIP-trp-T7</b> : pOSIP KO harboring a P <sub>trp</sub> integration cassette controlling the T7 polymerase; Km <sup>R</sup>	This study

(Continues)

TABLE 1 (Continued)

Plasmid	Description	Reference/source
pJL42	pCDF-1b plasmid with mCherry expressed from the T7 promoter; Sp <sup>R</sup>	This study
pJL83	<b>pCDF-mCherry</b> : pCDF-1b with <i>lacI</i> removed and mCherry expressed from the T7 promoter; Sp <sup>R</sup>	This study
pJL88	<b>pSEVA-mazF</b> : pSEVA27-sl3 plasmid with <i>mazF</i> expressed from the T7 promoter; Km <sup>R</sup>	This study
pJL89	<b>pCDF-mazF</b> : pCDF-1b plasmid with <i>lacI</i> removed and <i>mazF</i> expressed from the T7 promoter; Sp <sup>R</sup>	This study
pSEVA27-sl-serA <sup>mut</sup> CB TIR1	<b>pSEVA-serACB</b> : Expression of the serine operon from the T7 promoter; Km <sup>R</sup>	Rennig et al. (2019)

Note: Names used to refer to strains and plasmids in the main text are written in bold.

Abbreviation: *E. coli*: *Escherichia coli*.

dissolved in double-distilled water and autoclaved. The strains and plasmids that were used are listed in Table 1. Chemicals were purchased from Sigma-Aldrich (Taufkirchen, Germany). Restriction enzymes and polymerase chain reaction (PCR) polymerases were purchased from Thermo Fisher Scientific (Waltham, MA). USER enzyme was purchased from BioNordika (Herlev, Denmark).

## 2.2 | Plasmid and strain construction

Plasmids were purified using the Macherey Nagel plasmid purification kit (Dure, Germany). Cells were transformed using the TSS buffer method. Forty milliliters exponentially growing cells (OD ~0.4–0.8) were spun down and resuspended in 0.5–1 ml TSS buffer (100 g/L PEG 8000, 30 mM MgCl<sub>2</sub>, 20 ml dimethyl sulfide in LB). Cells were mixed with 100 ng plasmid and put on ice for 15 min. The plasmid-cell mix was subjected to heat shock (42°C, 1 min) and put on ice for 2 min. Cells were incubated for 1 hr at 37°C before plating on LB agar with appropriate antibiotics. All plasmids were constructed by USER cloning (Cavaleiro, Kim, Seppälä, Nielsen, & Nørholm, 2015; Nour-Eldin, Hansen, Nørholm, Jensen, & Halkier, 2006). Briefly, template DNA was amplified using Phusion U polymerase and primers containing a uracil complementary overhang. PCR products were purified and subjected to *DpnI* digestion. The backbone and the template fragments were mixed with USER enzyme and buffer to a volume of 10 µl and incubated for 20 min at 37°C followed by 20 min at 25°C. The mix was diluted with 10 µl water and 10 µl of the diluted mix was used for transformation into competent cells. All primers used in the study were ordered from Integrated DNA Technologies (Leuven, Belgium) and are listed in Table S1. The tryptophan repressor, operator and promoter and the *mazF* gene were amplified from the *Escherichia coli* genome. Integration of *trp*-T7 and *trp*\*-T7 was carried out according to the protocol by (St-Pierre et al., 2013). Briefly, the cassettes were cloned on a pOSIP vector, followed by integration at the attB-186(O) site. The antibiotic cassette was excised using an FLP recombinase.

## 2.3 | Microtiter plate cultivations

Single transformants were inoculated in 96-deep well plates in 500 µl M9 medium supplemented with 0.4% glucose, 0.02% YE, and 0.5 mM tryptophan. Cultures were grown overnight at 37°C, 250 rpm. Cells were washed and inoculated with a 1:100 inoculum ratio in microtiter plates with 150 µl M9 medium supplemented with 0.5% glucose and a specified amount of tryptophan or YE. The plates were sealed with oxygen-permeable film (Breathe-Easy sealing membrane, Sigma-Aldrich) and incubated at medium shaking, 37°C in an ELx808 Absorbance Reader (BioTek, Winooski, VT) for OD<sub>630</sub> measurements, or in a Synergy H1 Hybrid Multi-Mode Reader (BioTek) for OD<sub>630</sub> and fluorescence (excitation 587 nm, emission 617 nm) measurements.

## 2.4 | Flow cytometry

Single transformants were inoculated in 2 ml M9 medium supplemented with 0.4% glucose, 0.02% YE, and 0.5 mM tryptophan. Cultures were grown overnight at 37°C, 250 rpm. Cells were washed and inoculated with an inoculum ratio of 1:100 in a 24-deep well plate in 2.5 ml M9 medium supplemented with 0.4% glucose. The MG1655(DE3) strains were induced with 1 mM IPTG at OD<sub>630</sub> approximately 0.3–0.4. Samples were diluted appropriately and analyzed in a MACSQuant® VYB flow cytometer (Miltenyi Biotec, Cologne, Germany). The expression of mCherry was detected using a yellow laser (561 nm) and the 615/20 nm Y2 filter. Forward (FSC) and side (SSC) scatter was detected with the yellow laser and a 561/10 nm filter. The results were analyzed with FlowJo (Becton, Dickinson and Company, Franklin Lakes, NJ).

## 2.5 | Serine production in small batch fermentation

Preinoculums were prepared by inoculating single transformants in 2.5 ml M9 medium supplemented with 0.4% glucose, 0.02% YE, 0.5 mM tryptophan, and 2 mM glycine. Cultures were grown

overnight at 37°C, 250 rpm. Cells were washed and inoculated to an OD of 0.05 in 50 ml M9 medium supplemented with 0.4% glucose, 2 mM glycine and 0.04 or 0.5 mM tryptophan.

## 2.6 | Serine production in fed-batch fermentation

Medium for fed-batch fermentation was prepared as previously described (Mundhada et al., 2017), with an addition of 0.125% YE instead of 0.2% YE to the batch medium. A preculture was grown overnight at 37°C, 250 rpm, in 2xYT medium with 0.5 mM trp and 0.1% glucose. Cells were washed and the bioreactor was inoculated to an OD of 0.1. Oxygen saturation was maintained at 40% throughout the fermentation. Samples were taken regularly for analysis. A correlation of 0.374 between cell dry weight (CDW) and OD was used for CDW estimation and subsequent yield calculations (Mundhada et al., 2016).

## 2.7 | Analytical methods

Samples from the small batch and fed-batch fermentations were filtered and diluted appropriately. Glucose and byproducts were analyzed with high-performance liquid chromatography as previously described (Kildegaard et al., 2014). Serine was analyzed with LC-MS as previously described (Mundhada et al., 2016).

## 2.8 | Data availability

All data is available from the authors upon request.

# 3 | RESULTS AND DISCUSSION

## 3.1 | Design and construction of a trp-T7 expression system

To establish a high-expressing, inexpensive inducible system with low basal level expression, we wanted to combine the high level of gene expression achievable from the T7 polymerase with the tightly regulated trp operon. Using such a system, we hypothesized that the onset of gene expression could be tuned to a certain time point by regulating the amount of tryptophan added to the medium at the start of the cultivation (Figure 1a,b).

The trp operon consists of five structural genes encoding tryptophan synthase and is regulated by the TrpR. In growth medium with excess tryptophan, TrpR binds the *trpO*, blocking transcription of the operon from the *P<sub>trp</sub>*. The repression is further (~10-fold) enhanced by attenuation, a mechanism where the *trpL* forms RNA stem-loops that blocks transcription in the presence of tryptophan (Yanofsky, 1981). We constructed two integrative cassettes where the *E. coli* tryptophan regulation machinery was designed to control the T7 polymerase (Figure 1c).

The first variant contained the tryptophan repressor and the native tryptophan leader sequence, promoter, and operator. This variant is hereafter referred to as trp-T7. The second variant contained the tryptophan repressor, the tryptophan promoter and operator, but not the tryptophan leader sequence. This variant is hereafter referred to as trp\*-T7. The cassettes were cloned on a pOSIP integration vector (St-Pierre et al., 2013) and integrated into the attB-186(O) site in *E. coli* K12 MG1655, resulting in the two strains hereafter referred to as trp-T7 and a trp\*-T7.


Low-copy plasmid pSEVA27 (hereafter pSEVA) and high-copy plasmid pCDF-1b (hereafter pCDF) harboring *pT7* controlled by *lacO* were used as expression vectors to evaluate expression homogeneity and basal level expression, the timing of induction in different concentrations of tryptophan, and biochemical production from the expression system (Figure 1d). We decided to not replace the *lacO* site with a *trpO* site to facilitate cloning during future use of the system, as most T7-based expression vectors have a *lacO* site present. We further hypothesized that *trpR*-based repression of the genomic T7 polymerase would provide sufficient regulation of the system, as the trp operon is inherently under more strict regulation when compared with the lac operon. However, keeping *lacO* meant that we had to remove *lacI* from pCDF to enable target gene expression during growth on glucose.

## 3.2 | Evaluation of expression profiles in different concentrations of tryptophan and YE

For evaluation of the promoter system, strains with integrated trp-T7 and trp\*-T7 (that contains or lacks the *trpL*, respectively), were transformed with a low- or high-copy plasmid harboring mCherry under control of *pT7*, resulting in four different strains (Table 1 and Figure 1c,d). The induction profile of each strain was investigated using different starting concentrations of tryptophan and (tryptophan-containing) YE with mCherry as a reporter gene (Figure 2a). Strains were grown in 0–0.5 mM tryptophan or 0–1% YE, and OD and fluorescence was monitored for 20 hr.

Overall, the initial concentration of tryptophan and YE was inversely correlated to the fluorescence per OD and the initiation of mCherry expression (Figures 2c,d and S1), showing that tryptophan and YE could be used to control the induction of the GOI in both versions of the trp-T7 strains, and for both the high- and low-copy vector. The fluorescence and fluorescence per OD at different time points were generally higher with decreasing concentrations of tryptophan or YE, since expression was induced earlier (Figure 2b,c). An exception was trp-T7 with pCDF-mCherry, where fluorescence per OD was similar for all YE concentrations between 0% and 0.25% (Figures 2c and S1). As expected, growth was dependent on the amount of YE added to the medium (Figure S1).

For strains carrying the low-copy plasmid pSEVA-mCherry (Figure 2b), high initial concentrations of tryptophan and YE (0.5,




**FIGURE 1** The layout and function of the trp-T7 expression system. (a) Tryptophan-based repression and induction for the production of a protein or biochemical. (b) Repression and induction of the trp-T7 expression system is controlled by tryptophan concentration. When tryptophan is present in the growth medium it is bound by TrpR, which binds *trpO* and inhibits expression of T7 polymerase and the GOI. After tryptophan is consumed, the conformation of TrpR changes, leading to dissociation from *trpO*. T7 polymerase is produced and transcribes the plasmid-based GOI from *pT7*. (c) The layout of the genome integrated cassettes *trp*-T7 and *trp*\*-T7. (d) The plasmids with the different T7-transcribed genes that were used to test *trp*-T7-based protein and biochemical production in this study. GOI, gene of interest; *trpO*, tryptophan operator; TrpR, tryptophan repressor [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

0.25 mM, and 1%, 0.5%, and 0.25%, respectively) completely repressed mCherry expression in *trp*-T7, while low levels of expression could be seen after approximately 10–13 hr in *trp*\*-T7. This is expected, as the *trpL*-containing *trp*-T7 version of the promoter should exhibit a higher level of repression than *trp*\*-T7. With most concentrations of tryptophan and YE, the fluorescence per OD was only marginally higher in *trp*\*-T7 compared to *trp*-T7. However, for the strains carrying the high-copy plasmid pCDF-mCherry (Figure 2c), the fluorescence per OD was significantly higher in *trp*\*-T7, compared to *trp*-T7, for both *trp* and YE cultures. Low but significant levels of mCherry were observed in *trp*\*-T7 even at high initial concentrations of tryptophan and YE, while, for *trp*-T7, repression was similar to that seen in the low-copy plasmid strains. Interestingly, the fluorescence per OD in strains harboring the *trp*-T7 expression cassette was higher when mCherry was expressed from the low-copy pSEVA plasmid than from high-copy pCDF plasmid. This may be due to the higher metabolic burden resulting from maintaining multiple plasmid copies (Jones, Kim, & Keasling, 2000).

### 3.3 | Comparing expression strength and homogeneity to lac-T7

Next, we sought to investigate the expression heterogeneity of both promoter constructs and compare it to the commonly used lac-T7 expression system. A lac-T7 strain (*E. coli* MG1655 (DE3); Mundhada et al., 2016) was transformed with pSEVA-mCherry or pCDF-mCherry. Overnight cultures were inoculated in minimal medium without tryptophan. The lac-T7 cultures were induced with 1 mM IPTG at OD approximately 0.3–0.35. Expression levels were investigated with flow cytometry 8 and 24 hr after inoculation (Figure 3a).

For the pSEVA-mCherry strains, fluorescence was slightly lower for *trp*-T7 compared to *trp*\*-T7 after 8 hr ( $p = .027$ ), while no significant difference was seen after 24 hr ( $p = .12$ ) (Figures 3b and S2a). Fluorescence of the lac-T7 strain was on average 3.9- and 2.7-fold higher compared with the *trp*-T7 and *trp*\*-T7 strain, respectively. The lac-T7 strain also had a narrower distribution of fluorescence at the single-cell level. The broad fluorescence profile seen in the *trp* strains




**FIGURE 2** Experiment layout and fluorescence per OD after 7, 13, and 20 hr for strains harboring the trp-T7 or trp\*-T7 expression system. The strains were grown with different starting concentrations of tryptophan (trp) or yeast extract (YE). (a) Experiment scheme. (b) Fluorescence per OD for strains trp-T7 and trp\*-T7 with mCherry expressed from low-copy plasmid pSEVA. (c) Fluorescence per OD for strains trp-T7 and trp\*-T7 with mCherry expressed from high-copy plasmid pCDF. The different time points are shown in bright (7 hr), medium (13 hr), or dark (20 hr) shades of blue or orange for strains grown with tryptophan or YE, respectively. The fluorescence per OD was calculated as the average of three biological replicates. The standard deviations are shown as error bars [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

could be the result of heterogeneous transport and degradation rates of tryptophan, resulting in differences in the *trp* operon induction response (Yanofsky, Horn, & Gollnick, 1991), and such population heterogeneity could be improved by for example engineering constitutive expression of tryptophan permeases. After 24 hr, the difference between the *trp* and *lac* strains had decreased, with the *lac*-T7 strain having an average fluorescence that was 1.8- and 1.3-fold higher compared with the *trp*-T7 and *trp*\*-T7 strain, respectively. There was a subpopulation of low- and nonexpressing cells forming in the *lac*-T7 strain, while expression in the *trp* strains was homogenous (Figures 3b and 2c).

For the pCDF-mCherry strains, fluorescence was significantly higher for the *trp*\*-T7 compared with the *trp*-T7 strain after 8 hr of growth ( $p = .009$ ; Figures 3c and S2b). The *lac*-T7 was expressing higher levels of mCherry than both *trp* strains, with a 17.9- and 5.5-fold higher average fluorescence compared to *trp*-T7 and *trp*\*-T7, respectively. Similar to the pSEVA-mCherry cultures, the *lac*-T7

strain had a narrower expression profile compared to both *trp* strains. After 24 hr, the difference between the *trp* and *lac* strains had decreased slightly, with *lac*-T7 having 8.2- and 3.5-fold higher average fluorescence levels compared with *trp*-T7 and *trp*\*-T7, respectively (Figures 3c and S2b). However, mCherry expression in the *lac*-T7 strain was highly heterogeneous, with a large population not expressing or expressing only low amounts of the protein (Figures 3c and S2d). The expression heterogeneity of the *lac*-T7 system constitutes a significant problem when using the system in fed-batch or continuous production processes. It results in decreased productivity, especially as the subpopulation that stops producing the compound of interest usually quickly outgrows the producing population (Rugbjerg, Sarup-Lytzen, Nagy, & Sommer, 2018). Application of a less leaky version of the *lac*-T7 system or application of a constitutive *lac* permease could potentially improve population homogeneity of the IPTG-induced system, without affecting the expression levels negatively (Khlebnikov & Keasling, 2002).



**FIGURE 3** Histograms showing the fluorescence of strains *trp*-T7, *trp*\*-T7, and *lac*-T7 after 8 and 24 hr of growth. (a) Experiment scheme. (b) Fluorescence of strains with mCherry expressed from low-copy vector pSEVA. (c) Fluorescence of strains with mCherry expressed from high-copy vector pCDF. mCherry was induced by growth in tryptophan deplete medium (*trp*-T7, *trp*\*-T7) or by addition of 1 mM IPTG (*lac*-T7). Each histogram shown is a representative sample out of three biological replicates. IPTG, isopropyl  $\beta$ -D-1-thiogalactopyranoside [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



Overall, the average fluorescence was significantly lower for *trp*-T7 compared with the *lac*-T7 strains (Figures 3b,c and S2a,b), which may partly be explained by the tight regulation of the *trp* expression system. In addition, due to the *lacO* site present on the expression plasmids, mCherry expression is also repressed by endogenously expressed *lacI*. Comparison of mCherry fluorescence levels for *trp*-T7 and *trp*\*-T7 with or without IPTG induction showed that fluorescence per OD increased approximately 10% after IPTG addition for both pSEVA- and pCDF-based expression vectors (data not shown). Thus, higher protein expression levels could potentially be achieved by deleting the endogenous *lacI* gene.

### 3.4 | Evaluating the system for toxic protein production


Production of toxic proteins is challenging, as they can inhibit cell growth. An expression system with low basal level expression is preferred to avoid growth inhibition and decreased productivity when expressing such proteins. To investigate whether our constructs are suitable for toxic protein production, the *mazF* gene was cloned behind *pT7* on pSEVA and pCDF, resulting in plasmids pSEVA-*mazF* and pCDF-*mazF*. MazF is an ACA-specific endoribonuclease that stalls cell growth by ribosome independent messenger RNA cleavage (Muñoz-Gomez, Santos-Sierra, Berzal-Herranz, Lemmonier, & Mu, 2004; Zhang et al., 2003). The toxin has previously been used to demonstrate the suitability of inducible expression systems in *E. coli* (Guan et al., 2013). The *mazF* plasmids and respective control plasmids (no gene after *pT7*) were transformed to *trp*-T7 and *trp*\*-T7, resulting in eight different strains.

The experiment was carried out similarly as shown in Figure 2a; overnight cultures were inoculated in microtiter plates with 0–0.25 mM *trp* or 0–0.5% YE and growth was monitored for 24 hr (Figure 4a,b). At the highest tryptophan and the two highest YE concentrations, no significant difference could be seen in growth rate and final OD between the control strains and the strains harboring the *mazF* plasmids, with the exception of *trp*\*-T7 harboring pCDF-*mazF* where a slight decrease in growth rate could be seen also for the highest *trp* and YE concentrations (Figure 4b). At the lower concentrations, the MazF-expressing strains did not grow, grew with a much lower growth rate or reached a lower OD compared with the control strain for all constructs. Overall, the results show that the *trp*-T7 expression system is tight enough to be suitable for the expression of toxic proteins from high- and low-copy vectors in *E. coli*.


Interestingly, growth was resumed in the toxin-expressing strains toward the end of the microtiter plate cultivation. This phenomenon has been observed for the expression of other toxins in our lab and is possibly a result of single cells overcoming the toxic effect of MazF by mutating the toxin gene, the T7 polymerase or possibly by overexpressing the native antitoxin *mazE*. It is worth noting that growth was initiated at approximately the same timepoint in all cultures where growth was completely inhibited by toxin expression, indicating some common underlying mechanism.

### 3.5 | Applying *trp*-T7 for production of L-serine

L-serine is currently used in the pharmaceutical and cosmetics industry but has potential for extended industrial use as a building block for numerous chemicals (Werpy & Petersen, 2004). In previous



**FIGURE 4** Growth curves of strains harboring the *trp*-T7 expression systems and a *mazF*-expressing plasmid (pink lines) or the corresponding control plasmid (blue lines). A darker shade of each color represents a higher concentration of *trp* or YE added to the medium at the beginning of the cultivation. (a) Growth curves of strains *trp*-T7 and *trp*\*-T7 harboring *mazF* on pSEVA. (b) Growth curves of strains *trp*-T7 and *trp*\*-T7 harboring *mazF* on pCDF. The growth curves were calculated as an average of three biological replicates. YE, yeast extract [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



**FIGURE 5** Serine production in *trp*-T7-ser harboring the *trp*-T7 expression system and plasmid pSEVA-serACB. (a) Small batch fermentation. Growth of *trp*-T7-ser in 0.04 mM tryptophan (triangles, dark orange) and 0.5 mM tryptophan (triangles, blue). L-serine production of *trp*-T7-ser in 0.04 mM tryptophan (circles, orange) and 0.5 mM tryptophan (circles, bright blue). OD and serine concentrations were calculated as the average of three biological replicates. The standard deviations are shown as error bars. (b) Growth (orange) and production (blue) curves for fed-batch fermentation of *trp*-T7-ser. (c) Glucose consumed during the fed-batch fermentation of *trp*-T7-ser. (d) L-serine yield during the fed-batch fermentation of *trp*-T7-ser. (e) The productivity of the fed-batch fermentation of *trp*-T7-ser [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

studies, the L-serine pathway has been expressed from *pT7* by an integrated *lacUV5*-T7 polymerase induced by the addition of IPTG (Mundhada et al., 2016, 2017). To test L-serine production from our tryptophan repressible expression system, the *trp*-T7 cassette was integrated into an L-serine-tolerant strain of *E. coli* (Mundhada et al., 2017). We chose to only apply one of the cassettes, as mCherry production levels of *trp*-T7 and *trp*<sup>+</sup>-T7 were similar according to flow cytometry (Figure 3b). The strain was transformed with pSEVA27-sl-serA<sup>mut</sup>CB TIR1 (hereafter pSEVA-serACB) harboring an expression-optimized serine operon under control of *pT7* (Rennig et al., 2019), resulting in strain *trp*-T7-ser.

Overnight cultures of *trp*-T7-ser were inoculated in shake flasks with either 0.04 or 0.5 mM tryptophan to test the inducible and repressible function of the system, respectively. OD and serine were measured continuously for 24 hr. At the end of cultivation, serine titers reached 0.55 g/L (Figure 5a), slightly lower than the 0.62 g/L achieved when using *lac*-T7 (Rennig et al., 2019). A decrease in growth rate and final OD could be seen for the serine-producing strain, which may be due to carbon being utilized for serine production instead of growth.

To investigate the scalability of the process, we further tested the strain using fed-batch fermentation. Tryptophan was replaced by YE for initial repression of the serine pathway, as this is a more plausible alternative for large-scale fermentation. After calculating the tryptophan concentration in the YE and taking the larger cultivation volume in the bioreactor into account, 0.125% YE was added

to the medium before the start of cultivation. An overnight culture was inoculated in a 1 L bioreactor containing 250 ml medium supplemented with 0.125% YE. The glucose feed was started after 9 hr (at OD ~9) and growth and serine production were monitored for approximately 42 hr (Figure 5b). At the end of the fermentation, 26 g/L L-serine had been produced, approximately half of the 48 g/L reached when using *lac*-T7 (Rennig et al., 2019). The productivity was 0.73 g/Lh (Figure 5e), approximately 70% of that reached with the *lac*-T7 system. However, the same yield of L-serine per glucose (0.33 and 0.30 g/g for *trp*-T7 and *lac*-T7, respectively) was achieved; the lower titer and productivity may partly be explained by a decreased glucose uptake and growth rate in our *trp*-T7 strain (Figure 5c,d). Increasing the concentration of YE to 0.2%, which was used in the *lac*-T7 fermentation, might improve growth rate and glucose uptake but may also affect the timing of induction of the L-serine pathway. Another contributing factor could also be the lower expression levels from *trp*-T7 compared with *lac*-T7 that was observed previously (Figure 3b).

## 4 | CONCLUSIONS

Timing the induction of protein and product pathway expression can improve bioprocess performance and can serve as an important mode of control during fermentation. Low-cost expression systems are of special interest for large-scale fermentation and for the

production of low- or medium-value compounds. In this study, we demonstrated that the *E. coli* tryptophan promoter can be used to control T7-based expression of a GOI and that induction can be tuned by varying tryptophan or YE concentration in the start medium. Tryptophan depletion did not seem to impact growth in our experiments. However, for the production of other types of biochemicals or proteins, the growth rate and tryptophan uptake rate may be affected, which would impact the dynamics of the expression system. This could especially be the case if the expressed (heterologous) pathway enzymes or products have a high tryptophan content. In those cases, engineering approaches may have to be taken to tune the system and adapt to the new conditions.

Production levels were lower compared with using the lac-T7 expression system for both protein and biochemical production. Further improvement of the system could be made by engineering constitutive tryptophan uptake or using a *lacI* knockout strain to improve induction homogeneity or engineering the dynamic range using a mutant promoter library approach (Rohllhill et al., 2017). Overall, the low-cost and tight regulation of the trp-T7 system makes it an attractive alternative for large-scale production of biochemicals and proteins.

#### ACKNOWLEDGMENTS

The pOSIP KO and the PE-FLP plasmids were a gift from Drew Endy & Keith Shearwin (Addgene plasmids # 45985 and 45978, respectively). We would like to thank Hanne Bjerre Christensen and Lars Schrübbers for conducting the serine analysis. The authors would like to thank the Novo Nordisk Foundation (Grant number NNF16CC0020908 and NNF10CC1016517) for funding.

#### CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

#### ORCID

Jenny Landberg  <http://orcid.org/0000-0002-8094-0478>

Alex Toftgaard Nielsen  <http://orcid.org/0000-0001-6616-0187>

#### REFERENCES

- Bass, S. H., & Yansura, D. G. (2000). Application of the *E. coli* trp promoter. *Molecular Biotechnology*, 16(3), 253–260. <https://doi.org/10.1385/MB:16:3:253>
- Blazek, J., & Alper, H. S. (2013). Promoter engineering: Recent advances in controlling transcription at the most fundamental level. *Biotechnology Journal*, 8(1), 46–58. <https://doi.org/10.1002/biot.201200120>
- de Boer, H. A., Comstock, L. J., & Vasser, M. (1983). The tac promoter: A functional hybrid derived from the trp and lac promoters. *Proceedings of the National Academy of Sciences of the United States of America*, 80(1), 21–25. <https://doi.org/10.1073/pnas.80.1.21>
- Brewster, R. C., Jones, D. L., & Phillips, R. (2012). Tuning promoter strength through RNA polymerase binding site design in *Escherichia coli*. *PLoS Computational Biology*, 8(12):e1002811. <https://doi.org/10.1371/journal.pcbi.1002811>
- Brockman, I. M., & Prather, K. L. J. (2015). Dynamic metabolic engineering: New strategies for developing responsive cell factories. *Biotechnology Journal*, 10(9), 1360–1369. <https://doi.org/10.1002/biot.201400422>
- Brosius, J., Erfle, M., & Storella, J. (1985). Spacing of the –10 and –35 regions in the tac promoter. Effect on its in vivo activity. *Journal of Biological Chemistry*, 260(6), 3539–3541.
- Burg, J. M., Cooper, C. B., Ye, Z., Reed, B. R., Moreb, E. A., & Lynch, M. D. (2016). Large-scale bioprocess competitiveness: The potential of dynamic metabolic control in two-stage fermentations. *Current Opinion in Chemical Engineering*, 14, 121–136. <https://doi.org/10.1016/j.coche.2016.09.008>
- Cavaleiro, A. M., Kim, S. H., Seppälä, S., Nielsen, M. T., & Nørholm, M. H. H. (2015). Accurate DNA assembly and genome engineering with optimized uracil excision cloning. *ACS Synthetic Biology*, 4(9), 1042–1046. <https://doi.org/10.1021/acssynbio.5b00113>
- Chao, P., Law, S., Chen, P. T., & Hung, B. (2002). High production of heterologous proteins in *Escherichia coli* using the thermo-regulated T7 expression system. *Applied Microbiology and Biotechnology*, 58, 446–453. <https://doi.org/10.1007/s00253-001-0891-6>
- Chen, Y., Ho, J. M. L., Shis, D. L., Gupta, C., Long, J., Wagner, D. S., ... Bennett, M. R. (2018). Tuning the dynamic range of bacterial promoters regulated by ligand-inducible transcription factors. *Nature Communications*, 8, 64. <https://doi.org/10.1038/s41467-017-02473-5>
- Chevalet, L., Robert, A., Gueneau, F., Bonnefoy, J. Y., & Nguyen, T. (2000). Recombinant protein production driven by the tryptophan promoter is tightly controlled in ICONE 200, a new genetically engineered *E. coli* mutant. *Biotechnology and Bioengineering*, 69(4), 351–358. [https://doi.org/10.1002/1097-0290\(20000820\)69:4<351::AID-BIT1>3.0.CO;2-G](https://doi.org/10.1002/1097-0290(20000820)69:4<351::AID-BIT1>3.0.CO;2-G)
- Ellefson, J. W., Ledbetter, M. P., & Ellington, A. D. (2018). Directed evolution of a synthetic phylogeny of programmable Trp repressors article. *Nature Chemical Biology*, 14(4), 361–367. <https://doi.org/10.1038/s41589-018-0006-7>
- Elvin, C. M., Thompson, P. R., Argall, M. E., Philip Hendr, N., Stamford, P. J., Lilley, P. E., & Dixon, N. E. (1990). Modified bacteriophage lambda promoter vectors for overproduction of proteins in *Escherichia coli*. *Gene*, 87(1), 123–126. [https://doi.org/10.1016/0378-1119\(90\)90503-J](https://doi.org/10.1016/0378-1119(90)90503-J)
- Gronenborn, B. (1976). Overproduction of phage Lambda repressor under control of the lac promoter of *Escherichia coli*. *MGG Molecular & General Genetics*, 148(3), 243–250. <https://doi.org/10.1007/BF00332898>
- Guan, L., Liu, Q., Li, C., & Zhang, Y. (2013). Development of a Fur-dependent and tightly regulated expression system in *Escherichia coli* for toxic protein synthesis. *BMC Biotechnology*, 13, 25. <https://doi.org/10.1186/1472-6750-13-25>
- Gupta, A., Reizman, I. M. B., Reisch, C. R., & Prather, K. L. J. (2017). Dynamic regulation of metabolic flux in engineered bacteria using a pathway-independent quorum-sensing circuit. *Nature Biotechnology*, 35(3), 273–279. <https://doi.org/10.1038/nbt.3796>
- Gustavsson, M., & Lee, S. Y. (2016). Prospects of microbial cell factories developed through systems metabolic engineering. *Microbial Biotechnology*, 9(5), 610–617. <https://doi.org/10.1111/1751-7915.12385>
- Gvzdaitis, B. G., Beil, S., Kreibaum, U., Simutis, R., Havlik, I., Dors, M., & Schneider, F. (1994). Temperature control in fermenters: Application of neural nets and feedback control in breweries. *Journal of the Institute of Brewing*, 100(3–4), 99–104.
- Jones, K. L., Kim, S., & Keasling, J. D. (2000). Low-copy plasmids can perform as well as or better than high-copy plasmids for metabolic engineering of bacteria. *Metabolic Engineering*, 338(2), 328–338. <https://doi.org/10.1006/mben.2000.0161>
- Keasling, J. D. (2012). Synthetic biology and the development of tools for metabolic engineering. *Metabolic Engineering*, 14(3), 189–195. <https://doi.org/10.1016/j.mben.2012.01.004>
- Khlebnikov, A., & Keasling, J. D. (2002). Effect of lacY expression on homogeneity of induction from the P tac and P trc promoters by natural and synthetic inducers. *Biotechnology Progress*, 18, 672–674.
- Kildegaard, K. R., Hallström, B. M., Blicher, T. H., Sonnenschein, N., Jensen, N. B., Sherstyk, S., ... Borodina, I. (2014). Evolution reveals a

- glutathione-dependent mechanism of 3-hydroxypropionic acid tolerance. *Metabolic Engineering*, 26, 57–66. <https://doi.org/10.1016/j.ymben.2014.09.004>
- Kim, S. H., Cavaleiro, A. M., Rennig, M., & Nørholm, M. H. H. (2016). SEVA linkers: A versatile and automatable DNA backbone exchange standard for synthetic biology. *ACS Synthetic Biology*, 5(10), 1177–1181. <https://doi.org/10.1021/acssynbio.5b00257>
- Kim, E. M., Woo, H. M., Tian, T., Yilmaz, S., Javidpour, P., Keasling, J. D., & Lee, T. S. (2017). Autonomous control of metabolic state by a quorum sensing (QS)-mediated regulator for bisabolene production in engineered *E. coli*. *Metabolic Engineering*, 44(11), 325–336. <https://doi.org/10.1016/j.ymben.2017.11.004>
- Kincade, J. M., & deHaset, P. L. (1991). Bacteriophage lambda promoters pL and pR sequence determinants of in vivo activity and of sensitivity to the DNA gyrase inhibitor, coumermycin. *Gene*, 97(1), 7–12. [https://doi.org/10.1016/0378-1119\(91\)90003-T](https://doi.org/10.1016/0378-1119(91)90003-T)
- Liu, P., Zhu, X., Tan, Z., Zhang, X., & Ma, Y. (2016). Construction of *Escherichia coli* cell factories for production of organic acids and alcohols. In Q. Ye, J. Bao & J.-J. Zhong (Eds.), *Bioreactor engineering research and industrial applications I: Cell factories* (pp. 107–140). Berlin, Heidelberg: Springer. [https://doi.org/10.1007/10\\_2014\\_294](https://doi.org/10.1007/10_2014_294)
- Liu, X., Gupta, S. T. P., Bhimsaria, D., Reed, J. L., Rodriguez-Martinez, J. A., Ansari, A. Z., & Raman, S. (2019). De novo design of programmable inducible promoters. *Nucleic Acids Research*, 47(19), 10452–10463. <https://doi.org/10.1093/nar/gkz772>
- Mundhada, H., Schneider, K., Christensen, H. B., & Nielsen, A. T. (2016). Engineering of high yield production of L-serine in *Escherichia coli*. *Biotechnology and Bioengineering*, 113(4), 807–816. <https://doi.org/10.1002/bit.25844>
- Mundhada, H., Seoane, J. M., Schneider, K., Koza, A., Christensen, H. B., Klein, T., ... Nielsen, A. T. (2017). Increased production of L-serine in *Escherichia coli* through adaptive laboratory evolution. *Metabolic Engineering*, 39, 141–150. <https://doi.org/10.1016/j.ymben.2016.11.008>
- Muñoz-Gomez, A. J., Santos-Sierra, S., Berzal-Herranz, A., Lemmonier, M., & Mu, A. J. (2004). Insights into the specificity of RNA cleavage by the *Escherichia coli* MazF toxin. *FEBS Letters*, 567, 316–320. <https://doi.org/10.1016/j.febslet.2004.05.005>
- Nielsen, J., & Keasling, J. D. (2016). Engineering cellular metabolism. *Cell*, 164(6), 1185–1197. <https://doi.org/10.1016/j.cell.2016.02.004>
- Nour-Eldin, H. H., Hansen, B. G., Nørholm, M. H. H., Jensen, J. K., & Halkier, B. A. (2006). Advancing uracil-excision based cloning towards an ideal technique for cloning PCR fragments. *Nucleic Acids Research*, 34(18), 1–8. <https://doi.org/10.1093/nar/gkl635>
- Rennig, M., Mundhada, H., Wordofa, G. G., Gerngross, D., Wulff, T., Worberg, A., ... Nørholm, M. (2019). Industrializing a bacterial strain for L-serine production through translation initiation optimization. *ACS Synthetic Biology*, 8, 2347–2358. <https://doi.org/10.1021/acssynbio.9b00169>
- Rohlhill, J., Sandoval, N. R., & Papoutsakis, E. T. (2017). Sort-Seq approach to engineering a formaldehyde-inducible promoter for dynamically regulated *Escherichia coli* growth on methanol. *ACS Synthetic Biology*, 6, 1584–1595. <https://doi.org/10.1021/acssynbio.7b00114>. research-article.
- Rosano, G. L., & Ceccarelli, E. A. (2014). Recombinant protein expression in *Escherichia coli*: Advances and challenges. *Frontiers in Microbiology*, 5, 172. <https://doi.org/10.3389/fmicb.2014.00172>
- Rugbjerg, P., Sarup-Lytzen, K., Nagy, M., & Sommer, M. (2018). Synthetic addiction extends the productive life time of engineered *Escherichia coli* populations. *Proceedings of the National Academy of Sciences of the United States of America*, 115(10), 2347–2352. <https://doi.org/10.1073/pnas.1718622115>
- Saida, F., Uzan, M., Odaert, B., & Bontems, F. (2006). Expression of highly toxic genes in *E. coli*: Special strategies and genetic tools. *Current Protein and Peptide Science*, 7(1), 47–56. <https://doi.org/10.2174/138920306775474095>
- Somerville, R. L. (1988). The trp promoter of *Escherichia coli* and its exploitation in the design of efficient protein production systems. *Biotechnology and Genetic Engineering Reviews*, 6(1), 1–41. <https://doi.org/10.1080/02648725.1988.10647844>
- St-Pierre, F., Cui, L., Priest, D. G., Endy, D., Dodd, I. B., & Shearwin, K. E. (2013). One-step cloning and chromosomal integration of DNA. *ACS Synthetic Biology*, 2(9), 537–541. <https://doi.org/10.1021/sb400021j>
- Studier, F. W. (2005). Protein production by auto-induction in high density shaking cultures. *Protein Expression and Purification*, 41(1), 207–234. <https://doi.org/10.1016/j.pep.2005.01.016>
- Studier, F. W., & Moffatt, B. A. (1986). Use of bacteriophage T7 RNA polymerase to direct selective high-level expression of cloned genes. *Journal of Molecular Biology*, 189(1), 113–130. [https://doi.org/10.1016/0022-2836\(86\)90385-2](https://doi.org/10.1016/0022-2836(86)90385-2)
- Terpe, K. (2006). Overview of bacterial expression systems for heterologous protein production: From molecular and biochemical fundamentals to commercial systems. *Applied Microbiology and Biotechnology*, 72(2), 211–222. <https://doi.org/10.1007/s00253-006-0465-8>
- Valdez-Cruz, N. A., Caspeta, L., Pérez, N. O., Ramírez, O. T., & Trujillo-Roldán, M. A. (2010). Production of recombinant proteins in *E. coli* by the heat inducible expression system based on the phage lambda pL and/or pR promoters. *Microbial Cell Factories*, 9(18), 1–16.
- Werpy, T., & Petersen, G. (2004). *Top value added chemicals from biomass volume I—Results of screening for potential candidates from sugars and synthesis gas*. US Department of Energy Efficiency and Renewable Energy. <https://doi.org/10.2172/15008859>
- Yanofsky, C. (1981). Attenuation in the control of expression of bacterial operons. *Nature*, 289(5800), 751–758. <https://doi.org/10.1038/289751a0>
- Yanofsky, C., Horn, V., & Gollnick, P. (1991). Physiological studies of tryptophan transport and tryptophanase operon induction in *Escherichia coli*. *Journal of Bacteriology*, 173(19), 6009–6017. <https://doi.org/10.1128/jb.173.19.6009-6017.1991>
- Zargar, A., Quan, D. N., & Bentley, W. E. (2016). Enhancing intercellular coordination: Rewiring quorum sensing networks for increased protein expression through autonomous induction. *ACS Synthetic Biology*, 5, 923–928. <https://doi.org/10.1021/acssynbio.5b00261>
- Zhang, Y., Zhang, J., Hoefflich, K. P., Ikura, M., Qing, G., & Inouye, M. (2003). MazF cleaves cellular mRNAs specifically at ACA to block protein synthesis in *Escherichia coli*. *Molecular Cell*, 12(4), 913–923. [https://doi.org/10.1016/S1097-2765\(03\)00402-7](https://doi.org/10.1016/S1097-2765(03)00402-7)

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Landberg J, Mundhada H, Nielsen AT. An autoinducible trp-T7 expression system for production of proteins and biochemicals in *Escherichia coli*. *Biotechnology and Bioengineering*. 2020;117:1513–1524. <https://doi.org/10.1002/bit.27297>