

Received 20 October 2016

Accepted 24 October 2016

Edited by H. Stoeckli-Evans, University of  
Neuchâtel, Switzerland

**Keywords:** crystal structure; carbazole derivatives; phenylsulfonyl; benzofuran; thiophene; hydrogen bonding; C—H... $\pi$  interactions; offset  $\pi$ – $\pi$  interactions.

**CCDC reference:** 1473995

**Supporting information:** this article has supporting information at journals.iucr.org/e

# Crystal structures of two new carbazole derivatives: 12-(4-nitrophenyl)-7-phenylsulfonyl-7*H*-benzofuro-[2,3-*b*]carbazole and 2-methyl-4-(4-nitrophenyl)-9-phenylsulfonyl-9*H*-thieno[2,3-*b*]carbazole


K. Swaminathan,<sup>a</sup> P. Narayanan,<sup>a</sup> K. Sethusankar,<sup>a\*</sup> Velu Saravanan<sup>b</sup> and Arasambattu K. Mohanakrishnan<sup>b</sup>

<sup>a</sup>Department of Physics, RKM Vivekananda College (Autonomous), Chennai 600 004, India, and <sup>b</sup>Department of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India. \*Correspondence e-mail: ksethusankar@yahoo.co.in


The title compounds, C<sub>30</sub>H<sub>18</sub>N<sub>2</sub>O<sub>5</sub>S, (I), and C<sub>27</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>S<sub>2</sub>, (II), are carbazole derivatives with a phenylsulfonyl group and a nitrophenyl group attached to the carbazole moiety in identical positions in both molecules. A benzofuran ring system in (I) and a methylthiophene ring in (II) are fused with the respective carbazole moieties on the same sides. The mean plane of the carbazole ring system makes a dihedral angle of 3.17 (7)° with the benzofuran ring system in (I) and a dihedral angle of 3.39 (11)° with the methylthiophene ring in (II), implying that both fused units are essentially planar. The mean planes of the carbazole ring systems in both the compounds are almost orthogonal to the respective nitro-substituted phenyl rings, making dihedral angles of 75.64 (10) and 77.63 (12)° in compounds (I) and (II), respectively. In (I), the phenylsulfonyl ring system is positionally disordered with a refined occupancy ratio of 0.63 (2):0.37 (2). In both compounds, the molecular structures are stabilized by intramolecular C—H...O hydrogen bonds, generating *S*(6) ring motifs with the sulfone group O atoms. In the crystal of compound (I), molecules are linked by pairs of C—H...O hydrogen bonds, which generate *R*<sub>2</sub><sup>2</sup>(18) inversion dimers, and interconnected by *C*(14) chains running along the *c*-axis direction, whereas in compound (II), the C—H...O hydrogen bonds generate *R*<sub>4</sub><sup>3</sup>(37) ring motifs. In the crystals of both compounds, C—H...O hydrogen-bonded sheets are formed lying parallel to (10 $\bar{1}$ ). In addition, C—H... $\pi$  and offset  $\pi$ – $\pi$  interactions [intercentroid distance = 3.7158 (14) Å in (I) and 3.9040 (15) Å in (II)] are also present in the crystals of both compounds.

## 1. Chemical context

Carbazole and its derivatives are interesting compounds owing to their applications in pharmacy and molecular electronics. Carbazole derivatives exhibit various biological activities such as antitumor (Itoigawa *et al.*, 2000), anti-oxidative (Tachibana *et al.*, 2001), anti-inflammatory and antimutagenic (Ramsewak *et al.*, 1999). They also exhibit electroactivity and luminescence and are considered to be potential candidates for electronic applications, such as colour displays, organic semiconductors, laser and solar cells (Friend, *et al.* 1999). Tetrahydrocarbazole systems are present in the framework of a number of indole-type alkaloids of biological interest (Saxton, 1983). Carbazole-based heterocyclic polymer systems can be chemically or electrochemically polymerized to give products with a number of applications, such as rechargeable batteries (Sacak, 1999) and electrochromic displays (Santhanam & Sundaresan, 1986). This enables their



use as suitable building blocks for the design and synthesis of molecular glasses, which are widely studied as components of electroactive and photoactive materials (Zhang *et al.*, 2004). Against this background, the X-ray structure determination of the title compounds, (I) and (II), has been carried out to study their structural aspects and the results are presented here.



## 2. Structural commentary


The molecular structures of the title compounds, (I) and (II), are illustrated in Figs. 1 and 2, respectively. In both compounds, the carbazole ring systems (N1/C1–C12) are essentially planar with maximum deviations of 0.089 (3) and

0.089 (3) Å for atom C10 in compounds (I) and (II), respectively. In compound (I), the benzofuran moiety (O5/C10/C11/CC25–C30) is essentially planar with a maximum deviation of 0.021 (3) Å for atom C10 while the phenylsulfonyl ring system is positionally disordered with a refined occupancy factor of 0.63 (2): 0.37 (2).


The mean planes of the carbazole ring systems make dihedral angles of 3.17 (7) and 3.39 (11)°, respectively, with the benzofuran ring in (I) and the methylthiophene ring in (II), indicating that the ring systems they are essentially coplanar. The nitrophenyl rings in compounds (I) and (II) are inclined to the carbazole ring system by 75.64 (10) and 77.63 (12)°, respectively. The NO<sub>2</sub> groups are inclined to the benzene ring (C19–C24) to which they are attached by 9.8 (4)° in (I) and 9.3 (3)° in (II). The phenylsulfonyl ring (C13–C18) is almost normal to the nitro-substituted phenyl ring (C19–C24) with a dihedral angle of 84.7 (2)° in (I) and 83.98 (17)° in (II).

In both compounds, as a result of the electron-withdrawing character of the phenylsulfonyl group, the N–C<sub>sp<sup>2</sup></sub> bond lengths are longer than the mean value of 1.355 (14) Å for N–C bond lengths (CSD; Groom *et al.*, 2016). Atom S1 has a distorted tetrahedral geometry. The widening of the O1=S1=O2 angle and narrowing of the N1–S1–C13 angle from the ideal tetrahedral values are attributed to the Thorpe–Ingold effect (Bassindale, 1984). The widening of the angles may be due to the repulsive interaction between the two short S=O bonds.

The sums of the bond angles around atom N1 are 349.58° in (I) and 351.18° in (II), intermediate between *sp*<sup>2</sup> and *sp*<sup>3</sup>



**Figure 1**  
The molecular structure of compound (I), showing the atom labelling. Displacement ellipsoids are drawn at the 30% probability level. Intramolecular C2–H2···O1 and C9–H9···O2 hydrogen bonds, which generate two *S*(6) ring motifs, are shown as dashed lines (see Table 1). For the sake of clarity, the minor component of the disordered phenylsulfonyl ring has been omitted.



**Figure 2**  
The molecular structure of compound (II), showing the atom labelling. Displacement ellipsoids are drawn at the 30% probability level. Intramolecular C2–H2···O1 and C9–H9···O2 hydrogen bonds, which generate two *S*(6) ring motifs, are shown as dashed lines (see Table 2).

**Table 1**  
Hydrogen-bond geometry (Å, °) for (I).

$Cg_1$  is the centroid of the furan ring O5/C10/C11/C25/C30 and  $Cg_3$  is the centroid of the benzene ring C1–C6.

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
C2–H2···O1	0.93	2.41	3.015 (9)	122
C9–H9···O2	0.93	2.28	2.849 (9)	119
C14–H14···O4 <sup>i</sup>	0.93	2.55	3.296 (7)	138
C20–H20···O2 <sup>ii</sup>	0.93	2.53	3.454 (11)	172
C16–H16··· $Cg_1^{iii}$	0.93	2.76	3.676 (6)	169
C23–H23··· $Cg_3^{iv}$	0.93	2.99	3.908 (4)	169

Symmetry codes: (i)  $x - \frac{1}{2}, -y + \frac{5}{2}, z - \frac{1}{2}$ ; (ii)  $-x + 1, -y + 2, -z$ ; (iii)  $-x, -y + 2, -z$ ; (iv)  $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$ .


**Table 2**  
Hydrogen-bond geometry (Å, °) for (II).

$Cg_1$  is the centroid of the thiophene ring S2/C10/C11/C25/C26 and  $Cg_3$  is centroid of the benzene ring C1–C6.


$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
C2–H2···O1	0.93	2.38	2.970 (4)	121
C9–H9···O2	0.93	2.33	2.907 (3)	120
C3–H3···O4 <sup>i</sup>	0.93	2.48	3.356 (4)	157
C14–H14···O4 <sup>ii</sup>	0.93	2.58	3.335 (4)	139
C20–H20···O2 <sup>iii</sup>	0.93	2.53	3.368 (5)	150
C16–H16··· $Cg_1^{iv}$	0.93	2.89	3.812 (4)	172
C23–H23··· $Cg_3^v$	0.93	2.75	3.646 (4)	163

Symmetry codes: (i)  $x, y + 1, z$ ; (ii)  $x + \frac{1}{2}, -y - \frac{1}{2}, z + \frac{1}{2}$ ; (iii)  $-x + 1, -y, -z + 2$ ; (iv)  $-x + 2, -y, -z + 2$ ; (v)  $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$ .

hybridization. In both compounds, the molecular structure is stabilized by intramolecular C–H···O hydrogen bonds, which generate  $S(6)$  ring motifs with the sulfone oxygen atoms (Tables 1 and 2).




**Figure 3**  
The crystal packing of compound (I), viewed normal to the  $(10\bar{1})$  plane, showing C–H···O hydrogen bonds that generate  $R_2^2(18)$  inversion dimers and the C–H···O hydrogen bonds that generate  $C(14)$  chains running along the  $c$ -axis direction (see Table 1 for details). H atoms not involved in the hydrogen bonding and the benzofuran ring have been excluded for clarity.



**Figure 4**  
The crystal packing of compound (II), viewed along the  $ac$  diagonal, showing the intermolecular C–H···O hydrogen bonds (see Table 2), which generate  $R_2^2(37)$  ring motifs and form sheets lying parallel to the  $(10\bar{1})$  plane. H atoms not involved in hydrogen bonding have been excluded for clarity.

### 3. Supramolecular features

In the crystal of compound (I), molecules are linked *via* pairs of C20–H20···O2<sup>ii</sup> hydrogen bonds (Table 1), forming  $R_2^2(18)$  inversion dimers. The molecules are also interconnected by C14–H14···O4<sup>i</sup> hydrogen bonds, which generate  $C(14)$  chains. These interactions result in the formation of sheets parallel to  $(10\bar{1})$ . The crystal packing also features C16–H16··· $Cg_1^{iii}$  and C23–H23··· $Cg_3^{iv}$  interactions (Table 1 and Fig. 3;  $Cg_1$  and  $Cg_3$  are the centroids of the rings C10/C11/C25/30/O5 and C1–C6, respectively). There are also offset  $\pi$ – $\pi$  interactions present [ $Cg_4 \cdots Cg_4(-x + 1, -y + 2, -z) = 3.7158(14)$  Å, interplanar distance =  $3.472(1)$  Å, slippage =  $1.324(11)$  Å;  $Cg_4$  is the centroid of the C7–C12 ring]; see Table 1 and Fig. 3.



**Figure 5**  
The crystal packing of compound (II), viewed along the  $c$  axis, showing the C–H···O intermolecular hydrogen bonds which generate  $R_2^2(18)$  inversion dimers (see Table 2). H atoms not involved in hydrogen bonding have been excluded for clarity.

**Table 3**  
Experimental details.

	(I)	(II)
Crystal data		
Chemical formula	C <sub>30</sub> H <sub>18</sub> N <sub>2</sub> O <sub>5</sub> S	C <sub>27</sub> H <sub>18</sub> N <sub>2</sub> O <sub>4</sub> S <sub>2</sub>
<i>M<sub>r</sub></i>	518.52	498.55
Crystal system, space group	Monoclinic, <i>P</i> <sub>2</sub> <sub>1</sub> / <i>n</i>	Monoclinic, <i>P</i> <sub>2</sub> <sub>1</sub> / <i>n</i>
Temperature (K)	296	296
<i>a</i> , <i>b</i> , <i>c</i> (Å)	12.1347 (5), 12.0708 (5), 17.6391 (7)	12.5052 (8), 11.2594 (6), 17.0731 (9)
β (°)	108.617 (2)	102.914 (2)
<i>V</i> (Å <sup>3</sup> )	2448.50 (17)	2343.1 (2)
<i>Z</i>	4	4
Radiation type	Mo <i>K</i> α	Mo <i>K</i> α
μ (mm <sup>-1</sup> )	0.18	0.27
Crystal size (mm)	0.25 × 0.20 × 0.10	0.35 × 0.30 × 0.25
Data collection		
Diffractometer	Bruker Kappa APEXII CCD	Bruker Kappa APEXII CCD
Absorption correction	Multi-scan ( <i>SADABS</i> ; Bruker, 2008)	Multi-scan ( <i>SADABS</i> ; Bruker, 2008)
<i>T</i> <sub>min</sub> , <i>T</i> <sub>max</sub>	0.958, 0.982	0.911, 0.936
No. of measured, independent and observed [ <i>I</i> > 2σ( <i>I</i> )] reflections	32749, 4318, 2814	44472, 5100, 3714
<i>R</i> <sub>int</sub>	0.040	0.034
(sin θ/λ) <sub>max</sub> (Å <sup>-1</sup> )	0.595	0.639
Refinement		
<i>R</i> [ <i>F</i> <sup>2</sup> > 2σ( <i>F</i> <sup>2</sup> )], <i>wR</i> ( <i>F</i> <sup>2</sup> ), <i>S</i>	0.043, 0.131, 1.05	0.055, 0.163, 1.08
No. of reflections	4318	5100
No. of parameters	401	317
No. of restraints	140	0
H-atom treatment	H-atom parameters constrained	H-atom parameters constrained
Δρ <sub>max</sub> , Δρ <sub>min</sub> (e Å <sup>-3</sup> )	0.28, -0.21	0.44, -0.36

Computer programs: *APEX2* and *SAINT* (Bruker, 2008), *SHELXS97* and *SHELXL97* (Sheldrick, 2008), *ORTEP-3 for Windows* (Farrugia, 2012), *Mercury* (Macrae *et al.*, 2008) and *PLATON* (Spek, 2009).

In the crystal of compound (II), molecules are linked by C3—H3···O4<sup>i</sup> and C14—H14···O<sup>ii</sup><sub>4</sub> hydrogen bonds (Table 2), which result in the formation of *R*<sub>4</sub><sup>3</sup>(37) ring motifs (Fig. 4). The crystal packing also features pairs of C20—H20···O2<sup>iii</sup> hydrogen bonds, which generate *R*<sub>2</sub><sup>2</sup>(18) inversion dimers (Fig. 5), which are interconnected by C16—H16···Cg1<sup>iv</sup> and C23—H23···Cg3<sup>v</sup> interactions [Cg1 and Cg3 are the centroids of rings C10/C11/C25/C26/S2 and C1—C6, respectively]. These interactions result in the formation of sheets parallel to (10 $\bar{1}$ ). There are also offset π–π interactions present [Cg4···Cg4(−*x* + 1, −*y*, −*z* + 2) = 3.9040 (15) Å, interplanar distance = 3.791 (1) Å, slippage 0.932 Å; Cg4 is the centroid of the C7–C12, ring]; see Table 2 and Figs. 4 and 5.

#### 4. Database survey

A search of Cambridge Structural Database (CSD version 5.37; last update May 2016; Groom *et al.*, 2016) yielded four hits for 7*H*-[1] benzofuran[2,3-*b*]carbazole and 47 hits for 9-(phenylsulfonyl)-9*H*-carbazole. However, the compound 7-phenylsulfonyl-7*H*-benzofuran[2,3-*b*]carbazole (EYOFEE01; Panchatcharam *et al.*, 2011), which crystallizes in *P*<sub>2</sub><sub>1</sub>/*c* is the closest analogue of compound (I). The compound 2-methyl-9-(phenylsulfonyl)-9*H*-thieno[2,3-*b*]carbazole (IQOBIA; Sureshbabu *et al.*, 2011), which crystallizes in space group *P*<sub>2</sub><sub>1</sub>/*c*, is the closest analogue of compound (II). The crystal packing of the title compounds is stabilized by C—H···O, C—H···π and π–π interactions but the related structures

(EYOFEE01; Panchatcharam *et al.*, 2011) exhibit C—H···π and π–π interactions only.

#### 5. Synthesis and crystallization

**Compound (I):** A solution of [3-(4-nitrobenzoyl)-1-(phenylsulfonyl)-1*H*-indol-2-yl]methyl pivalate (0.1 g, 1.92 mmol), anhydrous SnCl<sub>4</sub> (0.06 g, 2.30 mmol) and benzofuran (0.027 g, 2.30 mmol) in dry DCE (10 ml) was stirred at room temperature under nitrogen for 3 h. After completion of the reaction (monitored by TLC), it was poured into ice–water (100 ml). The organic layer was separated and the aqueous layer was extracted with DCM (2 × 20 ml). The combined extract was washed with water (3 × 50ml) and dried (Na<sub>2</sub>SO<sub>4</sub>). Removal of the solvent by column chromatographic purification (silica gel; hexane–ethyl acetate, 8:2) yielded compound (I) as a colourless solid (0.073 g, 74%). Colourless block-like crystals were obtained by slow evaporation of a solution of (I) in ethyl acetate at room temperature (m.p. 589–591 K).

**Compound (II):** A solution of [3-(4-nitrobenzoyl)-1-(phenylsulfonyl)-*H*-indol-2-yl]methyl pivalate (0.1 g, 1.92 mmol), anhydrous SnCl<sub>4</sub> (0.06 g, 2.30 mmol) and 2-methylthiophene (0.024 g, 2.30 mmol) in dry DCE (10 ml) was stirred at room temperature under nitrogen atmosphere for 3 h. After the completion of the reaction (monitored by TLC), it was poured into ice–water (100 ml), the organic layer was separated and the aqueous layer was extracted with DCM (2 × 20ml). The combined extract was washed with water (3 ×

50 ml) and dried ( $\text{Na}_2\text{SO}_4$ ). Removal of the solvent by column chromatographic purification (silica gel; hexane–ethyl acetate, 8:2) yielded compound (II) as a colourless solid (0.067 g, 72%). Colourless block-like crystals were obtained by slow evaporation of a solution of (II) in ethyl acetate at room temperature (m.p. 531–533 K).

## 6. Refinement

Crystal data, data collection and structure refinement details for compounds (I) and (II) are summarized in Table 3. The positions of the hydrogen atoms were localized from the difference electron-density maps. The C-bound H atoms were treated as riding atoms: C–H = 0.93–0.96 Å with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C-methyl})$  and  $1.2U_{\text{eq}}(\text{C})$  for other H atoms. In compound (I), the phenylsulfonyl ring (C13–C18) is positionally disordered with a refined occupancy ratio of 0.63 (2): 0.37 (2). The bond distances of the disordered components were restrained using standard similarity restraint SADI [*SHELXL97*; Sheldrick, 2008] with s.u. of 0.01 Å. Ellipsoid displacement (SIMU and DELU) restraints were also applied to the disordered ring. The methyl groups were allowed to rotate, but not to tip, to best fit the electron density.

## Acknowledgements

The authors thank Dr Jagan and Dr Babu Varghese, Senior Scientific Officer, SAIF, IIT Madras, Chennai, India, for the data collection.

## References

- Bassindale, A. (1984). *The Third Dimension in Organic Chemistry*, ch. 1, p. 11. New York: John Wiley and Sons.
- Bruker (2008). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Friend, R. H., Gymer, R. W., Holmes, A. B., Burroughes, J. H., Marks, R. N., Taliani, C., Bradley, D. D. C., Dos Santos, D. A., Brédas, J. L., Lögdlund, M. & Salaneck, W. R. (1999). *Nature*, **397**, 121–128.
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
- Itoigawa, M., Kashiwada, Y., Ito, C., Furukawa, H., Tachibana, Y., Bastow, K. F. & Lee, K. H. (2000). *J. Nat. Prod.* **63**, 893–897.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Panchatcharam, R., Dhayalan, V., Mohanakrishnan, A. K., Chakkaravarthi, G. & Manivannan, V. (2011). *Acta Cryst.* **E67**, o2829.
- Ramsewak, R. S., Nair, M. G., Strasburg, G. M., DeWitt, D. L. & Nitiss, J. L. (1999). *J. Agric. Food Chem.* **47**, 444–447.
- Sacak, M. (1999). *J. Appl. Polym. Sci.* **74**, 1792–1796.
- Santhanam, K. S. V. & Sundaresan, N. S. (1986). *Indian J. Technol.* **24**, 417–422.
- Saxton, J. E. (1983). Editor. *Heterocyclic Compounds*, Vol. 25, *The Monoterpenoid Indole Alkaloids*, ch. 8 and 11. New York: Wiley.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Sureshbabu, R., Saravanan, V., Dhayalan, V. & Mohanakrishnan, A. K. (2011). *Eur. J. Org. Chem.* 922–935.
- Tachibana, Y., Kikuzaki, H., Lajis, N. H. & Nakatani, N. (2001). *J. Agric. Food Chem.* **49**, 5589–5594.
- Zhang, Q., Chen, J., Cheng, Y., Wang, L., Ma, D., Jing, X. & Wang, F. (2004). *J. Mater. Chem.* **14**, 895–900.

## supporting information

*Acta Cryst.* (2016). E72, 1739-1743 [https://doi.org/10.1107/S2056989016016996]

## Crystal structures of two new carbazole derivatives: 12-(4-nitrophenyl)-7-phenylsulfonyl-7*H*-benzofuro[2,3-*b*]carbazole and 2-methyl-4-(4-nitrophenyl)-9-phenylsulfonyl-9*H*-thieno[2,3-*b*]carbazole

**K. Swaminathan, P. Narayanan, K. Sethusankar, Velu Saravanan and Arasambattu K. Mohanakrishnan**

### Computing details

For both compounds, data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT* (Bruker, 2008); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008) and *PLATON* (Spek, 2009).

### (I) 12-(4-Nitrophenyl)-7-phenylsulfonyl-7*H*-benzofuro[2,3-*b*]carbazole

#### Crystal data

$C_{30}H_{18}N_2O_5S$

$M_r = 518.52$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 12.1347$  (5) Å

$b = 12.0708$  (5) Å

$c = 17.6391$  (7) Å

$\beta = 108.617$  (2)°

$V = 2448.50$  (17) Å<sup>3</sup>

$Z = 4$

$F(000) = 1072$

$D_x = 1.407$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 4318 reflections

$\theta = 2.4$ – $25.0$ °

$\mu = 0.18$  mm<sup>-1</sup>

$T = 296$  K

Block, colourless

$0.25 \times 0.20 \times 0.10$  mm

#### Data collection

Bruker Kappa APEXII CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega$  &  $\varphi$  scans

Absorption correction: multi-scan  
(SADABS; Bruker, 2008)

$T_{\min} = 0.958$ ,  $T_{\max} = 0.982$

32749 measured reflections

4318 independent reflections

2814 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.040$

$\theta_{\max} = 25.0$ °,  $\theta_{\min} = 2.4$ °

$h = -14 \rightarrow 11$

$k = -14 \rightarrow 14$

$l = -18 \rightarrow 20$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.043$   
 $wR(F^2) = 0.131$   
 $S = 1.05$   
 4318 reflections  
 401 parameters  
 140 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0594P)^2 + 0.7253P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.002$   
 $\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.21 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional R-factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and R-factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C8	0.36384 (19)	0.94124 (18)	0.02090 (14)	0.0501 (6)	
C9	0.3369 (2)	0.96522 (19)	-0.05953 (15)	0.0552 (6)	
H9	0.3249	0.9108	-0.0987	0.066*	
C10	0.3295 (2)	1.07565 (19)	-0.07650 (14)	0.0512 (6)	
C25	0.2981 (2)	1.22953 (19)	-0.14680 (14)	0.0540 (6)	
C26	0.2698 (2)	1.3006 (2)	-0.21190 (16)	0.0669 (7)	
H26	0.2531	1.2750	-0.2641	0.080*	
C27	0.2680 (2)	1.4115 (2)	-0.19423 (18)	0.0729 (8)	
H27	0.2483	1.4631	-0.2356	0.088*	
C28	0.2947 (2)	1.4475 (2)	-0.11625 (17)	0.0721 (8)	
H28	0.2932	1.5231	-0.1065	0.087*	
C29	0.3236 (2)	1.3757 (2)	-0.05248 (16)	0.0621 (7)	
H29	0.3418	1.4018	-0.0003	0.075*	
C30	0.32498 (19)	1.26319 (18)	-0.06814 (14)	0.0497 (6)	
C11	0.34632 (18)	1.16109 (17)	-0.02110 (13)	0.0465 (5)	
C12	0.37395 (18)	1.13576 (18)	0.06066 (13)	0.0480 (5)	
C7	0.38385 (18)	1.02299 (18)	0.08091 (13)	0.0488 (6)	
C6	0.4188 (2)	0.9654 (2)	0.15713 (15)	0.0558 (6)	
C5	0.4579 (3)	1.0012 (3)	0.23615 (17)	0.0798 (8)	
H5	0.4621	1.0765	0.2480	0.096*	
C4	0.4905 (3)	0.9234 (3)	0.29710 (19)	0.0967 (10)	
H4	0.5165	0.9465	0.3502	0.116*	
C3	0.4845 (3)	0.8114 (3)	0.2793 (2)	0.0917 (10)	
H3	0.5059	0.7604	0.3210	0.110*	
C2	0.4480 (2)	0.7735 (2)	0.20217 (19)	0.0751 (8)	

---

H2	0.4449	0.6981	0.1908	0.090*	
C1	0.4157 (2)	0.8517 (2)	0.14180 (15)	0.0575 (6)	
C19	0.38864 (19)	1.22457 (18)	0.12135 (13)	0.0501 (6)	
C20	0.4853 (2)	1.2920 (2)	0.14172 (16)	0.0698 (8)	
H20	0.5419	1.2810	0.1173	0.084*	
C21	0.5000 (2)	1.3751 (2)	0.19725 (17)	0.0749 (8)	
H21	0.5664	1.4190	0.2114	0.090*	
C22	0.4151 (2)	1.3916 (2)	0.23098 (15)	0.0656 (7)	
C23	0.3172 (3)	1.3273 (3)	0.21208 (18)	0.0837 (9)	
H23	0.2601	1.3400	0.2358	0.100*	
C24	0.3051 (2)	1.2436 (2)	0.15732 (17)	0.0726 (8)	
H24	0.2393	1.1989	0.1443	0.087*	
N1	0.38211 (16)	0.83465 (15)	0.05764 (12)	0.0563 (5)	
N2	0.4307 (3)	1.4799 (2)	0.29103 (16)	0.0916 (8)	
O5	0.30118 (15)	1.11572 (13)	-0.15385 (9)	0.0620 (5)	
O3	0.3507 (3)	1.5024 (3)	0.3147 (2)	0.1593 (14)	
O4	0.5242 (3)	1.5253 (2)	0.31592 (15)	0.1177 (9)	
S1	0.3009 (3)	0.72479 (18)	0.01555 (15)	0.0506 (10)	0.63 (2)
O1	0.3496 (10)	0.6265 (6)	0.0582 (5)	0.0755 (19)	0.63 (2)
O2	0.2837 (9)	0.7347 (7)	-0.0686 (2)	0.072 (2)	0.63 (2)
C13	0.1670 (5)	0.7463 (4)	0.0295 (4)	0.0473 (17)	0.63 (2)
C14	0.0952 (6)	0.8257 (4)	-0.0189 (4)	0.0530 (15)	0.63 (2)
H14	0.1180	0.8613	-0.0583	0.064*	0.63 (2)
C15	-0.0106 (5)	0.8520 (5)	-0.0086 (4)	0.0750 (19)	0.63 (2)
H15	-0.0586	0.9052	-0.0410	0.090*	0.63 (2)
C16	-0.0447 (4)	0.7989 (7)	0.0502 (4)	0.086 (2)	0.63 (2)
H16	-0.1154	0.8164	0.0572	0.103*	0.63 (2)
C17	0.0271 (5)	0.7194 (8)	0.0987 (4)	0.090 (3)	0.63 (2)
H17	0.0043	0.6839	0.1380	0.108*	0.63 (2)
C18	0.1329 (5)	0.6931 (6)	0.0883 (4)	0.075 (2)	0.63 (2)
H18	0.1809	0.6400	0.1207	0.090*	0.63 (2)
S1'	0.3118 (8)	0.7228 (6)	0.0149 (6)	0.102 (3)	0.37 (2)
O1'	0.379 (2)	0.6375 (16)	0.0646 (13)	0.133 (7)	0.37 (2)
O2'	0.2987 (18)	0.7264 (11)	-0.0681 (7)	0.095 (4)	0.37 (2)
C13'	0.1784 (11)	0.7417 (13)	0.0298 (10)	0.094 (5)	0.37 (2)
C14'	0.1010 (14)	0.8274 (11)	-0.0020 (12)	0.094 (4)	0.37 (2)
H14'	0.1214	0.8835	-0.0311	0.112*	0.37 (2)
C15'	-0.0070 (11)	0.8293 (15)	0.0097 (12)	0.115 (5)	0.37 (2)
H15'	-0.0588	0.8867	-0.0116	0.138*	0.37 (2)
C16'	-0.0375 (10)	0.746 (2)	0.0533 (10)	0.145 (6)	0.37 (2)
H16'	-0.1098	0.7468	0.0611	0.174*	0.37 (2)
C17'	0.0399 (14)	0.660 (2)	0.0851 (10)	0.151 (6)	0.37 (2)
H17'	0.0195	0.6037	0.1142	0.181*	0.37 (2)
C18'	0.1478 (12)	0.6579 (17)	0.0734 (11)	0.139 (5)	0.37 (2)
H18'	0.1996	0.6005	0.0947	0.167*	0.37 (2)

---



Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C8	0.0494 (13)	0.0414 (12)	0.0632 (16)	-0.0007 (10)	0.0232 (11)	0.0000 (11)
C9	0.0664 (16)	0.0426 (13)	0.0613 (16)	-0.0044 (11)	0.0270 (12)	-0.0069 (11)
C10	0.0543 (14)	0.0498 (14)	0.0523 (15)	-0.0023 (11)	0.0211 (11)	-0.0048 (11)
C25	0.0598 (15)	0.0401 (13)	0.0647 (16)	-0.0039 (11)	0.0237 (12)	-0.0032 (12)
C26	0.0830 (19)	0.0594 (16)	0.0586 (16)	-0.0087 (14)	0.0230 (14)	0.0026 (13)
C27	0.086 (2)	0.0544 (16)	0.076 (2)	-0.0041 (14)	0.0218 (15)	0.0146 (14)
C28	0.092 (2)	0.0445 (14)	0.080 (2)	-0.0022 (14)	0.0282 (16)	0.0051 (14)
C29	0.0735 (17)	0.0467 (14)	0.0677 (17)	-0.0039 (12)	0.0247 (13)	-0.0053 (13)
C30	0.0488 (13)	0.0450 (13)	0.0584 (15)	-0.0028 (10)	0.0215 (11)	-0.0050 (11)
C11	0.0460 (13)	0.0378 (12)	0.0584 (15)	-0.0020 (10)	0.0204 (11)	0.0011 (10)
C12	0.0445 (13)	0.0455 (13)	0.0551 (14)	-0.0038 (10)	0.0174 (10)	-0.0044 (11)
C7	0.0449 (13)	0.0458 (13)	0.0554 (14)	-0.0010 (10)	0.0155 (11)	0.0037 (11)
C6	0.0494 (14)	0.0566 (15)	0.0575 (16)	-0.0049 (12)	0.0118 (11)	0.0036 (12)
C5	0.091 (2)	0.0721 (19)	0.0633 (19)	-0.0111 (16)	0.0071 (15)	0.0048 (15)
C4	0.109 (3)	0.102 (3)	0.0600 (19)	-0.015 (2)	0.0007 (17)	0.0138 (18)
C3	0.086 (2)	0.087 (2)	0.088 (2)	0.0008 (18)	0.0086 (18)	0.036 (2)
C2	0.0679 (18)	0.0649 (18)	0.086 (2)	0.0024 (14)	0.0151 (15)	0.0206 (16)
C1	0.0469 (14)	0.0537 (15)	0.0702 (17)	0.0007 (11)	0.0162 (12)	0.0091 (13)
C19	0.0531 (14)	0.0442 (13)	0.0538 (14)	-0.0046 (11)	0.0180 (11)	-0.0009 (11)
C20	0.0643 (17)	0.0712 (18)	0.0826 (19)	-0.0174 (14)	0.0358 (14)	-0.0242 (15)
C21	0.0733 (18)	0.0747 (19)	0.0825 (19)	-0.0272 (15)	0.0332 (15)	-0.0247 (16)
C22	0.0803 (19)	0.0528 (15)	0.0639 (17)	-0.0056 (14)	0.0232 (14)	-0.0174 (13)
C23	0.0744 (19)	0.097 (2)	0.092 (2)	-0.0128 (17)	0.0439 (16)	-0.0337 (18)
C24	0.0674 (17)	0.0768 (18)	0.0816 (19)	-0.0217 (14)	0.0348 (15)	-0.0278 (15)
N1	0.0583 (12)	0.0431 (11)	0.0698 (14)	0.0006 (9)	0.0238 (10)	0.0027 (10)
N2	0.105 (2)	0.0823 (18)	0.0870 (19)	-0.0048 (18)	0.0303 (17)	-0.0308 (15)
O5	0.0860 (12)	0.0478 (10)	0.0554 (11)	-0.0062 (8)	0.0271 (9)	-0.0050 (8)
O3	0.132 (2)	0.172 (3)	0.187 (3)	0.000 (2)	0.068 (2)	-0.111 (2)
O4	0.142 (2)	0.0987 (18)	0.1132 (19)	-0.0397 (17)	0.0411 (16)	-0.0519 (15)
S1	0.0718 (16)	0.0251 (13)	0.0596 (17)	0.0034 (10)	0.0278 (11)	0.0003 (10)
O1	0.090 (5)	0.024 (2)	0.106 (4)	0.013 (3)	0.022 (3)	0.011 (2)
O2	0.106 (4)	0.075 (5)	0.049 (3)	-0.002 (3)	0.043 (3)	-0.018 (2)
C13	0.056 (3)	0.041 (3)	0.045 (4)	-0.015 (2)	0.016 (3)	0.000 (3)
C14	0.060 (3)	0.053 (3)	0.043 (3)	-0.007 (2)	0.011 (2)	-0.001 (2)
C15	0.059 (4)	0.086 (4)	0.072 (4)	-0.001 (3)	0.011 (3)	0.011 (3)
C16	0.057 (3)	0.117 (6)	0.089 (4)	-0.014 (3)	0.031 (3)	0.017 (4)
C17	0.077 (4)	0.125 (6)	0.077 (4)	-0.003 (4)	0.038 (3)	0.032 (4)
C18	0.073 (4)	0.087 (4)	0.065 (4)	-0.005 (3)	0.022 (3)	0.029 (3)
S1'	0.083 (4)	0.076 (5)	0.160 (7)	-0.015 (3)	0.058 (4)	-0.027 (4)
O1'	0.098 (9)	0.053 (7)	0.222 (13)	-0.004 (6)	0.014 (7)	-0.009 (6)
O2'	0.142 (10)	0.024 (4)	0.164 (10)	-0.026 (5)	0.110 (8)	-0.031 (5)
C13'	0.084 (8)	0.132 (11)	0.067 (10)	-0.020 (7)	0.027 (7)	0.000 (8)
C14'	0.073 (7)	0.139 (10)	0.075 (8)	-0.020 (6)	0.032 (6)	-0.024 (6)
C15'	0.062 (7)	0.167 (11)	0.110 (9)	-0.015 (7)	0.017 (7)	-0.001 (8)
C16'	0.113 (9)	0.187 (15)	0.153 (11)	-0.019 (9)	0.067 (9)	0.015 (11)

C17'	0.141 (10)	0.178 (14)	0.166 (10)	-0.015 (10)	0.095 (9)	0.024 (11)
C18'	0.119 (8)	0.183 (12)	0.130 (10)	-0.027 (8)	0.061 (8)	0.033 (9)

*Geometric parameters (Å, °)*

C8—C9	1.381 (3)	C21—H21	0.9300
C8—C7	1.410 (3)	C22—C23	1.368 (4)
C8—N1	1.426 (3)	C22—N2	1.472 (3)
C9—C10	1.363 (3)	C23—C24	1.372 (4)
C9—H9	0.9300	C23—H23	0.9300
C10—O5	1.383 (3)	C24—H24	0.9300
C10—C11	1.390 (3)	N1—S1'	1.645 (6)
C25—O5	1.381 (3)	N1—S1	1.677 (3)
C25—C30	1.381 (3)	N2—O3	1.204 (3)
C25—C26	1.386 (3)	N2—O4	1.209 (3)
C26—C27	1.377 (4)	S1—O1	1.428 (4)
C26—H26	0.9300	S1—O2	1.436 (4)
C27—C28	1.379 (4)	S1—C13	1.740 (3)
C27—H27	0.9300	C13—C14	1.3900
C28—C29	1.374 (3)	C13—C18	1.3900
C28—H28	0.9300	C14—C15	1.3900
C29—C30	1.387 (3)	C14—H14	0.9300
C29—H29	0.9300	C15—C16	1.3900
C30—C11	1.462 (3)	C15—H15	0.9300
C11—C12	1.406 (3)	C16—C17	1.3900
C12—C7	1.403 (3)	C16—H16	0.9300
C12—C19	1.485 (3)	C17—C18	1.3900
C7—C6	1.451 (3)	C17—H17	0.9300
C6—C5	1.390 (4)	C18—H18	0.9300
C6—C1	1.398 (3)	S1'—O2'	1.422 (7)
C5—C4	1.387 (4)	S1'—O1'	1.428 (8)
C5—H5	0.9300	S1'—C13'	1.736 (7)
C4—C3	1.384 (5)	C13'—C14'	1.3900
C4—H4	0.9300	C13'—C18'	1.3900
C3—C2	1.369 (4)	C14'—C15'	1.3900
C3—H3	0.9300	C14'—H14'	0.9300
C2—C1	1.383 (3)	C15'—C16'	1.3900
C2—H2	0.9300	C15'—H15'	0.9300
C1—N1	1.423 (3)	C16'—C17'	1.3900
C19—C24	1.376 (3)	C16'—H16'	0.9300
C19—C20	1.378 (3)	C17'—C18'	1.3900
C20—C21	1.373 (3)	C17'—H17'	0.9300
C20—H20	0.9300	C18'—H18'	0.9300
C21—C22	1.358 (4)		
C9—C8—C7	123.5 (2)	C23—C22—N2	119.3 (3)
C9—C8—N1	127.4 (2)	C22—C23—C24	118.6 (2)
C7—C8—N1	109.02 (19)	C22—C23—H23	120.7

C10—C9—C8	114.1 (2)	C24—C23—H23	120.7
C10—C9—H9	123.0	C23—C24—C19	121.2 (2)
C8—C9—H9	123.0	C23—C24—H24	119.4
C9—C10—O5	122.5 (2)	C19—C24—H24	119.4
C9—C10—C11	125.9 (2)	C1—N1—C8	107.16 (18)
O5—C10—C11	111.62 (19)	C1—N1—S1'	122.0 (4)
O5—C25—C30	112.2 (2)	C8—N1—S1'	123.3 (4)
O5—C25—C26	123.2 (2)	C1—N1—S1	120.53 (18)
C30—C25—C26	124.5 (2)	C8—N1—S1	121.89 (19)
C27—C26—C25	115.6 (2)	O3—N2—O4	122.6 (3)
C27—C26—H26	122.2	O3—N2—C22	118.7 (3)
C25—C26—H26	122.2	O4—N2—C22	118.7 (3)
C26—C27—C28	121.1 (3)	C25—O5—C10	105.33 (17)
C26—C27—H27	119.5	O1—S1—O2	120.9 (5)
C28—C27—H27	119.5	O1—S1—N1	109.8 (5)
C29—C28—C27	122.4 (3)	O2—S1—N1	105.2 (4)
C29—C28—H28	118.8	O1—S1—C13	107.3 (5)
C27—C28—H28	118.8	O2—S1—C13	107.5 (5)
C28—C29—C30	118.1 (2)	N1—S1—C13	105.2 (3)
C28—C29—H29	121.0	C14—C13—C18	120.0
C30—C29—H29	121.0	C14—C13—S1	116.5 (4)
C25—C30—C29	118.3 (2)	C18—C13—S1	123.4 (4)
C25—C30—C11	105.33 (19)	C13—C14—C15	120.0
C29—C30—C11	136.3 (2)	C13—C14—H14	120.0
C10—C11—C12	119.5 (2)	C15—C14—H14	120.0
C10—C11—C30	105.46 (19)	C16—C15—C14	120.0
C12—C11—C30	135.0 (2)	C16—C15—H15	120.0
C7—C12—C11	116.4 (2)	C14—C15—H15	120.0
C7—C12—C19	122.4 (2)	C15—C16—C17	120.0
C11—C12—C19	121.1 (2)	C15—C16—H16	120.0
C12—C7—C8	120.5 (2)	C17—C16—H16	120.0
C12—C7—C6	132.6 (2)	C16—C17—C18	120.0
C8—C7—C6	106.8 (2)	C16—C17—H17	120.0
C5—C6—C1	118.6 (2)	C18—C17—H17	120.0
C5—C6—C7	133.2 (2)	C17—C18—C13	120.0
C1—C6—C7	108.0 (2)	C17—C18—H18	120.0
C4—C5—C6	119.2 (3)	C13—C18—H18	120.0
C4—C5—H5	120.4	O2'—S1'—O1'	120.6 (13)
C6—C5—H5	120.4	O2'—S1'—N1	108.6 (7)
C3—C4—C5	120.3 (3)	O1'—S1'—N1	101.5 (11)
C3—C4—H4	119.8	O2'—S1'—C13'	110.4 (12)
C5—C4—H4	119.8	O1'—S1'—C13'	112.2 (12)
C2—C3—C4	121.9 (3)	N1—S1'—C13'	101.3 (6)
C2—C3—H3	119.0	C14'—C13'—C18'	120.0
C4—C3—H3	119.0	C14'—C13'—S1'	125.4 (10)
C3—C2—C1	117.4 (3)	C18'—C13'—S1'	114.5 (10)
C3—C2—H2	121.3	C13'—C14'—C15'	120.0
C1—C2—H2	121.3	C13'—C14'—H14'	120.0

C2—C1—C6	122.6 (3)	C15'—C14'—H14'	120.0
C2—C1—N1	128.4 (2)	C16'—C15'—C14'	120.0
C6—C1—N1	108.9 (2)	C16'—C15'—H15'	120.0
C24—C19—C20	118.2 (2)	C14'—C15'—H15'	120.0
C24—C19—C12	121.1 (2)	C15'—C16'—C17'	120.0
C20—C19—C12	120.7 (2)	C15'—C16'—H16'	120.0
C21—C20—C19	121.5 (2)	C17'—C16'—H16'	120.0
C21—C20—H20	119.2	C18'—C17'—C16'	120.0
C19—C20—H20	119.2	C18'—C17'—H17'	120.0
C22—C21—C20	118.4 (2)	C16'—C17'—H17'	120.0
C22—C21—H21	120.8	C17'—C18'—C13'	120.0
C20—C21—H21	120.8	C17'—C18'—H18'	120.0
C21—C22—C23	122.0 (2)	C13'—C18'—H18'	120.0
C21—C22—N2	118.7 (3)		
C7—C8—C9—C10	-0.9 (3)	C6—C1—N1—C8	-0.2 (2)
N1—C8—C9—C10	-177.3 (2)	C2—C1—N1—S1'	32.8 (5)
C8—C9—C10—O5	-178.72 (19)	C6—C1—N1—S1'	-150.8 (4)
C8—C9—C10—C11	0.0 (4)	C2—C1—N1—S1	37.9 (4)
O5—C25—C26—C27	178.4 (2)	C6—C1—N1—S1	-145.7 (2)
C30—C25—C26—C27	-0.9 (4)	C9—C8—N1—C1	175.9 (2)
C25—C26—C27—C28	1.0 (4)	C7—C8—N1—C1	-0.9 (2)
C26—C27—C28—C29	-0.5 (4)	C9—C8—N1—S1'	-33.9 (5)
C27—C28—C29—C30	-0.3 (4)	C7—C8—N1—S1'	149.2 (4)
O5—C25—C30—C29	-179.2 (2)	C9—C8—N1—S1	-39.2 (3)
C26—C25—C30—C29	0.1 (4)	C7—C8—N1—S1	144.0 (2)
O5—C25—C30—C11	-0.4 (3)	C21—C22—N2—O3	172.2 (3)
C26—C25—C30—C11	178.9 (2)	C23—C22—N2—O3	-9.1 (5)
C28—C29—C30—C25	0.5 (4)	C21—C22—N2—O4	-9.4 (4)
C28—C29—C30—C11	-177.8 (2)	C23—C22—N2—O4	169.3 (3)
C9—C10—C11—C12	0.2 (4)	C30—C25—O5—C10	0.9 (2)
O5—C10—C11—C12	178.98 (18)	C26—C25—O5—C10	-178.4 (2)
C9—C10—C11—C30	-177.9 (2)	C9—C10—O5—C25	177.8 (2)
O5—C10—C11—C30	0.9 (2)	C11—C10—O5—C25	-1.1 (2)
C25—C30—C11—C10	-0.3 (2)	C1—N1—S1—O1	-48.8 (5)
C29—C30—C11—C10	178.2 (3)	C8—N1—S1—O1	170.8 (4)
C25—C30—C11—C12	-178.0 (2)	S1'—N1—S1—O1	62 (5)
C29—C30—C11—C12	0.5 (5)	C1—N1—S1—O2	179.7 (5)
C10—C11—C12—C7	0.6 (3)	C8—N1—S1—O2	39.3 (5)
C30—C11—C12—C7	178.0 (2)	S1'—N1—S1—O2	-70 (5)
C10—C11—C12—C19	-177.9 (2)	C1—N1—S1—C13	66.4 (3)
C30—C11—C12—C19	-0.5 (4)	C8—N1—S1—C13	-74.0 (3)
C11—C12—C7—C8	-1.5 (3)	S1'—N1—S1—C13	177 (5)
C19—C12—C7—C8	177.0 (2)	O1—S1—C13—C14	-168.3 (6)
C11—C12—C7—C6	174.7 (2)	O2—S1—C13—C14	-37.0 (5)
C19—C12—C7—C6	-6.8 (4)	N1—S1—C13—C14	74.8 (3)
C9—C8—C7—C12	1.7 (3)	O1—S1—C13—C18	15.2 (7)
N1—C8—C7—C12	178.71 (19)	O2—S1—C13—C18	146.6 (6)

C9—C8—C7—C6	-175.3 (2)	N1—S1—C13—C18	-101.7 (4)
N1—C8—C7—C6	1.7 (2)	C18—C13—C14—C15	0.0
C12—C7—C6—C5	-2.4 (5)	S1—C13—C14—C15	-176.6 (5)
C8—C7—C6—C5	174.2 (3)	C13—C14—C15—C16	0.0
C12—C7—C6—C1	-178.3 (2)	C14—C15—C16—C17	0.0
C8—C7—C6—C1	-1.8 (3)	C15—C16—C17—C18	0.0
C1—C6—C5—C4	-1.2 (4)	C16—C17—C18—C13	0.0
C7—C6—C5—C4	-176.8 (3)	C14—C13—C18—C17	0.0
C6—C5—C4—C3	0.2 (5)	S1—C13—C18—C17	176.3 (5)
C5—C4—C3—C2	0.7 (5)	C1—N1—S1'—O2'	-175.6 (10)
C4—C3—C2—C1	-0.6 (5)	C8—N1—S1'—O2'	38.5 (11)
C3—C2—C1—C6	-0.4 (4)	S1—N1—S1'—O2'	112 (6)
C3—C2—C1—N1	175.6 (2)	C1—N1—S1'—O1'	-47.6 (11)
C5—C6—C1—C2	1.3 (4)	C8—N1—S1'—O1'	166.5 (10)
C7—C6—C1—C2	177.9 (2)	S1—N1—S1'—O1'	-120 (6)
C5—C6—C1—N1	-175.4 (2)	C1—N1—S1'—C13'	68.1 (8)
C7—C6—C1—N1	1.2 (3)	C8—N1—S1'—C13'	-77.8 (8)
C7—C12—C19—C24	-72.6 (3)	S1—N1—S1'—C13'	-4 (5)
C11—C12—C19—C24	105.9 (3)	O2'—S1'—C13'—C14'	-51.1 (13)
C7—C12—C19—C20	109.0 (3)	O1'—S1'—C13'—C14'	171.2 (15)
C11—C12—C19—C20	-72.6 (3)	N1—S1'—C13'—C14'	63.7 (11)
C24—C19—C20—C21	1.0 (4)	O2'—S1'—C13'—C18'	124.6 (12)
C12—C19—C20—C21	179.5 (3)	O1'—S1'—C13'—C18'	-13.0 (17)
C19—C20—C21—C22	-1.5 (4)	N1—S1'—C13'—C18'	-120.5 (9)
C20—C21—C22—C23	0.9 (5)	C18'—C13'—C14'—C15'	0.0
C20—C21—C22—N2	179.5 (3)	S1'—C13'—C14'—C15'	175.5 (12)
C21—C22—C23—C24	0.0 (5)	C13'—C14'—C15'—C16'	0.0
N2—C22—C23—C24	-178.5 (3)	C14'—C15'—C16'—C17'	0.0
C22—C23—C24—C19	-0.5 (5)	C15'—C16'—C17'—C18'	0.0
C20—C19—C24—C23	0.0 (4)	C16'—C17'—C18'—C13'	0.0
C12—C19—C24—C23	-178.5 (3)	C14'—C13'—C18'—C17'	0.0
C2—C1—N1—C8	-176.6 (2)	S1'—C13'—C18'—C17'	-176.0 (11)

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

Cg1 is the centroid of the furan ring O5/C10/C11/C25/C30 and Cg3 is the centroid of the benzene ring C1—C6.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C2—H2 $\cdots$ O1	0.93	2.41	3.015 (9)	122
C9—H9 $\cdots$ O2	0.93	2.28	2.849 (9)	119
C14—H14 $\cdots$ O4 <sup>i</sup>	0.93	2.55	3.296 (7)	138
C20—H20 $\cdots$ O2 <sup>ii</sup>	0.93	2.53	3.454 (11)	172
C16—H16 $\cdots$ Cg1 <sup>iii</sup>	0.93	2.76	3.676 (6)	169
C23—H23 $\cdots$ Cg3 <sup>iv</sup>	0.93	2.99	3.908 (4)	169

Symmetry codes: (i)  $x-1/2, -y+5/2, z-1/2$ ; (ii)  $-x+1, -y+2, -z$ ; (iii)  $-x, -y+2, -z$ ; (iv)  $-x+1/2, y+1/2, -z+1/2$ .

## (II) 2-Methyl-4-(4-nitrophenyl)-9-phenylsulfonyl-9H-thieno[2,3-b]carbazole

*Crystal data* $C_{27}H_{18}N_2O_4S_2$  $M_r = 498.55$ Monoclinic,  $P2_1/n$ Hall symbol:  $-P\ 2_1n$  $a = 12.5052\ (8)\ \text{\AA}$  $b = 11.2594\ (6)\ \text{\AA}$  $c = 17.0731\ (9)\ \text{\AA}$  $\beta = 102.914\ (2)^\circ$  $V = 2343.1\ (2)\ \text{\AA}^3$  $Z = 4$  $F(000) = 1032$  $D_x = 1.413\ \text{Mg m}^{-3}$ Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$ 

Cell parameters from 5100 reflections

 $\theta = 2.2\text{--}27.0^\circ$  $\mu = 0.27\ \text{mm}^{-1}$  $T = 296\ \text{K}$ 

Block, colourless

 $0.35 \times 0.30 \times 0.25\ \text{mm}$ *Data collection*

Bruker Kappa APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 $\omega$  &  $\varphi$  scansAbsorption correction: multi-scan  
(SADABS; Bruker, 2008) $T_{\min} = 0.911$ ,  $T_{\max} = 0.936$ 

44472 measured reflections

5100 independent reflections

3714 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.034$  $\theta_{\max} = 27.0^\circ$ ,  $\theta_{\min} = 2.2^\circ$  $h = -15 \rightarrow 15$  $k = -14 \rightarrow 14$  $l = -21 \rightarrow 21$ *Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.055$  $wR(F^2) = 0.163$  $S = 1.08$ 

5100 reflections

317 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0625P)^2 + 2.4597P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} = 0.001$  $\Delta\rho_{\max} = 0.44\ \text{e \AA}^{-3}$  $\Delta\rho_{\min} = -0.36\ \text{e \AA}^{-3}$ *Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional R-factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and R-factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.6060 (2)	0.1940 (2)	0.88312 (17)	0.0422 (6)
C2	0.5786 (3)	0.2854 (3)	0.8277 (2)	0.0559 (8)
H2	0.5843	0.3645	0.8438	0.067*
C3	0.5428 (3)	0.2548 (3)	0.7482 (2)	0.0630 (9)

H3	0.5242	0.3146	0.7100	0.076*
C4	0.5337 (3)	0.1379 (3)	0.7237 (2)	0.0631 (9)
H4	0.5082	0.1203	0.6695	0.076*
C5	0.5616 (3)	0.0466 (3)	0.77821 (17)	0.0508 (7)
H5	0.5562	-0.0321	0.7612	0.061*
C6	0.5983 (2)	0.0746 (2)	0.85962 (16)	0.0390 (6)
C7	0.6289 (2)	0.0029 (2)	0.93182 (15)	0.0353 (5)
C8	0.6520 (2)	0.0804 (2)	0.99817 (16)	0.0360 (6)
C9	0.6768 (2)	0.0413 (2)	1.07625 (16)	0.0393 (6)
H9	0.6917	0.0937	1.1194	0.047*
C10	0.6786 (2)	-0.0812 (2)	1.08733 (15)	0.0380 (6)
C11	0.6568 (2)	-0.1622 (2)	1.02287 (15)	0.0353 (5)
C12	0.6333 (2)	-0.1197 (2)	0.94342 (15)	0.0350 (5)
C13	0.8439 (2)	0.2964 (2)	0.98979 (16)	0.0444 (6)
C14	0.9114 (3)	0.2047 (3)	1.02414 (18)	0.0523 (7)
H14	0.8886	0.1523	1.0593	0.063*
C15	1.0124 (3)	0.1910 (4)	1.0062 (2)	0.0681 (9)
H15	1.0579	0.1287	1.0286	0.082*
C16	1.0456 (3)	0.2685 (5)	0.9557 (3)	0.0829 (12)
H16	1.1145	0.2595	0.9443	0.100*
C17	0.9798 (4)	0.3594 (5)	0.9214 (3)	0.0949 (15)
H17	1.0039	0.4117	0.8869	0.114*
C18	0.8765 (3)	0.3742 (4)	0.9378 (2)	0.0735 (11)
H18	0.8307	0.4354	0.9141	0.088*
C19	0.6151 (2)	-0.2052 (2)	0.87505 (15)	0.0354 (5)
C20	0.5183 (2)	-0.2687 (3)	0.85455 (19)	0.0537 (8)
H20	0.4651	-0.2592	0.8844	0.064*
C21	0.4998 (3)	-0.3460 (3)	0.7902 (2)	0.0554 (8)
H21	0.4339	-0.3872	0.7755	0.067*
C22	0.5802 (2)	-0.3609 (2)	0.74852 (16)	0.0430 (6)
C23	0.6776 (3)	-0.3019 (3)	0.7681 (2)	0.0584 (8)
H23	0.7316	-0.3145	0.7393	0.070*
C24	0.6946 (2)	-0.2227 (3)	0.83158 (19)	0.0531 (8)
H24	0.7602	-0.1808	0.8452	0.064*
C25	0.6646 (2)	-0.2867 (2)	1.05076 (15)	0.0384 (6)
H25	0.6526	-0.3528	1.0173	0.046*
C26	0.6920 (3)	-0.2909 (2)	1.13271 (18)	0.0481 (7)
C27	0.7108 (4)	-0.4006 (3)	1.1839 (2)	0.0753 (11)
H27A	0.6928	-0.4695	1.1504	0.113*
H27B	0.6651	-0.3981	1.2222	0.113*
H27C	0.7864	-0.4044	1.2118	0.113*
N1	0.63886 (18)	0.19995 (19)	0.96867 (14)	0.0411 (5)
N2	0.5598 (3)	-0.4437 (2)	0.68029 (16)	0.0575 (7)
O1	0.6654 (2)	0.41840 (18)	0.97873 (17)	0.0737 (7)
O2	0.7283 (2)	0.29254 (19)	1.09819 (13)	0.0611 (6)
O3	0.6358 (2)	-0.4681 (3)	0.64953 (16)	0.0853 (9)
O4	0.4680 (2)	-0.4842 (2)	0.65745 (15)	0.0765 (8)
S1	0.71501 (6)	0.31228 (6)	1.01429 (5)	0.0488 (2)

S2                    0.70682 (7)                    -0.15330 (7)                    1.17880 (5)                    0.0541 (2)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0382 (14)	0.0378 (14)	0.0513 (16)	0.0023 (11)	0.0117 (12)	0.0048 (12)
C2	0.0580 (19)	0.0376 (15)	0.070 (2)	0.0046 (14)	0.0099 (16)	0.0134 (14)
C3	0.068 (2)	0.056 (2)	0.062 (2)	0.0058 (16)	0.0080 (17)	0.0246 (16)
C4	0.076 (2)	0.064 (2)	0.0454 (17)	0.0001 (18)	0.0045 (16)	0.0116 (15)
C5	0.0608 (19)	0.0445 (16)	0.0440 (16)	-0.0026 (14)	0.0049 (14)	0.0012 (13)
C6	0.0371 (14)	0.0362 (13)	0.0443 (15)	-0.0005 (11)	0.0102 (11)	0.0033 (11)
C7	0.0329 (13)	0.0343 (13)	0.0390 (13)	-0.0026 (10)	0.0084 (10)	-0.0021 (10)
C8	0.0336 (13)	0.0288 (12)	0.0467 (15)	-0.0012 (10)	0.0113 (11)	-0.0034 (10)
C9	0.0426 (14)	0.0371 (14)	0.0381 (14)	-0.0032 (11)	0.0088 (11)	-0.0078 (11)
C10	0.0356 (13)	0.0427 (14)	0.0361 (14)	-0.0020 (11)	0.0085 (11)	-0.0002 (11)
C11	0.0311 (13)	0.0346 (13)	0.0407 (14)	-0.0019 (10)	0.0090 (10)	-0.0003 (10)
C12	0.0323 (13)	0.0332 (13)	0.0397 (13)	-0.0031 (10)	0.0087 (10)	-0.0028 (10)
C13	0.0502 (16)	0.0416 (15)	0.0404 (14)	-0.0121 (12)	0.0083 (12)	-0.0014 (12)
C14	0.0515 (18)	0.0549 (18)	0.0494 (17)	-0.0044 (14)	0.0086 (14)	0.0041 (14)
C15	0.0486 (19)	0.083 (3)	0.069 (2)	0.0032 (18)	0.0053 (16)	0.004 (2)
C16	0.053 (2)	0.120 (4)	0.078 (3)	-0.009 (2)	0.0188 (19)	0.008 (3)
C17	0.081 (3)	0.127 (4)	0.083 (3)	-0.025 (3)	0.033 (2)	0.037 (3)
C18	0.068 (2)	0.072 (2)	0.077 (2)	-0.0113 (19)	0.0093 (19)	0.031 (2)
C19	0.0390 (14)	0.0305 (12)	0.0373 (13)	-0.0020 (10)	0.0095 (11)	-0.0005 (10)
C20	0.0455 (16)	0.0565 (18)	0.0640 (19)	-0.0130 (14)	0.0229 (14)	-0.0197 (15)
C21	0.0454 (17)	0.0556 (18)	0.065 (2)	-0.0183 (14)	0.0112 (14)	-0.0174 (15)
C22	0.0514 (16)	0.0332 (13)	0.0427 (15)	0.0001 (12)	0.0073 (12)	-0.0086 (11)
C23	0.0526 (18)	0.067 (2)	0.0615 (19)	-0.0067 (15)	0.0251 (15)	-0.0215 (16)
C24	0.0452 (16)	0.0558 (18)	0.0626 (19)	-0.0183 (14)	0.0215 (14)	-0.0202 (15)
C25	0.0373 (14)	0.0386 (14)	0.0404 (14)	0.0032 (11)	0.0111 (11)	0.0083 (11)
C26	0.0526 (17)	0.0402 (15)	0.0526 (17)	0.0009 (13)	0.0141 (14)	0.0086 (13)
C27	0.107 (3)	0.054 (2)	0.067 (2)	0.007 (2)	0.024 (2)	0.0234 (17)
N1	0.0430 (13)	0.0302 (11)	0.0495 (13)	-0.0007 (9)	0.0090 (10)	-0.0027 (9)
N2	0.0718 (19)	0.0405 (14)	0.0556 (16)	0.0054 (13)	0.0043 (14)	-0.0111 (12)
O1	0.0785 (17)	0.0302 (11)	0.110 (2)	0.0068 (11)	0.0148 (14)	-0.0106 (12)
O2	0.0765 (15)	0.0538 (13)	0.0603 (13)	-0.0182 (11)	0.0305 (11)	-0.0233 (11)
O3	0.091 (2)	0.089 (2)	0.0765 (17)	0.0123 (16)	0.0200 (15)	-0.0388 (15)
O4	0.0852 (19)	0.0571 (15)	0.0761 (17)	-0.0124 (13)	-0.0053 (14)	-0.0266 (12)
S1	0.0555 (5)	0.0302 (3)	0.0627 (5)	-0.0035 (3)	0.0172 (4)	-0.0106 (3)
S2	0.0642 (5)	0.0521 (5)	0.0442 (4)	-0.0033 (4)	0.0081 (4)	0.0038 (3)

*Geometric parameters (Å, °)*

C1—C2	1.388 (4)	C15—H15	0.9300
C1—C6	1.400 (4)	C16—C17	1.361 (6)
C1—N1	1.428 (4)	C16—H16	0.9300
C2—C3	1.374 (5)	C17—C18	1.391 (6)
C2—H2	0.9300	C17—H17	0.9300



C3—C4	1.379 (5)	C18—H18	0.9300
C3—H3	0.9300	C19—C24	1.381 (4)
C4—C5	1.378 (4)	C19—C20	1.382 (4)
C4—H4	0.9300	C20—C21	1.381 (4)
C5—C6	1.399 (4)	C20—H20	0.9300
C5—H5	0.9300	C21—C22	1.365 (4)
C6—C7	1.452 (4)	C21—H21	0.9300
C7—C12	1.394 (3)	C22—C23	1.363 (4)
C7—C8	1.408 (3)	C22—N2	1.469 (4)
C8—C9	1.372 (4)	C23—C24	1.383 (4)
C8—N1	1.434 (3)	C23—H23	0.9300
C9—C10	1.391 (4)	C24—H24	0.9300
C9—H9	0.9300	C25—C26	1.365 (4)
C10—C11	1.408 (4)	C25—H25	0.9300
C10—S2	1.725 (3)	C26—C27	1.501 (4)
C11—C12	1.406 (4)	C26—S2	1.729 (3)
C11—C25	1.477 (3)	C27—H27A	0.9600
C12—C19	1.491 (3)	C27—H27B	0.9600
C13—C18	1.372 (4)	C27—H27C	0.9600
C13—C14	1.379 (4)	N1—S1	1.668 (2)
C13—S1	1.762 (3)	N2—O4	1.216 (4)
C14—C15	1.374 (5)	N2—O3	1.216 (4)
C14—H14	0.9300	O1—S1	1.419 (2)
C15—C16	1.356 (6)	O2—S1	1.422 (2)
C2—C1—C6	121.7 (3)	C16—C17—H17	119.9
C2—C1—N1	129.3 (3)	C18—C17—H17	119.9
C6—C1—N1	108.9 (2)	C13—C18—C17	118.6 (4)
C3—C2—C1	117.7 (3)	C13—C18—H18	120.7
C3—C2—H2	121.2	C17—C18—H18	120.7
C1—C2—H2	121.2	C24—C19—C20	119.0 (2)
C2—C3—C4	121.7 (3)	C24—C19—C12	120.7 (2)
C2—C3—H3	119.2	C20—C19—C12	120.2 (2)
C4—C3—H3	119.2	C21—C20—C19	120.6 (3)
C5—C4—C3	121.1 (3)	C21—C20—H20	119.7
C5—C4—H4	119.5	C19—C20—H20	119.7
C3—C4—H4	119.5	C22—C21—C20	118.7 (3)
C4—C5—C6	118.7 (3)	C22—C21—H21	120.6
C4—C5—H5	120.7	C20—C21—H21	120.6
C6—C5—H5	120.7	C23—C22—C21	122.3 (3)
C5—C6—C1	119.2 (3)	C23—C22—N2	119.4 (3)
C5—C6—C7	133.1 (3)	C21—C22—N2	118.3 (3)
C1—C6—C7	107.6 (2)	C22—C23—C24	118.6 (3)
C12—C7—C8	120.3 (2)	C22—C23—H23	120.7
C12—C7—C6	131.8 (2)	C24—C23—H23	120.7
C8—C7—C6	107.8 (2)	C19—C24—C23	120.7 (3)
C9—C8—C7	123.0 (2)	C19—C24—H24	119.6
C9—C8—N1	128.7 (2)	C23—C24—H24	119.6

C7—C8—N1	108.2 (2)	C26—C25—C11	110.3 (2)
C8—C9—C10	116.3 (2)	C26—C25—H25	124.9
C8—C9—H9	121.8	C11—C25—H25	124.9
C10—C9—H9	121.8	C25—C26—C27	126.6 (3)
C9—C10—C11	122.8 (2)	C25—C26—S2	114.4 (2)
C9—C10—S2	125.7 (2)	C27—C26—S2	119.1 (2)
C11—C10—S2	111.5 (2)	C26—C27—H27A	109.5
C12—C11—C10	119.7 (2)	C26—C27—H27B	109.5
C12—C11—C25	128.2 (2)	H27A—C27—H27B	109.5
C10—C11—C25	112.1 (2)	C26—C27—H27C	109.5
C7—C12—C11	117.8 (2)	H27A—C27—H27C	109.5
C7—C12—C19	122.3 (2)	H27B—C27—H27C	109.5
C11—C12—C19	119.9 (2)	C1—N1—C8	107.4 (2)
C18—C13—C14	120.7 (3)	C1—N1—S1	121.13 (18)
C18—C13—S1	120.6 (3)	C8—N1—S1	122.65 (18)
C14—C13—S1	118.7 (2)	O4—N2—O3	123.3 (3)
C15—C14—C13	119.7 (3)	O4—N2—C22	118.4 (3)
C15—C14—H14	120.2	O3—N2—C22	118.2 (3)
C13—C14—H14	120.2	O1—S1—O2	120.27 (15)
C16—C15—C14	119.8 (4)	O1—S1—N1	106.83 (14)
C16—C15—H15	120.1	O2—S1—N1	106.12 (12)
C14—C15—H15	120.1	O1—S1—C13	108.62 (15)
C15—C16—C17	121.0 (4)	O2—S1—C13	108.44 (14)
C15—C16—H16	119.5	N1—S1—C13	105.62 (12)
C17—C16—H16	119.5	C10—S2—C26	91.74 (14)
C16—C17—C18	120.2 (4)		
C6—C1—C2—C3	0.3 (5)	C7—C12—C19—C20	-106.1 (3)
N1—C1—C2—C3	-175.9 (3)	C11—C12—C19—C20	74.5 (3)
C1—C2—C3—C4	0.2 (5)	C24—C19—C20—C21	-1.8 (5)
C2—C3—C4—C5	-0.8 (6)	C12—C19—C20—C21	178.5 (3)
C3—C4—C5—C6	0.9 (5)	C19—C20—C21—C22	1.7 (5)
C4—C5—C6—C1	-0.3 (4)	C20—C21—C22—C23	-0.3 (5)
C4—C5—C6—C7	176.7 (3)	C20—C21—C22—N2	179.8 (3)
C2—C1—C6—C5	-0.3 (4)	C21—C22—C23—C24	-1.0 (5)
N1—C1—C6—C5	176.7 (2)	N2—C22—C23—C24	178.8 (3)
C2—C1—C6—C7	-178.0 (3)	C20—C19—C24—C23	0.4 (5)
N1—C1—C6—C7	-1.0 (3)	C12—C19—C24—C23	-179.9 (3)
C5—C6—C7—C12	1.2 (5)	C22—C23—C24—C19	1.0 (5)
C1—C6—C7—C12	178.4 (3)	C12—C11—C25—C26	178.4 (3)
C5—C6—C7—C8	-175.8 (3)	C10—C11—C25—C26	-0.4 (3)
C1—C6—C7—C8	1.5 (3)	C11—C25—C26—C27	-178.2 (3)
C12—C7—C8—C9	-1.6 (4)	C11—C25—C26—S2	1.0 (3)
C6—C7—C8—C9	175.7 (2)	C2—C1—N1—C8	176.8 (3)
C12—C7—C8—N1	-178.7 (2)	C6—C1—N1—C8	0.2 (3)
C6—C7—C8—N1	-1.4 (3)	C2—C1—N1—S1	-34.9 (4)
C7—C8—C9—C10	0.1 (4)	C6—C1—N1—S1	148.4 (2)
N1—C8—C9—C10	176.6 (2)	C9—C8—N1—C1	-176.1 (3)

C8—C9—C10—C11	0.3 (4)	C7—C8—N1—C1	0.7 (3)
C8—C9—C10—S2	-179.7 (2)	C9—C8—N1—S1	36.2 (4)
C9—C10—C11—C12	0.7 (4)	C7—C8—N1—S1	-146.90 (19)
S2—C10—C11—C12	-179.24 (19)	C23—C22—N2—O4	-170.9 (3)
C9—C10—C11—C25	179.6 (2)	C21—C22—N2—O4	9.0 (4)
S2—C10—C11—C25	-0.3 (3)	C23—C22—N2—O3	9.1 (4)
C8—C7—C12—C11	2.5 (4)	C21—C22—N2—O3	-171.0 (3)
C6—C7—C12—C11	-174.1 (3)	C1—N1—S1—O1	48.2 (2)
C8—C7—C12—C19	-176.8 (2)	C8—N1—S1—O1	-168.5 (2)
C6—C7—C12—C19	6.6 (4)	C1—N1—S1—O2	177.6 (2)
C10—C11—C12—C7	-2.1 (4)	C8—N1—S1—O2	-39.0 (2)
C25—C11—C12—C7	179.2 (2)	C1—N1—S1—C13	-67.3 (2)
C10—C11—C12—C19	177.3 (2)	C8—N1—S1—C13	76.0 (2)
C25—C11—C12—C19	-1.4 (4)	C18—C13—S1—O1	-5.6 (3)
C18—C13—C14—C15	0.0 (5)	C14—C13—S1—O1	174.2 (2)
S1—C13—C14—C15	-179.7 (3)	C18—C13—S1—O2	-137.9 (3)
C13—C14—C15—C16	0.8 (5)	C14—C13—S1—O2	41.9 (3)
C14—C15—C16—C17	-0.8 (7)	C18—C13—S1—N1	108.7 (3)
C15—C16—C17—C18	0.0 (7)	C14—C13—S1—N1	-71.5 (3)
C14—C13—C18—C17	-0.9 (6)	C9—C10—S2—C26	-179.2 (3)
S1—C13—C18—C17	178.9 (3)	C11—C10—S2—C26	0.8 (2)
C16—C17—C18—C13	0.8 (7)	C25—C26—S2—C10	-1.1 (2)
C7—C12—C19—C24	74.2 (4)	C27—C26—S2—C10	178.2 (3)
C11—C12—C19—C24	-105.2 (3)		

*Hydrogen-bond geometry (Å, °)*

Cg1 is the centroid of the thiophene ring S2/C10/C11/C25/C26 and Cg3 is centroid of the benzene ring C1—C6.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C2—H2...O1	0.93	2.38	2.970 (4)	121
C9—H9...O2	0.93	2.33	2.907 (3)	120
C3—H3...O4 <sup>i</sup>	0.93	2.48	3.356 (4)	157
C14—H14...O4 <sup>ii</sup>	0.93	2.58	3.335 (4)	139
C20—H20...O2 <sup>iii</sup>	0.93	2.53	3.368 (5)	150
C16—H16...Cg1 <sup>iv</sup>	0.93	2.89	3.812 (4)	172
C23—H23...Cg3 <sup>v</sup>	0.93	2.75	3.646 (4)	163

Symmetry codes: (i)  $x, y+1, z$ ; (ii)  $x+1/2, -y-1/2, z+1/2$ ; (iii)  $-x+1, -y, -z+2$ ; (iv)  $-x+2, -y, -z+2$ ; (v)  $-x+3/2, y-1/2, -z+3/2$ .