

RESEARCH ARTICLE

Cortical Gyrfication Patterns Associated with Trait Anxiety

Tara A. Miskovich, Walker S. Pedersen, Emily L. Belleau, Skyler Shollenbarger, Krista M. Lisdahl, Christine L. Larson*

Department of Psychology, University of Wisconsin-Milwaukee, Milwaukee, WI, United States of America

* larsoncl@uwm.edu



OPEN ACCESS

Citation: Miskovich TA, Pedersen WS, Belleau EL, Shollenbarger S, Lisdahl KM, Larson CL (2016) Cortical Gyrfication Patterns Associated with Trait Anxiety. *PLoS ONE* 11(2): e0149434. doi:10.1371/journal.pone.0149434

Editor: Xi-Nian Zuo, Institute of Psychology, Chinese Academy of Sciences, CHINA

Received: October 19, 2015

Accepted: January 30, 2016

Published: February 12, 2016

Copyright: © 2016 Miskovich et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant files are available from the Dryad Digital Repository (doi:[10.5061/dryad.96p8g](https://doi.org/10.5061/dryad.96p8g)).

Funding: This work was supported by the National Institutes of Health (MH086809). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Dispositional anxiety is a stable personality trait that is a key risk factor for internalizing disorders, and understanding the neural correlates of trait anxiety may help us better understand the development of these disorders. Abnormal cortical folding is thought to reflect differences in cortical connectivity occurring during brain development. Therefore, assessing gyrfication may advance understanding of cortical development and organization associated with trait anxiety. Previous literature has revealed structural abnormalities in trait anxiety and related disorders, but no study to our knowledge has examined gyrfication in trait anxiety. We utilized a relatively novel measure, the local gyrfication index (LGI), to explore differences in gyrfication as a function of trait anxiety. We obtained structural MRI scans using a 3T magnetic resonance scanner on 113 young adults. Results indicated a negative correlation between trait anxiety and LGI in the left superior parietal cortex, specifically the precuneus, reflecting less cortical complexity among those high on trait anxiety. Our findings suggest that aberrations in cortical gyrfication in a key region of the default mode network is a correlate of trait anxiety and may reflect disrupted local parietal connectivity.

Introduction

Dispositional anxiety is a stable personality trait with biological underpinnings [1] that is a risk factor for numerous psychiatric disorders, including anxiety disorders, depression, and substance use [2–4]. Understanding individual differences in brain structure associated with this personality trait has the potential to advance understanding of the etiology of these burdensome disorders.

Previous research has shown that trait anxiety and anxiety-related traits are associated with variation in cortical volume and thickness of regions implicated in anxiety-related processes. For instance, trait anxiety and neuroticism have both been linked to reduced thickness in the orbitofrontal cortex [1,5,6]. Behavioral inhibition, a construct similar to trait anxiety, has been associated with decreased gray matter volume in the medial orbitofrontal cortices and the precuneus [7]. The orbitofrontal cortex is important for various cognitive and emotional processes that are thought to be impaired in anxiety, such as extinction learning [8]. Additionally, both

the orbitofrontal cortex and precuneus have been identified as nodes in the default mode network (DMN), a neural network involved in self-referential or self-focused attentional processing [9–11]. This network is particularly relevant for internalizing disorders such as anxiety and depression, as DMN dysfunction may be a core neural substrate of pathological self-focused thought, such as rumination or worry [12–14].

As these studies indicate, trait anxiety is associated with abnormalities in cortical volume and thickness; however, to our knowledge no one has assessed the relationship between trait anxiety and cortical folding or gyrification. The process of gyrification begins in the second trimester in utero [15,16], and is thought to be largely complete by the third trimester [17]. Gyrification has been posited to reflect underlying structural connectivity because cortical folding allows for more efficient network wiring as surface area expands in the rapidly developing brain [18]. Initially, theories of gyrification argued the ratio of folding to surface area (the gyrification index) [19] remains relatively constant after the first few years of life, and therefore should primarily reflect early developmental, and genetic influences [17]. However, more recent evidence has pointed to developmental changes in gyrification throughout adolescence and adulthood [16,20–22]. Additionally, monozygotic twins have differences in gyrification patterns, suggesting a greater role for environmental factors influencing gyrification, more so than measures of cortical volume [23,24]. This is particularly true within more superficial sulci [25]. Therefore, differences in gyrification patterns may arise across a larger window of development than previously thought. Importantly, gyrification has been linked to measures of cortical connectivity, both functional [26] and structural [27]. Thus, abnormal cortical folding may provide insight into variation in cortical connectivity that may arise during cortical development.

Aberrant gyrification has been linked to various psychiatric disorders such as schizophrenia [28,29] and autism [30,31]. Abnormal morphology may, therefore, represent a disruption in neurodevelopment that puts one at risk for these and other disorders. Of most relevance for the current study, investigations across clinical samples of anxiety and depression in adults have revealed gyrification abnormalities in the precuneus and posterior cingulate, another key node in the posterior DMN [32–34]. Additionally, there is some evidence suggesting that gyrification abnormalities are linked to dysfunction in this network. Nixon and colleagues [33] found that individuals with depression displayed hypogyrfication in the precuneus, as well as increased task-based functional connectivity between the precuneus and anterior nodes of the DMN. Since the DMN is thought to subservise self-focused thought, excessive connectivity may reflect pathological self-focus such as worry [13,14]. Thus, research in this area suggests that there may be a link between gyrification and DMN dysfunction in internalizing disorders that may reflect a vulnerability to developing an anxiety or depressive disorder.

The aim of the current study was to investigate abnormalities in cortical folding that may be associated with trait anxiety, a core risk factor for some of the most prevalent forms of psychopathology, including anxiety disorders, depression, and substance use. Examining gyrification alongside trait anxiety may provide us with important clues about structural differences that arise during neurodevelopment that are associated with this important risk factor. We used a three-dimensional measurement of gyrification, the local gyrification index (LGI) [35], to assess differences in gyrification as a function of trait anxiety.

Materials and Methods

Participants

One hundred and twenty one individuals ranging from 18 to 35 (72 females) were recruited from the University of Wisconsin-Milwaukee student body and local community to participate

in the study. All aspects of this study were approved by both the Medical College of Wisconsin and the University of Wisconsin, Milwaukee (IRB# PRO00012620). Informed written consent was obtained from each participant in accordance with the University of Wisconsin-Milwaukee and Wisconsin Medical College Institutional Review Boards. Mean age of the sample was 21.9 ± 3.8 years. Subjects were excluded from the study if they were left handed, had a history of significant head trauma, contradiction to magnetic resonance imaging, history of a neurological disorder, psychotic disorder, or bipolar disorder, as assessed by the Structured Clinical Interview for DSM-IV Disorders [36].

Eight subjects were excluded from the final analyses. Six were dropped due to excessive motion during scanning, one was dropped due to technical issues with the scanner, and one was dropped due to the presence of manic episodes. Final data analyses included the remaining 113 subjects.

Quantifying Trait Anxiety

Participants completed the Trait version of the Spielberger State-Trait Anxiety Inventory (STAI) [37], a 20-item questionnaire that has high test-retest reliability as well as high internal consistency [38]. The mean STAI score was 39.7 ± 10.7 (range 21–66).

MRI Acquisition

High resolution spoiled gradient recalled (SPGR) images were acquired in a sagittal orientation (TR = 8.2ms; TE = 3.2ms; FOV = 24cm; flip angle = 12° ; voxel size (0.9375 x 0.9375 x 1mm) on a 3.0 Tesla short bore GE Signa Excite MRI system using a 12-channel head coil.

Cortical Reconstruction and LGI Maps

FreeSurfer 5.0 (www.nmr.mgh.harvard.edu/freesurfer) [39,40] installed on a Red Hat Enterprise Linux v.6.6 was used to reconstruct the cortical surface as a triangular mesh. The FreeSurfer reconstruction pipeline is described in more detail in [39], but consists of transformation to Talairach space, normalization of intensity, removal of non-brain tissue (skull), and segmentation of gray and white matter. In addition, we used manual editing to further correct the pial and white matter surfaces and remove skull and dura mater left over from skull-stripping.

The LGI, developed by Schaer and colleagues [35], was used to assess the ratio of buried to visible cortex, or amount of gyrification. This technique is a three-dimensional extension of the GI technique of Zilles and colleagues [19]. This new method is fully automated, and therefore provides a less subjective measure that takes into account the three-dimensional nature of the cortical folds and is not biased by the orientation of the slices [35]. In addition, the primary advantage is the ability to localize specific abnormalities in cortical folding, whereas previous methods only yielded a global measure of gyrification. The FreeSurfer LGI computation creates a triangulated mesh of the outer hull of the brain, calculates the LGI ratio between the pial surface and outer hull at the center of a 25 mm spherical region of interest, and propagates these values onto the pial surface, assigning LGI values to each vertex on the mesh surface [35,41].

Statistical Analyses

The relationship between trait anxiety and gyrification was assessed on surface maps at every vertex in the brain using the different offset, different slope (DODS) design provided in Qdec in FreeSurfer, with age and sex added as covariates. LGI values were first mapped to a normalized brain for each subject. We applied a smoothing kernel of 5 mm FWHM. To correct for multiple comparisons cluster correction was done using Monte Carlo simulation with 10,000

iterations. Using a vertex-wise threshold of $p < .05$, a minimum cluster size of 688.85 mm^2 was considered significant.

For clusters showing a significant relationship between LGI and trait anxiety, we extracted average LGI, along with surface area and cortical thickness. Although these measurements have been linked to gyrfication [20,21], they have different developmental trajectories [42], thus, examining these measurements jointly may provide a more comprehensive account of structural differences related to trait anxiety. Cortical thickness was defined as the distance between the gray and white matter boundary. This distance was calculated at each vertex and then averaged within significant clusters. Surface area was calculated by assigning the average area of surrounding triangles to each vertex and summing those values within significant clusters to obtain total surface area for the cluster. Multiple regressions, conducted in SPSS, were used to determine the relationship between LGI and surface area, cortical thickness, and trait anxiety. Age and sex were used as covariates in these analyses, and the results are presented in partial correlations.

Results

The vertex-wise analysis revealed a negative correlation between trait anxiety and LGI in the precuneus in the left hemisphere (cluster size = 2608.73 mm^2 , 5940 vertices, cluster corrected $p = .003$; Fig 1). There were no regions where LGI was positively associated with trait anxiety, and there was no moderating effect of sex. There was no relationship between anxiety and gyrfication in the right hemisphere.

We conducted follow up analyses to better understand how other features of the cortical surface may be related to or explain differences in gyrfication in the precuneus. Specifically, we correlated average surface area and cortical thickness with average LGI within the significant precuneus cluster. Controlling for the effects of sex and age, we found that surface area was positively associated with LGI ($r = 0.204$, $p = 0.032$; Fig 2A). Additionally, we found that cortical thickness was negatively correlated with LGI ($r = -0.253$, $p = 0.007$; Fig 2B). The relationships between these measurements and gyrfication are consistent with previous findings [20,21]. To determine whether differences in surface area and cortical thickness accounted for the relationship between LGI and trait anxiety, we assessed the relationship between average precuneus LGI and trait anxiety with average surface area and cortical thickness added as covariates (in addition to age and sex). We found that even after controlling for variance in surface area and cortical thickness, there was still a negative association between trait anxiety and LGI ($r = -0.315$, $p = 0.001$; Fig 2C).

Discussion

In the current study, we investigated differences in cortical gyrfication associated with dispositional anxiety. We found that there was less gyrfication in the left precuneus as dispositional anxiety increased. Although both cortical thickness and surface area were associated with gyrfication, the relationship between precuneus gyrfication and trait anxiety remained significant when controlling for these other cortical measurements. These findings suggest that there may be differences that arise during neurodevelopmental in the precuneus in those with dispositional anxiety, though longitudinal replication is warranted. Since theoretical [43] and empirical work [26,27,33] have linked gyrfication to underlying cortical connectivity, these aberrations in cortical morphology may be indicative of abnormalities in connectivity between the precuneus and corresponding networks.

Functionally, the precuneus is involved in broad attentional and memory processes, as well as self-reflection and self-focused thinking [44], and is a key node in the DMN [9–11]. Studies




Fig 1. Decreased precuneus gyrification is associated with trait anxiety. (A) Inflated and pial surface maps of the left hemisphere demonstrating decreased gyrification in the precuneus as a function of trait anxiety. Images on the left depict the medial view of the left hemisphere. Images on the right are a view from the top of the right hemisphere and are tilted 30 degrees to provide a better angle for viewing the cluster extent. (B) Scatterplot of the correlation between trait anxiety and average LGI within the precuneus cluster.

doi:10.1371/journal.pone.0149434.g001

of anxiety and anxiety disorders have found abnormalities in both activation and functional connectivity of the DMN [45]. Increased DMN activation has been also been linked to subjective reports of worry [13,14]. Additionally, several studies have found that in healthy individuals, precuneus activation is negatively correlated with activation of the amygdala [46,47]. Moreover, connectivity between these regions has also shown to be altered in social anxiety [48], panic disorder [49] and in youths with generalized anxiety disorder [50]. Thus, the precuneus may play a modulatory role in affective processing, which is disrupted in anxiety [7,47]. The current findings raise the possibility that structural abnormalities in the precuneus may underlie some of the functional abnormalities previously observed in anxiety in this region and related circuitry.

Our findings are consistent with previous studies showing precuneus structural abnormalities in anxiety-related personality dimensions and LGI-specific precuneus findings in clinical populations. Both behavioral inhibition [7] and harm avoidance, [51] have been associated with decreased precuneus gray matter volume. This relationship has also been detected in trait anxiety, but did not survive correction for multiple comparisons [52]. Additionally, although this is the first study to assess gyrification within dispositional anxiety, our findings are consistent with previous studies examining gyrification in clinical samples of anxiety and depression. Yoon and colleagues [32] demonstrated that hypogyrfication in the precuneus was associated with greater severity of symptoms in panic disorder, and Zhang and colleagues found precuneus/posterior cingulate cortex (PCC) hypogyrfication in individuals with depression. Nixon and colleagues [33] replicated the finding of decreased precuneus gyrification in depression and further showed that this hypogyrfication was associated with hyper-connectivity between the precuneus and the dorsal lateral prefrontal cortex. Their findings are consistent with the hypothesis that morphological abnormalities may reflect altered functional connectivity. The present study extends these previous findings by suggesting that abnormalities in precuneus cortical folding across these disorders may in part be due to the presence of these abnormalities in dispositional anxiety, a core risk factor that is common to these emotional disorders [2–4].




Fig 2. Correlations between LGI and other cortical measurements. (A) Scatterplot of the correlation between total surface area and average LGI within the precuneus. (B) Scatterplot demonstrating a negative correlation between average cortical thickness and average LGI within the precuneus. (C) Scatterplot demonstrating the correlation between trait anxiety and LGI when controlling for both cortical surface area and thickness.

doi:10.1371/journal.pone.0149434.g002

As posited above, the current findings may provide insight into the development of trait anxiety by demonstrating a structural correlate that may arise during neurodevelopment. Although earlier theories posit that the degree of brain gyrfication takes place primarily in utero and then stays constant through childhood and beyond [16,17], recent evidence has suggested that there are developmental changes in gyrfication throughout the lifespan [16,20–22]. Since the age of onset for various anxiety disorders ranges from early childhood to adulthood [53], it is hard to definitely understand when these cortical differences occur without assessing changes longitudinally. Additionally, the current study can only infer that these patterns may represent abnormalities in neurodevelopment, for we did not assess cortical changes across development or between individuals of different ages. However, structural abnormalities in the precuneus and posterior cingulate have been identified in pediatric samples, with anxious youths showing gray matter volume abnormalities [54,55], supporting the possibility that the current LGI findings represent aberrations in early neurodevelopment.

We also investigated how other aspects of cortical structure correlated with LGI in the precuneus in order to provide a broader picture of the cortical surface and better understand how these measurements are related to each other. Consistent with previous research [20,21], surface area had a positive relationship with LGI while cortical thickness had a negative relationship. While LGI was associated with thickness and surface area, these variables did not fully account for the relationship between LGI and anxiety. Since other cortical structural measures could not fully explain the relationship between LGI and trait anxiety, it is possible that other structural factors are influencing this relationship. For instance, certain theories of gyrfication have emphasized the role of white matter tension [43] or white matter constraints [56] in the development of gyrfication. Therefore, it is possible that the current findings reflect underlying differences in white matter strength. Further research on gyrfication may be useful in understanding the association between gyrfication abnormalities and the structural connectivity of broader neural networks in this population.

Conclusions

In conclusion, we found differences in cortical gyrfication associated with trait anxiety that could not be explained by surface area or thickness within a key DMN node. These findings are consistent with findings in clinical populations, highlighting the possibility that differences in local parietal connectivity are associated with a personality dimension that is a key risk factor for many of the most common forms of psychopathology. Although we did not investigate connectivity directly, future research may shed light on whether these structural differences are linked to disruptions in network connectivity. This structural aberration may in turn contribute to the emotional difficulties evident in high trait anxiety through influences on a primary node of the DMN subserving self-referential and self-conscious thought.

Author Contributions

Conceived and designed the experiments: TAM CLL. Performed the experiments: TAM CLL WSP ELB. Analyzed the data: TAM CLL WSP ELB SS. Contributed reagents/materials/analysis tools: CLL. Wrote the paper: TAM CLL WSP ELB SS KML.

References

1. Rauch SL, Milad MR, Orr SP, Quinn BT, Fischl B, Pitman RK. Orbitofrontal thickness, retention of fear extinction, and extraversion. *Neuroreport*. 2005; 16: 1909–1912. PMID: [16272877](#)
2. Kotov R, Gamez W, Schmidt F, Watson D. Linking “big” personality traits to anxiety, depressive, and substance use disorders: a meta-analysis. *Psychol Bull*. 2010; 136: 768. doi: [10.1037/a0020327](#) PMID: [20804236](#)

3. Lahey BB. Public health significance of neuroticism. *Am Psychol.* 2009; 64: 241. doi: [10.1037/a0015309](https://doi.org/10.1037/a0015309) PMID: [19449983](https://pubmed.ncbi.nlm.nih.gov/19449983/)
4. Blackford JU, Pine DS. Neural substrates of childhood anxiety disorders: a review of neuroimaging findings. *Child Adolesc Psychiatr Clin N Am.* 2012; 21: 501–525. doi: [10.1016/j.chc.2012.05.002](https://doi.org/10.1016/j.chc.2012.05.002) PMID: [22800991](https://pubmed.ncbi.nlm.nih.gov/22800991/)
5. Wright CI, Williams D, Feczko E, Barrett LF, Dickerson BC, Schwartz CE, et al. Neuroanatomical correlates of extraversion and neuroticism. *Cereb Cortex.* 2006; 16: 1809–1819. PMID: [16421327](https://pubmed.ncbi.nlm.nih.gov/16421327/)
6. Kühn S, Schubert F, Gallinat J. Structural correlates of trait anxiety: reduced thickness in medial orbitofrontal cortex accompanied by volume increase in nucleus accumbens. *J Affect Disord.* 2011; 134: 315–319. doi: [10.1016/j.jad.2011.06.003](https://doi.org/10.1016/j.jad.2011.06.003) PMID: [21705088](https://pubmed.ncbi.nlm.nih.gov/21705088/)
7. Fuentes P, Barrós-Loscertales A, Bustamante JC, Rosell P, Costumero V, Ávila C. Individual differences in the Behavioral Inhibition System are associated with orbitofrontal cortex and precuneus gray matter volume. *Cogn Affect Behav Neurosci.* 2012; 12: 491–498. doi: [10.3758/s13415-012-0099-5](https://doi.org/10.3758/s13415-012-0099-5) PMID: [22592859](https://pubmed.ncbi.nlm.nih.gov/22592859/)
8. Milad MR, Rauch SL. The role of the orbitofrontal cortex in anxiety disorders. *Ann N Y Acad Sci.* 2007; 1121: 546–561. PMID: [17698998](https://pubmed.ncbi.nlm.nih.gov/17698998/)
9. Raichle ME, Snyder AZ. A default mode of brain function: a brief history of an evolving idea. *Neuroimage.* 2007; 37: 1083–1090. PMID: [17719799](https://pubmed.ncbi.nlm.nih.gov/17719799/)
10. Raichle ME, MacLeod AM, Snyder AZ, Powers WJ, Gusnard DA, Shulman GL. A default mode of brain function. *Proc Natl Acad Sci U S A.* 2001; 98: 676–682. PMID: [11209064](https://pubmed.ncbi.nlm.nih.gov/11209064/)
11. Buckner RL, Andrews-Hanna JR, Schacter DL. The brain's default network. *Ann N Y Acad Sci.* 2008; 1124: 1–38. doi: [10.1196/annals.1440.011](https://doi.org/10.1196/annals.1440.011) PMID: [18400922](https://pubmed.ncbi.nlm.nih.gov/18400922/)
12. Berman MG, Peltier S, Nee DE, Kross E, Deldin PJ, Jonides J. Depression, rumination and the default network. *Soc Cogn Affect Neurosci.* 2011; 6: 548–555. doi: [10.1093/scan/nsq080](https://doi.org/10.1093/scan/nsq080) PMID: [20855296](https://pubmed.ncbi.nlm.nih.gov/20855296/)
13. Servaas MN, Riese H, Ormel J, Aleman A. The neural correlates of worry in association with individual differences in neuroticism. *Hum Brain Mapp.* 2014; 35: 4303–4315. doi: [10.1002/hbm.22476](https://doi.org/10.1002/hbm.22476) PMID: [24532549](https://pubmed.ncbi.nlm.nih.gov/24532549/)
14. Paulesu E, Sambugaro E, Torti T, Danelli L, Ferri F, Scialfa G, et al. Neural correlates of worry in generalized anxiety disorder and in normal controls: a functional MRI study. *Psychol Med.* 2010; 40: 117–124. doi: [10.1017/S0033291709005649](https://doi.org/10.1017/S0033291709005649) PMID: [19419593](https://pubmed.ncbi.nlm.nih.gov/19419593/)
15. Chi JG, Dooling EC, Gilles FH. Gyral development of the human brain. *Ann Neurol.* 1977; 1: 86–93. PMID: [560818](https://pubmed.ncbi.nlm.nih.gov/560818/)
16. White T, Su S, Schmidt M, Kao C, Sapiro G. The development of gyrfication in childhood and adolescence. *Brain Cogn.* 2010; 72: 36–45. doi: [10.1016/j.bandc.2009.10.009](https://doi.org/10.1016/j.bandc.2009.10.009) PMID: [19942335](https://pubmed.ncbi.nlm.nih.gov/19942335/)
17. Armstrong E, Schleicher A, Omran H, Curtis M, Zilles K. The ontogeny of human gyrfication. *Cereb Cortex.* 1995; 5: 56–63. PMID: [7719130](https://pubmed.ncbi.nlm.nih.gov/7719130/)
18. Laughlin SB, Sejnowski TJ. Communication in neuronal networks. *Science.* 2003; 301: 1870–1874. PMID: [14512617](https://pubmed.ncbi.nlm.nih.gov/14512617/)
19. Zilles K, Armstrong E, Schleicher A, Kretschmann H. The human pattern of gyrfication in the cerebral cortex. *Anat Embryol.* 1988; 179: 173–179. PMID: [3232854](https://pubmed.ncbi.nlm.nih.gov/3232854/)
20. Klein D, Rotarska-Jagiela A, Genc E, Sritharan S, Mohr H, Roux F, et al. Adolescent brain maturation and cortical folding: evidence for reductions in gyrfication. *PloS one.* 2014; 9: e84914. doi: [10.1371/journal.pone.0084914](https://doi.org/10.1371/journal.pone.0084914) PMID: [24454765](https://pubmed.ncbi.nlm.nih.gov/24454765/)
21. Hogstrom LJ, Westlye LT, Walhovd KB, Fjell AM. The structure of the cerebral cortex across adult life: age-related patterns of surface area, thickness, and gyrfication. *Cereb Cortex.* 2013; 23: 2521–2530. doi: [10.1093/cercor/bhs231](https://doi.org/10.1093/cercor/bhs231) PMID: [22892423](https://pubmed.ncbi.nlm.nih.gov/22892423/)
22. Alemán-Gómez Y, Janssen J, Schnack H, Balaban E, Pina-Camacho L, Alfaro-Almagro F, et al. The human cerebral cortex flattens during adolescence. *J Neurosci.* 2013; 33: 15004–15010. doi: [10.1523/JNEUROSCI.1459-13.2013](https://doi.org/10.1523/JNEUROSCI.1459-13.2013) PMID: [24048830](https://pubmed.ncbi.nlm.nih.gov/24048830/)
23. White T, Andreasen NC, Nopoulos P. Brain volumes and surface morphology in monozygotic twins. *Cereb Cortex.* 2002; 12: 486–493. PMID: [11950766](https://pubmed.ncbi.nlm.nih.gov/11950766/)
24. Bartley AJ, Jones DW, Weinberger DR. Genetic variability of human brain size and cortical gyral patterns. *Brain.* 1997; 120 (Pt 2): 257–269. PMID: [9117373](https://pubmed.ncbi.nlm.nih.gov/9117373/)
25. Lohmann GD, von Cramon DY, Steinmetz H. Sulcal variability of twins. *Cereb Cortex.* 1999; 9: 754–763. PMID: [10554998](https://pubmed.ncbi.nlm.nih.gov/10554998/)
26. Dauvermann MR, Mukherjee P, Moorhead WT, Stanfield AC, Fusar-Poli P, Lawrie SM, et al. Relationship between gyrfication and functional connectivity of the prefrontal cortex in subjects at high genetic risk of schizophrenia. *Curr Pharm Des.* 2012; 18: 434–442. PMID: [22239574](https://pubmed.ncbi.nlm.nih.gov/22239574/)

27. Schaer M, Ottet MC, Scariati E, Dukes D, Franchini M, Eliez S, et al. Decreased frontal gyrfication correlates with altered connectivity in children with autism. *Front Hum Neurosci*. 2013; 7: 750. doi: [10.3389/fnhum.2013.00750](https://doi.org/10.3389/fnhum.2013.00750) PMID: [24265612](https://pubmed.ncbi.nlm.nih.gov/24265612/)
28. Palaniyappan L, Mallikarjun P, Joseph V, White TP, Liddle PF. Folding of the prefrontal cortex in schizophrenia: regional differences in gyrfication. *Biol Psychiatry*. 2011; 69: 974–979. doi: [10.1016/j.biopsych.2010.12.012](https://doi.org/10.1016/j.biopsych.2010.12.012) PMID: [21257157](https://pubmed.ncbi.nlm.nih.gov/21257157/)
29. White T, Andreasen NC, Nopoulos P, Magnotta V. Gyrfication abnormalities in childhood-and adolescent-onset schizophrenia. *Biol Psychiatry*. 2003; 54: 418–426. PMID: [12915286](https://pubmed.ncbi.nlm.nih.gov/12915286/)
30. Wallace GL, Robustelli B, Dankner N, Kenworthy L, Giedd JN, Martin A. Increased gyrfication, but comparable surface area in adolescents with autism spectrum disorders. *Brain*. 2013; 136: 1956–1967. doi: [10.1093/brain/awt106](https://doi.org/10.1093/brain/awt106) PMID: [23715094](https://pubmed.ncbi.nlm.nih.gov/23715094/)
31. Hardan AY, Jou RJ, Keshavan MS, Varma R, Minshew NJ. Increased frontal cortical folding in autism: a preliminary MRI study. *Psychiatry Res Neuroimaging*. 2004; 131: 263–268. PMID: [15465295](https://pubmed.ncbi.nlm.nih.gov/15465295/)
32. Yoon S, Jun CS, Jeong HS, Lee S, Lim SM, Ma J, et al. Altered cortical gyrfication patterns in panic disorder: Deficits and potential compensation. *J Psychiatr Res*. 2013; 47: 1446–1454. doi: [10.1016/j.jpsychires.2013.06.015](https://doi.org/10.1016/j.jpsychires.2013.06.015) PMID: [23871448](https://pubmed.ncbi.nlm.nih.gov/23871448/)
33. Nixon NL, Liddle PF, Nixon E, Worwood G, Liotti M, Palaniyappan L. Biological vulnerability to depression: linked structural and functional brain network findings. *Br J Psychiatry*. 2014; 204: 283–289. doi: [10.1192/bjp.bp.113.129965](https://doi.org/10.1192/bjp.bp.113.129965) PMID: [24357570](https://pubmed.ncbi.nlm.nih.gov/24357570/)
34. Zhang Y, Yu C, Zhou Y, Li K, Li C, Jiang T. Decreased gyrfication in major depressive disorder. *Neuroreport*. 2009; 20: 378–380. doi: [10.1097/WNR.0b013e3283249b34](https://doi.org/10.1097/WNR.0b013e3283249b34) PMID: [19218876](https://pubmed.ncbi.nlm.nih.gov/19218876/)
35. Schaer M, Cuadra MB, Tamarit L, Lazeyras F, Eliez S, Thiran JP. A surface-based approach to quantify local cortical gyrfication. *IEEE Trans Med Imaging*. 2008; 27: 161–170. doi: [10.1109/TMI.2007.903576](https://doi.org/10.1109/TMI.2007.903576) PMID: [18334438](https://pubmed.ncbi.nlm.nih.gov/18334438/)
36. First MB, Spitzer RL, Gibbon M, Williams JBW. Structured Clinical Interview for DSM-IV Axis I Disorders, Research Version, Non-Patient Edition (SCID-I/NP). New York: New York State Psychiatric Institute, Biometrics Research; 2002.
37. Spielberger CD, Gorsuch RL, Lushene RE, Vagg PR, Jacobs GA. State-Trait Anxiety Inventory for Adults: Manual and Sample: Manual, Instrument and Scoring Guide. Palo Alto, CA: Consulting Psychologists Press; 1983.
38. Barnes LLB, Harp D, Jung WS. Reliability generalization of scores on the Spielberger state-trait anxiety inventory. *Educational and Psychological Measurement*. 2002; 62: 603–618.
39. Dale AM, Fischl B, Sereno MI. Cortical surface-based analysis: I. Segmentation and surface reconstruction. *Neuroimage*. 1999; 9: 179–194. PMID: [9931268](https://pubmed.ncbi.nlm.nih.gov/9931268/)
40. Fischl B, Sereno MI, Dale AM. Cortical surface-based analysis: II: Inflation, flattening, and a surface-based coordinate system. *Neuroimage*. 1999; 9: 195–207. PMID: [9931269](https://pubmed.ncbi.nlm.nih.gov/9931269/)
41. Schaer M, Cuadra MB, Schmansky N, Fischl B, Thiran JP, Eliez S. How to measure cortical folding from MR images: a step-by-step tutorial to compute local gyrfication index. *J Vis Exp*. 2012; 59: e3417. doi: [10.3791/3417](https://doi.org/10.3791/3417) PMID: [22230945](https://pubmed.ncbi.nlm.nih.gov/22230945/)
42. Raznahan A, Shaw P, Lalonde F, Stockman M, Wallace GL, Greenstein D, et al. How does your cortex grow? *J Neurosci*. 2011; 31: 7174–7177. doi: [10.1523/JNEUROSCI.0054-11.2011](https://doi.org/10.1523/JNEUROSCI.0054-11.2011) PMID: [21562281](https://pubmed.ncbi.nlm.nih.gov/21562281/)
43. Van Essen DC. A tension-based theory of morphogenesis and compact wiring in the central nervous system. *Nature*. 1997; 385: 313–318. PMID: [9002514](https://pubmed.ncbi.nlm.nih.gov/9002514/)
44. Cavanna AE, Trimble MR. The precuneus: a review of its functional anatomy and behavioural correlates. *Brain*. 2006; 129: 564–583. PMID: [16399806](https://pubmed.ncbi.nlm.nih.gov/16399806/)
45. Sylvester CM, Corbetta M, Raichle ME, Rodebaugh TL, Schlaggar BL, Sheline YI, et al. Functional network dysfunction in anxiety and anxiety disorders. *Trends Neurosci*. 2012; 35: 527–535. doi: [10.1016/j.tins.2012.04.012](https://doi.org/10.1016/j.tins.2012.04.012) PMID: [22658924](https://pubmed.ncbi.nlm.nih.gov/22658924/)
46. Roy AK, Shehzad Z, Margulies DS, Kelly AC, Uddin LQ, Gotimer K, et al. Functional connectivity of the human amygdala using resting state fMRI. *Neuroimage*. 2009; 45: 614–626. doi: [10.1016/j.neuroimage.2008.11.030](https://doi.org/10.1016/j.neuroimage.2008.11.030) PMID: [19110061](https://pubmed.ncbi.nlm.nih.gov/19110061/)
47. Zhang S, Li CS. Functional connectivity mapping of the human precuneus by resting state fMRI. *Neuroimage*. 2012; 59: 3548–3562. doi: [10.1016/j.neuroimage.2011.11.023](https://doi.org/10.1016/j.neuroimage.2011.11.023) PMID: [22116037](https://pubmed.ncbi.nlm.nih.gov/22116037/)
48. Hahn A, Stein P, Windischberger C, Weissenbacher A, Spindelegger C, Moser E, et al. Reduced resting-state functional connectivity between amygdala and orbitofrontal cortex in social anxiety disorder. *Neuroimage*. 2011; 56: 881–889. doi: [10.1016/j.neuroimage.2011.02.064](https://doi.org/10.1016/j.neuroimage.2011.02.064) PMID: [21356318](https://pubmed.ncbi.nlm.nih.gov/21356318/)

49. Pannekoek JN, Veer IM, van Tol MJ, van der Werff, Steven JA, Demenescu LR, Aleman A, et al. Aberrant limbic and salience network resting-state functional connectivity in panic disorder without comorbidity. *J Affect Disord.* 2013; 145: 29–35. doi: [10.1016/j.jad.2012.07.006](https://doi.org/10.1016/j.jad.2012.07.006) PMID: [22858265](https://pubmed.ncbi.nlm.nih.gov/22858265/)
50. Strawn JR, Bitter SM, Weber WA, Chu WJ, Whitsel RM, Adler C, et al. Neurocircuitry of generalized anxiety disorder in adolescents: a pilot functional neuroimaging and functional connectivity study. *Depress Anxiety.* 2012; 29: 939–947. doi: [10.1002/da.21961](https://doi.org/10.1002/da.21961) PMID: [22628125](https://pubmed.ncbi.nlm.nih.gov/22628125/)
51. Gardini S, Cloninger CR, Venneri A. Individual differences in personality traits reflect structural variance in specific brain regions. *Brain Res Bull.* 2009; 79: 265–270. doi: [10.1016/j.brainresbull.2009.03.005](https://doi.org/10.1016/j.brainresbull.2009.03.005) PMID: [19480986](https://pubmed.ncbi.nlm.nih.gov/19480986/)
52. Donzuso G, Cerasa A, Gioia MC, Caracciolo M, Quattrone A. The neuroanatomical correlates of anxiety in a healthy population: differences between the State-Trait Anxiety Inventory and the Hamilton Anxiety Rating Scale. *Brain and Behavior.* 2014; 4: 504–514. doi: [10.1002/brb3.232](https://doi.org/10.1002/brb3.232) PMID: [25161817](https://pubmed.ncbi.nlm.nih.gov/25161817/)
53. Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry.* 2005; 62: 593–602. PMID: [15939837](https://pubmed.ncbi.nlm.nih.gov/15939837/)
54. Strawn JR, Hamm L, Fitzgerald DA, Fitzgerald KD, Monk CS, Phan KL. Neurostructural abnormalities in pediatric anxiety disorders. *J Anxiety Disord.* 2015; 32: 81–88. doi: [10.1016/j.janxdis.2015.03.004](https://doi.org/10.1016/j.janxdis.2015.03.004) PMID: [25890287](https://pubmed.ncbi.nlm.nih.gov/25890287/)
55. Strawn JR, Chu WJ, Whitsel RM, Weber WA, Norris MM, Adler CM, et al. A pilot study of anterior cingulate cortex neurochemistry in adolescents with generalized anxiety disorder. *Neuropsychobiology.* 2013; 67: 224–229. doi: [10.1159/000347090](https://doi.org/10.1159/000347090) PMID: [23635944](https://pubmed.ncbi.nlm.nih.gov/23635944/)
56. Tallinen T, Chung JY, Biggins JS, Mahadevan L. Gyrfication from constrained cortical expansion. *Proc Natl Acad Sci U S A.* 2014; 111: 12667–12672. doi: [10.1073/pnas.1406015111](https://doi.org/10.1073/pnas.1406015111) PMID: [25136099](https://pubmed.ncbi.nlm.nih.gov/25136099/)