



Coronary Large Conductance Ca^{2+} -Activated K^+ Channel Dysfunction in Diabetes Mellitus

Tong Lu* and Hon-Chi Lee*

Department of Cardiovascular Medicine, Mayo Clinic, Rochester, MN, United States

OPEN ACCESS

Edited by:

Yoshiaki Suzuki,
Nagoya City University, Japan

Reviewed by:

Prosper N'Gouemo,
Howard University, United States
Craig Doupnik,
USF Health, United States

*Correspondence:

Hon-Chi Lee
lee.honchi@mayo.edu
Tong Lu
lu.tong@mayo.edu

Specialty section:

This article was submitted to
Membrane Physiology and
Membrane Biophysics,
a section of the journal
Frontiers in Physiology

Received: 30 July 2021

Accepted: 14 September 2021

Published: 21 October 2021

Citation:

Lu T and Lee H (2021) Coronary
Large Conductance Ca^{2+} -Activated
 K^+ Channel Dysfunction in
Diabetes Mellitus.
Front. Physiol. 12:750618.
doi: 10.3389/fphys.2021.750618

Diabetes mellitus (DM) is an independent risk of macrovascular and microvascular complications, while cardiovascular diseases remain a leading cause of death in both men and women with diabetes. Large conductance Ca^{2+} -activated K^+ (BK) channels are abundantly expressed in arteries and are the key ionic determinant of vascular tone and organ perfusion. It is well established that the downregulation of vascular BK channel function with reduced BK channel protein expression and altered intrinsic BK channel biophysical properties is associated with diabetic vasculopathy. Recent efforts also showed that diabetes-associated changes in signaling pathways and transcriptional factors contribute to the downregulation of BK channel expression. This manuscript will review our current understandings on the molecular, physiological, and biophysical mechanisms that underlie coronary BK channelopathy in diabetes mellitus.

Keywords: BK channel, diabetes mellitus, coronary arteries, blood vessels, regulation

INTRODUCTION

Diabetes mellitus (DM) has become a global epidemic. The incidence and the prevalence of DM have steadily increased over the past few decades. According to the WHO report in 2021, over 422 million people worldwide have DM with a prevalence of 8.6%, causing 1.6 million deaths annually.¹ Type 1 diabetes mellitus (T1DM) accounts for 5–10% of the total cases of DM and is caused by autoimmune-mediated destruction of pancreatic β -cells, leading to hyperglycemia and insulin dependence (Bluestone et al., 2010; Op De Beeck and Eizirik, 2016). Type 2 diabetes mellitus (T2DM) represents 90–95% of the total cases of DM and is caused by insulin resistance with hyperinsulinemia, hyperglycemia, and hyperlipidemia in most patients (Pandey et al., 2015; Halim and Halim, 2019).

Both T1DM and T2DM are intimately related to micro-vascular and macro-vascular diseases, including ischemic heart disease, cerebrovascular disease, and peripheral vascular disease, resulting in myocardial infarction, stroke, retinopathy, nephropathy, and neuropathy with organ and tissue damages in 70% of diabetic patients (Kurisu et al., 2003; Yeung et al., 2012; Beckman and Creager, 2016; Sorop et al., 2016). The clinical consequences of diabetic vascular complication are devastating. DM is the leading cause of end stage renal disease, new cases of blindness,

¹https://www.who.int/health-topics/diabetes#tab=tab_1

and non-traumatic lower extremity amputation, imposing global direct health expenditure of \$ 760 billion in 2019 with a projected \$ 825 billion by 2030 and \$ 845 billion by 2045 (Williams et al., 2020). Hence, it is critically important to understand the mechanisms of vascular dysregulation in DM so that better diagnostic and therapeutic approaches can be developed to treat diabetic vascular complications more effectively.


Ionic mechanisms play a central role in the regulation of vascular reactivity. Vascular large conductance Ca^{2+} -activated K^+ (BK) channels are major determinants of such regulation. BK channels are densely populated in vascular smooth muscle cells (SMCs), particularly in small resistance arteries, and provide tight regulation of vascular tone and tissue perfusion. It is well established that vascular BK channel expression and function are abnormal in DM. Diabetic patients are known to have worse cardiovascular events and outcome, with higher risks of ischemic heart disease and myocardial infarction (Kurusu et al., 2003; Yeung et al., 2012; Sorop et al., 2016). In this review, we will focus on recent findings in the coronary arterial SMCs, highlighting the diabetes-mediated changes in channel expression, function, and intrinsic properties, as well as the molecular mechanisms associated with these changes.

STRUCTURE AND FUNCTION OF VASCULAR BK CHANNELS

Cardiac perfusion is regulated by vasoactive agents released by the endothelium from mechanical sensing of luminal shear stress, including endothelium-derived relaxation factors (EDRF) and endothelium-derived hyperpolarizing factors (EDHF), the pharmacologic action of neuroendocrine factors, and the response of coronary arteriolar SMCs to intraluminal pressure (Goodwill et al., 2017). Functional vascular BK channels are composed of the pore-forming α -subunits (BK- α) and the accessory β 1-subunits

(BK- β 1) and/or γ 1-subunits (BK- γ 1; **Figure 1**; Knaus et al., 1994; Yan and Aldrich, 2012). Four BK- α and four BK- β 1 assemble to form a functional BK channel. The stoichiometry and interaction between BK- α and BK- γ 1 are currently unclear. BK- α is expressed ubiquitously on the cell surface and in mitochondrial membranes of excitable and non-excitable cells, while BK- β 1 is distributed in the cell membranes of excitable cells. BK- γ 1 is mainly found in the cell membrane of non-excitable cells (Singh et al., 2013; Li et al., 2016). BK- α (encoded by the *KCNMA1* gene) contains the structure of six transmembrane domains (S1–S6) of voltage-gated K^+ channels in which S1–S4 constitute the voltage-sensing domain (VSD) and the S5-P loop-S6 form the ion permeation domain, containing the conserved K^+ selectivity filter (TVGYG; Ma et al., 2006; Cui et al., 2009). In addition, the BK channel has a unique S0 segment unit in the extracellular N-terminus and a large C-terminal domain (CTD). The CTD has four cytosolic domains (S7–S10) with two regulators of K^+ conductance domains (RCK1 and RCK2) that contain two high-affinity Ca^{2+} binding sites (Wu and Marx, 2010; Yuan et al., 2010). One such site is the Ca^{2+} bowl (889-QFLDQDDDD-897) in RCK2 with a Ca^{2+} concentration at half-maximal effect (EC_{50}) in the 10^{-6} M range (Xia et al., 2002; Bao et al., 2004). The other site (D367/E535/R514) is located in RCK1 (**Figure 1**; Zeng et al., 2005; Zhang et al., 2010b). The RCK1s and RCK2s of four BK- α subunits form an octameric gating ring that connects to the VSD through a rigid linker (Yuan et al., 2010; Tao et al., 2017). Binding to intracellular free Ca^{2+} and membrane depolarization activate BK channels through allosteric changes in the gating ring.

In addition to Ca^{2+} - and voltage-dependent activation, BK- α activity is tightly regulated by its accessory subunits, BK- β and BK- γ (Li and Yan, 2016; Gonzalez-Perez and Lingle, 2019). Four isoforms of β subunits (BK- β 1–4, encoded by the *KCNMB1–4* genes) and γ subunits (BK- γ 1–4, encoded by the *LRRC26*, *LRRC38*, *LRRC52*, and *LRRC55* genes) have been cloned in mammalian cells (Li and Yan, 2016; Gonzalez-Perez and Lingle, 2019).



In vascular SMCs, BK- β 1 is the predominant vascular isoform. It contains two transmembrane domains (TM1 and TM2) with a relatively large extracellular loop that can reach the inner mouth of the BK- α channel pore and modulates the binding of iberiotoxin (IBTX) and the effects of fatty acids on BK channel activity (Torres et al., 2014). The TM1 is thought to interact with the S2 of an adjacent BK- α subunit and the TM2 with the S0 of another adjacent BK- α subunit (Liu et al., 2010). The presence of the BK- β 1 subunit enhances channel sensitivity to Ca^{2+} activation.

BK- γ 1 is also expressed in vascular SMCs (Evanson et al., 2014). BK- γ 1 shares the structure of the leucine-rich repeat (LRR) protein superfamily and contains an extracellular N-terminus with six LRRs, a single transmembrane domain, and a short intracellular C-terminus (Figure 1). The effects of BK- γ 1 on BK- α regulation can be reproduced by a 40-amino acid peptide containing the transmembrane domain of BK- γ 1, suggesting that this is an important structure in the regulation of BK channel physiology (Li et al., 2016). BK- γ 1 is known to enhance BK- α sensitivity to Ca^{2+} and voltage stimuli by magnitudes similar to those of BK- β 1, allowing BK channel activation in the physiological range of intracellular free Ca^{2+} concentrations and membrane potentials of vascular SMCs (Tanaka et al., 1997; Cox and Aldrich, 2000; Yan and Aldrich, 2012). In heterologous expression systems, BK- β and BK- γ subunits can co-exist in the same functional BK channel complex. Their effects on the intrinsic properties of the channel were additive, suggesting that the multiplicity of BK- β /BK- γ combinations would generate a range of BK channels with distinct functional properties according to the specific stoichiometry of the contributing subunits (Gonzalez-Perez et al., 2015). Since nothing is known about the role of BK- γ in the regulation of coronary BK channels in DM, this review will focus on the findings regarding BK- α and BK- β 1 pathophysiology in DM.

Intracellular Ca^{2+} homeostasis in vascular SMCs is regulated by the balance between sarcolemmal Ca^{2+} entry (L-type Ca^{2+} channels and the transient receptor potential channels; TRP, etc.), release of Ca^{2+} from the endoplasmic reticulum/sarcoplasmic reticulum, uptake of cytoplasmic Ca^{2+} into intracellular stores, and extrusion through the sarcolemmal Ca^{2+} pump and Na^+ / Ca^{2+} exchanger (Leopold, 2015). In vascular SMCs, BK channels link Ca^{2+} homeostasis with cellular excitability and regulate vascular tone through membrane hyperpolarization, providing a negative feedback mechanism on Ca^{2+} entry. BK channels are colocalized with L-type Ca^{2+} channels and TRPC/TRPV channels to form BK channel- Ca^{2+} signaling complexes in the sarcolemma of vascular SMCs, allowing channel regulation in the local cellular milieu (Earley et al., 2005; Kwan et al., 2009; Suzuki et al., 2013; Hashad et al., 2018). Activation of L-type Ca^{2+} channels and TRP channels in vascular SMCs produces Ca^{2+} sparklets and triggers Ca^{2+} release from the SR to generate Ca^{2+} sparks (Nelson and Quayle, 1995; Takeda et al., 2011). With a single channel conductance of ~ 300 pS, BK channels contribute to 50% of the total K^+ currents in coronary arterial SMCs (Wang et al., 2011; Sun et al., 2020). Activation of vascular BK channels by Ca^{2+} sparks/sparklets in their vicinity gives rise to spontaneous transient outward currents (STOCs),

which hyperpolarize the cellular membrane potentials, inactivate L-type Ca^{2+} channels and TRP channels, reduce intracellular Ca^{2+} concentrations, and lead to vasorelaxation (Nelson et al., 1995; Ledoux et al., 2006). In addition, BK channels are also expressed in vascular endothelial cells (ECs). Activation of endothelial BK channels may hyperpolarize adjacent SMCs, bestowing EDHF effects (Bryan et al., 2005; Hughes et al., 2010). Nevertheless, activation of BK channels contributes to more than 70% of total vasodilation induced by bradykinin (Miura et al., 1999) and 40% of total vasodilation induced by shear stress in human coronary resistance vessels (Lu et al., 2019).

CORONARY BK CHANNEL DYSFUNCTION IN DM

Both T1DM and T2DM are known to be independent risk factors for cardiovascular diseases, and cardiovascular diseases continue to be a leading cause of mortality in diabetic patients (Dhalla et al., 1985; Stone et al., 1989; Brindisi et al., 2010; Leon and Maddox, 2015). Although, the prevalence of cardiovascular disease in the general population has decreased by 35–40% over recent decades, such a decline has not been observed in patients with DM (Gregg et al., 2007; Beckman and Creager, 2016; Cefalu et al., 2018). Endothelial dysfunction has been recognized as the mechanism that underlies vascular pathology of DM. Subsequent findings confirm that vascular smooth muscle dysfunction is equally important in the pathophysiology of diabetic cardiovascular complications (Creager et al., 2003).

Impaired BK channel-induced vasodilation was first discovered in the cerebral arteries of fructose-rich diet-induced insulin-resistant rats (Dimitropoulou et al., 2002; Erdos et al., 2002). Patch clamp studies provided direct evidence of BK channel dysfunction in freshly isolated coronary arterial SMCs from Zucker diabetic fatty (ZDF) rats, a genetic animal model of T2DM (Lu et al., 2005). Abnormal vascular BK channel function was also found in other diabetic animal models, including streptozotocin (STZ)-induced T1DM rodents, db/db T2DM mice, high fat diet (HFD)-induced obesity/diabetic mice and swine (Dimitropoulou et al., 2002; Pietryga et al., 2005; Burnham et al., 2006; McGahon et al., 2007; Yang et al., 2007; Dong et al., 2008; Lu et al., 2008, 2010, 2012, 2016, 2017a; Borbouse et al., 2009; Navedo et al., 2010; Zhang et al., 2010a; Mori et al., 2011; Nystoriak et al., 2014; Yi et al., 2014). It is worth noting that diabetic vascular BK channel dysfunction is a common finding in most vascular beds, but the results can vary in different species, animal models, and disease status (Mokelke et al., 2003, 2005; Christ et al., 2004; Pietryga et al., 2005; Burnham et al., 2006; Davies et al., 2007; McGahon et al., 2007; Lu et al., 2008; Borbouse et al., 2009; Navedo et al., 2010; Mori et al., 2011; Rueda et al., 2013; Nystoriak et al., 2014; Nieves-Cintrón et al., 2017). It has been found that in freshly isolated coronary arterioles from patients with T2DM, BK channel sensitivity to Ca^{2+} and voltage activation was reduced, indicating that the intrinsic biophysical properties of BK channels were altered in diabetic patients (Figure 2; Lu et al., 2019).




FIGURE 2 | voltages in the presence of 200 nM free Ca^{2+} in freshly isolated coronary arteriolar smooth muscle cells (SMCs) from non-diabetic controls and T2DM patients. BK channel was activated by membrane depolarization with reduced effects in diabetes mellitus (DM). The dashed line indicates the closed state (c) of channel. Right panel: BK channel open probability and voltage (nPo-V) relationships were fitted using the Boltzmann equation. The maximal nPo and voltage at half of maximal channel activation ($V_{0.5}$) were significantly decreased in T2DM patients ($n=9$), compared with controls ($n=12$). BK channel unitary current amplitude plotted against membrane voltages (i-V curves) were fitted using a linear equation. The unitary conductance of BK channels was not different between controls and T2DM patients. Data are presented as mean \pm SEM. There was a significant decrease in BK channel maximal nPo and $V_{0.5}$ in diabetic patients. **(C)** Altered BK channel kinetics in the coronary arterioles of T2DM patients. Typical histograms of BK channel open and closed dwell-time durations are illustrated. Data were obtained from inside-out patches at +60 mV in the presence of 200 nM free Ca^{2+} in the bath solution. Dwell-time distributions were best fitted by the sum of exponential probability density functions with three open time constant components (the slow τ_{o1} , the intermediate τ_{o2} , and the fast τ_{o3}) and four closed time constant components (the very slow τ_{c1} , the slow τ_{c2} , the intermediate τ_{c3} , and the fast τ_{c4}). Dashed lines represent the distribution of exponential components determined by the logarithm likelihood ratio test. The values of each time constant component and its relative weight (in parentheses) are given above each histogram. This figure was adapted from published results with the permission of Cardiovascular Research (Lu et al., 2019).

This finding supports the observation that the BK channel response to Ca^{2+} sparks was diminished in human diabetic vessels. The significance of coronary BK channel dysfunction in DM is underscored by the finding that ischemia-reperfusion-mediated myocardial infarction is exacerbated in STZ-induced T1DM mouse hearts and can be reproduced in non-diabetic hearts after exposure to the BK channel specific inhibitor, iberiotoxin (IBTX; Lu et al., 2016). Since IBTX is membrane impermeable and cardiac myocytes do not have BK- α expression on the sarcolemma, this finding provides evidence of the role of coronary vascular BK channels on cardioprotection during ischemia-reperfusion insults, as well as the loss of its protection in DM.

Altered BK Channel Protein Expression in Diabetic Vessels

Altered coronary vascular BK channel expression is common in DM (Burnham et al., 2006; McGahon et al., 2007). However, diverse levels of vascular BK channel expression in DM have been observed. In most cases, the protein expressions of BK channels are downregulated in coronary arteries (Burnham et al., 2006; Dong et al., 2008; Lu et al., 2008, 2017a; Zhang et al., 2010a; Rueda et al., 2013; Nystoriak et al., 2014; Li et al., 2017), but it was reportedly increased, despite impaired BK channel function in the coronary arteries of Ossabaw miniature swine with metabolic syndrome (Borhouse et al., 2009). Recently, human BK channel expression was examined in coronary arterioles obtained from atrial biopsies of patients who underwent coronary artery bypass grafting surgery. Protein downregulation was found in both BK- α and BK- β 1 in patients with T2DM, compared to age-matched non-diabetic subjects (Lu et al., 2019). However, the mRNA levels of BK- β 1 were (McGahon et al., 2007) not reduced in the coronary arteries of STZ-induced T1DM rats (Zhang et al., 2010a), db/db T2DM mice (Li et al., 2017) and HFD-induced diabetic mice (Lu et al., 2017a). The varied reports of BK channel expression suggest that a complex assortment of mechanisms exist in the regulation of vascular BK channel expression and function in DM. Reduced BK channel expression leads to impaired Ca^{2+} sparks/STOCs coupling, albeit the Ca^{2+} spark amplitudes and intracellular Ca^{2+} concentrations are known to be elevated in diabetic vascular SMCs.

Impaired BK Channel Biophysical Properties and Kinetics in Coronary Arterial SMCs in DM

Ca^{2+} -activated K^{+} channel currents (I) are determined by the number of activated channels (N), open probability (P_o), and channel unitary conductance (i), where $I = N \cdot P_o \cdot i$. BK channel current density is reduced in the coronary arteries of T1DM and T2DM animal models and in humans with DM (Lu et al., 2005, 2008, 2010, 2012, 2016, 2017a, 2019; Pietryga et al., 2005; Burnham et al., 2006; McGahon et al., 2007; Dong et al., 2008; Zhang et al., 2010a; Nystoriak et al., 2014; Yi et al., 2014; Li et al., 2017; Nieves-Cintrón et al., 2017; Tang et al., 2017; Zhang et al., 2020). BK channels are activated by intracellular free Ca^{2+} concentration and by membrane depolarization (Cox et al., 1997; Lu et al., 2008), and these are impaired in DM (Lu et al., 2008, 2019). BK channel sensitivity to voltage- and Ca^{2+} -mediated activation can be measured by using inside-out patch clamp studies in which the excised cell membrane can be clamped to various voltages and the cytoplasmic surface of the cell membrane directly exposed to bath solutions containing various free Ca^{2+} concentrations. In freshly isolated coronary arterial SMCs of ZDF rats at 8 months after the development of hyperglycemia, BK channels had a rightward-shifted Ca^{2+} concentration-dependent curve, with increased EC_{50} for Ca^{2+} activation and decreased Ca^{2+} cooperativity, compared to those of Lean control rats (Lu et al., 2008). Moreover, BK channel activation by membrane depolarization was also abnormal in coronary arterial SMCs of ZDF rats. The channel open probability-voltage (Po-V) relationships were rightward and downward shifted, with the voltage at 50% maximal Po increased by 40 mV. These results indicate that a higher cytoplasmic Ca^{2+} concentration and a more depolarized membrane potential are required to activate BK channels in DM. Changes in the intrinsic free energy of Ca^{2+} -binding ($\Delta\Delta\text{Ca}^{2+}$) that contributes to BK channel activation can be estimated based on the shift of Po-V relationship from 0 to 1 μM free Ca^{2+} in Lean and ZDF rats using the equation: $\Delta\Delta\text{Ca}^{2+} = -\Delta(z e V_{0.5})$, where z is the number of equivalence charge movement, e is the elementary charge, and $V_{0.5}$ is the voltage at half maximal activation (Shi et al., 2002). There was a 62.3% decrease in the $\Delta\Delta\text{Ca}^{2+}$ in ZDF rats, suggesting a less favorable condition for Ca^{2+} binding to

vascular BK channel Ca^{2+} sensors in ZDF rats (Lu et al., 2008). Similar results were also observed in BK channels in freshly isolated coronary microvascular SMCs from the atrial appendages of patients with T2DM. Ca^{2+} - and voltage sensitivity were significantly impaired in diabetic patients, with the maximal BK channel activity to free Ca^{2+} and voltage activation reduced by 70 and 50%, respectively (Figure 2; Lu et al., 2019). Such dysregulation contributed to a 27.4% attenuation in shear stress-mediated coronary arteriolar vasorelaxation in diabetic patients compared with non-diabetic controls (Lu et al., 2019). In addition, single BK channel current amplitudes were unaltered in DM, indicating that the conductance property of vascular BK channels is normal in DM.

Vascular large conductance Ca^{2+} -activated K^+ channel gating kinetics contain multiple components of open and closed states and dwell-times (McManus and Magleby, 1988, 1991). In coronary arterial SMCs, the open and closed dwell-time histograms of single BK channels were best fitted with three open-time constants: fast (τ_{o1}), intermediate (τ_{o2}), and slow (τ_{o3}), along with four closed-time constants: fast (τ_{c1}), intermediate (τ_{c2}), slow (τ_{c3}), and very slow (τ_{c4}). DM affects both channel open dwell-times and channel closed dwell-times. The BK channel mean closed-time constant and the individual closed-time constants were significantly prolonged. At the same time, the channel mean open-time constant and individual open-time constants were significantly reduced in DM. These findings were seen in both ZDF rats and in diabetic patients (Lu et al., 2008, 2019). These changes in BK channel gating kinetics suggest that channel openings are abbreviated, and closures prolonged in DM, with reduced channel P_o and maximal activation. Hence, diabetes not only affects BK channel expression, but also alters the intrinsic biophysical properties of the channel.

KCBMA1 and KCNMB1 Variations Associated With Obesity and DM

Genome-wide association studies (GWASs) are a powerful tool to find genetic variations associated with diseases. Results from a few studies have shown a strong association between *KCNMA1* splicing variants and the incidence of obesity or DM. The results from case-control cohorts involving 4,838 obese and 5,827 control subjects suggested that the *KCNMA1* rs2116830*G variant was associated with obesity with a *p* value of 2.82×10^{-10} (Jiao et al., 2011). A recent study reported that a *de novo* missense variant in *KCNMA1* (c.1123G>A) was identified in an adult male patient with a plethora of developmental phenotypes including neonatal DM. This loss-of-function polymorphism (p. G375A) of BK channel is located in the S6 transmembrane domain of BK channel (Liang et al., 2019). In addition, it is well known that BK- α and BK- $\beta 1$ undergo extensive alternative pre-mRNA splicing and that these splice variants have significant changes in BK channel intrinsic properties and surface expression (Poulsen et al., 2009). However, the pathophysiological roles of BK channel variants in the development of BK channelopathy in DM are largely unexplored and warrant further investigation.

SIGNALING MOLECULES AND PATHWAYS MEDIATING VASCULAR BK CHANNEL DYSFUNCTION IN DM

Effects of Reactive Oxygen Species on Vascular BK Channel Redox Modification

Increased reactive oxygen species (ROS) production is a hallmark of diabetic pathophysiology, and the role of ROS on vascular dysfunction has been extensively reviewed (Inoguchi et al., 2003; Konior et al., 2014). ROS is represented by a group of highly reactive molecules that include superoxide anion ($\text{O}_2^{\cdot-}$), peroxide ion (O_2^{2-}), hydrogen peroxide (H_2O_2), and peroxynitrite (ONOO^-). In vascular SMCs, multiple enzymatic systems such as the NADPH oxidases (NOXs), xanthine oxidase (XO), nitric oxide synthases (NOS), and the mitochondrial electron transport chain are known to produce $\text{O}_2^{\cdot-}$ and H_2O_2 (Taniyama et al., 2004; Byon et al., 2016). The NOXs, in particular NOX1 and NOX4, are the most important because they are commonly expressed in vascular cells and are the major source of ROS generation in vessels (Clemens and Griendling, 2006; Konior et al., 2014; Burtenshaw et al., 2017). $\text{O}_2^{\cdot-}$ is converted to H_2O_2 by superoxide dismutases (SODs) or reacts with nitric oxide (NO) to form ONOO^- . H_2O_2 is further reduced to H_2O by catalase (CAT) and glutathione peroxidase (GPx; Taniyama and Griendling, 2003). Oxidative stress due to ROS production outweighing their scavenging is implicated in vascular dysfunction associated with T1DM and T2DM. It is well documented that elevated glucose increases the production of intracellular advanced glycation end-products (AGEs), stimulates the protein kinase C (PKC)-dependent activation of NOX1 and NOX4 (Inoguchi et al., 2000; Lu et al., 2006; Deluyker et al., 2017), and reduces the activity and bioavailability of antioxidant enzymes, such as SODs, GSH, CAT, and GPx, which results in higher ROS levels in both vascular ECs and SMCs in DM (Szaleczky et al., 1999; Lu et al., 2012; Tiwari et al., 2013).

Reactive oxygen species triggers many signaling pathways and promotes redox-mediated protein posttranslational modification. We found that redox modification is involved in BK channel dysfunction through hyperglycemia. High glucose culture of HEK293 cells stably expressing BK- α resulted in altered BK- α activity and channel kinetics that were mimicked by the effects of exogenously applied H_2O_2 in BK- α expressing cells cultured in normal glucose (Lu et al., 2006). A 1-week culture with 22 mM glucose markedly downregulated the protein expression of CAT and CuZn-SOD in HEK293 cells, leading to a 3.3-fold increase of H_2O_2 concentration to the 10^{-3} M range. Consequently, high glucose culture produced a 50% reduction of BK- α current density, prolonged the channel activation and deactivation time constants (τ_A and τ_D), and upward shifted the τ -V curve, indicating that BK- α activation is suppressed in high glucose conditions (Lu et al., 2006). The effects of high glucose on BK- α voltage-dependent activation were mimicked by acute exposure to 2 mM H_2O_2 . Furthermore, the cysteine residue at 911 (C911) in BK- α is particularly vulnerable to H_2O_2 -mediated regulation (Tang et al., 2001), and a single substitution of C911 by alanine (C911A) eliminated


most of the inhibitory effects of BK- α under high glucose conditions and to exogenously applied H₂O₂ (Lu et al., 2006). In addition, acute exposure to ONOO⁻ (5–100 μ M) significantly suppressed BK channel activity in vascular SMCs (Brzezinska et al., 2000; Liu et al., 2002), but did not alter BK- α voltage-dependent activation (Lu et al., 2006), suggesting that the molecular mechanisms underlying BK channel regulation by H₂O₂ and ONOO⁻ are different. Further studies revealed a 3- to 4-fold increase of 3-nitrotyrosine levels on BK- α protein in freshly isolated aortas from STZ-induced T1DM rats compared to non-diabetic controls, suggesting that ONOO⁻-induced modification of BK- α may be mediated through protein tyrosine nitration rather than protein oxidation (Lu et al., 2010). The precise amino acid residue(s) in BK- α modified by ONOO⁻ has not been identified. Nevertheless, an increase of ROS accumulation is the culprit for the development of BK channel dysfunction in DM.

Angiotensin II Signaling and Vascular BK Channel Regulation

Angiotensin II (Ang II) is an oligopeptide hormone, exerting its physiological and pathophysiological effects through binding to Ang II type 1 (AT1R) and type 2 (AT2R) receptors and activating their downstream signaling pathways (Dasgupta and Zhang, 2011). In vascular SMCs, where AT1R is predominantly expressed, Ang II causes vasoconstriction and promotes vascular wall remodeling (Ribeiro-Oliveira et al., 2008). In contrast, activation of AT2R produces vasodilatation and impairs vascular remodeling, effects opposite to those of AT1R (Danyel et al., 2013). AT1R is a G-protein-coupled receptor, which is coupled

to G α q, G β γ , G α i, and β -arrestin (Kawai et al., 2017; Wang et al., 2018). Binding of Ang II to AT1R in vascular SMCs activates G α q which in turn activates the phospholipase C (PLC)-dependent inositol-1,4,5-triphosphate (IP₃)/diacylglycerol (DAG)-mediated Ca²⁺ signaling cascades, causing an increase in protein kinase C (PKC) activity (De Gasparo et al., 2000; Touyz and Schiffrin, 2000). Activation of PKC β stimulates NOXs with ROS overproduction under hyperglycemic conditions (Inoguchi et al., 2000; Evcimen and King, 2007) and is a cause of impaired vascular BK channel function in diabetic vessels (Figure 3; Zhou et al., 2006; Lu et al., 2012; Zhang et al., 2020). In addition to redox-mediated modification of BK- α , it has been shown that PKC-induced serine phosphorylation at 695 (S695) and 1151 (S1151) in the C-terminus of BK- α inhibits BK channel current density by 50%, and S1151 phosphorylation by PKC also abolishes BK- α activation by protein kinase A (PKA) and protein kinase G (PKG; Zhou et al., 2001, 2010). On the other hand, the activity of tyrosine-protein kinase is regulated by G α i and β -arrestin upon AT1R stimulation, causing BK channel dysfunction (Ma et al., 2000; Alioua et al., 2002; Fessart et al., 2005; Tian et al., 2007). Another study reported that the C-terminus of AT1R physically interacts with the C-terminus of BK- α in heterologous expression system, and such protein–protein interaction between AT1R and BK- α directly inhibits BK- α activity, independent of G-protein mediated processes (Zhang et al., 2014).

However, AT1R expression, Ang II bioavailability, and tissue sensitivity to Ang II are upregulated in diabetic vessels (Arun et al., 2004; Kawai et al., 2017). The pathophysiological importance of Ang II-mediated BK channel regulation in diabetic coronary



arteries is supported by the evidence that cardiac infarct size induced by experimental ischemia/reperfusion in STZ-induced T1DM mice was twice as large as non-diabetic mice (Lu et al., 2016). The effects of DM on myocardial ischemia/reperfusion injury can be reproduced by infusion of 2 μ M Ang II or 0.1 μ M membrane impermeable BK channel inhibitor, IBTX, but attenuated by the BK channel activator, NS-1619 (Lu et al., 2016). Similar results were observed in Akita T1DM mice with exacerbated cardiovascular complications and cardiac and vascular dysfunction, from an imbalance of Ang II/AT1R signaling in DM (Patel et al., 2012). Most importantly, the pathological roles of Ang II signaling are supported by clinical outcomes showing that treatment with AT1R blockers and ACE inhibitors reduced cardiovascular complications and cardiovascular death in patients with DM by 25–30% (Niklason et al., 2004; Abuissa et al., 2005; Cheng et al., 2014; Lv et al., 2018).

Caveolae Compartmentation and Vascular BK Channel Subcellular Distribution

Caveolae, which are nonclathrin-coated, flask-shaped invaginations of plasma membrane lipid raft subdomains, are characterized by their signature structural protein caveolin, with caveolin-1 (Cav-1) predominantly expressed in the vasculature (Gratton et al., 2004; Krajewska and Maslowska, 2004). Caveolae have emerged as a central platform for signal transduction in many tissues through the interaction between the Cav scaffolding domain and protein partners that contain a Cav-binding motif (Φ x Φ xxxx Φ or Φ xxxx Φ xx Φ , where Φ is an aromatic amino acid, and x is any amino acid; Okamoto et al., 1998). Many signaling molecules that are associated with BK channel regulation, such as the β -adrenergic receptors (Bucci et al., 2004), AT1R (Ushio-Fukai and Alexander, 2006; Basset et al., 2009), NOX1 (Hilenski et al., 2004; Wolin, 2004), cellular tyrosin protein kinase Src (c-Src; Zundel et al., 2000; Lee et al., 2001), guanylyl cyclase (Linder et al., 2005; Vellecco et al., 2016), PKA (Heijnen et al., 2004; Linder et al., 2005), protein kinase B (PKB or AKT; Sedding et al., 2005), PKC (Zeydanli et al., 2011; Ringvold and Khalil, 2017), PKG (Linder et al., 2005), NOS (Garcia-Cardena et al., 1996; Vellecco et al., 2016), and prostacyclin (PGI₂) synthase (PGIS; Spisni et al., 2001), are found in the low buoyant density, caveolae-rich membrane fractions of vascular ECs and SMCs. The significance of Cav-1 on vascular physiology is demonstrated by findings in Cav-1 knockout (KO) mice that show constitutively activated eNOS with elevated NO production as well as a failure to maintain a constant vasocontractile tone, resulting in the development of cardiovascular pathologies (Drab et al., 2001; Razani et al., 2001). Overgeneration of NO facilitates the production of ONOO⁻ and contributes to vascular dysfunction with excessive H₂O₂ accumulation (Pacher et al., 2007).

The consensus sequence of the Cav-binding motif is present in BK- α , but not in BK- β 1. Indeed, only BK- α but not BK- β 1 is detected in the caveolae-rich fractions of SMCs (Lu et al., 2016). Moreover, BK- α is colocalized in the caveolae with other ion channels (Wang et al., 2005; Riddle et al., 2011; Howitt et al., 2012; Lu et al., 2016), especially those associated with Ca²⁺ spark/sparklet generation, such as L-type Ca²⁺ channels

(Suzuki et al., 2013; Saeki et al., 2019), T-type Ca²⁺ channels (Hashad et al., 2018), TRPV4 (Goedicke-Fritz et al., 2015; Lu et al., 2017b), TRPC1, TRPC3, and TRPC6 (Bergdahl et al., 2003; Adebisi et al., 2011; Grayson et al., 2017) in vascular ECs and SMCs. The close proximity of BK channels with Ca²⁺ entry molecules leads to Ca²⁺ spark-coupled STOCs. However, it has been reported that Cav-1 interacts with BK channels and inhibits BK channel activities in coronary ECs (Wang et al., 2005; Riddle et al., 2011). Cholesterol depletion by methyl- β -cyclodextrin and silencing of Cav-1 by small interference RNA enhance BK currents, while exposure to the scaffolding domain peptide of Cav-1 (AP-CAV) inhibits BK currents (Wang et al., 2005; Riddle et al., 2011). Hence, the presence of caveolae may exert an inhibitory effect on BK channel activity.

Increased Cav-1 expression has been found in most diabetic vessels (Hillman et al., 2001; Bucci et al., 2004; Pascariu et al., 2004; Elcioglu et al., 2010; Uyy et al., 2010; Li et al., 2014). Cav-1 expression is directly upregulated by the Forkhead Box O (FOXO) transcription factor (Sandri et al., 2004; Van Den Heuvel et al., 2005). The FOXO-3a phosphorylation levels are significantly reduced in STZ-induced T1DM rat arteries and in cultured human coronary arterial SMCs (Zhang et al., 2010a). This explains the underlying mechanism that leads to Cav-1 upregulation in DM (Figure 3). Furthermore, in STZ-induced T1DM rats, our results in co-immunoprecipitation experiments show that AT1R, c-Src, and BK- α are enriched in the low buoyant density, caveolae-rich membrane fractions of aortas, compared to non-diabetic rats (Lu et al., 2010). Infusion with Ang II (0.05 μ g/kg) results in markedly enhanced AT1R protein translocation to the low buoyant density fractions of aortas after 1h (83.4% of total membrane AT1R in STZ-induced T1DM rats vs. 28.5% in controls), suggesting enhanced AT1R translocation into caveolae-rich lipid rafts upon agonist activation in diabetic vessels, consistent with previous report in cultured vascular SMCs (Ishizaka et al., 1998). However, the precise mechanism underlying AT1R translocation is currently unclear. The levels of vascular BK- α protein oxidation, tyrosine phosphorylation, and tyrosine nitration are significantly increased in STZ-induced T1DM rats, likely due to the co-localization of NOS, NOX1 and c-Src in the caveolae. Since BK- α but not BK- β 1 is present in caveolae, BK- α translocation into the caveolae of arteries in STZ-induced T1DM mice may promote the physical dissociation of BK- α and BK- β 1 (Lu et al., 2016), which may explain the uncoupling of BK- α and BK- β 1 in diabetic vessels. A working framework has emerged in caveolae targeting of BK channel regulation, in which caveolae compartmentalize BK- α with AT1R, NOS, NOXs, and c-Src to form BK- α -receptor-enzyme microdomain complexes in vascular SMCs (Figure 3). Such caveolae compartmentation is enhanced in diabetic vessels, which facilitates the redox modification of BK- α . Of note, because BK- β 1 does not translocate into caveolae, such subcellular distribution of BK- α and BK- β 1 may contribute to BK- α and BK- β 1 functional uncoupling, thereby exacerbating BK channelopathy in diabetic vessels (Figure 3). Additionally, caveolae take part in endosomal trafficking and regulating surface expression of many membrane proteins (Elkin et al., 2016). Taking into account the consequences of upregulation

of caveolae formation in the vascular SMCs in DM, BK- α caveolae translocation may have important pathophysiological implications for vascular BK channel dysfunction in DM.

Ubiquitin Proteasome System and Vascular BK Channel Protein Degradation

Protein homeostasis with a balanced regulation between synthesis and degradation is essential for the maintenance of normal cellular function. Cellular proteins are degraded mainly through the lysosomes and the ubiquitin proteasome system (UPS; Ciechanover, 2005). Lysosomal protein degradation occurs through fusion with endocytotic vesicles. This mechanism of protein degradation is non-specific, and all proteins are digested indiscriminately at the same rate. UPS-mediated protein degradation accounts for 80–90% of protein degradation in mammalian cells and it is substrate-specific (Powell, 2006; Schapira et al., 2019). This process is facilitated by three distinct enzymatic steps that involve an ubiquitin-activating enzyme (E1), a ubiquitin-conjugating enzyme (E2), and a ubiquitin ligase (E3). E1 interacts with ubiquitin through an E1-ubiquitin thioester bond in an ATP-dependent manner. It transfers the activated ubiquitin molecule to a cysteine residue on the E2 enzyme to form an E2-ubiquitin thioester-linked intermediate. The E3 ligase facilitates transfer of the E2-ubiquitin moiety to the substrate protein *via* an amide bond between the carboxy terminus of ubiquitin and a lysine side chain of the substrate protein. The E3 ligase is substrate-specific, allowing repeated positioning of the distal end of ubiquitin molecule for ubiquitin chain assembly with high precision. The poly-ubiquitinated protein is then recognized for enzymatic degradation in the 26S proteasome (Powell, 2006; Schapira et al., 2019). Hence, the E3 reaction is critical for determining the turnover of specific proteins. There are 617 E3 ligases functionally annotated in the human genome (Li et al., 2008). It is known that F-box (FBXO) proteins are a key component of the Skp1-Cullin-F-box (SCF)^{FBXO}-type ubiquitin ligase complex (SCF^{FBXO}) and serve as sites for enzyme-substrate interaction (Kipreos and Pagano, 2000). FBXO proteins contain several functional domains such as the F-box domain, the LRRs, and the WD40 repeats for protein-protein interaction. Two muscle-specific FBXO proteins, FBXO-9 and FBXO-32 (also known as atrogen-1), have been found to be upregulated in diabetic vessels. They mediate BK- β 1 protein ubiquitination in coronary arterial SMCs (Zhang et al., 2010a). The molecular basis of FBXO-32 and BK- β 1 interaction was identified using site-directed mutagenesis and co-immunoprecipitation approaches, which showed that the PDZ-binding motif (ETSV) on BK- β 1 is critical for FBXO-32-dependent ubiquitination (Zhang et al., 2010a). Deletion of the consensus sequence of the PDZ-binding motif in BK- β 1 significantly decreases BK- β 1 protein ubiquitination (Figure 4; Zhang et al., 2010a). Activation of FBXO proteins reduces BK- β 1 expression, while knockdown of *FBXO* and proteasomal inhibition enhances BK- β 1 levels, suggesting that accelerated UPS-mediated degradation of BK- β 1 is an important mechanism of BK channel regulation in DM.

The muscle RING-finger protein 1 (MuRF1) is another E3 ligase involved in UPS-dependent vascular BK- β 1 degradation

(Yi et al., 2014). Nuclear factor- κ B (NF- κ B) sites in the MuRF1 promoter are required for transcriptional activation, while FOXO sites are not (Wu et al., 2014). Overexpression of MuRF1 downregulates BK- β 1 expression, impairs BK- β 1-mediated BK channel activity, and reduces BK channel-induced vasodilation in mouse coronary arteries. We found that the N-terminus of BK- β 1 and the coiled-coil region of MuRF1 are necessary for BK- β 1 and MuRF1 interaction (Yi et al., 2014). Importantly, the protein expressions of FBXO-9, FBXO-32, and MuRF1 are unregulated in the arteries of STZ-induced T1DM animals and in primary human coronary arterial SMCs cultured with high glucose (Zhang et al., 2010a, 2020; Lu et al., 2012; Yi et al., 2014). Such upregulation of FBXO expression is mediated through the suppression of PI3K/AKT-dependent phosphorylation in FOXO-3a, thereby promoting FOXO-3a nuclear translocation and binding to the consensus sequence [GTAAA(C/T)A] in the promoter of *Fbxo* gene, activating its transcription (Furuyama et al., 2000). However, activation of MuRF1 is due to an increase of NF- κ B-mediated *Trim63* (encoding MuRF1) transcription (Wu et al., 2014). In DM or hyperglycemia, the activity of AKT is reduced (Okon et al., 2005), while that of NF- κ B is augmented (Narayanan et al., 2014), thereby promoting FBXO and MuRF1 expression (Figure 4). Indeed, inhibition of PKC β activity by ruboxistaurin, NF- κ B activity by TPCA-1, and proteasomal activity by MG132 downregulates BK- β 1 ubiquitination, preserves BK- β 1 expression, and improves BK channel function in coronary arterial SMCs (Zhang et al., 2010a; Lu et al., 2012; Yi et al., 2014).

BK- α protein expression is also regulated by lysosome and UPS degradation (Wang et al., 2013; Liu et al., 2014; Leo et al., 2015; Song et al., 2018). It has been found that the CRL4A and its substrate cereblon (CRBN) complex (CRL4A^{CRBN}) serves as the ubiquitin ligase that interacts with the C-terminus of BK- α and induces BK- α protein degradation in neurons (Liu et al., 2014). A recent study reported that both CRBN and BK- α proteins were targeted by SCF^{FBXO-7} ubiquitin ligase complex for ubiquitination and proteolysis, controlling BK- α function and regulating the learning and memory processes in the brain (Song et al., 2018). However, the specific E3 ligase(s) responsible for BK- α protein ubiquitination in blood vessels is unknown, and how the BK- α -specific E3s are regulated in DM remains to be determined.

Effects of Nuclear Factor Erythroid-2-Related Factor 2 Signaling on Vascular BK Channel Expression

Nuclear factor erythroid-2-related factor 2 (Nrf2) plays a critical role in the maintenance of intracellular redox homeostasis by regulating multiple downstream antioxidant enzymes and phase II detoxifying enzymes, which include NADPH dehydrogenase quinone 1 (NQO1), glutathione-disulfide reductase (GSR), glutathione translocase (GSTA), thioredoxin (TXN), thioredoxin reductase 1 (TXNRD1), heme oxygenase-1 (HO-1), SODs, CAT, and GPx (Gao and Mann, 2009; Chen et al., 2014). In addition, Nrf2 negatively regulates the expression of NOXs (McSweeney et al., 2016). The function of Nrf2 is principally regulated by the kelch-like ECH-association protein 1 (Keap1), which mediates




FIGURE 4 | Regulation of BK channel expression by ubiquitin proteasome system (UPS) and nuclear factor erythroid-2-related factor 2 (Nrf2) signaling. FBXO and MuRF1 are the E3 ligases targeting BK-β1 protein degradation via the UPS in vascular SMCs. FBXO is one of downstream targets of FOXO-3a. FOXO-3a activity is negatively controlled by AKT-dependent phosphorylation, while MuRF1 expression is controlled by NF-κB/p65. Under baseline conditions, p65 is bound to an inhibitory subunit, IκB that keeps it sequestered in an inactive state in the cytoplasm. Phosphorylation of IκB by IκB kinase promotes IκB degradation through the UPS, which in turn releases p65 and facilitates nuclear translocation. Under hyperglycemic conditions, overproduction of ROS inhibits AKT and activates NF-κB/p65, which in turn promotes FBXO and MuRF1 expression, leading to BK-β1 ubiquitination and accelerated degradation in vascular SMCs. Nrf2 is the master regulator of the antioxidant response. Under normal conditions, each molecule of Nrf2 interacts with two molecules of Keap1 resulting in UPS-mediated degradation. ROS modifies specific cysteine residues in Keap1 and releases Nrf2 from binding with Keap1. The unbound Nrf2 translocates into the nucleus and binds to the promoter region of target genes. Nrf2 directly upregulates BK-α mRNA expression via binding to the promoter region of *KCNMA1*. However, BK-β1 mRNA expression is not regulated by Nrf2 but by other transcription factor(s). In DM, Nrf2 expression and function is significantly downregulated, leading to a decrease in BK-α expression through reduced transcription and a decrease in BK-β1 expression through accelerated UPS degradation. The symbols “u” and “p” represent protein ubiquitination and phosphorylation, respectively.

Nrf2 ubiquitination and subsequent proteasomal degradation (Canning et al., 2015; Suzuki and Yamamoto, 2015). In the nuclei, Nrf2 binds to the promoters of antioxidant response elements (AREs) and electrophile response elements (EpREs) through interaction with the Nrf2-binding motif [TGA(G/C)xxxGC], where x represents any amino acid (Chorley et al., 2012). Both the *KCNMA1* and *KCNMB1* genes contain the consensus sequences of Nrf2-binding motifs in their promoter regions. Using promoter luciferase reporter assays, we confirmed that Nrf2 binds to the ARE of the *KCNMA1* promoter, but not to that of *KCNMB1* promoter. Mutation of the Nrf2-binding motif in the *KCNMA1* promoter abolishes the transcription response to Nrf2 (Sun et al., 2020). In addition, adenoviral expression of Nrf2 significantly augmented the mRNA levels of BK-α and BK-β1 in coronary arterial SMCs (Lu et al., 2017a; Sun et al., 2020). These results suggest that Nrf2 facilitates BK-α mRNA expression through activation of *KCNMA1* transcription, whereas the stimulatory effect of Nrf2 on BK-β1 mRNA expression is indirect and may be achieved by activating other transcription factor(s) or signaling mechanisms that upregulate *KCNMB1* transcription and expression in vascular SMCs.

Nuclear factor erythroid-2-related factor 2 deficiency has been implicated in diabetic complications including those associated with the heart (Tan et al., 2011; Bai et al., 2013),


blood vessels (Ungvari et al., 2011; Miao et al., 2012; Li et al., 2017; Lu et al., 2017a), kidneys (Zheng et al., 2011; Cui et al., 2012), and the brain (Pu et al., 2018; Tarantini et al., 2018). The expression of Nrf2 and its downstream genes is slightly increased in the cardiovascular systems of STZ-induced T1DM mice at 2–3 months after the onset of hyperglycemia, but then becomes significantly downregulated at 5–6 months after the development of hyperglycemia (Tan et al., 2011; Miao et al., 2012; Bai et al., 2013), suggesting the burnout of an important redox protective mechanism in the advanced stages of DM. In db/db and HFD-induced diabetic mice 6 months after the development of hyperglycemia, BK channel activity and BK channel-mediated vasodilation in coronary arteries are impaired, accompanied by a remarkable reduction in Nrf2 and its associated antioxidant enzymes (Li et al., 2017; Lu et al., 2017a). Nrf2 KO mice show excessive ROS production, as well as diminished BK channel expression and function in vascular SMCs (Ashino et al., 2013; Sun et al., 2020). Both mRNA and protein expression of BK-α are downregulated, whereas BK-β1 proteins but not mRNA levels are decreased in the arterial SMCs of Nrf2 KO mice, consistent with the notion that Nrf2 regulates BK-α via transcription, and BK-β1 through posttranscriptional mechanisms (Figure 4; Sun et al., 2020). Administration of dimethyl formamide (DMF, an FDA-approved Nrf2 activator) preserves

BK channel protein expression, BK channel activity, and BK channel-mediated vasodilation in the coronary arteries of db/db and HDF-induced diabetic mice (Li et al., 2017; Lu et al., 2017a). Currently, Nrf2 activators such as DMF and sulforaphane (SFN) are being used in clinical trials for cardiovascular diseases and metabolic disorder (Yagishita et al., 2020), but it has not been administered for diabetic patients with coronary heart disease (Houghton, 2019). Whether the beneficial effects of Nrf2 activators observed in animal studies would translate into better outcomes in diabetic patients with cardiovascular complications needs to be determined.

Effects of Calcineurin-Nuclear Factor of Activated T Cells Cytoplasmic 3 Isoform Pathway on BK- β 1 Transcription

Nuclear factor of activated T cells cytoplasmic 3 isoform (NFATc3) belongs to the nuclear factor of activated T cells (NFAT) family of transcription factors that were originally discovered in resting T cells and is important in immune response (Rao et al., 1997). NFATc3 is also involved in the development of skeletal muscle and of the cardiovascular systems (Crabtree and Olson, 2002). The activity of NFATc3 is modulated by the Ca^{2+} /calmodulin-dependent phosphatase, calcineurin. Elevation of the intracellular Ca^{2+} concentration activates calmodulin and promotes its binding to calcineurin, leading to calcineurin activation. Activated calcineurin dephosphorylates NFATc3, which in turn induces NFATc3 nuclear translocation. Calcineurin binds to the scaffolding protein A-kinase anchoring protein 150 (AKAP150), corresponding to AKAP79 in humans, which also anchors PKA and L-type Ca^{2+} channel to form a dynamic Ca^{2+} signaling complex (Oliveria et al., 2007). AKAP79/150 strongly suppresses PKA-mediated L-type Ca^{2+} channel phosphorylation and is required for the activation of NFAT by local Ca^{2+} influx through L-type channels (Oliveria et al., 2007).

Nuclear factor of activated T cells share a conserved DNA-binding domain that specifically binds to the DNA core sequence [(A/T)GGAAA] at the promoter region of target genes, activating gene transcription (Rao et al., 1997). Human and mouse *KCNMA1* and *KCNMB1* contain at least one NFAT-binding motif in their promoters. Inhibition of vascular BK channels by NFATc3 has been reported, while upregulation of NFATc3 expression by Ang II results in decreased BK channel activity in mouse arteries due to the downregulation of BK- β 1 mRNA expression (Nieves-Cintrón et al., 2007). The effects of NFATc3 on BK channel activity and BK- β 1 mRNA expression are abolished by calcineurin inhibitors, FK506 and cyclosporin A, in the presence of Ang II, a finding that has been confirmed in NFATc3 KO mice (Nieves-Cintrón et al., 2007). AKAP150 also participates in NFATc3-mediated BK channel downregulation in HFD-induced diabetic mice (Figure 5; Nystoriak et al., 2014). In HFD-induced diabetic mice, the activity of the AKAP150-NFATc3 signaling pathway is upregulated, contributing to impaired BK channel function with reduced BK- β 1 expression and increased vascular tone in the mesenteric arteries. However, in AKAP150 KO mice with HFD consumption, the deleterious effects of HFD on BK channels are not observed



(Nystoriak et al., 2014). Recently, *in vivo* administration of a NFATc3 inhibitor (A285222, Abbott Labs) in Akita T1DM mice is found to improve vascular endothelial function, enhance eNOS activity and NO production, reduce endothelin-1 secretion, lower blood pressure, and improve survival (Garcia-Vaz et al., 2020). The beneficial effects of NFATc3 inhibitors on coronary BK channel function in DM warrant further investigation.

Arachidonic Acid and Its Metabolites on BK Channel Regulation

Arachidonic acid (AA), a polyunsaturated omega-6 fatty acid, is abundant in normal human diet and in membrane phospholipids. It is an important precursor to a wide range of bioactive mediators and eicosanoids that regulate a multitude of essential functions in the body (Tallima and El Ridi, 2018). AA is metabolized by three major enzyme systems: It is converted by 12-lipoxygenase (12-LOX) into leukotrienes and 12-hydroxyeicosatetraenoic acid (12-HETE), by cytochrome P-450 (CYP-450) epoxygenase into epoxyeicosatrienoic acids (EETs), and by cyclooxygenases (COX) into prostaglandins, including PGI_2 and thromboxane A2 (TXA_2 ; Brash, 2001; Vila, 2004). Additionally, AA can be metabolized by CYP-450 omega-hydroxylase to produce 20-hydroxyeicosatetraenoic acid (20-HETE).

Arachidonic acid (Lu et al., 2005; Kur et al., 2014; Martin et al., 2014, 2021) and its metabolites (EETs, PGI_2 , 12-HETE, and 20-HETE; Li and Campbell, 1997; Yamaki et al., 2001; Zhang et al., 2001; Zink et al., 2001; Lauterbach et al., 2002; Morin et al., 2007) are known to activate vascular BK channels and promote vasodilation through endothelium-dependent

hyperpolarization mechanisms. Direct exposure to 10 μ M AA robustly increases BK channel activity in inside-out excised patches from human umbilical arterial SMCs, suggesting activation of BK channels directly by AA (Martin et al., 2021). Extracellular application of AA results in BK channel activation and hyperpolarization of resting membrane potentials in vascular SMCs (Kur et al., 2014; Martin et al., 2021). These changes can be blocked by LOX, CYP, and COX inhibitors, suggesting that AA metabolites affect BK channels. The effects of AA on BK channels require the presence of BK- β 1 (Sun et al., 2007; Martin et al., 2021).

The activation of vascular BK channels by PGI₂ is associated with cAMP-dependent, PKA-mediated phosphorylation. EETs and their metabolites dihydroxyicosatrienoic acids (DHETs) are also potent BK channel activators and vasodilators, including the human coronary microvessels and internal mammary arteries (Quilley et al., 1997; Archer et al., 2003; Feletou and Vanhoutte, 2006; Larsen et al., 2006). Several different mechanisms of EET- and DHET-mediated BK channel activation have been proposed, including direct activation (Wu et al., 2000; Lu et al., 2001), ADP-ribosylation of G α (Fukao et al., 2001; Li et al., 2002), and stimulation of PKA-mediated phosphorylation (Dimitropoulou et al., 2007; Imig et al., 2008). However, AA-induced vasodilation of coronary arterioles *via* BK channel activity is impaired in high glucose conditions and DM (Lu et al., 2005; Zhou et al., 2005, 2006; Yousif and Benter, 2007; Tsai et al., 2011). PGI₂ and EET levels are decreased in patients with cardiovascular diseases (Theken et al., 2012; Mokhtar et al., 2013; Schuck et al., 2013) and DM (Lane et al., 1982; Kazama et al., 1987; Migdalis et al., 2001; Dufлот et al., 2019). As a result of these findings, AA metabolites and analogues have been developed as potential therapeutic agents for cardiovascular diseases and diabetic vascular complications (Campbell et al., 2017; Wang et al., 2021).

FUTURE DIRECTIONS IN DIABETIC BK CHANNEL RESEARCH

Studies of the regulation of BK channel function and expression have greatly advanced our understanding on the role of BK channels in diabetic cardiovascular complications. DM involves a plethora of signaling abnormalities including those pertaining to insulin, ROS generation, Ang II signaling, and Ca²⁺ regulation.

REFERENCES

- Abuissa, H., Jones, P. G., Marso, S. P., and O'keefe, J. H. (2005). Angiotensin-converting enzyme inhibitors or angiotensin receptor blockers for prevention of type 2 diabetes: a meta-analysis of randomized clinical trials. *J. Am. Coll. Cardiol.* 46, 821–826. doi: 10.1016/j.jacc.2005.05.051
- Adebisi, A., Narayanan, D., and Jaggar, J. H. (2011). Caveolin-1 assembles type 1 inositol 1,4,5-trisphosphate receptors and canonical transient receptor potential 3 channels into a functional signaling complex in arterial smooth muscle cells. *J. Biol. Chem.* 286, 4341–4348. doi: 10.1074/jbc.M110.179747
- Alioua, A., Mahajan, A., Nishimaru, K., Zarei, M. M., and Stefani, E. (2002). Coupling of c-Src to large conductance voltage- and Ca²⁺-activated K⁺ channels as a new mechanism of agonist-induced vasoconstriction. *Proc. Natl. Acad. Sci. U. S. A.* 99, 14560–14565. doi: 10.1073/pnas.222348099

Thus, it is not surprising that DM affects vascular BK channel expression and function in many different ways, including transcription, translation, post-translation, surface trafficking, and channel degradation. Whether surface trafficking dysregulation of BK channel subunits contributes to BK channelopathy of the vascular SMCs in DM is unknown. Moreover, BK channels do not exist as isolated proteins but are assembled in membrane microdomains of vascular ECs and SMCs. Studies of BK channel organization by scaffolding proteins in close proximity with receptors, enzymes, and Ca²⁺ sources in blood vessels will provide further insights into BK channel physiology and into the molecular mechanisms underlying BK channelopathy in DM. In addition, our knowledge on BK- γ 1 in diabetic BK channel dysregulation is very limited. Little is known about the regulation of vascular BK- γ 1 expression and function in hyperglycemia and DM. Since the results of BK channel pathology from diabetic animal models are diverse, it is critical to study vascular BK channel biology and dysfunction using human tissues, which serve as the gold standard for diabetic BK channel research.

Ca²⁺-activated K⁺ channels are important regulators of vascular physiology and are critical determinants coronary circulation and cardioprotection. Preservation of BK channel expression and activities protects vascular function in DM. Hence, a better understanding of BK channelopathy and prevention of BK channel abnormalities in DM may lead to better vascular therapeutics and care for patients with DM.

AUTHOR CONTRIBUTIONS

TL and HL wrote the manuscript and critically reviewed the final version of the manuscript. All authors contributed to the article and approved the submitted version.

FUNDING

This work was supported by grants from the National Institute of Health (RO1 HL-080118 and RO1 HL-074180), American Diabetes Association (ADA-JFA-07-39, ADA 1-12-BS-119, ADA 1-16-IBS-195, and ADA 1-18-IBS-210), and the Mayo Clinic.

- Archer, S. L., Gragasin, F. S., Wu, X., Wang, S., and Mcmurtry, S. (2003). Endothelium-derived hyperpolarizing factor in human internal mammary artery is 11,12-epoxyicosatrienoic acid and causes relaxation by activating smooth muscle BK(Ca) channels. *Circulation* 107, 769–776. doi: 10.1161/01.CIR.0000047278.28407.C2
- Arun, K. H., Kaul, C. L., and Ramarao, P. (2004). High glucose concentration augments angiotensin II mediated contraction via AT1 receptors in rat thoracic aorta. *Pharmacol. Res.* 50, 561–568. doi: 10.1016/j.phrs.2004.06.001
- Ashino, T., Yamamoto, M., Yoshida, T., and Numazawa, S. (2013). Redox-sensitive transcription factor Nrf2 regulates vascular smooth muscle cell migration and neointimal hyperplasia. *Arterioscler. Thromb. Vasc. Biol.* 33, 760–768. doi: 10.1161/ATVBAHA.112.300614
- Bai, Y., Cui, W., Xin, Y., Miao, X., and Barati, M. T. (2013). Prevention by sulforaphane of diabetic cardiomyopathy is associated with up-regulation

- of Nrf2 expression and transcription activation. *J. Mol. Cell. Cardiol.* 57, 82–95. doi: 10.1016/j.yjmc.2013.01.008
- Bao, L., Kaldany, C., Holmstrand, E. C., and Cox, D. H. (2004). Mapping the BKCa channel's "Ca²⁺ bowl": side-chains essential for Ca²⁺ sensing. *J. Gen. Physiol.* 123, 475–489. doi: 10.1085/jgp.200409052
- Basset, O., Deffert, C., Foti, M., Bedard, K., and Jaquet, V. (2009). NADPH oxidase 1 deficiency alters caveolin phosphorylation and angiotensin II-receptor localization in vascular smooth muscle. *Antioxid. Redox Signal.* 11, 2371–2384. doi: 10.1089/ars.2009.2584
- Beckman, J. A., and Creager, M. A. (2016). Vascular complications of diabetes. *Circ. Res.* 118, 1771–1785. doi: 10.1161/CIRCRESAHA.115.306884
- Bergdahl, A., Gomez, M. F., Dreja, K., Xu, S. Z., and Adner, M. (2003). Cholesterol depletion impairs vascular reactivity to endothelin-1 by reducing store-operated Ca²⁺ entry dependent on TRPC1. *Circ. Res.* 93, 839–847. doi: 10.1161/01.RES.0000100367.45446.A3
- Bluestone, J. A., Herold, K., and Eisenbarth, G. (2010). Genetics, pathogenesis and clinical interventions in type 1 diabetes. *Nature* 464, 1293–1300. doi: 10.1038/nature08933
- Borbouse, L., Dick, G. M., Asano, S., Bender, S. B., and Dincer, U. D. (2009). Impaired function of coronary BK(Ca) channels in metabolic syndrome. *Am. J. Physiol. Heart Circ. Physiol.* 297, H1629–H1637. doi: 10.1152/ajpheart.00466.2009
- Brash, A. R. (2001). Arachidonic acid as a bioactive molecule. *J. Clin. Invest.* 107, 1339–1345. doi: 10.1172/JCI13210
- Brindisi, M. C., Bouillet, B., Verges, B., and Halimi, S. (2010). Cardiovascular complications in type 1 diabetes mellitus. *Diabetes Metab.* 36, 341–344. doi: 10.1016/j.diabet.2010.06.002
- Bryan, R. M., You, J., Golding, E. M., and Marrelli, S. P. (2005). Endothelium-derived hyperpolarizing factor: a cousin to nitric oxide and prostacyclin. *Anesthesiology* 102, 1261–1277. doi: 10.1097/00000542-200506000-00028
- Brzezinska, A. K., Gebremedhin, D., Chilian, W. M., Kalyanaraman, B., and Elliott, S. J. (2000). Peroxynitrite reversibly inhibits Ca(2+)-activated K(+) channels in rat cerebral artery smooth muscle cells. *Am. J. Physiol. Heart Circ. Physiol.* 278, H1883–H1890. doi: 10.1152/ajpheart.2000.278.6.H1883
- Bucci, M., Rovietto, F., Brancaleone, V., Lin, M. I., and Di Lorenzo, A. (2004). Diabetic mouse angiopathy is linked to progressive sympathetic receptor deletion coupled to an enhanced caveolin-1 expression. *Arterioscler. Thromb. Vasc. Biol.* 24, 721–726. doi: 10.1161/01.ATV.0000122362.44628.09
- Burnham, M. P., Johnson, I. T., and Weston, A. H. (2006). Reduced Ca²⁺-dependent activation of large-conductance Ca²⁺-activated K⁺ channels from arteries of type 2 diabetic Zucker diabetic fatty rats. *Am. J. Physiol. Heart Circ. Physiol.* 290, H1520–H1527. doi: 10.1152/ajpheart.00827.2005
- Burtenshaw, D., Hakimjavadi, R., Redmond, E. M., and Cahill, P. A. (2017). Nox, reactive oxygen species and regulation of vascular cell fate. *Antioxidants* 6:90. doi: 10.3390/antiox6040090
- Byon, C. H., Heath, J. M., and Chen, Y. (2016). Redox signaling in cardiovascular pathophysiology: a focus on hydrogen peroxide and vascular smooth muscle cells. *Redox Biol.* 9, 244–253. doi: 10.1016/j.redox.2016.08.015
- Campbell, W. B., Imig, J. D., Schmitz, J. M., and Falck, J. R. (2017). Orally active epoxyeicosatrienoic acid analogs. *J. Cardiovasc. Pharmacol.* 70, 211–224. doi: 10.1097/FJC.0000000000000523
- Canning, P., Sorrell, F. J., and Bullock, A. N. (2015). Structural basis of Keap1 interactions with Nrf2. *Free Radic. Biol. Med.* 88, 101–107. doi: 10.1016/j.freeradbiomed.2015.05.034
- Cefalu, W. T., Kaul, S., Gerstein, H. C., Holman, R. R., and Zinman, B. (2018). Cardiovascular outcomes trials in type 2 diabetes: where do we go from here? Reflections from a diabetes care editors' expert forum. *Diabetes Care* 41, 14–31. doi: 10.2337/dci17-0057
- Chen, J., Zhang, Z., and Cai, L. (2014). Diabetic cardiomyopathy and its prevention by nrf2: current status. *Diabetes Metab. J.* 38, 337–345. doi: 10.4093/dmj.2014.38.5.337
- Cheng, J., Zhang, W., Zhang, X., Han, F., and Li, X. (2014). Effect of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers on all-cause mortality, cardiovascular deaths, and cardiovascular events in patients with diabetes mellitus: a meta-analysis. *JAMA Intern. Med.* 174, 773–785. doi: 10.1001/jamainternmed.2014.348
- Chorley, B. N., Campbell, M. R., Wang, X., Karaca, M., and Sambandan, D. (2012). Identification of novel NRF2-regulated genes by ChIP-seq: influence on retinoid X receptor alpha. *Nucleic Acids Res.* 40, 7416–7429. doi: 10.1093/nar/gks409
- Christ, G. J., Day, N., Santizo, C., Sato, Y., and Zhao, W. (2004). Intracorporal injection of hSlo cDNA restores erectile capacity in STZ-diabetic F-344 rats in vivo. *Am. J. Physiol. Heart Circ. Physiol.* 287, H1544–H1553. doi: 10.1152/ajpheart.00792.2003
- Ciechanover, A. (2005). Intracellular protein degradation: from a vague idea thru the lysosome and the ubiquitin-proteasome system and onto human diseases and drug targeting. *Cell Death Differ.* 12, 1178–1190. doi: 10.1038/sj.cdd.4401692
- Clempus, R. E., and Griendling, K. K. (2006). Reactive oxygen species signaling in vascular smooth muscle cells. *Cardiovasc. Res.* 71, 216–225. doi: 10.1016/j.cardiores.2006.02.033
- Cox, D. H., and Aldrich, R. W. (2000). Role of the beta1 subunit in large-conductance Ca(2+)-activated K(+) channel gating energetics. Mechanisms of enhanced Ca(2+) sensitivity. *J. Gen. Physiol.* 116, 411–432. doi: 10.1085/jgp.116.3.411
- Cox, D. H., Cui, J., and Aldrich, R. W. (1997). Allosteric gating of a large conductance Ca-activated K+ channel. *J. Gen. Physiol.* 110, 257–281. doi: 10.1085/jgp.110.3.257
- Crabtree, G. R., and Olson, E. N. (2002). NFAT signaling: choreographing the social lives of cells. *Cell* 109, S67–S79. doi: 10.1016/s0092-8674(02)00699-2
- Creager, M. A., Luscher, T. F., Cosentino, F., and Beckman, J. A. (2003). Diabetes and vascular disease: pathophysiology, clinical consequences, and medical therapy: part I. *Circulation* 108, 1527–1532. doi: 10.1161/01.CIR.0000091257.27563.32
- Cui, W., Bai, Y., Miao, X., Luo, P., and Chen, Q. (2012). Prevention of diabetic nephropathy by sulforaphane: possible role of Nrf2 upregulation and activation. *Oxidative Med. Cell. Longev.* 2012:821936. doi: 10.1155/2012/821936
- Cui, J., Yang, H., and Lee, U. S. (2009). Molecular mechanisms of BK channel activation. *Cell. Mol. Life Sci.* 66, 852–875. doi: 10.1007/s00018-008-8609-x
- Danyel, L. A., Schmerler, P., Paulis, L., Unger, T., and Steckelings, U. M. (2013). Impact of AT2-receptor stimulation on vascular biology, kidney function, and blood pressure. *Integr. Blood Press. Control* 6, 153–161. doi: 10.2147/IBPC.S34425
- Dasgupta, C., and Zhang, L. (2011). Angiotensin II receptors and drug discovery in cardiovascular disease. *Drug Discov. Today* 16, 22–34. doi: 10.1016/j.drudis.2010.11.016
- Davies, K. P., Zhao, W., Tar, M., Figueroa, J. C., and Desai, P. (2007). Diabetes-induced changes in the alternative splicing of the slo gene in corporal tissue. *Eur. Urol.* 52, 1229–1237. doi: 10.1016/j.eururo.2006.11.028
- De Gasparo, M., Catt, K. J., Inagami, T., Wright, J. W., and Unger, T. (2000). International union of pharmacology. XXIII. The angiotensin II receptors. *Pharmacol. Rev.* 52, 415–472.
- Deluyker, D., Evens, L., and Bito, V. (2017). Advanced glycation end products (AGEs) and cardiovascular dysfunction: focus on high molecular weight AGEs. *Amino Acids* 49, 1535–1541. doi: 10.1007/s00726-017-2464-8
- Dhalla, N. S., Pierce, G. N., Innes, I. R., and Beamish, R. E. (1985). Pathogenesis of cardiac dysfunction in diabetes mellitus. *Can. J. Cardiol.* 1, 263–281.
- Dimitropoulou, C., Han, G., Miller, A. W., Molero, M., and Fuchs, L. C. (2002). Potassium (BK(Ca)) currents are reduced in microvascular smooth muscle cells from insulin-resistant rats. *Am. J. Physiol. Heart Circ. Physiol.* 282, H908–H917. doi: 10.1152/ajpheart.00382.2001
- Dimitropoulou, C., West, L., Field, M. B., White, R. E., and Reddy, L. M. (2007). Protein phosphatase 2A and Ca²⁺-activated K⁺ channels contribute to 11,12-epoxyeicosatrienoic acid analog mediated mesenteric arterial relaxation. *Prostaglandins Other Lipid Mediat.* 83, 50–61. doi: 10.1016/j.prostaglandins.2006.09.008
- Dong, L., Zheng, Y. M., Van Riper, D., Rathore, R., and Liu, Q. H. (2008). Functional and molecular evidence for impairment of calcium-activated potassium channels in type-1 diabetic cerebral artery smooth muscle cells. *J. Cereb. Blood Flow Metab.* 28, 377–386. doi: 10.1038/sj.cbfm.9600536
- Drab, M., Verkade, P., Elger, M., Kasper, M., and Lohn, M. (2001). Loss of caveolae, vascular dysfunction, and pulmonary defects in caveolin-1 gene-disrupted mice. *Science* 293, 2449–2452. doi: 10.1126/science.1062688
- Dufлот, T., Moreau-Grange, L., Roche, C., Jacob, M., and Wils, J. (2019). Altered bioavailability of epoxyeicosatrienoic acids is associated with conduit artery endothelial dysfunction in type 2 diabetic patients. *Cardiovasc. Diabetol.* 18:35. doi: 10.1186/s12933-019-0843-z
- Earley, S., Heppner, T. J., Nelson, M. T., and Brayden, J. E. (2005). TRPV4 forms a novel Ca²⁺ signaling complex with ryanodine receptors

- and BKCa channels. *Circ. Res.* 97, 1270–1279. doi: 10.1161/01.RES.0000194321.60300.d6
- Elicioglu, K. H., Kabasakal, L., Cetinel, S., Conturk, G., and Sezen, S. F. (2010). Changes in caveolin-1 expression and vasoreactivity in the aorta and corpus cavernosum of fructose and streptozotocin-induced diabetic rats. *Eur. J. Pharmacol.* 642, 113–120. doi: 10.1016/j.ejphar.2010.05.049
- Elkin, S. R., Lakoduk, A. M., and Schmid, S. L. (2016). Endocytic pathways and endosomal trafficking: a primer. *Wien. Med. Wochenschr.* 166, 196–204. doi: 10.1007/s10354-016-0432-7
- Erdos, B., Miller, A. W., and Busija, D. W. (2002). Alterations in KATP and KCa channel function in cerebral arteries of insulin-resistant rats. *Am. J. Physiol. Heart Circ. Physiol.* 283, H2472–H2477. doi: 10.1152/ajpheart.00516.2002
- Evanson, K. W., Bannister, J. P., Leo, M. D., and Jaggar, J. H. (2014). LRRC26 is a functional BK channel auxiliary gamma subunit in arterial smooth muscle cells. *Circ. Res.* 115, 423–431. doi: 10.1161/CIRCRESAHA.115.303407
- Evcimen, N. D., and King, G. L. (2007). The role of protein kinase C activation and the vascular complications of diabetes. *Pharmacol. Res.* 55, 498–510. doi: 10.1016/j.phrs.2007.04.016
- Feletou, M., and Vanhoutte, P. M. (2006). Endothelium-derived hyperpolarizing factor: where are we now? *Arterioscler. Thromb. Vasc. Biol.* 26, 1215–1225. doi: 10.1161/01.ATV.0000217611.81085.c5
- Fessart, D., Simaan, M., and Laporte, S. A. (2005). C-Src regulates clathrin adapter protein 2 interaction with beta-arrestin and the angiotensin II type 1 receptor during clathrin-mediated internalization. *Mol. Endocrinol.* 19, 491–503. doi: 10.1210/me.2004-0246
- Fukao, M., Mason, H. S., Kenyon, J. L., Horowitz, B., and Keef, K. D. (2001). Regulation of BK(Ca) channels expressed in human embryonic kidney 293 cells by epoxyeicosatrienoic acid. *Mol. Pharmacol.* 59, 16–23. doi: 10.1124/mol.59.1.16
- Furuyama, T., Nakazawa, T., Nakano, I., and Mori, N. (2000). Identification of the differential distribution patterns of mRNAs and consensus binding sequences for mouse DAF-16 homologues. *Biochem. J.* 349, 629–634. doi: 10.1042/bj3490629
- Gao, L., and Mann, G. E. (2009). Vascular NAD(P)H oxidase activation in diabetes: a double-edged sword in redox signalling. *Cardiovasc. Res.* 82, 9–20. doi: 10.1093/cvr/cvp031
- Garcia-Cardena, G., Fan, R., Stern, D. F., Liu, J., and Sessa, W. C. (1996). Endothelial nitric oxide synthase is regulated by tyrosine phosphorylation and interacts with caveolin-1. *J. Biol. Chem.* 271, 27237–27240. doi: 10.1074/jbc.271.44.27237
- Garcia-Vaz, E., Mcneilly, A. D., Berglund, L. M., Ahmad, A., and Gallagher, J. R. (2020). Inhibition of NFAT signaling restores microvascular endothelial function in diabetic mice. *Diabetes* 69, 424–435. doi: 10.2337/db18-0870
- Goedicke-Fritz, S., Kaistha, A., Kacik, M., Markert, S., and Hofmeister, A. (2015). Evidence for functional and dynamic microcompartmentation of Cav-1/TRPV4/K(Ca) in caveolae of endothelial cells. *Eur. J. Cell Biol.* 94, 391–400. doi: 10.1016/j.ejcb.2015.06.002
- Gonzalez-Perez, V., and Lingle, C. J. (2019). Regulation of BK channels by beta and gamma subunits. *Annu. Rev. Physiol.* 81, 113–137. doi: 10.1146/annurev-physiol-022516-034038
- Gonzalez-Perez, V., Xia, X. M., and Lingle, C. J. (2015). Two classes of regulatory subunits coassemble in the same BK channel and independently regulate gating. *Nat. Commun.* 6:8341. doi: 10.1038/ncomms9341
- Goodwill, A. G., Dick, G. M., Kiel, A. M., and Tune, J. D. (2017). Regulation of coronary blood flow. *Compr. Physiol.* 7, 321–382. doi: 10.1002/cphy.c160016
- Gratton, J. P., Bernatchez, P., and Sessa, W. C. (2004). Caveolae and caveolins in the cardiovascular system. *Circ. Res.* 94, 1408–1417. doi: 10.1161/01.RES.0000129178.56294.17
- Grayson, T. H., Murphy, T. V., and Sandow, S. L. (2017). Transient receptor potential canonical type 3 channels: interactions, role and relevance: a vascular focus. *Pharmacol. Ther.* 174, 79–96. doi: 10.1016/j.pharmthera.2017.02.022
- Gregg, E. W., Gu, Q., Cheng, Y. J., Narayan, K. M., and Cowie, C. C. (2007). Mortality trends in men and women with diabetes, 1971 to 2000. *Ann. Intern. Med.* 147, 149–155. doi: 10.7326/0003-4819-147-3-200708070-00167
- Halim, M., and Halim, A. (2019). The effects of inflammation, aging and oxidative stress on the pathogenesis of diabetes mellitus (type 2 diabetes). *Diabetes Metab. Syndr.* 13, 1165–1172. doi: 10.1016/j.dsx.2019.01.040
- Hashad, A. M., Harraz, O. F., Brett, S. E., Romero, M., and Kassmann, M. (2018). Caveolae link CaV3.2 channels to BKCa-mediated feedback in vascular smooth muscle. *Arterioscler. Thromb. Vasc. Biol.* 38, 2371–2381. doi: 10.1161/ATVBAHA.118.311394
- Heijnen, H. F., Waaijenborg, S., Crapo, J. D., Bowler, R. P., and Akkerman, J. W. (2004). Colocalization of eNOS and the catalytic subunit of PKA in endothelial cell junctions: a clue for regulated NO production. *J. Histochem. Cytochem.* 52, 1277–1285. doi: 10.1177/002215540405201004
- Hilenski, L. L., Clempus, R. E., Quinn, M. T., Lambeth, J. D., and Griendling, K. K. (2004). Distinct subcellular localizations of Nox1 and Nox4 in vascular smooth muscle cells. *Arterioscler. Thromb. Vasc. Biol.* 24, 677–683. doi: 10.1161/01.ATV.0000112024.13727.2c
- Hillman, N., Cox, S., Noble, A. R., and Gallagher, P. J. (2001). Increased numbers of caveolae in retinal endothelium and pericytes in hypertensive diabetic rats. *Eye* 15, 319–325. doi: 10.1038/eye.2001.103
- Houghton, C. A. (2019). Sulforaphane: its “coming of age” as a clinically relevant nutraceutical in the prevention and treatment of chronic disease. *Oxidative Med. Cell. Longev.* 2019:2716870. doi: 10.1155/2019/2716870
- Howitt, L., Grayson, T. H., Morris, M. J., Sandow, S. L., and Murphy, T. V. (2012). Dietary obesity increases NO and inhibits BKCa-mediated, endothelium-dependent dilation in rat cremaster muscle artery: association with caveolins and caveolae. *Am. J. Physiol. Heart Circ. Physiol.* 302, H2464–H2476. doi: 10.1152/ajpheart.00965.2011
- Hughes, J. M., Riddle, M. A., Paffett, M. L., Bosc, L. V. G., and Walker, B. R. (2010). Novel role of endothelial BKCa channels in altered vasoreactivity following hypoxia. *Am. J. Physiol. Heart Circ. Physiol.* 299, H1439–H1450. doi: 10.1152/ajpheart.00124.2010
- Imig, J. D., Dimitropoulou, C., Reddy, D. S., White, R. E., and Falck, J. R. (2008). Afferent arteriolar dilation to 11, 12-EET analogs involves PP2A activity and Ca²⁺-activated K⁺ channels. *Microcirculation* 15, 137–150. doi: 10.1080/10739680701456960
- Inoguchi, T., Li, P., Umeda, F., Yu, H. Y., and Kakimoto, M. (2000). High glucose level and free fatty acid stimulate reactive oxygen species production through protein kinase C--dependent activation of NAD(P)H oxidase in cultured vascular cells. *Diabetes* 49, 1939–1945. doi: 10.2337/diabetes.49.11.1939
- Inoguchi, T., Sonta, T., Tsubouchi, H., Etoh, T., and Kakimoto, M. (2003). Protein kinase C-dependent increase in reactive oxygen species (ROS) production in vascular tissues of diabetes: role of vascular NAD(P)H oxidase. *J. Am. Soc. Nephrol.* 14, S227–S232. doi: 10.1097/01.ASN.0000077407.90309.65
- Ishizaka, N., Griendling, K. K., Lassegue, B., and Alexander, R. W. (1998). Angiotensin II type 1 receptor: relationship with caveolae and caveolin after initial agonist stimulation. *Hypertension* 32, 459–466. doi: 10.1161/01.HYP.32.3.459
- Jiao, H., Arner, P., Hoffstedt, J., Brodin, D., and Dubern, B. (2011). Genome wide association study identifies KCNMA1 contributing to human obesity. *BMC Med. Genet.* 4:51. doi: 10.1186/1755-8794-4-51
- Kawai, T., Forrester, S. J., O'Brien, S., Baggett, A., and Rizzo, V. (2017). AT1 receptor signaling pathways in the cardiovascular system. *Pharmacol. Res.* 125, 4–13. doi: 10.1016/j.phrs.2017.05.008
- Kazama, Y., Kanemaru, Y., Noguchi, T., and Onaya, T. (1987). Reevaluation of circulating prostacyclin and thromboxane in diabetes. *Prostaglandins Leukot. Med.* 26, 115–122. doi: 10.1016/0262-1746(87)90106-5
- Kipreos, E. T., and Pagano, M. (2000). The F-box protein family. *Genome Biol.* 1:REVIEWS3002. doi: 10.1186/gb-2000-1-5-reviews3002
- Knaus, H. G., Garcia-Calvo, M., Kaczorowski, G. J., and Garcia, M. L. (1994). Subunit composition of the high conductance calcium-activated potassium channel from smooth muscle, a representative of the mSlo and slowpoke family of potassium channels. *J. Biol. Chem.* 269, 3921–3924. doi: 10.1016/S0021-9258(17)41720-0
- Konior, A., Schramm, A., Czesnikiewicz-Guzik, M., and Guzik, T. J. (2014). NADPH oxidases in vascular pathology. *Antioxid. Redox Signal.* 20, 2794–2814. doi: 10.1089/ars.2013.5607
- Krajewska, W. M., and Maslowska, I. (2004). Caveolins: structure and function in signal transduction. *Cell. Mol. Biol. Lett.* 9, 195–220.
- Kur, J., Mcgahon, M. K., Fernandez, J. A., Scholfield, C. N., and Mcgeown, J. G. (2014). Role of ion channels and subcellular Ca²⁺ signaling in arachidonic acid-induced dilation of pressurized retinal arterioles. *Invest. Ophthalmol. Vis. Sci.* 55, 2893–2902. doi: 10.1167/iovs.13-13511
- Kurusu, S., Inoue, I., Kawagoe, T., Ishihara, M., and Shimatani, Y. (2003). Diabetes mellitus is associated with insufficient microvascular reperfusion

- following revascularization for anterior acute myocardial infarction. *Intern. Med.* 42, 554–559. doi: 10.2169/internalmedicine.42.554
- Kwan, H. Y., Shen, B., Ma, X., Kwok, Y. C., and Huang, Y. (2009). TRPC1 associates with BK(Ca) channel to form a signal complex in vascular smooth muscle cells. *Circ. Res.* 104, 670–678. doi: 10.1161/CIRCRESAHA.108.188748
- Lane, L. S., Jansen, P. D., Lahav, M., and Rudy, C. (1982). Circulating prostacyclin and thromboxane levels in patients with diabetic retinopathy. *Ophthalmology* 89, 763–766. doi: 10.1016/S0161-6420(82)34729-6
- Larsen, B. T., Miura, H., Hatoum, O. A., Campbell, W. B., and Hammock, B. D. (2006). Epoxyeicosatrienoic and dihydroxyeicosatrienoic acids dilate human coronary arterioles via BK(Ca) channels: implications for soluble epoxide hydrolase inhibition. *Am. J. Physiol. Heart Circ. Physiol.* 290, H491–H499. doi: 10.1152/ajpheart.00927.2005
- Lauterbach, B., Barbosa-Sicard, E., Wang, M. H., Honeck, H., and Kargel, E. (2002). Cytochrome P450-dependent eicosapentaenoic acid metabolites are novel BK channel activators. *Hypertension* 39, 609–613. doi: 10.1161/hy0202.103293
- Ledoux, J., Werner, M. E., Brayden, J. E., and Nelson, M. T. (2006). Calcium-activated potassium channels and the regulation of vascular tone. *Physiology* 21, 69–78. doi: 10.1152/physiol.00040.2005
- Lee, H., Woodman, S. E., Engelman, J. A., Volonte, D., and Galbiati, F. (2001). Palmitoylation of caveolin-1 at a single site (Cys-156) controls its coupling to the c-Src tyrosine kinase: targeting of dually acylated molecules (GPI-linked, transmembrane, or cytoplasmic) to caveolae effectively uncouples c-Src and caveolin-1 (TYR-14). *J. Biol. Chem.* 276, 35150–35158. doi: 10.1074/jbc.M104530200
- Leo, M. D., Bulley, S., Bannister, J. P., Kuruvilla, K. P., and Narayanan, D. (2015). Angiotensin II stimulates internalization and degradation of arterial myocyte plasma membrane BK channels to induce vasoconstriction. *Am. J. Phys. Cell Phys.* 309, C392–C402. doi: 10.1152/ajpcell.00127.2015
- Leon, B. M., and Maddox, T. M. (2015). Diabetes and cardiovascular disease: epidemiology, biological mechanisms, treatment recommendations and future research. *World J. Diabetes* 6, 1246–1258. doi: 10.4239/wjcd.v6.i13.1246
- Leopold, J. A. (2015). Vascular calcification: mechanisms of vascular smooth muscle cell calcification. *Trends Cardiovasc. Med.* 25, 267–274. doi: 10.1016/j.tcm.2014.10.021
- Li, W., Bengtson, M. H., Ulbrich, A., Matsuda, A., and Reddy, V. A. (2008). Genome-wide and functional annotation of human E3 ubiquitin ligases identifies MULAN, a mitochondrial E3 that regulates the organelle's dynamics and signaling. *PLoS One* 3:e1487. doi: 10.1371/journal.pone.0003918
- Li, P. L., and Campbell, W. B. (1997). Epoxyeicosatrienoic acids activate K⁺ channels in coronary smooth muscle through a guanine nucleotide binding protein. *Circ. Res.* 80, 877–884. doi: 10.1161/01.RES.80.6.877
- Li, Q., Guan, X., Yen, K., Zhang, J., and Yan, J. (2016). The single transmembrane segment determines the modulatory function of the BK channel auxiliary gamma subunit. *J. Gen. Physiol.* 147, 337–351. doi: 10.1085/jgp.201511551
- Li, Y., Wang, X. L., Sun, X., Chai, Q., and Li, J. (2017). Regulation of vascular large-conductance calcium-activated potassium channels by Nrf2 signalling. *Diab. Vasc. Dis. Res.* 14, 353–362. doi: 10.1177/1479164117703903
- Li, X., Xing, W., Wang, Y., Mi, C., and Zhang, Z. (2014). Upregulation of caveolin-1 contributes to aggravated high-salt diet-induced endothelial dysfunction and hypertension in type 1 diabetic rats. *Life Sci.* 113, 31–39. doi: 10.1016/j.lfs.2014.07.027
- Li, Q., and Yan, J. (2016). Modulation of BK channel function by auxiliary beta and gamma subunits. *Int. Rev. Neurobiol.* 128, 51–90. doi: 10.1016/bs.irn.2016.03.015
- Li, P. L., Zhang, D. X., Ge, Z. D., and Campbell, W. B. (2002). Role of ADP-ribose in 11,12-EET-induced activation of K(Ca) channels in coronary arterial smooth muscle cells. *Am. J. Physiol. Heart Circ. Physiol.* 282, H1229–H1236. doi: 10.1152/ajpheart.00736.2001
- Liang, L., Li, X., Moutton, S., Vergano, S. A. S., and Cogne, B. (2019). De novo loss-of-function KCNMA1 variants are associated with a new multiple malformation syndrome and a broad spectrum of developmental and neurological phenotypes. *Hum. Mol. Genet.* 28, 2937–2951. doi: 10.1093/hmg/ddz117
- Linder, A. E., McCluskey, L. P., Cole, K. R. 3rd, Lanning, K. M., and Webb, R. C. (2005). Dynamic association of nitric oxide downstream signaling molecules with endothelial caveolin-1 in rat aorta. *J. Pharmacol. Exp. Ther.* 314, 9–15. doi: 10.1124/jpet.105.083634
- Liu, G., Niu, X., Wu, R. S., Chudasama, N., and Yao, Y. (2010). Location of modulatory beta subunits in BK potassium channels. *J. Gen. Physiol.* 135, 449–459. doi: 10.1085/jgp.201010417
- Liu, Y., Terata, K., Chai, Q., Li, H., and Kleinman, L. H. (2002). Peroxynitrite inhibits Ca²⁺-activated K⁺ channel activity in smooth muscle of human coronary arterioles. *Circ. Res.* 91, 1070–1076. doi: 10.1161/01.RES.0000046003.14031.98
- Liu, J., Ye, J., Zou, X., Xu, Z., and Feng, Y. (2014). CRL4A(CRBN) E3 ubiquitin ligase restricts BK channel activity and prevents epileptogenesis. *Nat. Commun.* 5:3924. doi: 10.1038/ncomms4924
- Lu, T., Chai, Q., Jiao, G., Wang, X. L., and Sun, X. (2019). Downregulation of BK channel function and protein expression in coronary arteriolar smooth muscle cells of type 2 diabetic patients. *Cardiovasc. Res.* 115, 145–153. doi: 10.1093/cvr/cvy137
- Lu, T., Chai, Q., Yu, L., D'uscio, L. V., and Katusic, Z. S. (2012). Reactive oxygen species signaling facilitates FOXO-3a/FBXO-dependent vascular BK channel beta1 subunit degradation in diabetic mice. *Diabetes* 61, 1860–1868. doi: 10.2337/db11-1658
- Lu, T., He, T., Katusic, Z. S., and Lee, H. C. (2006). Molecular mechanisms mediating inhibition of human large conductance Ca²⁺-activated K⁺ channels by high glucose. *Circ. Res.* 99, 607–616. doi: 10.1161/01.RES.0000243147.41792.93
- Lu, T., Jiang, B., Wang, X. L., and Lee, H. C. (2016). Coronary arterial BK channel dysfunction exacerbates ischemia/reperfusion-induced myocardial injury in diabetic mice. *Appl. Physiol. Nutr. Metab.* 41, 992–1001. doi: 10.1139/apnm-2016-0048
- Lu, T., Katakam, P. V., Vanrollins, M., Weintraub, N. L., and Spector, A. A. (2001). Dihydroxyeicosatrienoic acids are potent activators of Ca(2+)-activated K(+) channels in isolated rat coronary arterial myocytes. *J. Physiol.* 534, 651–667. doi: 10.1111/j.1469-7793.2001.t01-1-00651.x
- Lu, T., Sun, X., Li, Y., Chai, Q., and Wang, X. L. (2017a). Role of Nrf2 signaling in the regulation of vascular BK channel beta1 subunit expression and BK channel function in high-fat diet-induced diabetic mice. *Diabetes* 66, 2681–2690. doi: 10.2337/db17-0181
- Lu, T., Wang, X. L., Chai, Q., Sun, X., and Sieck, G. C. (2017b). Role of the endothelial caveolae microdomain in shear stress-mediated coronary vasorelaxation. *J. Biol. Chem.* 292, 19013–19023. doi: 10.1074/jbc.M117.786152
- Lu, T., Wang, X. L., He, T., Zhou, W., and Kaduce, T. L. (2005). Impaired arachidonic acid-mediated activation of large-conductance Ca²⁺-activated K⁺ channels in coronary arterial smooth muscle cells in Zucker diabetic fatty rats. *Diabetes* 54, 2155–2163. doi: 10.2337/diabetes.54.7.2155
- Lu, T., Ye, D., He, T., Wang, X. L., and Wang, H. L. (2008). Impaired Ca²⁺-dependent activation of large-conductance Ca²⁺-activated K⁺ channels in the coronary artery smooth muscle cells of Zucker diabetic fatty rats. *Biophys. J.* 95, 5165–5177. doi: 10.1529/biophysj.108.138339
- Lu, T., Zhang, D. M., Wang, X. L., He, T., and Wang, R. X. (2010). Regulation of coronary arterial BK channels by caveolae-mediated angiotensin II signaling in diabetes mellitus. *Circ. Res.* 106, 1164–1173. doi: 10.1161/CIRCRESAHA.109.209767
- Lv, X., Zhang, Y., Niu, Y., Song, Q., and Zhao, Q. (2018). Comparison of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers on cardiovascular outcomes in hypertensive patients with type 2 diabetes mellitus: a PRISMA-compliant systematic review and meta-analysis. *Medicine* 97:e0256. doi: 10.1097/MD.00000000000010256
- Ma, Y. C., Huang, J., Ali, S., Lowry, W., and Huang, X. Y. (2000). Src tyrosine kinase is a novel direct effector of G proteins. *Cell* 102, 635–646. doi: 10.1016/S0092-8674(00)00086-6
- Ma, Z., Lou, X. J., and Horrigan, F. T. (2006). Role of charged residues in the S1-S4 voltage sensor of BK channels. *J. Gen. Physiol.* 127, 309–328. doi: 10.1085/jgp.200509421
- Martin, P., Moncada, M., Castillo, K., Orsi, F., and Ducca, G. (2021). Arachidonic acid effect on the allosteric gating mechanism of BK (Sl α) channels associated with the beta1 subunit. *Biochim. Biophys. Acta Biomembr.* 1863:183550. doi: 10.1016/j.bbmem.2021.183550
- Martin, P., Moncada, M., Enrique, N., Asuaje, A., and Capuccino, J. M. V. (2014). Arachidonic acid activation of BKCa (Sl α) channels associated to the beta1-subunit in human vascular smooth muscle cells. *Pflugers Arch.* 466, 1779–1792. doi: 10.1007/s00424-013-1422-x
- McGahon, M. K., Dash, D. P., Arora, A., Wall, N., and Dawicki, J. (2007). Diabetes downregulates large-conductance Ca²⁺-activated potassium beta 1

- channel subunit in retinal arteriolar smooth muscle. *Circ. Res.* 100, 703–711. doi: 10.1161/01.RES.0000260182.36481.c9
- McManus, O. B., and Magleby, K. L. (1988). Kinetic states and modes of single large-conductance calcium-activated potassium channels in cultured rat skeletal muscle. *J. Physiol.* 402, 79–120. doi: 10.1113/jphysiol.1988.sp017195
- McManus, O. B., and Magleby, K. L. (1991). Accounting for the Ca(2+)-dependent kinetics of single large-conductance Ca(2+)-activated K+ channels in rat skeletal muscle. *J. Physiol.* 443, 739–777. doi: 10.1113/jphysiol.1991.sp018861
- McSweeney, S. R., Warabi, E., and Siow, R. C. (2016). Nrf2 as an endothelial mechanosensitive transcription factor: going with the flow. *Hypertension* 67, 20–29. doi: 10.1161/HYPERTENSIONAHA.115.06146
- Miao, X., Bai, Y., Sun, W., Cui, W., and Xin, Y. (2012). Sulforaphane prevention of diabetes-induced aortic damage was associated with the up-regulation of Nrf2 and its down-stream antioxidants. *Nutr. Metab.* 9:84. doi: 10.1186/1743-7075-9-84
- Migdalís, I. N., Kalogeropoulou, K., Iliopoulou, V., Triantafílu, P., and Charalabides, J. (2001). Plasma levels of endothelin, lipid peroxides and prostacyclin in diabetic patients with macroangiopathy. *Diabetes Res. Clin. Pract.* 54, 129–136. doi: 10.1016/S0168-8227(01)00275-3
- Miura, H., Liu, Y., and Gutterman, D. D. (1999). Human coronary arteriolar dilation to bradykinin depends on membrane hyperpolarization: contribution of nitric oxide and Ca²⁺-activated K⁺ channels. *Circulation* 99, 3132–3138. doi: 10.1161/01.CIR.99.24.3132
- Mokelke, E. A., Dietz, N. J., Eckman, D. M., Nelson, M. T., and Sturek, M. (2005). Diabetic dyslipidemia and exercise affect coronary tone and differential regulation of conduit and microvessel K⁺ current. *Am. J. Physiol. Heart Circ. Physiol.* 288, H1233–H1241. doi: 10.1152/ajpheart.00732.2004
- Mokelke, E. A., Hu, Q., Song, M., Toro, L., and Reddy, H. K. (2003). Altered functional coupling of coronary K⁺ channels in diabetic dyslipidemic pigs is prevented by exercise. *J. Appl. Physiol.* 95, 1179–1193. doi: 10.1152/jappphysiol.00972.2002
- Mokhtar, S. S., Vanhoutte, P. M., Leung, S. W. S., Yusof, M. I., Sulaiman, W. A. W., Saad, A. Z. M., et al. (2013). Reduced expression of prostacyclin synthase and nitric oxide synthase in subcutaneous arteries of type 2 diabetic patients. *Tohoku J. Exp. Med.* 231, 217–222. doi: 10.1620/tjem.231.217
- Mori, A., Suzuki, S., Sakamoto, K., Nakahara, T., and Ishii, K. (2011). BMS-191011, an opener of large-conductance Ca²⁺-activated potassium channels, dilates rat retinal arterioles in vivo. *Biol. Pharm. Bull.* 34, 150–152. doi: 10.1248/bpb.34.150
- Morin, C., Sirois, M., Echave, V., Gomes, M. M., and Rousseau, E. (2007). Functional effects of 20-HETE on human bronchi: hyperpolarization and relaxation due to BKCa channel activation. *Am. J. Phys. Lung Cell. Mol. Phys.* 293, L1037–L1044. doi: 10.1152/ajplung.00145.2007
- Narayanan, A., Amaya, M., Voss, K., Chung, M., and Benedict, A. (2014). Reactive oxygen species activate NFκB (p65) and p53 and induce apoptosis in RVFV infected liver cells. *Virology* 449, 270–286. doi: 10.1016/j.virol.2013.11.023
- Navedo, M. F., Takeda, Y., Nieves-Cintrón, M., Molkentin, J. D., and Santana, L. F. (2010). Elevated Ca²⁺ sparklet activity during acute hyperglycemia and diabetes in cerebral arterial smooth muscle cells. *Am. J. Phys. Cell Phys.* 298, C211–C220. doi: 10.1152/ajpcell.00267.2009
- Nelson, M. T., Cheng, H., Rubart, M., Santana, L. F., and Bonev, A. D. (1995). Relaxation of arterial smooth muscle by calcium sparks. *Science* 270, 633–637. doi: 10.1126/science.270.5236.633
- Nelson, M. T., and Quayle, J. M. (1995). Physiological roles and properties of potassium channels in arterial smooth muscle. *Am. J. Phys.* 268, C799–C822. doi: 10.1152/ajpcell.1995.268.4.C799
- Nieves-Cintrón, M., Amberg, G. C., Nichols, C. B., Molkentin, J. D., and Santana, L. F. (2007). Activation of NFATc3 down-regulates the beta1 subunit of large conductance, calcium-activated K⁺ channels in arterial smooth muscle and contributes to hypertension. *J. Biol. Chem.* 282, 3231–3240. doi: 10.1074/jbc.M608822200
- Nieves-Cintrón, M., Syed, A. U., Buonarati, O. R., Rigor, R. R., and Nystoriak, M. A. (2017). Impaired BKCa channel function in native vascular smooth muscle from humans with type 2 diabetes. *Sci. Rep.* 7:14058. doi: 10.1038/s41598-017-14565-9
- Niklason, A., Hedner, T., Niskanen, L., Lanke, J., and Captopril Prevention Project Study Group (2004). Development of diabetes is retarded by ACE inhibition in hypertensive patients—a subanalysis of the captopril prevention project (CAPPP). *J. Hypertens.* 22, 645–652. doi: 10.1097/00004872-200403000-00029
- Nystoriak, M. A., Nieves-Cintrón, M., Nygren, P. J., Hinke, S. A., and Nichols, C. B. (2014). AKAP150 contributes to enhanced vascular tone by facilitating large-conductance Ca²⁺-activated K⁺ channel remodeling in hyperglycemia and diabetes mellitus. *Circ. Res.* 114, 607–615. doi: 10.1161/CIRCRESAHA.114.302168
- Okamoto, T., Schlegel, A., Scherer, P. E., and Lisanti, M. P. (1998). Caveolins, a family of scaffolding proteins for organizing "preassembled signaling complexes" at the plasma membrane. *J. Biol. Chem.* 273, 5419–5422. doi: 10.1074/jbc.273.10.5419
- Okon, E. B., Chung, A. W., Rauniyar, P., Padilla, E., and Tejerina, T. (2005). Compromised arterial function in human type 2 diabetic patients. *Diabetes* 54, 2415–2423. doi: 10.2337/diabetes.54.8.2415
- Oliveria, S. F., Dell'acqua, M. L., and Sather, W. A. (2007). AKAP79/150 anchoring of calcineurin controls neuronal L-type Ca²⁺ channel activity and nuclear signaling. *Neuron* 55, 261–275. doi: 10.1016/j.neuron.2007.06.032
- Op De Beeck, A., and Eizirik, D. L. (2016). Viral infections in type 1 diabetes mellitus—why the beta cells? *Nat. Rev. Endocrinol.* 12, 263–273. doi: 10.1038/nrendo.2016.30
- Pacher, P., Beckman, J. S., and Liaudet, L. (2007). Nitric oxide and peroxynitrite in health and disease. *Physiol. Rev.* 87, 315–424. doi: 10.1152/physrev.00029.2006
- Pandey, A., Chawla, S., and Guchhait, P. (2015). Type-2 diabetes: current understanding and future perspectives. *IUBMB Life* 67, 506–513. doi: 10.1002/iub.1396
- Pascariu, M., Bendayan, M., and Ghitescu, L. (2004). Correlated endothelial caveolin overexpression and increased transcytosis in experimental diabetes. *J. Histochem. Cytochem.* 52, 65–76. doi: 10.1177/002215540405200107
- Patel, V. B., Bodiga, S., Basu, R., Das, S. K., and Wang, W. (2012). Loss of angiotensin-converting enzyme-2 exacerbates diabetic cardiovascular complications and leads to systolic and vascular dysfunction: a critical role of the angiotensin II/AT1 receptor axis. *Circ. Res.* 110, 1322–1335. doi: 10.1161/CIRCRESAHA.112.268029
- Pietryga, M., Brazert, J., Wender-Ozegowska, E., Biczysko, R., and Dubiel, M. (2005). Abnormal uterine doppler is related to vasculopathy in pregestational diabetes mellitus. *Circulation* 112, 2496–2500. doi: 10.1161/CIRCULATIONAHA.104.492843
- Poulsen, A. N., Wulf, H., Hay-Schmidt, A., Jansen-Olesen, I., and Olesen, J. (2009). Differential expression of BK channel isoforms and beta-subunits in rat neuro-vascular tissues. *Biochim. Biophys. Acta* 1788, 380–389. doi: 10.1016/j.bbame.2008.10.001
- Powell, S. R. (2006). The ubiquitin-proteasome system in cardiac physiology and pathology. *Am. J. Physiol. Heart Circ. Physiol.* 291, H1–H19. doi: 10.1152/ajpheart.00062.2006
- Pu, D., Zhao, Y., Chen, J., Sun, Y., and Lv, A. (2018). Protective effects of sulforaphane on cognitive impairments and AD-like lesions in diabetic mice are associated with the upregulation of Nrf2 transcription activity. *Neuroscience* 381, 35–45. doi: 10.1016/j.neuroscience.2018.04.017
- Quilley, J., Fulton, D., and McGiff, J. C. (1997). Hyperpolarizing factors. *Biochem. Pharmacol.* 54, 1059–1070. doi: 10.1016/S0006-2952(97)00039-7
- Rao, A., Luo, C., and Hogan, P. G. (1997). Transcription factors of the NFAT family: regulation and function. *Annu. Rev. Immunol.* 15, 707–747. doi: 10.1146/annurev.immunol.15.1.707
- Razani, B., Engelman, J. A., Wang, X. B., Schubert, W., and Zhang, X. L. (2001). Caveolin-1 null mice are viable but show evidence of hyperproliferative and vascular abnormalities. *J. Biol. Chem.* 276, 38121–38138. doi: 10.1074/jbc.M105408200
- Ribeiro-Oliveira, A., Nogueira, A. I., Pereira, R. M., Boas, W. W., and Dos Santos, R. A. (2008). The renin-angiotensin system and diabetes: an update. *Vasc. Health Risk Manag.* 4, 787–803. doi: 10.2147/VHRM.S1905
- Riddle, M. A., Hughes, J. M., and Walker, B. R. (2011). Role of caveolin-1 in endothelial BKCa channel regulation of vasoreactivity. *Am. J. Phys. Cell Phys.* 301, C1404–C1414. doi: 10.1152/ajpcell.00013.2011
- Ringvold, H. C., and Khalil, R. A. (2017). Protein kinase C as regulator of vascular smooth muscle function and potential target in vascular disorders. *Adv. Pharmacol.* 78, 203–301. doi: 10.1016/bs.apha.2016.06.002
- Rueda, A., Fernandez-Velasco, M., Benitah, J. P., and Gomez, A. M. (2013). Abnormal Ca²⁺ spark/STOC coupling in cerebral artery smooth muscle cells of obese type 2 diabetic mice. *PLoS One* 8:e53321. doi: 10.1371/journal.pone.0053321
- Saeki, T., Suzuki, Y., Yamamura, H., Takeshima, H., and Imaizumi, Y. (2019). A junctophilin-caveolin interaction enables efficient coupling between ryanodine

- receptors and BKCa channels in the Ca(2+) microdomain of vascular smooth muscle. *J. Biol. Chem.* 294, 13093–13105. doi: 10.1074/jbc.RA119.008342
- Sandri, M., Sandri, C., Gilbert, A., Skurk, C., and Calabria, E. (2004). Foxo transcription factors induce the atrophy-related ubiquitin ligase atrogin-1 and cause skeletal muscle atrophy. *Cell* 117, 399–412. doi: 10.1016/S0092-8674(04)00400-3
- Schapiro, M., Calabrese, M. F., Bullock, A. N., and Crews, C. M. (2019). Targeted protein degradation: expanding the toolbox. *Nat. Rev. Drug Discov.* 18, 949–963. doi: 10.1038/s41573-019-0047-y
- Schuck, R. N., Theken, K. N., Edin, M. L., Caughey, M., and Bass, A. (2013). Cytochrome P450-derived eicosanoids and vascular dysfunction in coronary artery disease patients. *Atherosclerosis* 227, 442–448. doi: 10.1016/j.atherosclerosis.2013.01.034
- Sedding, D. G., Hermsen, J., Seay, U., Eickelberg, O., and Kummer, W. (2005). Caveolin-1 facilitates mechanosensitive protein kinase B (Akt) signaling in vitro and in vivo. *Circ. Res.* 96, 635–642. doi: 10.1161/01.RES.0000160610.61306.0f
- Shi, J., Krishnamoorthy, G., Yang, Y., Hu, L., and Chaturvedi, N. (2002). Mechanism of magnesium activation of calcium-activated potassium channels. *Nature* 418, 876–880. doi: 10.1038/nature00941
- Singh, H., Lu, R., Bopassa, J. C., Meredith, A. L., and Stefani, E. (2013). MitoBK(Ca) is encoded by the Kcnma1 gene, and a splicing sequence defines its mitochondrial location. *Proc. Natl. Acad. Sci. U. S. A.* 110, 10836–10841. doi: 10.1073/pnas.1302028110
- Song, T., Liang, S., Liu, J., Zhang, T., and Yin, Y. (2018). CRL4 antagonizes SCFFbxo7-mediated turnover of cereblon and BK channel to regulate learning and memory. *PLoS Genet.* 14:e1007165. doi: 10.1371/journal.pgen.1007165
- Sorop, O., Van Den Heuvel, M., Van Ditzhuijzen, N. S., De Beer, V. J., and Heinonen, I. (2016). Coronary microvascular dysfunction after long-term diabetes and hypercholesterolemia. *Am. J. Physiol. Heart Circ. Physiol.* 311, H1339–H1351. doi: 10.1152/ajpheart.00458.2015
- Spisni, E., Griffoni, C., Santi, S., Riccio, M., and Marulli, R. (2001). Colocalization prostacyclin (PGI2) synthase-caveolin-1 in endothelial cells and new roles for PGI2 in angiogenesis. *Exp. Cell Res.* 266, 31–43. doi: 10.1006/excr.2001.5198
- Stone, P. H., Muller, J. E., Hartwell, T., York, B. J., and Rutherford, J. D. (1989). The effect of diabetes mellitus on prognosis and serial left ventricular function after acute myocardial infarction: contribution of both coronary disease and diastolic left ventricular dysfunction to the adverse prognosis. The MILIS study group. *J. Am. Coll. Cardiol.* 14, 49–57. doi: 10.1016/0735-1097(89)90053-3
- Sun, X., Qian, L. L., Li, Y., Pfeifer, T. M., and Wang, X. L. (2020). Regulation of KCNMA1 transcription by Nrf2 in coronary arterial smooth muscle cells. *J. Mol. Cell. Cardiol.* 140, 68–76. doi: 10.1016/j.yjmcc.2020.03.001
- Sun, X., Zhou, D., Zhang, P., Moczydlowski, E. G., and Haddad, G. G. (2007). Beta-subunit-dependent modulation of hSlo BK current by arachidonic acid. *J. Neurophysiol.* 97, 62–69. doi: 10.1152/jn.00700.2006
- Suzuki, T., and Yamamoto, M. (2015). Molecular basis of the Keap1-Nrf2 system. *Free Radic. Biol. Med.* 88, 93–100. doi: 10.1016/j.freeradbiomed.2015.06.006
- Suzuki, Y., Yamamura, H., Ohya, S., and Imaizumi, Y. (2013). Caveolin-1 facilitates the direct coupling between large conductance Ca²⁺-activated K⁺ (BKCa) and Cav1.2 Ca²⁺ channels and their clustering to regulate membrane excitability in vascular myocytes. *J. Biol. Chem.* 288, 36750–36761. doi: 10.1074/jbc.M113.511485
- Szaleczky, E., Prechl, J., Feher, J., and Somogyi, A. (1999). Alterations in enzymatic antioxidant defence in diabetes mellitus—a rational approach. *Postgrad. Med. J.* 75, 13–17. doi: 10.1136/pgmj.75.879.13
- Takeda, Y., Nystoriak, M. A., Nieves-Cintrón, M., Santana, L. F., and Navedo, M. F. (2011). Relationship between Ca²⁺ sparklets and sarcoplasmic reticulum Ca²⁺ load and release in rat cerebral arterial smooth muscle. *Am. J. Physiol. Heart Circ. Physiol.* 301, H2285–H2294. doi: 10.1152/ajpheart.00488.2011
- Tallima, H., and El Ridi, R. (2018). Arachidonic acid: physiological roles and potential health benefits—a review. *J. Adv. Res.* 11, 33–41. doi: 10.1016/j.jare.2017.11.004
- Tan, Y., Ichikawa, T., Li, J., Si, Q., and Yang, H. (2011). Diabetic downregulation of Nrf2 activity via ERK contributes to oxidative stress-induced insulin resistance in cardiac cells in vitro and in vivo. *Diabetes* 60, 625–633. doi: 10.2337/db10-1164
- Tanaka, Y., Meera, P., Song, M., Knaus, H. G., and Toro, L. (1997). Molecular constituents of maxi KCa channels in human coronary smooth muscle: predominant alpha + beta subunit complexes. *J. Physiol.* 502, 545–557. doi: 10.1111/j.1469-7793.1997.545bj.x
- Tang, X. D., Daggett, H., Hanner, M., Garcia, M. L., and Mcmanus, O. B. (2001). Oxidative regulation of large conductance calcium-activated potassium channels. *J. Gen. Physiol.* 117, 253–274. doi: 10.1085/jgp.117.3.253
- Tang, X., Qian, L. L., Wang, R. X., Yao, Y., and Dang, S. P. (2017). Regulation of coronary arterial large conductance Ca²⁺-activated K⁺ channel protein expression and function by n-3 polyunsaturated fatty acids in diabetic rats. *J. Vasc. Res.* 54, 329–343. doi: 10.1159/000479870
- Taniyama, Y., and Griendling, K. K. (2003). Reactive oxygen species in the vasculature: molecular and cellular mechanisms. *Hypertension* 42, 1075–1081. doi: 10.1161/01.HYP.0000100443.09293.4F
- Taniyama, Y., Ushio-Fukai, M., Hitomi, H., Rocic, P., and Kingsley, M. J. (2004). Role of p38 MAPK and MAPKAPK-2 in angiotensin II-induced Akt activation in vascular smooth muscle cells. *Am. J. Phys. Cell Phys.* 287, C494–C499. doi: 10.1152/ajpcell.00439.2003
- Tao, X., Hite, R. K., and Mackinnon, R. (2017). Cryo-EM structure of the open high-conductance Ca(2+)-activated K(+) channel. *Nature* 541, 46–51. doi: 10.1038/nature20608
- Tarantini, S., Valcarcel-Ares, M. N., Yabluchanskiy, A., Tucsek, Z., and Hertelendy, P. (2018). Nrf2 deficiency exacerbates obesity-induced oxidative stress, neurovascular dysfunction, blood-brain barrier disruption, neuroinflammation, amyloidogenic gene expression, and cognitive decline in mice, mimicking the aging phenotype. *J. Gerontol. A Biol. Sci. Med. Sci.* 73, 853–863. doi: 10.1093/gerona/glx177
- Theken, K. N., Schuck, R. N., Edin, M. L., Tran, B., and Ellis, K. (2012). Evaluation of cytochrome P450-derived eicosanoids in humans with stable atherosclerotic cardiovascular disease. *Atherosclerosis* 222, 530–536. doi: 10.1016/j.atherosclerosis.2012.03.022
- Tian, L., Mcclafferty, H., Chen, L., and Shipston, M. J. (2007). Reversible tyrosine protein phosphorylation regulates large conductance voltage- and calcium-activated potassium (BK) channels via cortactin. *J. Biol. Chem.* 283, 3067–3076. doi: 10.1074/jbc.M706826200
- Tiwari, B. K., Pandey, K. B., Abidi, A. B., and Rizvi, S. I. (2013). Markers of oxidative stress during diabetes mellitus. *J. Biomark.* 2013:378790. doi: 10.1155/2013/378790
- Torres, Y. P., Granados, S. T., and Latorre, R. (2014). Pharmacological consequences of the coexpression of BK channel alpha and auxiliary beta subunits. *Front. Physiol.* 5:383. doi: 10.3389/fphys.2014.00383
- Touyz, R. M., and Schiffrin, E. L. (2000). Signal transduction mechanisms mediating the physiological and pathophysiological actions of angiotensin II in vascular smooth muscle cells. *Pharmacol. Rev.* 52, 639–672.
- Tsai, S. H., Hein, T. W., Kuo, L., and Yang, V. C. (2011). High glucose impairs EDHF-mediated dilation of coronary arterioles via reduced cytochrome P450 activity. *Microvasc. Res.* 82, 356–363. doi: 10.1016/j.mvr.2011.09.008
- Ungvari, Z., Bailey-Downs, L., Gautam, T., Jimenez, R., and Losonczy, G. (2011). Adaptive induction of NF-E2-related factor-2-driven antioxidant genes in endothelial cells in response to hyperglycemia. *Am. J. Physiol. Heart Circ. Physiol.* 300, H1133–H1140. doi: 10.1152/ajpheart.00402.2010
- Ushio-Fukai, M., and Alexander, R. W. (2006). Caveolin-dependent angiotensin II type 1 receptor signaling in vascular smooth muscle. *Hypertension* 48, 797–803. doi: 10.1161/01.HYP.0000242907.70697.5d
- Uyy, E., Antohe, F., Ivan, L., Haraba, R., and Radu, D. L. (2010). Upregulation of caveolin-1 expression is associated with structural modifications of endothelial cells in diabetic lung. *Microvasc. Res.* 79, 154–159. doi: 10.1016/j.mvr.2009.11.008
- Van Den Heuvel, A. P., Schulze, A., and Burgering, B. M. (2005). Direct control of caveolin-1 expression by FOXO transcription factors. *Biochem. J.* 385, 795–802. doi: 10.1042/BJ20041449
- Vellecco, V., Mitidieri, E., Gargiulo, A., Brancaleone, V., and Matassa, D. (2016). Vascular effects of linagliptin in non-obese diabetic mice are glucose-independent and involve positive modulation of the endothelial nitric oxide synthase (eNOS)/caveolin-1 (CAV-1) pathway. *Diabetes Obes. Metab.* 18, 1236–1243. doi: 10.1111/dom.12750
- Vila, L. (2004). Cyclooxygenase and 5-lipoxygenase pathways in the vessel wall: role in atherosclerosis. *Med. Res. Rev.* 24, 399–424. doi: 10.1002/med.10065
- Wang, R. X., Chai, Q., Lu, T., and Lee, H. C. (2011). Activation of vascular BK channels by docosahexaenoic acid is dependent on cytochrome P450 epoxygenase activity. *Cardiovasc. Res.* 90, 344–352. doi: 10.1093/cvr/cvq411
- Wang, J., Hanada, K., Gareri, C., and Rockman, H. A. (2018). Mechanoactivation of the angiotensin II type 1 receptor induces beta-arrestin-biased signaling through Gα_q coupling. *J. Cell. Biochem.* 119, 3586–3597. doi: 10.1002/jcb.26552

- Wang, Z., Subramanya, A. R., Satlin, L. M., Pastor-Soler, N. M., and Carattino, M. D. (2013). Regulation of large-conductance Ca^{2+} -activated K^{+} channels by WNK4 kinase. *Am. J. Phys. Cell Phys.* 305, C846–C853. doi: 10.1152/ajpcell.00133.2013
- Wang, B., Wu, L., Chen, J., Dong, L., and Chen, C. (2021). Metabolism pathways of arachidonic acids: mechanisms and potential therapeutic targets. *Signal Transduct. Target. Ther.* 6:94. doi: 10.1038/s41392-020-00443-w
- Wang, X. L., Ye, D., Peterson, T. E., Cao, S., and Shah, V. H. (2005). Caveolae targeting and regulation of large conductance Ca^{2+} -activated K^{+} channels in vascular endothelial cells. *J. Biol. Chem.* 280, 11656–11664. doi: 10.1074/jbc.M410987200
- Williams, R., Karuranga, S., Malanda, B., Saeedi, P., and Basit, A. (2020). Global and regional estimates and projections of diabetes-related health expenditure: results from the international diabetes federation diabetes atlas, 9th edition. *Diabetes Res. Clin. Pract.* 162:108072. doi: 10.1016/j.diabres.2020.108072
- Wolin, M. S. (2004). Subcellular localization of Nox-containing oxidases provides unique insight into their role in vascular oxidant signaling. *Arterioscler. Thromb. Vasc. Biol.* 24, 625–627. doi: 10.1161/01.ATV.0000117201.14603.5d
- Wu, C. L., Cornwell, E. W., Jackman, R. W., and Kandarian, S. C. (2014). NF- κ B but not FoxO sites in the MuRF1 promoter are required for transcriptional activation in disuse muscle atrophy. *Am. J. Phys. Cell Phys.* 306, C762–C767. doi: 10.1152/ajpcell.00361.2013
- Wu, S. N., Li, H. F., and Chiang, H. T. (2000). Actions of epoxyeicosatrienoic acid on large-conductance Ca^{2+} -activated K^{+} channels in pituitary GH(3) cells. *Biochem. Pharmacol.* 60, 251–262. doi: 10.1016/S0006-2952(00)00317-8
- Wu, R. S., and Marx, S. O. (2010). The BK potassium channel in the vascular smooth muscle and kidney: alpha- and beta-subunits. *Kidney Int.* 78, 963–974. doi: 10.1038/ki.2010.325
- Xia, X. M., Zeng, X., and Lingle, C. J. (2002). Multiple regulatory sites in large-conductance calcium-activated potassium channels. *Nature* 418, 880–884. doi: 10.1038/nature00956
- Yagishita, Y., Gattbonton-Schwager, T. N., Mccallum, M. L., and Kensler, T. W. (2020). Current landscape of NRF2 biomarkers in clinical trials. *Antioxidants* 9:716. doi: 10.3390/antiox9080716
- Yamaki, F., Kaga, M., Horinouchi, T., Tanaka, H., and Koike, K. (2001). MaxiK channel-mediated relaxation of Guinea-pig aorta following stimulation of IP receptor with beraprost via cyclic AMP-dependent and -independent mechanisms. *Naunyn Schmiedebergs Arch Pharmacol.* 364, 538–550. doi: 10.1007/s002100100485
- Yan, J., and Aldrich, R. W. (2012). BK potassium channel modulation by leucine-rich repeat-containing proteins. *Proc. Natl. Acad. Sci. U. S. A.* 109, 7917–7922. doi: 10.1073/pnas.1205435109
- Yang, Y., Jones, A. W., Thomas, T. R., and Rubin, L. J. (2007). Influence of sex, high-fat diet, and exercise training on potassium currents of swine coronary smooth muscle. *Am. J. Physiol. Heart Circ. Physiol.* 293, H1553–H1563. doi: 10.1152/ajpheart.00151.2007
- Yeung, C. Y., Lam, K. S., Li, S. W., Lam, K. F., and Tse, H. F. (2012). Sudden cardiac death after myocardial infarction in type 2 diabetic patients with no residual myocardial ischemia. *Diabetes Care* 35, 2564–2569. doi: 10.2337/dc12-0118
- Yi, F., Wang, H., Chai, Q., Wang, X., and Shen, W. K. (2014). Regulation of large conductance Ca^{2+} -activated K^{+} (BK) channel beta1 subunit expression by muscle RING finger protein 1 in diabetic vessels. *J. Biol. Chem.* 289, 10853–10864. doi: 10.1074/jbc.M113.520940
- Yousif, M. H., and Benter, I. F. (2007). Role of cytochrome P450 metabolites of arachidonic acid in regulation of corporal smooth muscle tone in diabetic and older rats. *Vasc. Pharmacol.* 47, 281–287. doi: 10.1016/j.vph.2007.08.002
- Yuan, P., Leonetti, M. D., Pico, A. R., Hsiung, Y., and Mackinnon, R. (2010). Structure of the human BK channel Ca^{2+} -activation apparatus at 3.0 Å resolution. *Science* 329, 182–186. doi: 10.1126/science.1190414
- Zeng, X. H., Xia, X. M., and Lingle, C. J. (2005). Divalent cation sensitivity of BK channel activation supports the existence of three distinct binding sites. *J. Gen. Physiol.* 125, 273–286. doi: 10.1085/jgp.200409239
- Zeydanli, E. N., Kandilci, H. B., and Turan, B. (2011). Doxycycline ameliorates vascular endothelial and contractile dysfunction in the thoracic aorta of diabetic rats. *Cardiovasc. Toxicol.* 11, 134–147. doi: 10.1007/s12012-011-9107-1
- Zhang, D. M., He, T., Katusic, Z. S., Lee, H. C., and Lu, T. (2010a). Muscle-specific F-box only proteins facilitate BK channel β 1 subunit downregulation in vascular smooth muscle cells of diabetes mellitus. *Circ. Res.* 107, 1454–1459. doi: 10.1161/CIRCRESAHA.110.228361
- Zhang, G., Huang, S. Y., Yang, J., Shi, J., and Yang, X. (2010b). Ion sensing in the RCK1 domain of BK channels. *Proc. Natl. Acad. Sci. U. S. A.* 107, 18700–18705. doi: 10.1073/pnas.1010124107
- Zhang, Z., Li, M., Lu, R., Alioua, A., and Stefani, E. (2014). The angiotensin II type 1 receptor (AT1R) closely interacts with large conductance voltage- and Ca^{2+} -activated K^{+} (BK) channels and inhibits their activity independent of G-protein activation. *J. Biol. Chem.* 289, 25678–25689. doi: 10.1074/jbc.M114.595603
- Zhang, Y., Oltman, C. L., Lu, T., Lee, H. C., and Dellsperger, K. C. (2001). EET homologs potentially dilate coronary microvessels and activate BK(Ca) channels. *Am. J. Physiol. Heart Circ. Physiol.* 280, H2430–H2440. doi: 10.1152/ajpheart.2001.280.6.H2430
- Zhang, Z. Y., Qian, L. L., Wang, N., Miao, L. F., and Ma, X. (2020). Glucose fluctuations promote vascular BK channels dysfunction via PKC α /NF- κ B/MuRF1 signaling. *J. Mol. Cell. Cardiol.* 145, 14–24. doi: 10.1016/j.yjmcc.2020.05.021
- Zheng, H., Whitman, S. A., Wu, W., Wondrak, G. T., and Wong, P. K. (2011). Therapeutic potential of Nrf2 activators in streptozotocin-induced diabetic nephropathy. *Diabetes* 60, 3055–3066. doi: 10.2337/db11-0807
- Zhou, X. B., Arntz, C., Kamm, S., Motejlek, K., and Sausbier, U. (2001). A molecular switch for specific stimulation of the BKCa channel by cGMP and cAMP kinase. *J. Biol. Chem.* 276, 43239–43245. doi: 10.1074/jbc.M104202200
- Zhou, W., Wang, X. L., Kaduce, T. L., Spector, A. A., and Lee, H. C. (2005). Impaired arachidonic acid-mediated dilation of small mesenteric arteries in Zucker diabetic fatty rats. *Am. J. Physiol. Heart Circ. Physiol.* 288, H2210–H2218. doi: 10.1152/ajpheart.00704.2004
- Zhou, W., Wang, X. L., Lamping, K. G., and Lee, H. C. (2006). Inhibition of protein kinase C β protects against diabetes-induced impairment in arachidonic acid dilation of small coronary arteries. *J. Pharmacol. Exp. Ther.* 319, 199–207. doi: 10.1124/jpet.106.106666
- Zhou, X. B., Wulfsen, I., Utku, E., Sausbier, U., and Sausbier, M. (2010). Dual role of protein kinase C on BK channel regulation. *Proc. Natl. Acad. Sci. U. S. A.* 107, 8005–8010. doi: 10.1073/pnas.0912029107
- Zink, M. H., Oltman, C. L., Lu, T., Katakam, P. V., and Kaduce, T. L. (2001). 12-lipoxygenase in porcine coronary microcirculation: implications for coronary vasoregulation. *Am. J. Physiol. Heart Circ. Physiol.* 280, H693–H704. doi: 10.1152/ajpheart.2001.280.2.H693
- Zundel, W., Swiersz, L. M., and Giaccia, A. (2000). Caveolin 1-mediated regulation of receptor tyrosine kinase-associated phosphatidylinositol 3-kinase activity by ceramide. *Mol. Cell. Biol.* 20, 1507–1514. doi: 10.1128/MCB.20.5.1507-1514.2000

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Lu and Lee. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.