

## RESEARCH ARTICLE

# Complete sequences of epidermin and nukacin encoding plasmids from oral-derived *Staphylococcus epidermidis* and their antibacterial activity

Kenta Nakazono<sup>1</sup> , Mi Nguyen-Tra Le<sup>2,3</sup> , Miki Kawada-Matsuo<sup>2,3\*</sup> , Noy Kimheang<sup>2</sup>, Junzo Hisatsune<sup>3,4</sup>, Yuichi Oogai<sup>5</sup>, Masanobu Nakata<sup>5</sup>, Norifumi Nakamura<sup>1</sup>, Motoyuki Sugai<sup>3,4</sup>, Hitoshi Komatsuzawa<sup>2,3</sup>

**1** Department of Oral and Maxillofacial surgery, Field of Maxillofacial Rehabilitation, Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima, Japan, **2** Department of Bacteriology, Hiroshima University Graduate School of Biomedical and Health Sciences, Hiroshima, Japan, **3** Project Research Centre for Nosocomial Infectious Diseases, Hiroshima University, Hiroshima, Japan, **4** Antimicrobial Resistance Research Center, National Institute of Infectious Diseases, Higashi Murayama, Japan, **5** Department of Oral Microbiology, Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima, Japan

 These authors contributed equally to this work.

\* [mmatsuo@hiroshima-u.ac.jp](mailto:mmatsuo@hiroshima-u.ac.jp)



## OPEN ACCESS

**Citation:** Nakazono K, Le MN-T, Kawada-Matsuo M, Kimheang N, Hisatsune J, Oogai Y, et al. (2022) Complete sequences of epidermin and nukacin encoding plasmids from oral-derived *Staphylococcus epidermidis* and their antibacterial activity. PLoS ONE 17(1): e0258283. <https://doi.org/10.1371/journal.pone.0258283>

**Editor:** Jean-San Chia, College of Medicine, National Taiwan University, TAIWAN

**Received:** September 22, 2021

**Accepted:** December 22, 2021

**Published:** January 18, 2022

**Copyright:** © 2022 Nakazono et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All complete plasmid files are available from the NCBI database (accession numbers OK031036 and OK031035).

**Funding:** Research Program on Emerging and Re-emerging Infectious Diseases from the Japan Agency for Medical Research and Development (AMED) under grant number 21fk0108604j0001. Grant in Aid for Scientific Research (C) (Grant No: 21K09858A) from the Ministry of Education,

## Abstract

*Staphylococcus epidermidis* is a commensal bacterium in humans. To persist in the bacterial flora of the host, some bacteria produce antibacterial factors such as the antimicrobial peptides known as bacteriocins. In this study, we tried to isolate bacteriocin-producing *S. epidermidis* strains. Among 150 *S. epidermidis* isolates from the oral cavities of 287 volunteers, we detected two bacteriocin-producing strains, KSE56 and KSE650. Complete genome sequences of the two strains confirmed that they carried the epidermin-harboring plasmid pEpi56 and the nukacin IVK45-like-harboring plasmid pNuk650. The amino acid sequence of epidermin from KSE56 was identical to the previously reported sequence, but the epidermin synthesis-related genes were partially different. The prepeptide amino acid sequences of nukacin KSE650 and nukacin IVK45 showed one mismatch, but both mature peptides were entirely similar. pNuk650 was larger and had an additional seven ORFs compared to pIVK45. We then investigated the antibacterial activity of the two strains against several skin and oral bacteria and found their different activity patterns. In conclusion, we report the complete sequences of 2 plasmids coding for bacteriocins from *S. epidermidis*, which were partially different from those previously reported. Furthermore, this is the first report to show the complete sequence of an epidermin-carrying plasmid, pEpi56.

## Introduction

Staphylococci are classified into two groups, *Staphylococcus aureus* and coagulase -negative staphylococci (CoNS) due to their clinical importance. CoNS are abundant colonizers on the

Culture, Sports, Sciences, and Technology of Japan.

**Competing interests:** The authors have declared that no competing interests exist.

skin and are considered to contribute to the maintenance of skin integrity and homeostasis [1–3]. CoNS assist in immune activity to prevent pathogen colonization by inducing antimicrobial peptides from the epithelium, by direct production of antibacterial factors such as phenol-soluble modulins (PSMs) and bacteriocins [4–6]. Therefore, the colonization of CoNS provides several benefits to the host. However, CoNS are commonly isolated in clinical cultures and considered to be major nosocomial pathogens in humans [7, 8]. CoNS are often isolated from blood and indwelling medical implants such as intravascular catheters and urinary catheters, leading to opportunistic infectious diseases. In addition, most clinical isolates of *Staphylococcus epidermidis* carry the genes encoding for antibiotic resistance and biofilm formation, which significantly challenge current antibiotic therapy [9, 10].

In the oral cavity, oral bacterial flora is composed of a great diversity of bacterial species. Many oral indigenous bacteria, including oral streptococci, are known to produce antimicrobial factors such as bacteriocins and hydrogen peroxide [11–15]. Bacteriocins exhibit a wide range of antimicrobial activity against the bacterial species that make up the oral flora [16]. Therefore, bacteriocins are thought to be involved in the exclusion and symbiosis of other bacteria in the oral cavity. *S. epidermidis* is also found in oral cavity [17, 18]. Some *S. epidermidis* are known to produce antimicrobial peptides known as bacteriocins, including epidermin [19–21], Pep5 [21–23], epilancin K7 [21, 24], epilancin 15X [25, 26], epicidin 280 [27] and Nukacin IVK-45 [28]. These bacteriocins are known to be lantibiotics containing specific amino acids such as lanthionine,  $\beta$ -methyllanthionine, and dehydrated amino acids [11–13]. However, there are no reports about the bacteriocin produced by *S. epidermidis* isolated from the oral cavity.

So far, there have been many reports on bacteriocins produced by oral isolates of streptococcal species [11–13, 16] but very few reports on other oral bacterial species. To understand the meaning of bacteriocins for bacterial flora formation, more information about bacteriocins produced by many oral bacterial species is required. In this study, we focused on the bacteriocins of oral-derived *S. epidermidis* to understand the antibacterial activity against oral and skin bacteria. We examined 150 *S. epidermidis* strains isolated from the oral cavity and investigated their bacteriocin-producing activity. As a result, we found two strains that produced epidermin and nukacin IVK45. We performed the complete-genome analysis of these two strains and identified the plasmids harboring the epidermin or nukacin IVK45-like bacteriocin gene clusters. The nucleotide sequences of these plasmids were not entirely similar to the previously reported sequences. Additionally, we evaluated the antibacterial activity of these two bacteriocins against the skin and oral commensal bacteria.

## Materials and methods

### Bacterial strains and growth conditions

*S. epidermidis* clinical isolates were grown in trypticase soy broth (TSB) (Becton, Dickinson and Company [BD], Franklin Lakes, NJ, USA) at 37°C. The *Staphylococcus aureus* MW2 strain and *braRS*-inactivated mutant were obtained previously [29]. Other bacteria used in this study are listed in Table 1. Staphylococcal strains and *Micrococcus luteus* were grown in TSB at 37°C and 30°C, respectively. Streptococcal strains were grown in TSB at 37°C with 5% CO<sub>2</sub>. *Cutibacterium acnes* was grown on sheep blood agar at 37°C anaerobically. *Corynebacterium* and *Rothia mucilanginosa* were grown at 37°C in R medium and BHI (BD) aerobically, respectively. The composition of R medium is as follows: 1g of bacto peptone (BD), 0.5g of yeast extract (BD), 0.5g of malt extract (BD), 0.5g of casamino acids (BD), 0.2g of beef extract (BD), 0.2g of glycerol, 5mg of Tween 80, 0.1g of MgSO<sub>4</sub> in 100 ml distilled water. When necessary, tetracycline (5  $\mu$ g/ml) was added to the medium.

Table 1. Strains used in this study.

Strains	Character	origin
<i>Staphylococcus epidermidis</i>		
KSE1	Wild type	This study
KSE3	Wild type	This study
KSE56	Wild type	This study
KSE650	Wild type	This study
KSE56-	KSE56 plasmid deleted	This study
KSE650-	KSE650 plasmid deleted	This study
<i>Staphylococcus warneri</i> ISK-1	Wild type	
<i>Staphylococcus hominis</i> JCM31912	Wild type	Riken BRC <sup>1</sup>
<i>Staphylococcus haemolyticus</i> JCM2416	Wild type	Riken BRC <sup>1</sup>
<i>Staphylococcus capitis</i> subsp. <i>capitis</i> JCM2420	Wild type	Riken BRC <sup>1</sup>
<i>Staphylococcus simulans</i> JCM2424	Wild type	Riken BRC <sup>1</sup>
<i>Cutibacterium acnes</i> JCM6425	Wild type	Riken BRC <sup>1</sup>
<i>Corynebacterium accolens</i> JCM8331	Wild type	Riken BRC <sup>1</sup>
<i>Corynebacterium pseudodiphtheriticum</i> JCM1320	Wild type	Riken BRC <sup>1</sup>
<i>Rothia mucilaginosa</i> JCM10910	Wild type	Riken BRC <sup>1</sup>
<i>Micrococcus luteus</i> JCM1464	Wild type	Riken BRC <sup>1</sup>
<i>Streptococcus mutans</i> UA159	Wild type	[30]
<i>Streptococcus sanguinis</i> GTC217	Wild type	Gifu University
<i>Streptococcus salivarius</i> GTC215	Wild type	Gifu University
<i>Streptococcus gordonii</i> JCM12995	Wild type	Riken BRC <sup>1</sup>
<i>Staphylococcus aureus</i>		
COL	Wild type	[31]
RN4220	NCTS8325 derivative	[32]
MW2	clinical strain, methicillin-resistant ( <i>mecA</i> +)	[33]
ΔTCS16	<i>braRS</i> inactivation in MW2, Tcr2	[29]

1. Japan Collection of Microorganisms.

2. Tetracycline resistance.

<https://doi.org/10.1371/journal.pone.0258283.t001>

## Isolation of *Staphylococcus epidermidis* from the oral cavity

*S. epidermidis* strains were isolated from the oral cavities of 287 volunteers. Saliva collected from the oral cavity was plated on No.110 medium (Eiken Chemical Co. Ltd, Tokyo, Japan) and incubated for 2 days at 37°C. The strains were picked from a single white colony on the agar and further investigated by PCR with specific primers for *S. epidermidis* (forward primer: GGCAAATT TGTGGGTCAAGA, reverse primer: TGGCTAATGGTTTGTACCA). Isolated *S. epidermidis* strains were replated on TSB containing 2% agar (TSA) medium. The isolated strains were then replated again on TSA to pick up a single colony and finally, *S. epidermidis* confirmed by PCR was used in this study. Clinical isolates were designated as KSE strains. Saliva collection and *S. epidermidis* isolation were approved by the Ethical Committee of the Kagoshima University Graduate School of Medical and Dental Sciences (No. 701) and the Ethical Committee for Epidemiology of Hiroshima University (E-1998). Written informed consent was obtained from all participants. All methods were performed in accordance with the approved guidelines and regulations.

## Screening of bacteriocin-producing *S. epidermidis*

To investigate bacteriocin production among *S. epidermidis* strains, we performed a direct assay using *S. aureus* MW2 *braRS* knockout mutant as an indicator strain because this mutant

showed increased susceptibility to several bacteriocins [34]. Overnight cultures of *S. epidermidis* strains were spotted on a TSA plate and cultured at 37°C for 24 h. Then, 3.5 ml of prewarmed half-strength TSB soft agar (1%) containing *braRS* knockout mutant cells ( $10^7$  cells/ml) were poured over the TSA plate. The plates were incubated at 37°C for 24 h. The strains which showed the growth inhibition zones surrounding *S. epidermidis* strain were picked up. The strains were reconfirmed for bacteriocin production by the direct assay again.

### Complete genome sequences of bacteriocin-producing *S. epidermidis* strains

To perform whole-genome sequencing of *S. epidermidis* strains, DNA was extracted from each strain. *S. epidermidis* cells grown overnight in 5 ml TSB were collected and then suspended in 0.5 ml of CS buffer (100 mM Tris-HCl [pH 7.5], 150 mM NaCl, 10 mM EDTA) containing lysostaphin (Sigma-Aldrich, St. Louis, MO, USA) (final concentration: 50 µg/ml) and RNase (Nippon Gene, Tokyo, Japan) (final: 20 µg/ml). After incubation at 37°C for 1 h, proteinase K (Nacalai Tesque, Kyoto, Japan) (final: 150 µg/ml) and SDS (final 1%) were added, followed by incubation at 55°C for 5 h. After treatment with phenol followed by phenol-chloroform, DNA was precipitated by ethanol. Whole-genome sequencing (WGS) of *S. epidermidis* strains was performed using the Illumina MiSeq sequencing platform, followed by annotation with Rapid Annotation using Subsystem Technology (RAST) version 2.0 [35]. After confirming the presence of bacteriocin genes using WGS, long-read sequencing by MinION (Oxford Nanopore Technologies, UK) was carried out to determine the complete sequences of the chromosomes and plasmids of these strains. Hybrid assembly of Illumina short reads and MinION long reads was performed with Unicycler v0.4.8. The complete sequences of plasmids harboring bacteriocin genes were selected, including epidermin-carrying plasmid pEpi56 and nukacin-carrying plasmid pNuk650. Each plasmid was compared with publicly available plasmids or gene clusters, including the *epiY<sup>2</sup>-epiP* gene cluster (X62386), *epiG-epiT<sup>2</sup>* gene cluster (U77778), and pIVK45 (accession number KP702950).

### Accession numbers

The complete plasmids carrying epidermin (pEpi56) and nukacin (pNuk650) have been deposited in the NCBI database under accession numbers OK031036 and OK031035, respectively.

### Identification of epidermin and nukacin KSE650 produced by *S. epidermidis*

To identify the bacteriocin, we purified the bacteriocin from two *S. epidermidis* strains. Overnight cultures (500 ml) of *S. epidermidis* KSE56 and KSE650 were centrifuged at 4,000 x g for 15 min. Macro-Prep cationic resin (1.5 ml) (Bio rad, USA) was added to the supernatant and stirred for 12 h. The resin was collected into an open column, then washed three times with 10 ml of 25 mM ammonium acetate (pH 7.5). To elute the bacteriocin, the resin was treated with 500 µl of 5% acetic acid. This elution was repeated 10 times. After each fraction was evaporated completely, the samples were dissolved in 50 µl of distilled water. Each solution was tested for antibacterial activity against *M. luteus*. Overnight cultures of *M. luteus* (100 µl) were inoculated on TSA plates. Then, 5 µl of each solution was spotted on TSA. After overnight incubation at 37°C, growth inhibition was observed. Samples with antibacterial activity were subjected to HPLC chromatography using an Octadecyl C18 column. After equilibrating the column with 0.1% TFA water, the sample was injected. Thereafter, a linear gradient of 0 to 60% acetonitrile

for 30 min was applied to the column. Each peak was fractionated, and the samples were evaporated, then dissolved with 50  $\mu$ l of distilled water. Subsequently, the antibacterial activity of each fraction was tested with the method above. ESI-MS analysis was performed by LTQ Orbitrap XL (Thermo Fisher Scientific, USA).

### Isolation of the strain curing bacteriocin-encoded plasmid

Plasmid deletion in KSE56 and KSE650 was performed with the method described elsewhere [36]. Overnight cultures of KSE56 or KSE650 were inoculated into 5 ml of fresh TSB and incubated at 37°C with shaking. When the OD<sub>660</sub> reached 0.5, acriflavine was added at a concentration of 25  $\mu$ g/ml. After incubation for 12 h, the culture was diluted and plated on TSA. After 24 h of incubation at 37°C, colonies were picked, replated on TSA and then incubated at 37°C for 24 h. Next, 0.75% soft agar (3.5 ml) containing *Bacillus coagulans* (200  $\mu$ l of overnight culture) was poured on that plate and incubated at 37°C for 24 h. The strains with no inhibitory zone were picked. Finally, PCR was performed using specific primers for *S. epidermidis*-specific genes and bacteriocin genes coding for nukacin KSE650 or epidermin.

### Susceptibility tests

Two methods were used for the evaluation of bacteriocins. A direct assay was performed with a previously described method [34]. An overnight culture of the bacteriocin-producing strain was spotted on a TSA plate and cultured at 37°C for 24 h. Then, 3.5 ml of prewarmed half-strength TSB soft agar (1%) containing indicator bacterial cells ( $10^7$  cells/ml) was poured over the TSA plate. The plates were incubated at 37°C for 16 h. The diameters of the growth inhibition zones surrounding the bacteriocin-producing strains were measured in three directions. Three independent experiments were performed, and the average diameter was calculated.

Another method was to evaluate the minimal antibacterial dose of purified bacteriocins. Purified epidermin and nukacin KSE650 were adjusted to 0.5 mg/ml. The bacteriocin solution underwent 2-fold serial dilutions (2-fold to 128-fold dilution). Then, 3.5 ml of prewarmed half-strength TSB soft agar (1%) containing bacterial cells ( $10^7$  cells/ml) was poured over the TSA plate. Thereafter, 2  $\mu$ l of the bacteriocin solutions with serial dilution (1  $\mu$ g to 0.03  $\mu$ g) were spotted on the plate. After the incubation at 37°C for 16 h, the minimum antibacterial dose for the growth inhibition zones was determined.

### Co-culture of *S. epidermidis* with *M. luteus*

For analysis of the proportion of each bacterium (*S. epidermidis* and *M. luteus*) in co-culture by qPCR, we first set up the method for the calculation of bacterial cell number by qPCR. A single overnight culture of the bacterium was first adjusted to OD<sub>660</sub> = 1.0, and then a 10-fold serial dilution was performed in 500  $\mu$ l of lysis buffer. After heating at 95°C for 15 min, samples were centrifuged at 15,000 x rpm for 10 min. Using the supernatant, qPCR was performed with the respective specific primers. For *S. epidermidis*, the forward and reverse primers used were GGCAAATTTGTGGGTCAAGA and TGGCTAATGGTTTGTACCA, respectively. For *M. luteus*, the forward and reverse primers were GGGTTGCGATACTGTGAGGT and TTCGGGTGTTACCGACTTTC, respectively. Finally, the linear relationship between bacterial cell number and cut off value (Ct value) was constructed in each bacterium. Overnight cultures of *S. epidermidis* KSE1 (no bacteriocin production), KSE56, KSE650 and *M. luteus* were adjusted to OD<sub>660</sub> = 1.0, and the bacterial culture was diluted to 10-fold. Next, 100  $\mu$ l of *S. epidermidis* culture and *M. luteus* were mixed thoroughly. A small portion (20  $\mu$ l) of mixed culture was spotted on TSA. After overnight incubation at 37°C, the bacterial colonies growing on agar plates were scraped and suspended in 500  $\mu$ l of lysis buffer. After heating at 95°C for 15 min, the



bacterial suspension was centrifuged at 15,000 x rpm for 10 min and the culture supernatant was stocked as the template for quantitative PCR (qPCR). qPCR was performed using appropriate specific primers to determine the cell number of each bacterium in the co-culture samples. Finally, the proportion of 2 bacterial species was determined. Three independent experiments were performed. Post hoc multiple comparisons were made using Tukey's test.

## Results

### Isolation of *S. epidermidis* that produced bacteriocin


From 287 volunteers, 150 *S. epidermidis* strains (52.3%) were isolated from the oral cavity. Among 150 *S. epidermidis* strains, 2 strains showing a clear inhibitory zone against the *S. aureus* MW2 *braRS* inactivated mutant were identified by the direct method (Fig 1).

### Nucleotide sequence of epidermin-encoding plasmid

The size of the entire plasmid, pEpi56, is 64,386 bp, with 81 ORFs (Fig 2A and Table 2). The plasmid contains epidermin synthesis genes (*epiA* coding for epidermin KSE56, modification genes *epiBCD*, processing genes *epiP*, export genes *epiHT*, immunity genes *epiGEF*, and regulatory gene *epiQ*), replication-related genes, and other genes including the genes coding for hypothetical proteins (Table 2). Compared with epidermin-related genes in the Tü3298 strain [19] *epiT*, which codes for an exporter, was intact in pEpi56, while a gene disrupted into two fragments (*epiT'* and *epiT''* or *epiY* and *epiY'*) was found in the Tü3298 strain (Figs 2B and S1). The nucleotide sequence of *epiA* in KSE56 showed 2 mismatches with that of the Tü3298 strain (S2 Fig). However, the amino acid sequence of epidermin KSE56 showed 100% identity with that in the Tü3298 strain.


### Nucleotide sequence of nukacin-encoding plasmid

The size of the entire plasmid, pNuk650, was 26,160 bp, with 29 open reading frames (ORFs). The plasmid contained nukacin KSE650 synthesis genes (*nukA* coding for prepeptide nukacin KSE650, posttranslational modification enzyme genes *nukM*, processing and secretion transporter genes *nukT*, and immunity protein genes *nukFEGH*), replication-related genes, and



**Fig 1. Direct assay of bacteriocin-producing *S. epidermidis* against *braRS*-inactivated *S. aureus*.** The antibacterial activity of bacteriocin-producing *S. epidermidis* was evaluated by the direct assay using *S. aureus* MW2 *braRS*-inactivated mutant.

<https://doi.org/10.1371/journal.pone.0258283.g001>



**Fig 2. Gene map of the epidermin-carrying plasmid in KSE56.** (a) Epidermin-encoding plasmid from KSE56 (pEpi56). ORFs are shown as arrows, indicating the orientation of transcription. The arrow numbers indicate the ORF number displayed in Table 2. Colors indicate the classification of gene function. (b) Bacteriocin-coding region (KSE56 epidermin). The bacteriocin-coding region from pEpi56 was compared with pTu32 *epiP*-Y<sup>+</sup> (accession number X62386) and pTu32 *epiT*-G (accession number U77778). Striped blue arrows indicate truncated *epiT*.

<https://doi.org/10.1371/journal.pone.0258283.g002>

other genes including genes coding for hypothetical proteins (Fig 3A and Table 3). Compared to the plasmid pIVK45 (21,840 bp), which carried the gene coding for nukacin IVK45 [28] pNuk650 was larger with a higher number of ORFs (Fig 3A). The amino acid sequence of nukacin KSE650 showed similarity to nukacin IVK45 with one mismatch at the 4<sup>th</sup> position, but displayed lower similarity to nukacin ISK-1 with 10 mismatches [36, 37] (Fig 3B). The mature peptide of nukacin KSE650 showed a perfect match with nukacin IVK45 and 5 mismatches with nukacin ISK-1.

### Identification of epidermin KSE56 and nukacin KSE650

Epidermin KSE56 and nukacin KSE650 were purified from the culture supernatant of KSE56 and KSE650, respectively. After applying the sample purified by Macro Prep resin to Octadecyl C18 column, peak fractions in both samples were collected and each peak fraction was checked

Table 2. Genes in pEpi56.

No.	Location (bp)	Size (aa) <sup>a</sup>	Translation signal <sup>b</sup>	Homologue as determined by BLAST and/or FASTA					
				Source	Description(s)	Identity (%)	Overlap (aa) <sup>c</sup>	Accession no.	Note
1	190–1191	333	<u>GAGGTTTTTTATT</u> <b>ATG</b>	<i>S. epidermidis</i>	replication initiator protein A	99	333/338	WP_002498716.1	
2	1423–1983	186	<u>AAGGAGTAATAAAA</u> <b>ATG</b>	<i>S. epidermidis</i>	TIGR00730 family Rossman fold protein	99	186/186	WP_158171994.1	
3	2300–2515	71	-	<i>S. epidermidis</i>	hypothetical protein	67	48/78	MBM0824966.1	
4	2889–3014	41	<u>GGAGAATAATTAATAAACCCGTTACAAAATA</u> <u>AGCAATATCTATAAGTTTTTTAAAAATAAAA</u> <u>ATTCTAAAAATATGTAAGT</u> <b>ATG</b>	<i>S. epidermidis</i> SK135	ATP-binding cassette domain-containing protein	100	41/41	EFA87131.1	
5	3507–3695	62	<u>GAGTTAGACCAATAAATGAAACGAAAAAAC</u> <u>AATTG</u> <b>TTG</b>	<i>S. epidermidis</i>	hypothetical protein	100	62/62	MBC8789835.1	
6	4346–4513	55	<u>GGAGGCATTGT</u> <b>CATG</b>	<i>S. epidermidis</i>	hypothetical protein	100	55/55	WP_002498713.1	
7	4819–5685	288	<u>GGAGTGATATAT</u> <b>ATG</b>	<i>S. epidermidis</i>	RepB family plasmid replication initiator protein	99	287/288	WP_203085279.1	
8	5791–5934	47	<u>GGAGACATAAAAAGTT</u> <b>ATG</b>	<i>S. epidermidis</i>	hypothetical protein	100	47/47	WP_002498711.1	
9	6397–7026	209	<u>GAGTAAT</u> <b>CATG</b>	<i>S. epidermidis</i>	ABC transporter, ATP-binding protein	100	209/209	EJD97739.1	
10	7029–9071	680	<u>AGGTATTTATACAT</u> <b>ATG</b>	<i>S. epidermidis</i> NIHLM040	bacteriocin-associated integral membrane protein	100	680/680	EJD97738.1	
11	9165–9557	130	<u>GGAGGATTAAGTT</u> <b>GATG</b>	<i>S. epidermidis</i> NIHLM040	bacteriocin, lactococcin 972 family	100	130/130	EJD97736.1	
12	9743–10105	120	<u>GAGAATTATACAAAA</u> <b>ATG</b>	<i>S. epidermidis</i>	DUF3139 domain-containing protein	100	120/120	WP_002498706.1	
13	10304–10669	121	<u>GAGGACATACATTAGATATTTGG</u> <b>TTG</b>	<i>S. epidermidis</i> NIHLM040	IS431mec, transposase	100	121/121	EJD97734.1	
14	10732–10884	50	<u>GGAGTCTTCTGT</u> <b>ATG</b>	<i>S. epidermidis</i> NIHLM040	hypothetical protein	100	50/50	EJD97733.1	
15	11171–12556	461	<u>GAGGTGCTAT</u> <b>ATG</b>	<i>S. epidermidis</i> NIHLM040	putative epidermin leader peptide-processing serine protease EpiP	100	461/461	EJD97732.1	<i>epiP</i>
16	12567–13184	205	<u>GGAATAAAA</u> <b>ATG</b>	<i>S. epidermidis</i>	winged helix family transcriptional regulator	100	205/205	MBM0752529.1	<i>epiQ</i>
17	13181–13726	181	<u>GGAGGAATAAGAT</u> <b>ATG</b>	<i>S. epidermidis</i> NIHLM040	epidermin decarboxylase	100	181/181	EJD97730.1	<i>epiD</i>
18	13742–14992	416	<u>GGATGGTT</u> <b>GTTG</b>	<i>S. epidermidis</i> NIHLM040	putative epidermin biosynthesis protein EpiC	100	416/416	EJD97729.1	<i>epiC</i>

(Continued)



Table 2. (Continued)

No.	Location (bp)	Size (aa) <sup>a</sup>	Translation signal <sup>b</sup>	Homologue as determined by BLAST and/or FASTA					
				Source	Description(s)	Identity (%)	Overlap (aa) <sup>c</sup>	Accession no.	Note
19	14985–17945	986	<u>GAGGTGAAATAGAA</u> <b>TTG</b>	<i>S. epidermidis</i> NIHLM040	thiopeptide-type bacteriocin biosynthesis domain protein	100	986/986	EJD97728.1	<i>epiB</i>
20	18011–18169	52	<u>AGGAGTGTTTAAA</u> <b>ATG</b>	<i>S. epidermidis</i> NIHLM040	lantibiotic epidermin	100	52/52	EJD97726.1	<i>epiA</i>
21	18419–19969	516	<u>GGACTAATATTGAGT</u> <b>TTG</b>	<i>S. epidermidis</i>	ABC transporter ATP-binding protein/permease	100	516/516	WP_002498696.1	<i>epiT'</i>
22	19985–20977	330	<u>GAGATAAGGGAGATATA</u> <b>ATG</b>	<i>S. epidermidis</i>	YdcF family protein	100	330/330	WP_032605946.1	<i>epiH</i>
23	21136–21831	231	<u>GGAGGAATAATTCT</u> <b>TTG</b>	<i>S. epidermidis</i>	lantibiotic protection ABC transporter ATP-binding protein	100	231/231	WP_002498693.1	<i>epiF</i>
24	21833–22597	254	<u>GGAATAATA</u> <b>ATG</b>	<i>S. epidermidis</i>	lantibiotic immunity ABC transporter MutE/EpiE family permease subunit	100	254/254	WP_002498692.1	<i>epiE</i>
25	22587–23279	230	<u>GGAATATAA</u> <b>ATG</b>	<i>S. epidermidis</i>	epidermin immunity protein F	100	230/230	WP_002498691.1	<i>epiG</i>
26	23432–24034	200	<u>GAGGTGGAAATCA</u> <b>ATG</b>	<i>S. epidermidis</i> NIHLM040	putative transposon DNA-invertase Bin3	100	200/200	EJD97719.1	
27	24455–26071	538	<u>GGAGGAAGAAAAA</u> <b>ATG</b>	<i>S. epidermidis</i> NIHLM040	ABC transporter, ATP-binding protein	100	538/538	EJD97718.1	
28	26621–27463	280	<u>GGAGCATTAATT</u> <b>ATG</b>	<i>S. epidermidis</i>	hypothetical protein	100	280/280	WP_002498688.1	
29	27952–28383	143	<u>AAGGAGTCTTCTGT</u> <b>ATG</b>	<i>S. epidermidis</i> NIHLM040	IS431mec, transposase family protein	100	143/143	EJD97715.1	
30	28376–28627	83	<u>AGGCACCTTCAACGAAGGTAGCA</u> <b>ATG</b>	<i>S. epidermidis</i> NIHLM040	IS431mec, transposase family protein	100	83/83	EJD97714.1	
31	28733–29455	240	<u>GGAGTGTAAGCT</u> <b>TTG</b>	<i>S. epidermidis</i>	peptide ABC transporter permease	100	240/240	WP_002498749.1	
32	29472–30107	211	<u>GGAGCTGTAAACA</u> <b>TTG</b>	<i>S. epidermidis</i> NIHLM040	ABC transporter, ATP-binding protein	100	211/211	EJD97793.1	
33	30389–30484	31	<u>GGAGAGATTAA</u> <b>ATG</b>	<i>S. epidermidis</i> NIHLM040	hypothetical protein	100	31/31	EJD97792.1	
34	30495–30665	56	<u>AGGTTAATTTT</u> <b>ATG</b>	<i>S. epidermidis</i>	hypothetical protein	100	56/56	TID00490.1	
35	30897–31535	212	<u>AGGTTCAAGATGAAAACAAAGAA</u> <b>ATG</b>	<i>S. epidermidis</i> NIHLM040	hypothetical protein	100	212/212	EJD97791.1	
36	31698–32063	121	<u>GAGGAGAGAACTTTTAAA</u> <b>ATG</b>	<i>S. epidermidis</i> NIHLM040	hypothetical protein	100	121/121	EJD97790.1	
37	32230–32406	58	<u>GGAGTGATTTA</u> <b>ATG</b>	<i>S. epidermidis</i> NIHLM040	hypothetical protein	100	58/58	EJD97789.1	

(Continued)

Table 2. (Continued)

No.	Location (bp)	Size (aa) <sup>a</sup>	Translation signal <sup>b</sup>	Homologue as determined by BLAST and/or FASTA					
				Source	Description(s)	Identity (%)	Overlap (aa) <sup>c</sup>	Accession no.	Note
38	32573–34183	536	<u>GG</u> AAGGATTATT <b>ATG</b>	<i>S. epidermidis</i>	DNA mismatch repair protein MutS	100	536/536	WP_002498743.1	
39	34762–35058	98	<u>GG</u> ATTG <b>AA</b> TG	<i>S. epidermidis</i>	replication initiation protein	100	98/98	MBF2337202.1	
40	35232–35510	92	<u>GG</u> AGAGATT <b>AA</b> ATG	<i>S. epidermidis</i>	hypothetical protein	100	92/92	WP_002498740.1	
41	35521–35691	56	<u>GG</u> ATTT <b>TT</b> ATG	<i>S. epidermidis</i>	hypothetical protein	100	56/56	WP_099800689.1	
42	36232–36369	45	<u>GG</u> AG ACATAAGAAGGT <b>ATG</b>	<i>S. epidermidis</i>	hypothetical protein	100	45/45	MBM6015004.1	
43	36517–36732	71	<u>GG</u> AAATGACACATCTTAAATCGACATATTC <b>CA</b> AAA ATATGTTTAGAATACTGGTT <b>AC</b> ATG	<i>S. epidermidis</i>	hypothetical protein	100	71/71	WP_002498738.1	
44	37358–37726	122	<u>GAG</u> ACGTCT <b>ATG</b>	<i>S. epidermidis</i> NIHLM040	hypothetical protein	100	122/122	EJD97781.1	
45	37880–38335	151	-	<i>S. epidermidis</i>	putative plasmid recombination enzyme	100	151/151	TID00443.1	
46	38651–38905	84	<u>GG</u> AGTTCCTTT <b>AA</b> ATG	<i>S. epidermidis</i>	hypothetical protein	100	84/84	EJD97779.1	
47	38927–39067	46	<u>GG</u> AAGATGAAATAGTCCT <b>AA</b> TG	<i>S. epidermidis</i>	hypothetical protein	100	46/46	WP_151520775.1	
48	39102–40481	459	<u>GG</u> AGGTATGATAG <b>ATG</b>	<i>S. epidermidis</i> NIHLM040	drug resistance MFS transporter, drug:H + antiporter-2 family	100	459/459	EJD97777.1	
49	40630–41637	335	<u>GG</u> AGCGATGG <b>AA</b> ATG	<i>S. epidermidis</i>	tryptophan—tRNA ligase	100	335/335	WP_002498732.1	
50	41862–42590	242	<u>AAG</u> GAGAATA <b>AA</b> CAATG	<i>S. epidermidis</i> NIHLM040	ABC transporter permease	100	242/242	EJD97775.1	
51	42594–43457	287	<u>AAG</u> GAGAATA <b>AA</b> ATG	<i>S. epidermidis</i> NIHLM040	ABC transporter, ATP-binding protein	100	287/287	EJD97774.1	
52	43704–44525	273	<u>GG</u> AGGATTT <b>TT</b> ATG	<i>S. epidermidis</i> NIHLM040	transcriptional regulator, LysR family	100	273/273	EJD97773.1	
53	44678–45817	379	<u>GAG</u> GATGGGATA <b>AA</b> TATG	<i>S. epidermidis</i> NIHLM040	MFS transporter	100	379/379	EJD97772.1	
54	46236–46613	125	<u>GG</u> AAAAGAGT <b>AA</b> ATG	<i>S. epidermidis</i> NIHLM040	hypothetical protein	96	125/125	EJE04311.1	
55	46649–47338	229	<u>GG</u> AGACGATA <b>AT</b> GTG	<i>S. epidermidis</i> NIHLM040	ABC transporter, ATP-binding protein	100	229/229	EJD97770.1	
56	47346–48107	253	<u>GG</u> AGGAATGAAGCA <b>AT</b> TATG	<i>S. epidermidis</i>	ABC transporter permease	99	253/253	WP_002503830.1	
57	48465–48857	130	-	<i>S. epidermidis</i> NIHLM040	hypothetical protein	100	130/130	EJD97768.1	
58	48948–49919	323	<u>GG</u> AGAA <b>AT</b> TATG	<i>S. epidermidis</i>	DUF418 domain-containing protein	99	323/323	WP_095694513.1	
59	49974–50108	44	<u>GG</u> AAGG <b>AT</b> TG	<i>S. epidermidis</i>	hypothetical protein	100	44/44	EFA87101.1	

(Continued)

Table 2. (Continued)

No.	Location (bp)	Size (aa) <sup>a</sup>	Translation signal <sup>b</sup>	Homologue as determined by BLAST and/or FASTA					
				Source	Description(s)	Identity (%)	Overlap (aa) <sup>c</sup>	Accession no.	Note
60	50567–50722	51	-	<i>S. epidermidis</i>	hypothetical protein	100	51/51	MBC2926404.1	
61	51633–52454	273	<u>AGGTGTGATTTAA<b>ATG</b></u>	<i>S. epidermidis</i>	relaxase MobL	99	273/273	WP_161382396.1	
62	52466–52849	127	<u>GGAGGAATAAA<b>ATG</b></u>	<i>S. epidermidis</i> NIHLM040	hypothetical protein	100	127/127	EJD97765.1	
63	52851–53129	92	<u>GGAAATGATTTTT<b>TTG</b></u>	<i>S. epidermidis</i> NIHLM040	hypothetical protein	100	92/92	EJD97764.1	
64	54078–54224	48		<i>S. epidermidis</i>	hypothetical protein	100	48/48	WP_002456268.1	
65	54621–54800	59	<u>GGAGGCTTATAC<b>ATG</b></u>	<i>S. epidermidis</i> NIHLM040	CsbD family protein	100	59/59	EJD97762.1	
66	54833–55231	132	<u>GAGGTGTTTGTAT<b>ATG</b></u>	<i>S. epidermidis</i>	YolD-like family protein	100	132/132	WP_002498728.1	
67	55394–55651	85	-	<i>S. epidermidis</i> NIHLM040	prevent-host-death family protein	100	85/85	EJD97760.1	
68	55651–55917	88	-	<i>S. epidermidis</i> NIHLM040	addiction module toxin, Txe/YoeB family	100	88/88	EJD97759.1	
69	55934–56104	56	<u>GGAGGACTCGTTA<b>ATG</b></u>	<i>S. epidermidis</i>	hypothetical protein	100	56/56	KAB2267008.1	
70	56465–56689	74		<i>S. epidermidis</i>	putative glycoside hydrolase	100	74/74	QRX38739.1	
71	57190–57546	118	<u>GGAGGTTGTATGT<b>ATG</b></u>	<i>S. epidermidis</i> NIHLM040	hypothetical protein	100	118/118	EJD97756.1	
72	57860–58408	182	-	<i>S. epidermidis</i> NIHLM040	putative resolvase	100	182/182	EJD97755.1	
73	59658–60926	422	<u>GGAGAATTTAATA<b>ATG</b></u>	<i>S. epidermidis</i>	penicillin-binding protein PBP4	99	422/422	WP_002498725.1	
74	61202–61603	133	-	<i>S. epidermidis</i>	transposase DNA-binding domain protein	100	133/133	TID00494.1	
75	61744–61926	60	<u>GAGTCGTTTAG<b>ATG</b></u>	<i>S. epidermidis</i>	transposase	98	60/60	WP_203079065.1	
76	61958–62188	76	<u>GAGGTGTATTGAC<b>ATG</b></u>	<i>S. epidermidis</i> NIHLM040	hypothetical protein	99	76/76	EJD97751.1	
77	62255–62407	50	<u>GGAGGAATTTAA<b>ATG</b></u>	<i>S. epidermidis</i> NIHLM040	hypothetical protein	100	50/50	EJD97750.1	
78	62434–62595	53	<u>GGAGGCGGGAAA<b>ATG</b></u>	<i>S. epidermidis</i>	BH0509 family protein	100	53/53	EJD97749.1	
79	62670–62909	79	<u>GGAGGAAGATA<b>ATG</b></u>	<i>S. epidermidis</i>	hypothetical protein	100	79/79	WP_002498719.1	
80	63024–63272	82	<u>GGAGGTATCAAGGTT<b>ATG</b></u>	<i>S. epidermidis</i>	CopG family transcriptional regulator	100	82/82	MBM0752797.1	
81	63390–64280	296	-	<i>S. epidermidis</i>	ParA family protein	100	268/296	WP_002498717.1	

<sup>a</sup> aa, amino acids.

<sup>b</sup> Bold letters indicate start codons. Underlines indicate putative ribosome binding sites complementary to the 3' end of the 16s rRNA.

<sup>c</sup> Overlap is indicated as the number of overlapping amino acids/total number of amino acids.

<https://doi.org/10.1371/journal.pone.0258283.t002>



Table 3. Genes in pNuk650.

No.	Location (bp)	Size (aa) <sup>a</sup>	Translation signal <sup>b</sup>	Homologue as determined by BLAST and/or FASTA				
				Source	Description(s)	Identity (%)	Overlap (aa) <sup>c</sup>	Accession no.
1	413–541	42	GGAAAAGATATCC <b>ATG</b>	<i>S. epidermidis</i>	RepB (pAQZ2)	83	42/42	AZL87916
2	680–850	56	-	<i>S. epidermidis</i>	replication protein	91	56/56	WP_194378762
3	976–1911	311	GGAAAGAGGTTTATATT <b>ATG</b>	<i>S. epidermidis</i>	replication initiator protein A	100	311/311	WP_194378689
4	2467–3261	264	AGGAGGTATTATT <b>TTG</b>	<i>S. epidermidis</i>	ParA family protein	100	264/264	WP_172686110
5	3258–3467	69	GAGGGTGT <b>GTG</b>	<i>S. epidermidis</i>	plasmid replication associated protein, putative transcriptional regulator	98	66/69	AKQ51589
6	3821–3994	57	AGGGGGTATTATA <b>ATG</b>	<i>S. epidermidis</i> (pIVK45)	NukA	98	57/57	AKQ51579
7	4068–4250	60	AGGTACGCGTTTTTAAATTGTATAT <b>ATG</b>	<i>S. epidermidis</i>	transposase family protein	92	38/60	MBV5159007
8	4256–4393	45	GAGACC <b>ATG</b>	<i>S. epidermidis</i>	hypothetical protein	100	45/45	WP_194378692
9	4605–4844	79	-	<i>S. epidermidis</i>	transposase	100	74/79	WP_172686114
10	5583–6326	247	GAGTGAATTAT <b>ATG</b>	<i>S. epidermidis</i>	LytTR family transcriptional regulator DNA-binding domain-containing protein	100	247/247	WP_194378694
11	6570–9323	917	AGGAGAGGTTGTATAT <b>ATG</b>	<i>S. epidermidis</i> (pIVK45)	NukM	100	917/917	AKQ51580
12	9345–11429	694	AGGTGAATACAA <b>TTG</b>	<i>S. epidermidis</i> (pIVK45)	NukT	99	694/694	KP702950
13	11442–12350	302	AGGAGGTTCAAT <b>TTATG</b>		NukF	99	302/302	AKQ51583
14	12351–13103	250	GGAAAGGAATATTTATA <b>AAATG</b>	<i>S. epidermidis</i> (pIVK45)	NukE	99	250/250	AKQ51582
15	13100–13837	245	AAGGAGAGATTTAT <b>CTTG</b>	<i>S. epidermidis</i> (pIVK45)	NukG	88	245/245	AKQ51591
16	13844–14122	92	GAGGATTAATAACTA <b>ATG</b>	<i>S. epidermidis</i> (pIVK45)	NukH	100	92/92	AKQ51584
17	14444–14623	59	-	<i>S. epidermidis</i>	replication initiator protein A, partial	88	59/272	WP_064595943
18	14790–14930	46	GGATAACAAAATAACATCAACACAA <b>ATGTCACG</b> ATTTCAATAATATAGC <b>ATG</b>	<i>S. epidermidis</i>	hypothetical protein	98	46/46	WP_172686106
19	15014–15157	47	GGAAATGATAAATTC <b>AACTTTTCTTCCGATC</b> ATTAATAAAA <b>ATAAATG</b>		no significant similarity found			
20	15425–16423	332	TAAGGTGTCGAATCTAAATA <b>AAACTGGGGGC</b> TTTTTT <b>ATG</b>	<i>S. epidermidis</i>	protein rep	98	332/332	WP_145461985
21	17100–17483	127	AGGGGTTTTTTT <b>ATG</b>	<i>S. epidermidis</i> IS-K	bacterial transcription activator, effector-binding domain protein	99	127/127	EID36019
22	17957–18664	235	GAGAGGTGTTTTTTTATGTCTGGTGAACAGT AGTATATAGAA <b>ATG</b>	<i>S. epidermidis</i>	RepB family plasmid replication initiator protein	100	235/235	WP_194378685
23	18712–19323	203	AGGAGTAGTTT <b>ATG</b>	<i>S. epidermidis</i>	helix-turn-helix domain-containing protein	99	203/203	WP_194378686
24	19890–20699	269	GGAGAGAAATATATAT <b>TTG</b>	<i>S. epidermidis</i>	CPBP family intramembrane metalloprotease	100	269/269	WP_168429436
25	20725–21039	104	GAGGTGTAAAA <b>ATG</b>	<i>S. epidermidis</i>	helix-turn-helix domain-containing protein	99	104/104	WP_002455864
26	21312–22928	538	AGGATATT <b>ATG</b>	<i>S. epidermidis</i>	MutS family DNA mismatch repair protein	99	538/538	WP_194378687
27	23374–25119	581	AGGTGAAGTTAAA <b>AGTG</b>	<i>S. epidermidis</i>	AIPR family protein	100	581/581	WP_194378688

(Continued)



Table 3. (Continued)

No.	Location (bp)	Size (aa) <sup>a</sup>	Translation signal <sup>b</sup>	Homologue as determined by BLAST and/or FASTA				
				Source	Description(s)	Identity (%)	Overlap (aa) <sup>c</sup>	Accession no.
28	25145–25853	202	<u>GGAATCA</u> <b>ATG</b>	<i>S. epidermidis</i> (pIVK45)	Sin recombinase	100	202/202	AKQ51586
29	25976–26077	33	<u>AAGGAGGAATACT</u> <b>ATG</b>	<i>S. epidermidis</i>	NAD-dependent epimerase/dehydratase family protein	100	33/33	WP_172686124


<sup>a</sup> aa, amino acids.

<sup>b</sup> Bold letters indicate start codons. Underlines indicate putative ribosome binding sites complementary to the 3' end of the 16s rRNA.

<sup>c</sup> Overlap is indicated as the number of overlapping amino acids/total number of amino acid.


<https://doi.org/10.1371/journal.pone.0258283.t003>

activity (>20 mm diameter inhibitory zone) against *M. luteus*, and an activity (>5 mm diameter) against *R. mucilaginosa*, *C. pseudodiphtheriticum*, *S. haemolyticus*, *S. captis*, *S. hominis*, *S. simulans*, and *S. saprophyticus*. KSE56 also showed an antibacterial activity against *S. epidermidis* without bacteriocin production (KSE1, 10, 12, 16), plasmid-curing KSE56 and plasmid-



**Fig 4. Purification of epidermin and nukacin KSE650 by reverse phase-HPLC and mass determination by ESI-MS.** (a) RP-HPLC chromatogram of epidermin and nukacin KSE650. The arrow shows the peak corresponding to epidermin (upper) or nukacin KSE650 (lower). (b) Mass determination of epidermin (upper) or nukacin KSE650 (lower) by ESI-MS. Several isotopic peaks in each mass/charge (m/z) state.

<https://doi.org/10.1371/journal.pone.0258283.g004>



**Fig 5. Antibacterial activity of KSE56, KSE650, and their plasmid-deleted strains.** Direct assays were performed using KSE56, KSE650, and their plasmid-deleted strains. *S. hominis* was used as an indicator strain.

<https://doi.org/10.1371/journal.pone.0258283.g005>

curing KSE650. The inhibitory zone was not observed in *S. epidermidis* KSE56, *S. epidermidis* KSE650, *C. accolens*, *S. warneri* ISK-1, and *S. aureus* strains. Regarding oral streptococci, KSE56 showed a strong activity against *S. salivarius* and *S. gordonii*, and modest activity against *S. mutans* and *S. sanguinis*.

The nukacin KSE650-producing strain KSE650, showed strong antibacterial activity (>20 mm diameter) against *M. luteus*, *S. capitis*, and *S. simulans*, and an activity (>5 mm diameter) against *C. pseudodiphtheriticum*, *R. mucilaginosa*, *S. haemolyticus*, *S. hominis*, and *S. saprophyticus*. KSE650 also showed an antibacterial activity against *S. epidermidis* without bacteriocin production (KSE1, 10, 12, 16), plasmid-curing KSE56 and plasmid-curing KSE650. The inhibitory zone was not observed in *S. epidermidis* KSE56, *S. epidermidis* KSE650, *C. accolens*, *S. warneri* ISK-1, and *S. aureus* strains. Regarding oral streptococci, KSE650 showed activity against *S. salivarius* and *S. gordonii*, and no activity against *S. mutans* and *S. sanguinis*. Compared to the nukacin ISK-1-producing *S. warneri* strain, *S. warneri* showed stronger activity against commensal and oral bacteria except for *S. capitis* and *S. simulans*. Notably, *S. warneri* ISK-1 showed activity against the *S. aureus* strain.

We also checked the antibacterial activity using purified epidermin and nukacin KSE650 (Table 5). The antibacterial pattern against each bacterium was similar to the results of the direct assay.

### Co-culture of *S. epidermidis* with *M. luteus*

Co-cultures of *S. epidermidis* KSE1 (bacteriocin negative), KSE56, and KSE650 with *M. luteus* JCM1464 were analyzed. *M. luteus* was utilized as an indicator bacterium in co-culture assay because in the direct method, KSE56 and KSE650 showed a significant antibacterial effect against *M. luteus*. In co-culture with *M. luteus*, the proportion of *S. epidermidis* KSE1 was 46.2%, while the proportions of KSE56 and KSE650 were 70.4% and 79.8%, respectively (Fig 6).

### Discussion

In this study, we tried to isolate *S. epidermidis* strains that produced bacteriocin. We used the *S. aureus* MW2 *braRS*-inactivated mutant as the indicator strain for screening. We previously

**Table 4. Antibacterial activity of KSE56 and KSE650 against various bacterial species.**

Indicator strains	Halo size (mm)		
	KSE56	KSE650	<i>S. warneri</i>
<i>Corynebacterium pseudodiphtheriticum</i> JCM1320	10.0±0.8	10.7±0.5	11.7±0.5
<i>Corynebacterium accolens</i> JCM8331	-	-	11.3±0.5
<i>Micrococcus luteus</i> JCM1464	31.7±1.2	27.0±0	33.0±0
<i>Rothia mucilaginosa</i> JCM10910	8.7±0.5	8.0±0	13.0±0
<i>Cutibacterium acnes</i> JCM6425	15.0±0.8	-	-
<i>Staphylococcus haemolyticus</i> JCM2416	11.7±0.6	13.3±0.5	16.0±0.8
<i>Staphylococcus capitis</i> JCM2420	11.3±0.6	27.3±0.5	17.3±0.5
<i>Staphylococcus simulans</i> JCM2424	13.7±0.6	28.7±0.5	22.7±0.5
<i>Staphylococcus saprophyticus</i> JCM20595	13.0±0	12.3±0.5	13.3±0.5
<i>Staphylococcus hominis</i> JCM31912	15.3±0.6	16.3±0.5	21.7±0.5
<i>Staphylococcus epidermidis</i> KSE1	12.3±0.5	7.0±0.8	N.D. <sup>2</sup>
<i>Staphylococcus epidermidis</i> KSE10	12.0±0	7.3±0.5	N.D.
<i>Staphylococcus epidermidis</i> KSE12	17.0±0.8	9.7±0.5	N.D.
<i>Staphylococcus epidermidis</i> KSE16	14.3±0.5	8.7±0.5	N.D.
<i>Staphylococcus epidermidis</i> KSE56	-	-	-
<i>Staphylococcus epidermidis</i> KSE650	-	-	-
<i>Staphylococcus epidermidis</i> KSE56 plasmid-deleted	20.3±0.5	11.3±0.5	N.D.
<i>Staphylococcus epidermidis</i> KSE650 plasmid-deleted	11.0±0	11.7±0.5	N.D.
<i>Staphylococcus warneri</i> ISK-1	-	-	-
<i>Staphylococcus aureus</i> MW2	-	-	11.3±0.5
<i>Staphylococcus aureus</i> COL	-	-	11.0±0
<i>Staphylococcus aureus</i> RN4220 (MSSA)	-	-	10.7±0.5
<i>Streptococcus mutans</i> UA159	15.0±0.8	-	-
<i>Streptococcus sanguinis</i> GTC217	12.0±0	-	10.3±0.9
<i>Streptococcus salivarius</i> GTC215	27.7±0.5	12.3±0.5	18.3±0.5
<i>Streptococcus gordonii</i> JCM12995	29.0±0	17.0±0	23.0±0

"-" and "N.D." represent "no inhibitory zone" and "Not determined", respectively.

<https://doi.org/10.1371/journal.pone.0258283.t004>

reported that BraRS was involved in resistance to several bacteriocins including nisin A, nukacin ISK-1 and bacitracin [34]; therefore, a *braRS*-inactivated mutant increased susceptibility to these bacteriocins. Nisin A and nukacin ISK-1 are lantibiotics that act against lipid II molecules, which are responsible for cell wall biosynthesis, and subsequently, form a pour complex [38]. In addition, it was reported that many gram-positive bacteria, including staphylococci, streptococci, bacilli, lactococci and enterococci, produced lantibiotics that bind to lipid II [12, 19–27, 39, 40]. Therefore, the *braRS*-inactivated mutant is a good indicator strain to screen lipid II-binding lantibiotics. Finally, we identified 2 strains that produce epidermin and nukacin IVK45-like bacteriocins. Whole genome analysis of the 2 strains revealed that both genes were located on the plasmids (S2A and 4 Figs).

Epidermin was first identified in the *S. epidermidis* Tü3298 strain [19, 41]. In the Tü3298 strain, epidermin is located on the plasmid, pTu32. Recently, the whole genome sequence of the Tü3298 strain was determined [42], but the entire plasmid sequence of pEpi56 was not reported. Therefore, our study is the first to report the complete nucleotide sequence of epidermin harboring plasmids. Additionally, the epidermin-producing strain identified in this study was the second strain, following the Tü3298 strain. The nucleotide sequence of the *epiA* coding epidermin showed 2 mismatches between the two strains, but the amino acid sequence was


**Table 5. Minimum antibacterial dose of purified epidermin and nukacin KSE650.**

Indicator strains	Minimum antibacterial dose ( $\mu\text{g}$ )	
	Epidermin	Nukacin KSE650
<i>Corynebacterium pseudodiphtheriticum</i> JCM1320	2	2
<i>Corynebacterium accolens</i> JCM8331	> 2	> 2
<i>Micrococcus luteus</i> JCM1464	< 0.03	< 0.03
<i>Rothia mucilaginosa</i> JCM10910	0.5	1
<i>Staphylococcus haemolyticus</i> JCM2416	0.125	0.25
<i>Staphylococcus capitis</i> JCM2420	0.125	< 0.03
<i>Staphylococcus simulans</i> JCM2424	0.125	< 0.03
<i>Staphylococcus saprophyticus</i> JCM20595	0.5	0.25
<i>Staphylococcus hominis</i> JCM31912	0.06	0.25
<i>Staphylococcus epidermidis</i> KSE1	0.5	1
<i>Staphylococcus aureus</i> MW2	2	2
<i>Streptococcus mutans</i> UA159	1	> 2
<i>Streptococcus sanguinis</i> GTC217	0.5	1
<i>Streptococcus salivarius</i> GTC215	0.06	0.25
<i>Streptococcus gordonii</i> JCM12995	0.06	0.125

<https://doi.org/10.1371/journal.pone.0258283.t005>


similar. When the epidermin synthesis genes were compared between the 2 strains, *epiT* showed a significant difference (Fig 2B). *epiT* in KSE56 was intact, while this gene in Tu3298 was disrupted into 2 genes, *epiT'* and *epiT''* in Tü3298.

EpiT is involved in the secretion of the peptide. In previous reports that demonstrated the antibacterial activity of epidermin in Tü3298 [19–21], epidermin was correctly modified and secreted externally. However, Peschel A et al reported that the introduction of intact *gdmT*, encoding the secretion protein for gallidermin, which was close to epidermin in Tü3298, increased the production of epidermin in culture supernatant [43]. Therefore, the secretion activity of *epiT'/T''* is considered to be partial, while the intact *epiT* gene in KSE56 may be responsible for full secretion of the epidermin peptide.



**Fig 6. The proportion of *S. epidermidis* KSE1, KSE56, and KSE650 in co-culture with *M. luteus*.** Co-culture assays were performed according to the method described in the Materials and methods. Post hoc multiple comparisons were made using Tukey's test.

<https://doi.org/10.1371/journal.pone.0258283.g006>



**Fig 7. Structure of nukacin ISK-1 and nukacin KSE650.** The mature peptide sequences of nukacin ISK-1 and nukacin KSE650 are shown. The deduced calculated mass of mature nukacin KSE650 is consistent with that observed by ESI-MS. The structure is identical to that of nukacin ISK-1, except for the residues indicated by gray circles. Dhb, Ala-S-Ala, and Abu-S-Ala indicate dehydrobutyrine, lanthionine, and 3-methylanthionine respectively.

<https://doi.org/10.1371/journal.pone.0258283.g007>

Nukacin IVK-1 was first identified in *S. warneri* [37]. Since then, nukacin ISK-1 like bacteriocins have been identified in *S. epidermidis* [28], *S. hominis* [44], and *S. simulans* [45]. The amino acid sequence of KSE650 shows a high similarity with that of IVK45 by only one mismatch in the entire peptide, and 100% match with the mature peptide. Comparison of the plasmid between the two strains showed that KSE650 was larger than Tü3298, but the composition and the order of nukacin-related genes were identical (Fig 2A). The larger size of pNuk650 was due to the insertion of an approximately 8 kbp fragment, which was detected in pNuk650 but not in pIVK45 (Fig 3A, red arrows).

The antibacterial activity of these peptides against skin and oral commensal bacteria (oral streptococci) showed different patterns. In particular, the epidermin-producing strain (KSE56) had antibacterial activity against oral streptococci, while nukacin-producing strains had less activity. Interestingly, comparing nukacin ISK-1 and nukacin KSE650 suggested that 5 amino acid differences (Fig 7) were responsible for the different activities against several bacteria used in this study. Previously, it was reported that the structure of ring A in nukacin ISK-1 binds to the pyrophosphate moiety of lipid II, the precursor for cell wall peptidoglycan biosynthesis, and ring C was also associated with the binding of the isoprene chain [46]. Since lipid II molecules are widely conserved among gram positive bacteria, the different antibacterial activities between nukacin ISK-1 and nukacin KSE650 are influenced by the other molecules specific to each bacterial species. Furthermore, it is noteworthy that epidermin and nukacin KSE650 showed no inhibitory zone against *S. epidermidis* KSE650 and KSE56, respectively, while epidermin and nukacin KSE650 showed an activity against plasmid-curing KSE650 and plasmid-curing KSE56, respectively (Table 4). Although the immunity factors for epidermin and nukacin KSE650 were EpiFEG and NukFEG/NukH, respectively, which could be found in a respective plasmid, our results indicate that these immunity factors showed a cross-resistance to another bacteriocin. We previously reported that BraRS and ApsRS, TCSs, are involved in resistance to nisin A and nukacin ISK-1 [34]. Since *S. epidermidis* also possesses TCSs with similarity to BraRS and ApsRS, *S. epidermidis* TCSs may be involved in the resistance to epidermin and nukacin KSE650.



In conclusion, we determined the complete sequence of two plasmids encoding epidermin and nukacin KSE650 in *S. epidermidis* isolated from the oral cavity. *S. epidermidis* is the major commensal bacterium in human skin and the oral cavity. Based on our findings of the direct assay and co-culture assay, it is speculated that bacteriocins produced by *S. epidermidis* affect the bacterial composition of the host flora, including the skin, nasal and oral flora. However, in this study, we focused on the isolation of lantibiotic-producing strains using a *braRS*-inactivated strain as the indicator. Therefore, it is possible that *S. epidermidis* also produces other types of bacteriocins. Further studies are required to demonstrate the influence of *S. epidermidis* bacteriocins on the formation of bacterial flora.

## Supporting information

### **S1 Fig. Comparison of amino acid sequences of EpiT between the KSE56 and Tü3298 strains.**

(PDF)

### **S2 Fig. Comparison of nucleotide (A) and amino acid sequences (B) of epiA between the KSE56 and Tü3298 strains.**

(PDF)

## Acknowledgments

We thank Dr. Tomoko Amimoto, the Natural Science Center for Basic Research and Development (N-BARD), Hiroshima University for the measurement of ESI-MS analysis.

## Author Contributions

**Conceptualization:** Miki Kawada-Matsuo, Norifumi Nakamura, Hitoshi Komatsuzawa.

**Data curation:** Kenta Nakazono, Mi Nguyen-Tra Le.

**Formal analysis:** Kenta Nakazono, Mi Nguyen-Tra Le.

**Funding acquisition:** Miki Kawada-Matsuo, Motoyuki Sugai, Hitoshi Komatsuzawa.

**Investigation:** Kenta Nakazono, Mi Nguyen-Tra Le, Miki Kawada-Matsuo, Noy Kimheang, Junzo Hisatsune, Yuichi Oogai.

**Methodology:** Kenta Nakazono, Miki Kawada-Matsuo, Norifumi Nakamura, Motoyuki Sugai, Hitoshi Komatsuzawa.

**Project administration:** Miki Kawada-Matsuo, Norifumi Nakamura, Hitoshi Komatsuzawa.

**Resources:** Miki Kawada-Matsuo, Masanobu Nakata, Norifumi Nakamura, Motoyuki Sugai, Hitoshi Komatsuzawa.

**Software:** Mi Nguyen-Tra Le.

**Supervision:** Miki Kawada-Matsuo, Masanobu Nakata, Norifumi Nakamura, Motoyuki Sugai, Hitoshi Komatsuzawa.

**Validation:** Miki Kawada-Matsuo, Masanobu Nakata, Norifumi Nakamura, Hitoshi Komatsuzawa.

**Visualization:** Miki Kawada-Matsuo, Masanobu Nakata, Hitoshi Komatsuzawa.

**Writing – original draft:** Kenta Nakazono, Mi Nguyen-Tra Le, Miki Kawada-Matsuo.

**Writing – review & editing:** Mi Nguyen-Tra Le, Miki Kawada-Matsuo, Yuichi Oogai, Masanobu Nakata, Norifumi Nakamura, Motoyuki Sugai, Hitoshi Komatsuzawa.

## References

1. Parlet CP, Brown MM, Horswill AR. Commensal Staphylococci Influence *Staphylococcus aureus* Skin Colonization and Disease. *Trends Microbiol.* 2019; 27: 497–507. <https://doi.org/10.1016/j.tim.2019.01.008> PMID: 30846311
2. Grice EA, Kong HH, Conlan S, Deming CB, Davis J, Young AC, et al. Topographical and temporal diversity of the human skin microbiome. *Science.* 2009; 324: 1190–1192. <https://doi.org/10.1126/science.1171700> PMID: 19478181
3. Byrd AL, Belkaid Y, Segre JA. The human skin microbiome. *Nat Rev Microbiol.* 2018; 16: 143–155. <https://doi.org/10.1038/nrmicro.2017.157> PMID: 29332945
4. Nguyen TH, Park MD, Otto M. Host Response to *Staphylococcus epidermidis* Colonization and Infections. *Front Cell Infect Microbiol.* 2017; 7: 90. <https://doi.org/10.3389/fcimb.2017.00090> PMID: 28377905
5. Lai Y, Cogen AL, Radek KA, Park HJ, Macleod DT, Leichtle A, et al. Activation of TLR2 by a small molecule produced by *Staphylococcus epidermidis* increases antimicrobial defense against bacterial skin infections. *J Invest Dermatol.* 2010; 130: 2211–2221. <https://doi.org/10.1038/jid.2010.123> PMID: 20463690
6. Scharschmidt TC, Vasquez KS, Truong H-A, Gearty S V, Pauli ML, Nosbaum A, et al. A Wave of Regulatory T Cells into Neonatal Skin Mediates Tolerance to Commensal Microbes. *Immunity.* 2015; 43: 1011–1021. <https://doi.org/10.1016/j.immuni.2015.10.016> PMID: 26588783
7. Rogers KL, Fey PD, Rupp ME. Coagulase-negative staphylococcal infections. *Infect Dis Clin North Am.* 2009; 23: 73–98. <https://doi.org/10.1016/j.idc.2008.10.001> PMID: 19135917
8. Uçkay I, Pittet D, Vaudaux P, Sax H, Lew D, Waldvogel F. Foreign body infections due to *Staphylococcus epidermidis*. *Ann Med.* 2009; 41: 109–119. <https://doi.org/10.1080/07853890802337045> PMID: 18720093
9. Otto M. *Staphylococcus epidermidis*—the “accidental” pathogen. *Nat Rev Microbiol.* 2009; 7: 555–567. <https://doi.org/10.1038/nrmicro2182> PMID: 19609257
10. Miragaia M, Couto I, de Lencastre H. Genetic diversity among methicillin-resistant *Staphylococcus epidermidis* (MRSE). *Microb Drug Resist.* 2005; 11: 83–93. <https://doi.org/10.1089/mdr.2005.11.83> PMID: 15910220
11. Cotter PD, Hill C, Ross RP. Bacteriocins: developing innate immunity for food. *Nat Rev Microbiol.* 2005; 3: 777–788. <https://doi.org/10.1038/nrmicro1273> PMID: 16205711
12. Jack RW, Tagg JR, Ray B. Bacteriocins of gram-positive bacteria. *Microbiol Rev.* 1995; 59: 171–200. <https://doi.org/10.1128/mr.59.2.171-200.1995> PMID: 7603408
13. Nissen-Meyer J, Nes IF. Ribosomally synthesized antimicrobial peptides: their function, structure, biogenesis, and mechanism of action. *Arch Microbiol.* 1997; 167: 67–77. PMID: 9133328
14. Ryan CS, Kleinberg I. Bacteria in human mouths involved in the production and utilization of hydrogen peroxide. *Arch Oral Biol.* 1995; 40: 753–763. [https://doi.org/10.1016/0003-9969\(95\)00029-o](https://doi.org/10.1016/0003-9969(95)00029-o) PMID: 7487577
15. García-Mendoza A, Liébana J, Castillo AM, de la Higuera A, Piédrola G. Evaluation of the capacity of oral streptococci to produce hydrogen peroxide. *J Med Microbiol.* 1993; 39: 434–439. <https://doi.org/10.1099/00222615-39-6-434> PMID: 8246261
16. Watanabe A, Kawada-Matsuo M, Le MN-T, Hisatsune J, Oogai Y, Nakano Y, et al. Comprehensive analysis of bacteriocins in *Streptococcus mutans*. *Sci Rep.* 2021; 11: 12963. <https://doi.org/10.1038/s41598-021-92370-1> PMID: 34155274
17. Vieira AR, Hiller NL, Powell E, Kim LH-J, Spirk T, Modesto A, et al. Profiling microorganisms in whole saliva of children with and without dental caries. *Clin Exp Dent Res.* 2019; 5: 438–446. <https://doi.org/10.1002/cre2.206> PMID: 31452955
18. McManus BA, Daly B, Polyzois I, Wilson P, Brennan GI, Fleming TE, et al. Comparative Microbiological and Whole-Genome Analysis of *Staphylococcus aureus* Populations in the Oro-Nasal Cavities, Skin and Diabetic Foot Ulcers of Patients With Type 2 Diabetes Reveals a Possible Oro-Nasal Reservoir for Ulcer Infection. *Front Microbiol.* 2020; 11: 748. <https://doi.org/10.3389/fmicb.2020.00748> PMID: 32425909
19. Allgaier H, Jung G, Werner RG, Schneider U, Zähler H. Epidermin: sequencing of a heterodetic tetracyclic 21-peptide amide antibiotic. *Eur J Biochem.* 1986; 160: 9–22. <https://doi.org/10.1111/j.1432-1033.1986.tb09933.x> PMID: 3769923

20. Schnell N, Entian KD, Schneider U, Götz F, Zähler H, Kellner R, et al. Prepeptide sequence of epidermin, a ribosomally synthesized antibiotic with four sulphide-rings. *Nature*. 1988; 333: 276–278. <https://doi.org/10.1038/333276a0> PMID: 2835685
21. Bierbaum G, Götz F, Peschel A, Kupke T, van de Kamp M, Sahl HG. The biosynthesis of the lantibiotics epidermin, gallidermin, Pep5 and epilancin K7. *Antonie Van Leeuwenhoek*. 1996; 69: 119–127. <https://doi.org/10.1007/BF00399417> PMID: 8775972
22. Kaletta C, Entian KD, Kellner R, Jung G, Reis M, Sahl HG. Pep5, a new lantibiotic: structural gene isolation and prepeptide sequence. *Arch Microbiol*. 1989; 152: 16–19. <https://doi.org/10.1007/BF00447005> PMID: 2764678
23. Weil HP, Beck-Sickinger AG, Metzger J, Stevanovic S, Jung G, Josten M, et al. Biosynthesis of the lantibiotic Pep5. Isolation and characterization of a prepeptide containing dehydroamino acids. *Eur J Biochem*. 1990; 194: 217–223. <https://doi.org/10.1111/j.1432-1033.1990.tb19446.x> PMID: 2253617
24. van de Kamp M, Horstink LM, van den Hooven HW, Konings RN, Hilbers CW, Frey A, et al. Sequence analysis by NMR spectroscopy of the peptide lantibiotic epilancin K7 from *Staphylococcus epidermidis* K7. *Eur J Biochem*. 1995; 227: 757–771. <https://doi.org/10.1111/j.1432-1033.1995.tb20199.x> PMID: 7867636
25. Ekkelenkamp MB, Hanssen M, Danny Hsu S-T, de Jong A, Milatovic D, Verhoef J, et al. Isolation and structural characterization of epilancin 15X, a novel lantibiotic from a clinical strain of *Staphylococcus epidermidis*. *FEBS Lett*. 2005; 579: 1917–1922. <https://doi.org/10.1016/j.febslet.2005.01.083> PMID: 15792796
26. Velásquez JE, Zhang X, van der Donk WA. Biosynthesis of the antimicrobial peptide epilancin 15X and its N-terminal lactate. *Chem Biol*. 2011; 18: 857–867. <https://doi.org/10.1016/j.chembiol.2011.05.007> PMID: 21802007
27. Heidrich C, Pag U, Josten M, Metzger J, Jack RW, Bierbaum G, et al. Isolation, characterization, and heterologous expression of the novel lantibiotic epicidin 280 and analysis of its biosynthetic gene cluster. *Appl Environ Microbiol*. 1998; 64: 3140–3146. <https://doi.org/10.1128/AEM.64.9.3140-3146.1998> PMID: 9726851
28. Janek D, Zipperer A, Kulik A, Krismer B, Peschel A. High Frequency and Diversity of Antimicrobial Activities Produced by Nasal *Staphylococcus* Strains against Bacterial Competitors. *PLoS Pathog*. 2016; 12: e1005812. <https://doi.org/10.1371/journal.ppat.1005812> PMID: 27490492
29. Matsuo M, Kato F, Oogai Y, Kawai T, Sugai M, Komatsuzawa H. Distinct two-component systems in methicillin-resistant *Staphylococcus aureus* can change the susceptibility to antimicrobial agents. *The Journal of antimicrobial chemotherapy*. 2010. pp. 1536–1537. <https://doi.org/10.1093/jac/dkq141> PMID: 20430791
30. Murchison HH, Barrett JF, Cardineau GA, Curtiss R 3rd. Transformation of *Streptococcus mutans* with chromosomal and shuttle plasmid (pYA629) DNAs. *Infect Immun*. 1986; 54: 273–282. <https://doi.org/10.1128/iai.54.2.273-282.1986> PMID: 3021626
31. Kornblum J, Hartman BJ, Novick RP, Tomasz A. Conversion of a homogeneously methicillin-resistant strain of *Staphylococcus aureus* to heterogeneous resistance by Tn551-mediated insertional inactivation. *Eur J Clin Microbiol*. 1986; 5: 714–718. <https://doi.org/10.1007/BF02013311> PMID: 3026802
32. Kreiswirth BN, Löfdahl S, Betley MJ, O'Reilly M, Schlievert PM, Bergdoll MS, et al. The toxic shock syndrome exotoxin structural gene is not detectably transmitted by a prophage. *Nature*. 1983; 305: 709–712. <https://doi.org/10.1038/305709a0> PMID: 6226876
33. Four pediatric deaths from community-acquired methicillin-resistant *Staphylococcus aureus*—Minnesota and North Dakota, 1997–1999. *MMWR Morb Mortal Wkly Rep*. 1999; 48: 707–710. PMID: 21033181
34. Kawada-Matsuo M, Yoshida Y, Zendo T, Nagao J, Oogai Y, Nakamura Y, et al. Three Distinct Two-Component Systems Are Involved in Resistance to the Class I Bacteriocins, Nukacin ISK-1 and Nisin A, in *Staphylococcus aureus*. Otto M, editor. *PLoS One*. 2013; 8: e69455. <https://doi.org/10.1371/journal.pone.0069455> PMID: 23894484
35. Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, et al. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res*. 2014; 42: D206–14. <https://doi.org/10.1093/nar/gkt1226> PMID: 24293654
36. Aso Y, Koga H, Sashihara T, Nagao J-I, Kanemasa Y, Nakayama J, et al. Description of complete DNA sequence of two plasmids from the nukacin ISK-1 producer, *Staphylococcus warneri* ISK-1. *Plasmid*. 2005; 53: 164–178. <https://doi.org/10.1016/j.plasmid.2004.08.003> PMID: 15737403
37. Aso Y, Sashihara T, Nagao J-I, Kanemasa Y, Koga H, Hashimoto T, et al. Characterization of a gene cluster of *Staphylococcus warneri* ISK-1 encoding the biosynthesis of and immunity to the lantibiotic, nukacin ISK-1. *Biosci Biotechnol Biochem*. 2004; 68: 1663–1671. <https://doi.org/10.1271/bbb.68.1663> PMID: 15322349

38. Islam MR, Nagao J-I, Zendo T, Sonomoto K. Antimicrobial mechanism of lantibiotics. *Biochem Soc Trans.* 2012; 40: 1528–1533. <https://doi.org/10.1042/BST20120190> PMID: 23176511
39. Vogel V, Spellerberg B. Bacteriocin Production by Beta-Hemolytic Streptococci. *Pathog (Basel, Switzerland).* 2021;10. <https://doi.org/10.3390/pathogens10070867> PMID: 34358017
40. Barbosa J, Caetano T, Mendo S. Class I and Class II Lanthipeptides Produced by *Bacillus* spp. *J Nat Prod.* 2015; 78: 2850–2866. <https://doi.org/10.1021/np500424y> PMID: 26448102
41. Schnell N, Engelke G, Augustin J, Rosenstein R, Ungermann V, Götz F, et al. Analysis of genes involved in the biosynthesis of lantibiotic epidermin. *Eur J Biochem.* 1992; 204: 57–68. <https://doi.org/10.1111/j.1432-1033.1992.tb16605.x> PMID: 1740156
42. Moran JC, Horsburgh MJ. Whole-Genome Sequence of *Staphylococcus epidermidis* Tü3298. *Genome Announc.* 2016;4. <https://doi.org/10.1128/genomeA.00112-16> PMID: 26966218
43. Peschel A, Schnell N, Hille M, Entian KD, Götz F. Secretion of the lantibiotics epidermin and gallidermin: sequence analysis of the genes *gdmT* and *gdmH*, their influence on epidermin production and their regulation by EpiQ. *Mol Gen Genet.* 1997; 254: 312–318. <https://doi.org/10.1007/s004380050421> PMID: 9150266
44. Wilaipun P, Zendo T, Okuda K, Nakayama J, Sonomoto K. Identification of the nukacin KQU-131, a new type-A(II) lantibiotic produced by *Staphylococcus hominis* KQU-131 isolated from Thai fermented fish product (Pla-ra). *Biosci Biotechnol Biochem.* 2008; 72: 2232–2235. <https://doi.org/10.1271/bbb.80239> PMID: 18685189
45. Ceotto H, Holo H, da Costa KFS, Nascimento J dos S, Salehian Z, Nes IF, et al. Nukacin 3299, a lantibiotic produced by *Staphylococcus simulans* 3299 identical to nukacin ISK-1. *Vet Microbiol.* 2010; 146: 124–131. <https://doi.org/10.1016/j.vetmic.2010.04.032> PMID: 20627619
46. Fujinami D, Mahin A-A, Elsayed KM, Islam MR, Nagao J-I, Roy U, et al. The lantibiotic nukacin ISK-1 exists in an equilibrium between active and inactive lipid-II binding states. *Commun Biol.* 2018; 1: 150. <https://doi.org/10.1038/s42003-018-0150-3> PMID: 30272026