

RESEARCH ARTICLE

Transcranial direct current stimulation in attention-deficit hyperactivity disorder: A meta-analysis of neuropsychological deficits

Mohammad Ali Salehinejad^{1,2}*, Miles Wischnewski³, Vahid Nejati^{4,5}, Carmelo M. Vicario^{2,6}, Michael A. Nitsche^{2,7}

1 Ruhr-University Bochum, International Graduate School of Neuroscience, Bochum, Germany, **2** Leibniz Research Centre for Working Environment and Human Factors, Department of Psychology and Neurosciences, Dortmund, Germany, **3** Donders Institute for Brain, Cognition, and Behaviour, Radboud University Nijmegen, The Netherlands, **4** Department of Psychology, Shahid Beheshti University, Tehran, Iran, **5** Department of Psychology, University of Regensburg, Regensburg, Germany, **6** University of Messina, Department of Scienze Cognitive della Formazione e degli Studi Culturali, Messina, Italy, **7** University Medical Hospital Bergmannsheil, Department of Neurology, Bochum, Germany

* These authors contributed equally to this work.

* salehinejad@ifado.de



OPEN ACCESS

Citation: Salehinejad MA, Wischnewski M, Nejati V, Vicario CM, Nitsche MA (2019) Transcranial direct current stimulation in attention-deficit hyperactivity disorder: A meta-analysis of neuropsychological deficits. *PLoS ONE* 14(4): e0215095. <https://doi.org/10.1371/journal.pone.0215095>

Editor: Alessio Avenanti, University of Bologna, ITALY

Received: February 4, 2019

Accepted: March 26, 2019

Published: April 12, 2019

Copyright: © 2019 Salehinejad et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: Publication of this article was funded by the Open Access Fund of the Leibniz Association. MAS receives support from the National Organization of Science, Research & Technology, Deputy of Scholarship and Students Affairs, Iran, grant number: 95000171. The funder had no role in the study design, data collection, and analysis,

Abstract

Transcranial direct current stimulation (tDCS) is a promising method for altering cortical excitability with clinical implications in neuropsychiatric diseases. Its application in neurodevelopmental disorders especially attention-deficit hyperactivity disorder (ADHD), is in early stage and promising but its effectiveness has not been systematically examined yet. We conducted a meta-analysis on the effectiveness of tDCS on the most studied neuropsychological symptoms of ADHD, which is the first reported meta-analysis of tDCS studies on ADHD. Data from 10 randomized controlled studies (including 11 separate experiments) targeting inhibitory control, and/or working memory (WM) in ADHD were included. Results show that overall tDCS significantly improved inhibitory control. Sub-analyses further show that dorsolateral prefrontal cortex (dlPFC) (but not right inferior frontal gyrus) tDCS and anodal (but not cathodal) tDCS significantly improved inhibitory control with a small effect size. Anodal dlPFC-tDCS had the largest significant effect on inhibitory control with a small-to-medium effect size. Additionally, a significant improving effect of tDCS on inhibitory control accuracy (but not response time) and WM speed (but not accuracy) were found. Overall, this meta-analysis supports a beneficial effect of tDCS on inhibitory control and WM in ADHD with a small-to-medium effect size. TDCS seems to be a promising method for improving neuropsychological and cognitive deficits in ADHD. However, there might be a dissociation between neuropsychological deficits and clinical symptoms of ADHD and therefore, the significance of this meta-analysis for clinical purposes is limited. Future studies should systematically evaluate the role of inter-individual factors (i.e., ADHD subtype, types of the deficit) and stimulation parameters (i.e., site, polarity, intensity, duration, repetition rate) on tDCS efficacy in ADHD population and examine whether benefits are long-term.

decision to publish or preparation of the manuscript.

Competing interests: MAN is a member of the Scientific Advisory Board of Neuroelectronics. All other authors declare no competing interests. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

Introduction

Attention-deficit hyperactivity disorder (ADHD) is the most common neurodevelopmental disorder with an estimate of 11% prevalence in school-age children [1]. ADHD is primarily characterized by symptoms of inattention, hyperactivity, impulsivity [2] and various cognitive dysfunctions [3] that often persist into adolescence and adulthood. Apart from symptoms of hyperactivity and inattentiveness, a wide range of cognitive deficits is observed in individuals with ADHD such as problems in attention, inhibitory control working memory (WM), planning, problem-solving and executive functions [4–7]. Executive dysfunctions, especially inhibitory control and working memory, are pervasive and influential in ADHD pathophysiology to the extent that ADHD was labeled a disorder of cognitive control [7, 8] but, cognitive deficits in ADHD are heterogeneous [9–11].

A precise description of ADHD neuropathology is difficult to delineate due to its neuropsychological heterogeneity [12, 13], and substantial overlap between ADHD and typically developing children [14]. However, based on neuroimaging and neuropsychological findings relatively distinct brain regions can be identified to account for ADHD hallmark symptoms. Poor inhibitory control resulting from executive dysfunctions (i.e., inhibition-based model) and impulse-control deficits leading to hyperactivity (i.e., motivational dysfunction model) are two influential theories for neural foundations of ADHD [15, 16]. These major symptoms have been linked to frontostriatal circuitry and prefrontal cortex (PFC) abnormalities and can be diverged in the “prefrontal hypothesis of ADHD” [17]. According to this hypothesis, prefrontal regions including the dorsolateral PFC (dlPFC), orbitofrontal cortex (OFC) and inferior frontal gyrus (IFG) have altered activity in ADHD. Specifically, hypoactivity of the bilateral dlPFC, right inferior frontal gyrus (rIFG) and OFC, as well as smaller prefrontal volumes, are associated with behavioral deficits in ADHD [18–22]. Additionally, an adaptive increase of posterior parietal activity accompanies the respective hypoactivity of frontostriatal regions during performance of executive tasks [19]. These findings suggest that ADHD symptoms/causes stem from disturbances in large-scale brain networks [23, 24] and structural, functional and neurochemical abnormalities in cortical and subcortical structures [13, 17, 25], which should be considered in therapeutic interventions.

Pharmacological and non-pharmacological interventions have been among the most commonly used and recommended treatment options in ADHD. Pharmacological interventions can broadly be divided into stimulant and non-stimulant medications [26, 27]. Non-pharmacological or psychological interventions can be categorized into behavioral interventions, cognitive training and neurofeedback training [27, 28]. Stimulant medications have been so far the most effective psychopharmacological treatment and adherence to treatment is reported relatively high. Although current treatment options especially pharmacological interventions have been relatively effective and successful, the efficacy of these treatment options has been inconsistent in ADHD [27]. For example, the efficacy of pharmacological treatments is not long-lasting enough [29] and suffers from the partial response or non-response [30], adverse effects and relatively poor adherence [31]. On the other hand, neurofeedback training is found to have limited effects on ADHD when rated by blind evaluators [32]. Due to the heterogeneity of ADHD pathophysiology, symptoms and treatment response, there might be a need for new treatment or complementary treatment options, especially regarding interventions based on recent findings of the neuropathology of ADHD.

The significant role of brain abnormalities in ADHD symptomology encouraged researchers to probe new treatment options that target ADHD symptoms by modulating, altering and remediating deficient neurocognitive brain functions. Transcranial direct current stimulation (tDCS) is a non-invasive, painless, and well-tolerated brain stimulation technique that has

been gaining increased popularity in human neuroscience research in both healthy and clinical populations during last decade [33–35]. It induces a weak direct current applied to the scalp which modulates cortical excitability by shifting resting membrane potential, [36] and can induce both, acute and neuroplastic alterations of cortical excitability at the macroscopic level [37]. In the motor cortex, it has been suggested that anodal stimulation increases cortical excitability while cathodal stimulation decreases it [38] although such polarity-dependent effect is not always linear and could be affected by external and individual difference factors especially stimulation parameters such as current density, polarity, stimulation duration and/ or geometrical montage of electrodes [39, 40].

TDCS has been applied for modulating a variety of cognitive functions [41–44] or improving symptoms/deficits in healthy and clinical populations [45] (for an overview see [46, 47]). Its application in neurodevelopmental populations, especially ADHD, has gained attention in recent years [48, 49] while the efficacy of the method in ADHD is still unclear and needs further research. Different stimulation parameters and montages have been used in previous studies and different symptoms and deficits have been targeted by tDCS which does not evaluate the potential of this method in ADHD. Moreover, most clinical tDCS studies so far have focused on adult populations and there is a great need for more research into their therapeutic uses in pediatric patients [49, 50]. Considering the fact that pediatric brains still have to go through various stages of neurodevelopment and have accelerated neuroplasticity compared to adults [51], tDCS could be considered a potential therapy for some pediatric disorders, particularly when there are no other safe viable alternatives [52]. That being said, knowledge about tDCS effects in ADHD, is still relatively limited and warrants further investigation especially through systematic review or meta-analytic studies.

To date, meta-analysis or specific systematic review is available for the effects of tDCS on ADHD. The present meta-analysis, therefore, aims to investigate the effectiveness of tDCS on neuropsychological deficits in ADHD. Moreover, we did further sub-analyses on the available studies in order to (1) evaluate overall and deficit-specific efficacy of tDCS in ADHD, (2) identify the brain regions that are prominently involved in ADHD pathophysiology tDCS response, and (3) examine stimulation parameters that are important for tDCS efficacy in ADHD. Since inhibitory control and WM deficiencies are among the major neuropsychological deficits of ADHD, [53] and mediated by the genetic risk for ADHD, [54], we performed a meta-analysis on the effects of tDCS on inhibitory control and WM performance in respective patients.

Method

Our meta-analysis follows the guidelines of the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) [55] which consists of a checklist intended to facilitate preparation and reporting review/meta-analysis studies by identifying, selecting, and critically appraising relevant research, and collecting and analyzing data from the included studies. A brief version of the PRISMA checklist is available as supporting information (See [S1 Table](#)). The following steps are based on the PRISMA protocol.

Eligibility criteria

To ensure a high level of methodological accuracy, only peer-reviewed published studies were included in our analysis. Only studies on humans using tDCS in experimental designs with the control condition (i.e., sham stimulation and/or baseline control) with reported full-text in English were retained. Studies were included if (1) they had a randomized placebo- (sham⁵⁰) controlled or baseline-controlled design, (2) measured at least inhibitory control, and/or working memory performance, (3) reported data (text, figures, tables, appendices) which contained

effect size values or mean and standard deviation values in order to calculate effect sizes. Participants in the studies were required to be between 3 to 18 years of age (for childhood and adolescence ADHD) and 18 to 65 years of age (for adult ADHD) and to have a clinical diagnosis of ADHD (any subtype) based on the Diagnostic and Statistical Manual of Mental Disorders (DSM) or meet accepted cut-offs on validated ADHD symptom rating scales. Studies which included patients under medication for ADHD treatment in either the control or the active arm were not excluded.

Search strategy and study selection

A comprehensive literature search was conducted, including the PubMed database, Web of Science, Medline, and Scopus with the final search updated on January 2019. Search terms included attention-deficit hyperactivity disorder OR ADHD OR attention disorders AND transcranial direct current stimulation OR tDCS OR transcranial electrical stimulation OR tES (see Fig 1). Furthermore, a manual search was carried out over the reference sections of retrieved studies and review articles. After removal of duplicates, the full text was assessed and studies that were eligible according to the above-mentioned inclusion criteria were selected. The final search identified a total of 241 articles with 1 additional article identified by reference lists and recent reviews. Records were reduced to 28 articles after removing duplications. A content expert examined each title and abstract for eligibility. 18 articles were excluded for not meeting the inclusion criteria. Thus 10 articles remained for full-text assessment and data extraction. 8 of these studies investigated inhibitory control and 5 studies explored tDCS effects on WM performance. Articles were screened and selected independently by the first and second authors. Respective study characteristics are summarized in Tables 1 and 2.

Outcome variables

We limited our focus on neuropsychological variables that were targeted in most ADHD studies using tDCS. Based on the most commonly available data, inhibitory control and WM were identified as the variables for which the data were extracted. Inhibitory control and WM are central components of ADHD executive dysfunctions, as proposed by causal models of ADHD, [14, 53, 56] and encompass the most consistently impaired domains in ADHD [8]. Moreover, inhibitory control and WM are major neuropsychological deficits from the neuroscientific perspective, including the prefrontal-striatal model of ADHD [57]. For the meta-analysis on inhibitory control, the following tasks were included: (1) Go/No-Go, (2) Stop Signal Task (SST), (3) Flanker, (4) Stroop, (5) Continuous Performance Test (CPT), and (6) Neuropsychological Development Assessment (NEPSY II). In the Go/No-Go task, inhibitory process is reflected by the ability to inhibit a motor action in the case of specific number of “No-Go” trial [58]. Similarly, in the SST, participants should stop responding to “Go” trials after the stop signal is presented which shows the ability to inhibitory control. In the Stroop task, inhibitory control is reflected by the ability to suppress the meaning of a written word and focus on the color [59]. For the Flanker task response, inhibition is measured by identifying the target defined by its location while ignoring one or more distracting items that flank the targets in the same or opposite direction [60]. In the CPT, which is primarily a test of attention, the errors of commission and omission reflect inhibitory control ability [61]. Finally, in NEPSY-II inhibitory control is measured by looking at a series of shapes or arrows and naming the shape or direction or an alternative response, depending on the color or shape of the arrow [62]. For the meta-analysis on WM, the following tasks were included: 1) N-back, 2) Digit Span, and 3) Corsi block-tapping test. In the N-back task, subjects need to identify a stimulus that repeats the one presented “n” items before its onset. In the digit span task, subjects are read or shown




Fig 1. (A) Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram of selection of studies, (B) Bias assessment in individual studies. tDCS = Transcranial Direct Current Stimulation; ADHD = attention-deficit hyperactivity disorder; IC = Inhibitory Control; WM = Working Memory.

<https://doi.org/10.1371/journal.pone.0215095.g001>

a list of digits and asked to recall them in order. In the Corbi Cubes test, the subject repeats sequences of touches in different cubes (either forward or backward) [63].

Risk of assessment bias

Quality assessment of the included studies was performed using the Cochrane Collaboration’s tool for risk of bias in randomized trials [64]. For each study, the authors judged the risk of selection bias, performance bias, detection bias, attrition bias, reporting bias, and other biases. Risk of bias was categorized as low, high or uncertain. The final rating was established through consensus with the involvement of the senior author. The assessment of each study and the percentage of bias are presented in Fig 1B.

Data extraction and statistical analysis

Hedges’ g was used as a measure of effect size. This effect size reflects the tDCS-induced difference compared to 1) sham, 2) baseline or 3) sham and baseline (i.e. posttest–pretest tDCS vs

Table 1. Characteristic of studies included in meta-analysis for the effects of tDCS on inhibitory control.

#	Authors	N	Mean age	tDCS montage (target/reference)	Intensity	Duration	Polarity	On-/off-line	Control	Task	Outcome	Hedges' g									
1	Allenby et al (2018)	37	37.17 (range 18–56)	F3/Fp2 (25 cm ² both)	2 mA	3 days x 20 min	Anodal	Offline	Baseline + sham	CPT	False positive errors	0.42									
											True positive errors	-0.06									
											Response time	-0.11									
				F3/Fp2 (25 cm ² both)	2 mA	3 days x 20 min	Anodal	Offline	Baseline + sham	SST	Reaction time	-0.18									
2	Bandeira et al (2016)	9	11.1 ± 2.8	F3/Fp2 (35 cm ² both)	2 mA	5 days x 30 min	Anodal	Offline	Baseline	NEPSY II	Total errors	0.12									
											Completion time	0.54									
3	Breitling et al (2016)	21	14.33 (range 13–17)	F8/mastoid (35 cm ² both)	1 mA	20 min	Anodal	Online	Sham	Flanker task	Omission errors	-0.11									
											Comission errors	0.46									
											Reaction time	-0.14									
														Reaction time variability	0.13						
								F8/mastoid (35 cm ² both)	1 mA	20 min	Cathodal	Online	Sham	Flanker task	Omission errors	-0.60					
														Comission errors	0.17						
4	Cosmo et al (2015)	30	31.8 ± 11.6	F3/F4 (35 cm ² both)	1 mA	20 min	Anodal	Offline	Sham	Go/No-go task (letters)	Correct responses	-0.39									
											Omission errors	1.17									
											Comission errors	-0.15									
																			Go/No-go task (fruits)	Correct responses	1.09
																				Omission errors	0.44
																				Comission errors	0.24
5	Munz et al (2015)	14	12.3 ± 1.4	F3+F4/both mastoids (0.5 cm ² all)	0–0.25 mA (oscillatory)	5 x 5 min	Anodal	Offline	Sham	Go/No-go task	Reaction time	0.88									
											Reaction time variability	0.83									
6	Nejati et al (2017) experiment 1	15	10 ± 2.2	F3/F4 (25 cm ² both)	1 mA	15 min	Anodal	Offline	Sham	Go/No-go task	Go accuracy	0.14									
											No-go accuracy	0.56									
											Reaction time	-0.41									
																		Stroop task	Accuracy	1.11	
								Reaction time	0.98												
7	Nejati et al (2017) experiment 2	10	9 ± 1.8	F3/Fp2 (25 cm ² both)	1 mA	15 min	Anodal	Offline	Sham	Go/No-go task	Go accuracy	0.41									
											No-go accuracy	0.57									
											Reaction time	-0.20									

(Continued)

Table 1. (Continued)

#	Authors	N	Mean age	tDCS montage (target/reference)	Intensity	Duration	Polarity	On-/off-line	Control	Task	Outcome	Hedges' g
				F3/Fp2 (25 cm ² both)	1 mA	15 min	Cathodal	Offline	Sham	Go/No-go task	Go Accuracy	0.41
											No-go accuracy	1.04
											Reaction time	-0.20
8	Soltaninejad et al (2015)	20	Range 15–17	F3/Fp2 (35 cm ² both)	1.5 mA	8 min	Anodal	Offline	Sham	Go/No-go task	Go accuracy	-0.05
											No-go accuracy	0.03
											Reaction time	0.23
										Stroop task	Accuracy	0.57
											Reaction time	0.23
				F3/Fp2 (35 cm ² both)	1.5 mA	8 min	Cathodal	Offline	Sham	Go/No-go task	Go accuracy	-0.54
											No-go accuracy	0.73
											Reaction time	-0.02
										Stroop task	Accuracy	0.33
											Reaction time	0.11
											Reaction time	0.02
9	Sotnikova et al (2017)	13	14.33 ± 1.3	F3 (13 cm ²)/Cz (35 cm ²)	1 mA	30 min	Anodal	Online	Sham	Go/No-go	accuracy (hits + correct rejections/ total number of stimuli)	-1.03
											Reaction time	0.16
											Reaction time variability	-0.14

tDCS = transcranial direct current stimulation; F3 = left dlPFC; F4 = right dlPFC; F8 = inferior frontal gyrus; Fp1 = left supraorbital area; Fp2 = right supraorbital area; online = task performance during tDCS; offline = task performance after tDCS; CPT = Conners Continuous Performance Task; SST = Stop Signal Task (SST).

<https://doi.org/10.1371/journal.pone.0215095.t001>

posttest–pretest sham). When the effect size Cohen's *d* was reported, the effect sizes were extracted directly from the selected articles. Otherwise, *d* was calculated using means and pooled standard error of mean, which were gathered from the results section, figures or tables. From these effect sizes, Hedges' *g* values were calculated to correct for effect inflation due to small sample sizes [65]. In this meta-analysis, positive values reflect a tDCS-induced increase of inhibitory control or WM performance, whereas negative values indicate a tDCS-induced decrease of inhibitory control or WM performance compared to sham/baseline values. A weighted average was calculated from these Hedges' *g* values to determine the cumulative effect size (\bar{E}) and 95% confidence intervals. Subsequently, the *Z*-statistic and *p*-value were calculated to investigate whether \bar{E} differed significantly from zero.

For the main investigation on the effects of tDCS on inhibitory control in ADHD patients, three meta-analytic steps were performed. First, an unsigned analysis, in which the general effect of frontal tDCS, independent of polarity was analyzed. This analysis included 46 extracted effect size values. We further explored the effect of electrode montage by comparing tDCS setups that primarily targeted the dlPFC (38 extracted effect size values) or the rIFG (8 extracted effect size values). Second, a signed analysis was performed in which the effects of anodal (34 extracted effect size values) and cathodal tDCS (12 extracted effect size values) were

Table 2. Characteristic of studies included in the meta-analysis for the effects of tDCS on working memory.

#	Authors	N	Mean age	tDCS montage (target/reference)	Intensity	Duration	Polarity	On-/off-line	Control	Task	Outcome	Hedges' g		
1	Bandeira et al (2016)	9	11.1 ± 2.8	F3/Fp2 (35 cm ² both)	2 mA	5 days x 30 min	Anodal	Offline	Baseline	Digit span forward	Amount	-0.87		
										Digit span backward	Amount	-0.40		
										Corsi cube forward	Amount	-0.45		
										Corsi cube backward	Amount	0.08		
2	Nejati et al (2017) experiment 1	15	10 ± 2.2	F3/F4 (25 cm ² both)	1 mA	15 min	Anodal	Offline	Sham	1-back task	Accuracy	0.15		
											Reaction time	1.70		
3	Nejati et al (2017) experiment 2	10	9 ± 1.8	F3/Fp2 (25 cm ² both)	1 mA	15 min	Anodal	Offline	Sham	1-back task	Accuracy	1.14		
												Reaction time	0.82	
												Accuracy	0.53	
												Reaction time	0.52	
4	Prehn-Kristensen et al (2014)	12	12.1 (range 10–14)	F3+F4/both mastoids (0.5 cm ² all)	0–0.25 mA (oscillatory)	5 x 5 min	Anodal	Offline	Baseline + sham	Digit span	Amount	-0.61		
5	Soff et al (2017)	15	14.2 ± 1.2	F3 (3.14 cm ²)/ Cz (12.5 cm ²)	1 mA	5 days x 20 min	Anodal	Offline	Baseline + sham	QB (1-back) task	QB score (errors and reaction time)	0.50		
6	Sotnikova et al (2017)	13	14.33 ± 1.3	F3 (13 cm ²)/ Cz (35 cm ²)	1 mA	30 min	Anodal	Online	Sham	1-back task	Accuracy	-0.99		
												Reaction time	-0.05	
												Reaction time variability	0.18	
												2-back task	Accuracy	-1.14
												Reaction time	0.65	
	Reaction time variability	1.06												

tDCS = transcranial direct current stimulation; F3 = left dlPFC; F4 = right dlPFC; Fp2 = right supraorbital area; online = task performance during tDCS; offline = task performance after tDCS; QbTest = Quantified Behavior Test.

<https://doi.org/10.1371/journal.pone.0215095.t002>

determined separately. This analysis was then followed by an exploration of tDCS montage. Finally, since the majority of outcome measurements investigated both, accuracy (i.e., amount of correct responses or error rate) and speed (i.e. reaction time or reaction time variability) on inhibitory control tasks, separate analyses were performed for these variables (27 and 19 extracted effect size values respectively). For the WM analyses, not enough observations could be gathered for cathodal tDCS stimulation (n = 2) and therefore no separate analyses were performed.

For all analyses, the sample distribution was checked for normality by the Kolmogorov-Smirnov test. Additionally, total heterogeneity (Qtotal) was tested against the χ^2 distribution with n-1 degrees of freedom to determine whether the variance of effect sizes was greater than to be expected from sampling error [65]. To further investigate possible effects of publication

bias, the fail-safe number, based on the Rosenthal method ($\alpha < 0.05$), was calculated, which gives an indication of the amount of null findings that are needed to render the cumulative effect non-significant [66]. Data were analyzed using MetaWin 2.1[67] and IBM SPSS 22.0. All statistical tests were tested against a significance level of $\alpha \leq 0.05$ (two-tailed).

Results

Overview

In total, 11 separate experiments from 10 studies published from 2014—April 2018 were included in this meta-analysis. In 8 of 10 studies (or 9 of 11 experiments), effects of tDCS on inhibitory control was investigated [6, 68–74] and in 5 of 10 studies (or 6 of 11 experiments), tDCS effects on WM were explored [6, 69, 74–76]. In what follows, we describe the effect size obtained from studies to examine the efficacy of tDCS in ADHD major neuropsychological symptoms (Please see Fig 2). Brief details of each study are also summarized in Tables 1 and 2.

Risk of bias

The authors' judgment on the risk of bias is displayed in Fig 1B. Overall, the risk of bias was low. However, uncertainty about the risk does exist, at least to a certain extent, for selection bias, performance bias, and detection bias. Especially, for determination of the latter, information was missing in the majority of studies. A selective reporting bias was observed for the study of Munz et al. [72] since accuracy data for the experimental conditions are not reported in the manuscript, nor in any appendices. In all studies, blinding of participants was reported, but blinding of experimenters was not reported in two studies [70, 73]. We need to note that true blinding procedure of participants requires debriefing at the end of treatment about the active or sham mode of stimulation confirmed by statistical analysis. This was reported or confirmed only in three included studies implicating that blinding of participants should be interpreted with caution. All studies used randomization to allocate participants to different experimental conditions.

Effects of tDCS on inhibitory control in ADHD patients

A significant cumulative effect size (\bar{E}) of 0.197 ($Z = 2.76$, $p = 0.006$) was observed for a general tDCS effect on inhibitory control, taking polarity not into account Kolmogorov-Smirnov's test of normality showed that the distribution of the effect sizes was not significantly different from a normal distribution (lower bound $p = 0.20$) and total heterogeneity of the effect sizes was not significant ($Q_{total} = 45.21$, $p = 0.463$). The fail-safe number indicated that 221 unpublished null-findings would be required to render the effect non-significant. Exploration of montage showed that only dlPFC stimulation (l-dlPFC and bilateral) ($\bar{E} = 0.243$, $Z = 2.92$, $p = 0.004$), but not rIFG stimulation ($\bar{E} = 0.005$, $Z = 0.04$, $p = 0.971$) yielded a significant increase of accuracy rates in inhibitory control task performance.

Subsequently, polarity-dependent effects were investigated. Studies using anodal tDCS showed a significant \bar{E} of 0.232 ($Z = 2.72$, $p = 0.007$), with a fail-safe number of 171 showing that anodal tDCS significantly improved inhibitory control. This sample was distributed normally (lower bound $p = .20$) and showed no significant heterogeneity ($Q_{total} = 33.05$, $p = 0.465$). As for the stimulation polarity-independent analysis, this effect was driven by studies using a left and bilateral dlPFC montage ($\bar{E} = 0.255$, $Z = 2.66$, $p = 0.008$), whereas the rIFG montage did not yield a significant effect ($\bar{E} = 0.084$, $Z = 0.33$, $p = 0.745$). In contrast to anodal tDCS, cathodal tDCS did not show a significant overall effect ($\bar{E} = 0.089$, $Z = 0.63$, $p = 0.530$).




Fig 2. Meta-analysis and forest plot results including Hedges' g and 95% confidence interval and Cumulative effect size of tDCS on inhibitory control (top) and working memory (down).

<https://doi.org/10.1371/journal.pone.0215095.g002>

Table 3. Meta-analysis results for the effects of tDCS on inhibitory control in ADHD patients.

Analysis	N	Cumulative effect size				Normality		Heterogeneity		
		\bar{E}	95% CI	Z	p-value	Fail-safe number	KS test	p-value	Qtotal	p-value
Polarity-independent										
All studies	46	0.197	0.057–0.336	2.758	0.0058	221	0.088	LB 0.200	45.21	0.463
dIPFC only	38	0.243	0.080–0.406	2.922	0.0035	97	0.077	LB 0.200	36.87	0.475
rIFG only	8	0.005	-0.261–0.271	0.037	0.9705	0	0.195	LB 0.200	6.411	0.493
Polarity-dependent										
Anodal tDCS	34	0.232	0.065–0.400	2.723	0.0065	171	0.095	LB 0.200	33.05	0.465
dIPFC only	30	0.255	0.067–0.443	2.658	0.0079	56	0.085	LB 0.200	28.98	0.466
rIFG only	4	0.084	-0.422–0.589	0.325	0.7452	0	¹		2.25	0.523
Cathodal tDCS	12	0.089	-0.189–0.367	0.628	0.5300	0	0.133	LB 0.200	11.27	0.421
dIPFC only	8	0.194	-0.212–0.600	0.937	0.3488	0	0.114	LB 0.200	7.06	0.423
rIFG only	4	-0.075	-0.635–0.486	-0.263	0.7926	0	¹		3.01	0.390
Speed vs Accuracy										
Accuracy	27	0.250	0.042–0.459	2.356	0.0185	36	0.096	LB 0.200	26.30	0.447
Speed	19	0.093	-0.075–0.261	1.085	0.2779	0	0.208	0.031	18.08	0.451

tDCS = Transcranial Direct Current Stimulation; dIPFC = dorsolateral prefrontal cortex; LB = lower bound; rIFG = right inferior frontal gyrus; \bar{E} = cumulative effect size; CI = Confidence interval; KS = Kolmogorov-Smirnov’s test of normality; Qtotal = total heterogeneity represented by Cohen’s Q; Significant results are highlighted in bold. dIPFC refers to either left dIPFC or bilateral dIPFC (for detailed information refer to Tables 1 and 2 under tDCS montage column). 1KS test could not be performed because of too small sample size

<https://doi.org/10.1371/journal.pone.0215095.t003>

Finally, an analysis was performed separating outcomes measures that focused on accuracy or amount of errors compared to the speed of response. The results showed that tDCS significantly increased accurate responses in inhibitory control tasks ($\bar{E} = 0.250, Z = 2.36, p = 0.019$). No significant cumulative effect was found for speed ($\bar{E} = 0.093, Z = 1.09, p = 0.278$). For this last analysis, a deviation from normality was observed ($p = 0.031$). Results are summarized in Table 3.

Effects of tDCS on working memory in ADHD patients

No significant cumulative effect was observed for tDCS on working memory, without taking polarity into account ($\bar{E} = 0.160, Z = 0.81, p = 0.418$). Also, no effect of tDCS was observed when only studies with an anodal montage were included ($\bar{E} = 0.116, Z = 0.53, p = 0.600$). However, when separating outcomes for accuracy and speed, a significant effect of tDCS on speed was observed. TDCS led to a faster response time ($\bar{E} = 0.681, Z = 2.43, p = 0.015$), with a fail-safe number of 17. The sample was normally distributed (lower bound $p = .20$) and no significant heterogeneity was seen ($Q_{total} = 5.88, p = 0.437$). These results should be interpreted with caution, given the low sample size ($N = 7$). Moreover, results showed that tDCS did have no significant effect on accuracy of working memory task performance ($\bar{E} = -0.187, Z = -0.76, p = 0.446$). Results are shown in Table 4.

Discussion

ADHD is a major neurodevelopmental disorder with remarkable heterogeneity in symptoms, etiologies, and treatment response. Recent studies have emphasized on executive dysfunctions and brain functional abnormalities underlying ADHD pathophysiology [14, 20, 54, 57] suggesting that modulation of cortical activity in the involved brain regions via tDCS might

Table 4. Meta-analysis results for the effects of tDCS on working memory in ADHD patients.

Analysis	N	Cumulative effect size				Normality		Heterogeneity		
		\bar{E}	95% CI	Z	p-value	Fail-safe number	KS test	p-value	Qtotal	p-value
Polarity independent										
All studies	18	0.160	-0.227–0.547	0.811	0.4176	0	0.111	> 0.200	17.36	0.430
Polarity-dependent										
Anodal tDCS	16	0.116	-0.317–0.549	0.525	0.5996	0	0.107	> 0.200	15.46	0.419
Speed vs Accuracy										
Accuracy	11	-0.187	-0.668–0.295	-0.762	0.4461	0	0.163	> 0.200	10.09	0.433
Speed	7	0.681	0.131–1.231	2.427	0.0152	17	0.130	> 0.200	5.88	0.437

tDCS = Transcranial Direct Current Stimulation; \bar{E} = cumulative effect size; CI = Confidence interval; KS = Kolmogorov-Smirnov’s test of normality; Qtotal = total heterogeneity represented by Cohen’s Q; Significant results are highlighted in bold

<https://doi.org/10.1371/journal.pone.0215095.t004>

ameliorate neuropsychological symptoms [77]. Exploring the impact of tDCS especially on ADHD neuropsychological deficits has significantly increased in the last few years. However, no meta-analysis about tDCS efficacy in ADHD is available. We conducted a meta-analysis of randomized, placebo-or-baseline-controlled trials of tDCS application in ADHD, which is the first meta-analysis for ADHD patients.

Overall, results of this meta-analysis show that tDCS improved inhibitory control and WM in ADHD. Further sub-analyses yielded the following findings: (1) tDCS has an overall significant cumulative effect on inhibitory control in ADHD with a small effect size, (2) when the targeted brain region is taken into account, only tDCS over the dlPFC had a significant effect on inhibitory control (small-to-medium effect size), but not tDCS over the rIFG, (3) when stimulation polarity was taken into account, only anodal, but not cathodal tDCS had a significant effect on inhibitory control, (4) when both polarity and targeted region are taken into account, only anodal tDCS of the dlPFC had a significant effect on inhibitory control with a small-to-medium effect size, (5) and when analyzing inhibitory control outcomes separately, tDCS had a significant cumulative effect on accuracy, but not speed (i.e., reaction time). With regard to WM (6), tDCS had an overall significant effect only on performance speed but not on accuracy. In what follows we discuss tDCS effects on these two major target deficits, the stimulation parameters that could yield to larger effects, and finally clinical and methodological implications for future studies.

tDCS effects on inhibitory control in ADHD

The significant overall effect of tDCS on inhibitory control in ADHD is in line with a previous meta-analysis about the effects of this technique on a variety of cognitive functions (e.g., WM, executive functions, language) [78–80] and neuropsychiatric populations [80–82]. Moreover, the small effects size of the overall effect of tDCS found in our meta-analysis is consistent with small-to-medium reported effect sizes in previous meta-analyses about the effectiveness of tDCS [78, 82–84]. However, the effects of tDCS on inhibitory control were associated with larger effect size when target area (i.e., electrode positioning) and stimulation polarity (anodal vs cathodal) were taken into account.

Anodal dlPFC tDCS had the largest effect size (small-to-medium effect) on inhibitory control in ADHD populations, whereas anodal rIFG tDCS had no significant effect. For stimulation over the dlPFC, both left and right dlPFC were targeted in different studies but the left dlPFC was almost always stimulated with the target electrode. First of all, this is in line with

previous findings about significant involvement of the dlPFC in ADHD symptoms, especially executive dysfunctions [16, 18]. Inhibitory control is a major component of dlPFC-supported executive functioning [85, 86] which is impaired in ADHD. The dlPFC is hypoactive in ADHD, which is associated with insufficient suppression of dlPFC signaling not only in the resting state, but also during the performance of cognitive tasks [20, 87]. Recent meta-analyses of neuroimaging studies about functional abnormalities in ADHD populations showed a bilateral hypoactivity of the dlPFC [88] and reduced left medial frontal cortex activation [89] in ADHD populations during inhibitory control, WM and attention task performance. This hypoactivity of the dlPFC is assumed to underlie the attentional deficit, impaired inhibitory control and executive dysfunctions in ADHD and therefore, the pathophysiological rationale for the therapeutic application of tDCS is to increase dlPFC activation with anodal stimulation. It further supports the “cognitive dysfunction or inhibition-based model” of ADHD which suggests that inhibition-based executive deficits are a core deficit in ADHD [56, 90].

Secondly, it highlights the importance of stimulation site for the tDCS effects on inhibitory control. Our findings showed that dlPFC tDCS and not rIFG tDCS is effective. Moreover, the target area of the reference electrode impacts on tDCS effects on inhibitory control. TDCS effects on inhibitory control were more effective when two specific regions were stimulated by the target and reference electrodes. The left dlPFC-right OFC montage was the most effective montage and all of the four experiments which used this protocol reported improved performance. In contrast, the left dlPFC-right dlPFC tDCS montage, which was used in two studies, did not improve any inhibitory control outcome measure. In two studies that targeted only one prefrontal region (i.e., left dlPFC-Cz tDCS, left/right dlPFC- mastoid tDCS), inhibitory control was also improved (see Table 5). Thirdly and in addition to the stimulation site, the results of this meta-analysis demonstrate the importance of stimulation polarity in the tDCS effects on inhibitory processes in ADHD. Anodal tDCS, especially over dlPFC yielded the largest effect size for improving inhibitory control in ADHD. Even the two studies that reported significant effects of cathodal tDCS on response inhibition, argued that cathodal tDCS over the left dlPFC was probably effective in improving inhibitory control in ADHD by increasing activity of the right dlPFC through the inhibitory link between contralateral dlPFC regions via the transcallosal connections. That being said, further systematic investigations are still required to more robustly specify the contribution of potential stimulation sites and polarity on response inhibition in ADHD populations.

Additionally, the findings suggest that heterogeneity of ADHD subtypes and brain abnormalities should not be neglected in interpreting the tDCS effects. This heterogeneity could explain why rIFG tDCS was not as effective as dlPFC tDCS in the present analysis. For example, both, rIFG and left/right dlPFC abnormalities are shown to be associated with inhibitory control deficits in ADHD [13, 18, 53]. However, a differential involvement of the dorsal/ventral divisions within the lateral PFC is suggested in ADHD pathophysiology. Dorsal regions (i.e., dlPFC) are suggested to support inhibitory control of cognitive processes, while more ventral regions (e.g., rIFG) support motor domains of inhibitory control [18, 91]. This shows that ADHD subtypes (i.e., inattention type vs hyperactive type) are important for variations in tDCS effects and implicates that the rIFG tDCS might be well-suited for those groups of patients with more hyperactive than inattentive symptoms. Moreover, dlPFC has a global contribution to executive functioning and response inhibition regardless of modality [42]. This global contribution of the dlPFC in all aspects of response inhibition might explain the significant implication of the dlPFC, but not rIFG in ADHD inhibitory processes. Furthermore, ADHD is a disorder with disturbances in large-scale inter-related brain networks [23, 24]. The dlPFC is a crucial part of the prefrontal network and compared to other prefrontal regions, has many connections with other cortical/subcortical areas [92, 93]. It is possible that dlPFC

Table 5. Characteristics of the stimulation site and polarity in the studies investigated tDCS effects on inhibitory control and working memory.

Inhibitory control				
#	Authors	Stimulation target (site)	polarity	Performance improvement
1	Breitling et al (2016)	right IFG—left mastoid	Anodal	No
2	Soltaninejad et al (2015)	Left dlPFC—right supraorbital area (OFC)	Anodal	No
		Left dlPFC—right supraorbital area (OFC)	Cathodal	Yes
3	Allenby et al (2018)	Left dlPFC—right supraorbital area (OFC)	Anodal	Yes
4	Bandeira et al (2016)	Left dlPFC—right supraorbital area (OFC)	Anodal	Yes
5	Nejati et al (2017) Exp 2	Left dlPFC—right supraorbital area (OFC)	Anodal	No
		Left dlPFC—right supraorbital area (OFC)	Cathodal	Yes
6	Nejati et al (2017) Exp 1	Left dlPFC—right dlPFC	Anodal	No
7	Cosmo et al (2015)	Left dlPFC—right dlPFC	Anodal	No
8	Sotnikova et al (2017)	Left dlPFC—vertex	Anodal	Yes
9	Munz et al (2015)	Left dlPFC—mastoid	Anodal	Yes (only RT)
		right dlPFC—mastoid	Anodal	Yes (only RT)
Working memory				
1	Soff et al. (2017)	Left dlPFC—vertex	Anodal	Yes (both accuracy and RT)
2	Sotnikova et al (2017)	Left dlPFC—vertex	Anodal	Yes (both accuracy and RT)
3	Nejati et al (2017) Exp 1	Left dlPFC—right dlPFC	Anodal	Yes (only RT)
4	Nejati et al (2017) Exp 2	Left dlPFC—right supraorbital area (OFC)	Anodal	Yes (both accuracy and RT)
		Left dlPFC—right supraorbital area (OFC)	Cathodal	No
5	Bandeira et al (2016)	Left dlPFC—right supraorbital area (OFC)	Anodal	No
6	Prehn-Kristensen et al (2014)	Left dlPFC / right dlPFC—mastoid	Anodal	Yes

tDCS = transcranial direct current stimulation; dlPFC = dorsolateral prefrontal cortex; IFG = inferior frontal gyrus; OFC = orbitofrontal cortex.

<https://doi.org/10.1371/journal.pone.0215095.t005>

stimulation effectively hits the entire prefrontal network including the dlPFC-rIFG sub-network, whereas, rIFG stimulation has more restricted effects. Such a global contribution of the dlPFC to other prefrontal region functions has been shown for various executive functions [42] which might be true for ADHD too.

tDCS and WM performance in ADHD

Our meta-analysis showed that tDCS does not have an overall significant effect on WM performance accuracy in ADHD, although it led to significantly faster response time. To be more specific, it was shown that anodal left dlPFC tDCS, but not cathodal stimulation improved response time during WM performance in ADHD. This finding should be discussed considering physiological processes underlying inhibitory control and WM, task characteristics, and results of previous meta-analyses about tDCS effects on WM. First of all, this result shows a dissociation of tDCS effects on inhibitory control and WM. Such a dissociation has also been shown for neuropsychological deficits in ADHD [53]. Inhibitory control is an executive control function [42] and requires executive attention to inhibit a prepotent response and monitoring, whereas memory tasks can deploy attentional resources [53]. Different effects of tDCS on inhibitory control and WM thus could be attributed to different cognitive processes underlying WM and inhibitory control as well as target area-specific effects of tDCS, which need to be studied in future systematically. Characteristics of the respective WM tasks can also affect the results. For example, performance on the n-back task can still be faster while accuracy cannot due to ceiling effects in case of near-optimal accuracy already without intervention.

Another factor that could be a source of variance of the tDCS effects on WM in the included studies is the difference of stimulation sites which is an important feature of tDCS efficacy. The stimulation montages of the studies on WM included anodal left dlPFC-right OFC tDCS (two experiments), left dlPFC-right dlPFC tDCS (two experiments), and left dlPFC-vertex tDCS (two experiments). Different montages used in these studies can explain the observed effects and may explain why some studies demonstrated no effects on WM accuracy. In four of the experiments, the reference electrode was positioned over regions that are involved in cognitive performance (i.e., right OFC and right dlPFC). In this case, it might be speculated that the cathodal return electrode compromises performance, and thus partially antagonizes anodal tDCS effects over the target region. The other two experiments used unilateral protocols that targeted the mere region of interest without positioning the cathodal return electrode over a potentially involved region and interestingly, in these studies both WM accuracy and RT improved, and prolonged effects outlasting the stimulation were also reported. Therefore, an electrode positioning which avoids such antagonistic effects might be advantageous.

Recent meta-analyses and review articles about the effects of tDCS on cognitive functioning showed mixed results regarding memory tasks. For example, a recent systematic review of 188 trials reported significant effects of anodal tDCS on RT only for executive functioning tasks, but not memory tasks in both, healthy and neuropsychiatric populations [80]. Similar to our finding, another meta-analysis of 12 tDCS WM trails found that tDCS significantly improved RT, but not accuracy [78]. A more recent meta-analysis about the effects of tDCS on WM (measured with n-back and digit span tasks) found a significant effect of anodal tDCS on accuracy and RT for offline, but not online tasks in healthy subjects; in neuropsychiatric populations, tDCS had a significant effect on WM accuracy (both offline and online) but not RT [84]. Finally, a recent meta-analysis of 61 studies investigated the effect of tDCS on cognitive tasks in relation to task and stimulation parameters. They found that in contrast to offline-task performance, online-task performance resulted in increased accuracy in clinical populations [94]. Interestingly, in 5 out of 6 studies included in our meta-analysis for WM, offline task performance was measured in ADHD individuals. Thus, the non-significant effect of tDCS on WM accuracy we found could be due to offline task performance in the majority of studies, which might be less efficient. Nevertheless, the results of the effect of tDCS on WM in ADHD should be treated with caution due to the limited number of qualified studies that did not allow us to disentangle polarity- and area-dependent effect of tDCS.

Clinical and methodological implications

This meta-analysis found that modulation of the activity of brain areas involved in the pathophysiology of ADHD with tDCS has a significant effect on response inhibition and WM, two major executive dysfunctions in ADHD [14, 53, 54]. This suggests that tDCS over the dlPFC might be suited as a potential therapeutic approach in ADHD, similar to other neuropsychiatric syndromes which are responsive to this method [45]. Increasing evidence from clinical and cognitive neuroscience describes ADHD as a disorder with functional/structural abnormalities, especially of the frontostriatal circuitry and PFC. Accordingly, modulation of the involved brain areas with non-invasive brain stimulation is a promising approach for improving neuropsychological symptoms in ADHD patients. The efficacy of this method might, however, depend on ADHD subtypes/symptoms and stimulation parameters, which need to be investigated systematically in future studies.

The findings of this meta-analysis have some important clinical and methodological implications to be considered for future tDCS studies in ADHD. The results of the included studies showed beneficial effects of tDCS in ADHD, and at the same time indicate that effects of tDCS

on ADHD deficits (namely inhibitory control and WM) are strongly dependent on 1) individual and inter-individual factors (e.g., type of symptoms/deficits) and 2) also stimulation parameters (i.e., site or cortical target, polarity, intensity, duration, and repetition rate). For example, heterogeneous ADHD subtypes in the analyzed studies (i.e., inattentive or hyperactive) might partially explain the non-significant effect of anodal rIFG tDCS on response inhibition, since dlPFC activity contributes to response inhibition in all ADHD subtypes (especially individuals with inattentive symptoms), while the rIFG is assumed to be specifically involved in ADHD individuals with more hyperactivity subtypes. Therefore, the inter-individual variability is an important factor which generally affects the efficacy of tDCS [95] and might be specifically relevant for the treatment of ADHD patients, that vary in subtypes. Furthermore, findings of those studies that investigated tDCS effects on other ADHD deficits showed that for those EF domains that involve motivational and emotional processing, tDCS over both, prefrontal and frontopolar areas is more effective compared to dlPFC-only tDCS. For example, the Wisconsin Card Sorting Test (WCST) is a well-documented measure of EF and primarily measures cognitive flexibility. This task involves aspects of both, hot (e.g., task switching, disinhibition, WM) and cold EFs (i.e., executive control, inhibition) and results showed that only when both, dlPFC and OFC were stimulated, cognitive flexibility improved. Therefore, the common stimulation protocols in ADHD that primarily target dorsolateral frontostriatal networks may not be ideal for all executive dysfunction in ADHD, especially hot executive dysfunctions. This highlights the necessity of symptom-driven protocols implicating that for each specific impairment (e.g., response inhibition, WM, selective attention, interference control, cognitive flexibility, etc.) a specific stimulation montage that targets the most relevant cortical region would be most appropriate.

Another important implication regarding the clinical efficacy of tDCS in ADHD concerns stimulation parameters (i.e., intensity, duration, repetition rate, polarity, and site). The stimulation intensities applied in the included studies ranged from 1 mA (seven experiments) to 2 mA (two experiments) and two experiments applied tDCS with 1.5 mA intensity. Findings from other clinical populations showed that higher intensities of stimulation can result in more prominent symptom improvement, for example in tinnitus [96], or cognitive impairment in Parkinson's disease [97]. This is in further accordance with physiological findings of tDCS about larger tDCS after-effects in motor cortex plasticity as a result of tDCS with higher intensities [98]. This was supported by the findings from two studies conducted on adult ADHD. The study with 2 mA intensity reported significant improvement in inhibitory control performance [68] while the study with 1 mA intensity reported no significant effects on Go/No-Go task performance [71] despite the fact both studies targeted left dlPFC with anodal tDCS. Stimulation intensity has, therefore, an impact on the effectiveness of tDCS, which is not necessarily linear [99] and should be considered in clinical applications.

With regard to stimulation duration, the length of stimulation duration varied from 8 min to 30 min and the longest duration of stimulation reported was five consecutive days of 30-min stimulation (Tables 1 & 2). The effects of different stimulation durations were not addressed systematically in ADHD studies so far. Longer stimulation duration, similar to stimulation intensity, could be an important factor to improve the efficacy of tDCS effects, including clinical effects. Prolongation of stimulation duration for increasing efficacy of the intervention works similarly well as enhancing stimulation intensity. One advantage of enhancing stimulation duration compared to intensity might be that this does not increase the probability of side effects like itching and tingling [33, 100], which might be specifically relevant for studies in children. Findings from other neuropsychiatric disorders show that repeated sessions of tDCS are more effective in reducing symptoms [33]. All studies included in this meta-analysis except one [69] examined effects of tDCS after one single session. Higher

clinical efficacy might be achieved with a longer duration of stimulation, and repetitive interventions, as shown for example in non-invasive brain stimulation studies in depression [101–103].

Stimulation polarity is another important parameter that determines the efficacy of tDCS. The physiological mechanism of tDCS effects, depending on the polarity, is the induction of LTP- and LTD-like plasticity [36, 104]. Anodal tDCS induces LTP-like plasticity, based on sub-threshold depolarization effects on membrane potentials, and respective enhancement of spontaneous neuronal activity, while cathodal tDCS has antagonistic effects [105]. In accordance with the pathophysiological foundation of ADHD that includes underactivation of the lateral and inferior PFC, anodal tDCS-generated excitability enhancement is conceptually a more promising approach in ADHD. The findings of this meta-analysis are in accordance with this assumption and showed that anodal, but not cathodal tDCS improved inhibitory control and WM in ADHD which could be attributed to the stimulation-induced compensation for regional cortical hypoactivity as well as alteration of functional cortical network connectivity following anodal tDCS that improves cognitive performance [106]. Nevertheless, this does not mean that cathodal tDCS is not of clinical interest in ADHD. Functional abnormalities in ADHD are not limited to hyperactivation of specific regions but also hyper-sensitiveness of other regions, especially those involved in motivational and emotional processing [11]. Moreover, findings from cathodal stimulation of the motor cortex showed that motor cortex excitability alterations induced by tDCS are intensity-dependent and nonlinear [99] which is not well studied for non-motor functions. It might be that cathodal tDCS over regions pathologically hyperactive in the disease might also be beneficial, which has not been systematically explored so far.

The cortical target area or stimulation site, another important parameter, contributes to the clinical efficacy of tDCS in ADHD. The contribution of this parameter was specifically clear in the studies which investigated tDCS effects on ADHD WM deficits. Only when the potentially involved cortical region (left dlPFC in WM) was stimulated with the target electrode, both WM accuracy and RT significantly improved and the effects were long-lasting. In these cases, the reference electrode was placed over a non-contributing area (i.e., vertex). In contrast, when the reference electrode targeted homologous regions of the contralateral hemisphere (i.e., right dlPFC or right supraorbital) the effects were not present or reduced to RT improvement only. A comparable pattern but in a different way was observed for tDCS effects on inhibitory control. Here, tDCS effects were more effective when two specific potentially involved regions were stimulated by the target and reference electrodes. The left dlPFC-right OFC montage was the most effective montage and all of the four experiments which used this protocol reported improved performance. In contrast, the left dlPFC-right dlPFC tDCS montage, which was used in two studies, did not improve any inhibitory control outcome measure. This demonstrates that the target area or areas significantly contribute to the tDCS effectiveness in ADHD.

To conclude, results from this meta-analysis support the cognitive benefits of tDCS in ADHD. However, for these cognitive benefits to be clinically useful, effects need to be sustained for longer durations. All of the studies included in this meta-analysis applied tDCS in an acute and short-term mode with no follow-up examination which does not allow to infer long-term efficacy of tDCS in improving ADHD neuropsychological deficits. Similarly, measuring inhibitory control and WM was done either during stimulation or right/shortly after intervention which hinders any robust conclusions about long-term effects of tDCS. One specific important note here is the dissociation between neuropsychological deficits and clinical symptoms of ADHD, which means improvement in inhibitory control and WM after or during tDCS does not say too much about the improvement of clinical symptoms. Moreover, applying statistically-proven blinding procedures, which was not reported in most of the included studies in this meta-analysis, is important for making firm conclusions about the

clinical efficacy of tDCS. Therefore, the results of this meta-analysis should be interpreted with caution, especially when it comes to the clinical efficacy of tDCS in ADHD. Nevertheless, we need to note that the studies included in this meta-analysis are actually the first tDCS studies on ADHD that primarily aimed to investigate whether tDCS has any beneficial effects on ADHD symptoms or not. Despite short-term tDCS-induced cognitive benefits in ADHD, the clinical efficacy of this technique needs further investigation in follow-up designs and in comparison to other interventions. Moreover, it requires stimulation protocols optimization using the information about the individual neural activity associated with deficits and task execution (individual level) and general stimulation parameters [49].

Safety of tDCS

There are unique issues concerning the safety, applicability, and ethics of tDCS application in pediatric populations which is mainly due to limited available data from children compared to the adult population [52]. However, the general safety of tDCS with standard protocols has been proven by a large body of evidence in recent years and a recent review concluded that application of conventional tDCS in human trials has not yet produced any reports of a Serious Adverse Effect or irreversible injury [107]. In pediatric populations, no severe adverse events have been reported, and even in children with epilepsy, seizures do not seem to worsen with tDCS [52]. The most frequently reported side-effects within studies included in that systematic review were headache, itchiness, and redness at the site of the stimulation. Similarly, no significant side-effects were reported in the studies included in our meta-analysis. This is important due to some concerns about the application of tDCS in children. One specific concern is about current intensity due to children's thinner skulls and the smaller distance between scalp and brain. Computational models of current flow within the brain suggest that about 50% of the current strength applied in adults result in respective effects in children [52] implicating that 0.5 mA applied in children results in similar physiological effects as 1 mA in adults [108]. From the studies included in this meta-analysis, eight studies applied a current intensity of 1 mA or lower, two studies applied 1.5 mA and one study applied 2 mA, and no significant adverse effect was reported (See Tables 1 & 2). However, the caveat still stands that studies included in this meta-analysis were not designed as safety studies and mostly included single session tDCS interventions. Therefore, systematic safety monitoring, especially for the clinical application of tDCS, is recommended. Moreover, it is important to keep in mind that the developing brain has "sensitive" or "critical" periods where the effects of interventions affecting the brain could be stronger than usual. This suggests that the risk to induce maladaptive neural plasticity due to tDCS might be high which necessitates the priority of dose-finding studies and longitudinal monitoring of tDCS-induced neuroplasticity in pediatric ADHD population [49].

Limitations and future directions

Despite promising results and novelty, this meta-analysis has some limitations to be considered. First, despite that our meta-analysis found a significant effect of tDCS on neuropsychological symptoms (namely inhibitory control and WM) in ADHD populations, the efficacy of this method on other executive dysfunctions, and clinical symptoms cannot be directly derived from these results. Inhibitory control and WM were chosen because they are two common deficits examined in most ADHD-tDCS studies which allowed us to run a meta-analytic study. Secondly, although we showed that anodal tDCS over the dlPFC has significant effects on IC in ADHD, no effects were seen for cathodal tDCS and rIFG tDCS. However, given the small number of studies investigating cathodal/rIFG tDCS, these results have to be interpreted with caution. With increasing interest in the application of tDCS in ADHD, future research might

be able to disentangle subtle protocol differences for ADHD subtypes. Thirdly, adaptation of stimulation protocols based on symptom subtypes, ADHD subgroups, specific cognitive deficits, and neuroanatomical differences is lacking in currently available studies and therefore, it is difficult to rate the potential of tDCS in ADHD.

Future tDCS studies should systematically investigate the following lines of research, to reveal the real clinical potential of tDCS for ADHD treatment. First, preliminary data suggest beneficial effects of tDCS on other cognitive dysfunctions in ADHD including cognitive flexibility, problem-solving, selective attention, and hyperactivity symptoms involved in ADHS [6, 70, 109]. Second, based on the findings of this meta-analysis, future tDCS studies on ADHD populations, with a statistically-proven blinding procedure (especially double-blinded trials), are recommended to target the stimulation area based on the symptoms subtypes. The above-mentioned stimulation parameters are other important factors that might affect the efficacy of the intervention and should be explored systematically. Lastly, future studies, especially those with clinical implications, require examining the long-term efficacy of tDCS in ADHD. With the expected increasing number of tDCS studies in ADHD populations, future meta-analytic studies might be able to deliver a more realistic picture about the effectiveness of tDCS in different subtypes of ADHD. It is lastly of note that other modalities of electrical stimulation such as transcranial alternating current stimulation (tACS), which has been shown to have neuroplastic effects beyond its impact on brain oscillations [110], or transcranial random noise stimulation (tRNS) can be a potentially interesting avenue for future research in ADHD, which is associated with abnormal brain oscillations [24].

Conclusion

The findings of this meta-analysis of tDCS interventions in ADHD suggest an improvement of neuropsychological deficits (i.e., inhibitory control and WM) by tDCS. Stimulation polarity and target area are relevant for the efficacy of tDCS in ADHD. Anodal dlPFC tDCS had a significantly superior effect on inhibitory control compared to cathodal/sham stimulation and anodal rIFG tDCS. TDCS significantly increased response accuracy of inhibitory control performance and decreased response time in WM tasks. Although our findings suggest improving effects of tDCS in ADHD neuropsychological deficits, the clinical utility of tDCS cannot be firmly rated with the currently available findings. Application of this method as a therapeutic intervention will require optimizing stimulation protocols based on general stimulation parameters and individual and inter-individual factors for improvement of clinical efficacy, exploration of clinical symptoms in addition to surrogate parameters, and achievement of sustained clinical benefits by tDCS over longer durations of time. Thus, future research is needed to more thoroughly explore and refine optimal stimulation parameters required for tDCS-based cognitive improvement and implementing robust experimental designs in different ADHD subtypes. Broadly speaking, the potential for tDCS as a non-invasive brain stimulation technique to safely improve neuroplasticity and treat neurological and neurodevelopmental disorders is encouraging. Future studies utilizing tDCS will further increase our understanding of neural networks and how to treat their pathological states in ADHD and other neurodevelopmental disorders including autism and learning disabilities.

Supporting information

S1 Table. The Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) 2009 checklist.

(DOC)

Acknowledgments

Publication of this article was funded by the Open Access Fund of the Leibniz Association.

Author Contributions

Conceptualization: Mohammad Ali Salehinejad.

Formal analysis: Miles Wischnewski.

Funding acquisition: Michael A. Nitsche.

Methodology: Mohammad Ali Salehinejad, Miles Wischnewski.

Supervision: Vahid Nejati, Carmelo M. Vicario, Michael A. Nitsche.

Visualization: Miles Wischnewski.

Writing – original draft: Mohammad Ali Salehinejad, Miles Wischnewski.

Writing – review & editing: Mohammad Ali Salehinejad, Michael A. Nitsche.

References

1. Visser SN, Danielson ML, Bitsko RH, Holbrook JR, Kogan MD, Ghandour RM, et al. Trends in the Parent-Report of Health Care Provider-Diagnosed and Medicated Attention-Deficit/Hyperactivity Disorder: United States, 2003–2011. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2014; 53(1):34–46.e2. <https://doi.org/10.1016/j.jaac.2013.09.001> PMID: 24342384
2. American Psychiatric Association. Diagnostic and statistical manual of mental disorders (DSM-5): American Psychiatric Pub; 2013.
3. Castellanos FX, Sonuga-Barke EJS, Milham MP, Tannock R. Characterizing cognition in ADHD: beyond executive dysfunction. *Trends in Cognitive Sciences*. 2006; 10(3):117–23. <https://doi.org/10.1016/j.tics.2006.01.011> PMID: 16460990
4. Hudec KL, Alderson RM, Patros CHG, Lea SE, Tarle SJ, Kasper LJ. Hyperactivity in boys with attention-deficit/hyperactivity disorder (ADHD): The role of executive and non-executive functions. *Research in Developmental Disabilities*. 2015;45–46:103–9. <http://dx.doi.org/10.1016/j.ridd.2015.07.012>.
5. Cortese S, Ferrin M, Brandeis D, Buitelaar J, Daley D, Dittmann RW, et al. Cognitive Training for Attention-Deficit/Hyperactivity Disorder: Meta-Analysis of Clinical and Neuropsychological Outcomes From Randomized Controlled Trials. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2015; 54(3):164–74. <https://doi.org/10.1016/j.jaac.2014.12.010>.
6. Nejati V, Salehinejad MA, Nitsche MA, Najian A, Javadi A-H. Transcranial Direct Current Stimulation Improves Executive Dysfunctions in ADHD: Implications for Inhibitory Control, Interference Control, Working Memory, and Cognitive Flexibility. *Journal of Attention Disorders*. 2017;0(0):1087054717730611. <https://doi.org/10.1177/1087054717730611> PMID: 28938852.
7. Kasper LJ, Alderson RM, Hudec KL. Moderators of working memory deficits in children with attention-deficit/hyperactivity disorder (ADHD): A meta-analytic review. *Clinical Psychology Review*. 2012; 32(7):605–17. <https://doi.org/10.1016/j.cpr.2012.07.001> PMID: 22917740
8. Willcutt EG, Doyle AE, Nigg JT, Faraone SV, Pennington BF. Validity of the Executive Function Theory of Attention-Deficit/Hyperactivity Disorder: A Meta-Analytic Review. *Biological Psychiatry*. 2005; 57(11):1336–46. <https://doi.org/10.1016/j.biopsych.2005.02.006> PMID: 15950006
9. Roy A, Oldehinkel AJ, Hartman CA. Cognitive Functioning in Adolescents with Self-Reported ADHD and Depression: Results from a Population-Based Study. *Journal of Abnormal Child Psychology*. 2017; 45(1):69–81. <https://doi.org/10.1007/s10802-016-0160-x> PMID: 27138748
10. Roberts M, Ashinoff BK, Castellanos FX, Carrasco M. When attention is intact in adults with ADHD. *Psychonomic Bulletin & Review*. 2017. <https://doi.org/10.3758/s13423-017-1407-4> PMID: 29181782
11. Rubia K. Cognitive Neuroscience of Attention Deficit Hyperactivity Disorder (ADHD) and Its Clinical Translation. *Frontiers in Human Neuroscience*. 2018; 12(100). <https://doi.org/10.3389/fnhum.2018.00100> PMID: 29651240
12. Fair DA, Bathula D, Nikolas MA, Nigg JT. Distinct neuropsychological subgroups in typically developing youth inform heterogeneity in children with ADHD. *Proceedings of the National Academy of Sciences*. 2012; 109(17):6769–74.

13. Samea F, Soluki S, Nejati V, Zarei M, Cortese S, Eickhoff SB, et al. Brain alterations in children/adolescents with ADHD revisited: A neuroimaging meta-analysis of 96 structural and functional studies. *Neuroscience & Biobehavioral Reviews*. 2019; 100:1–8. <https://doi.org/10.1016/j.neubiorev.2019.02.011>.
14. Nigg JT. Neuropsychologic Theory and Findings in Attention-Deficit/Hyperactivity Disorder: The State of the Field and Salient Challenges for the Coming Decade. *Biological Psychiatry*. 2005; 57(11):1424–35. <https://doi.org/10.1016/j.biopsych.2004.11.011> PMID: 15950017
15. Rubio B, Boes AD, Laganieri S, Rotenberg A, Jeurissen D, Pascual-Leone A. Noninvasive Brain Stimulation in Pediatric Attention-Deficit Hyperactivity Disorder (ADHD): A Review. *Journal of Child Neurology*. 2016; 31(6):784–96. <https://doi.org/10.1177/0883073815615672> PMID: 26661481.
16. Zhu Y, Jiang X, Ji W. The Mechanism of Cortico-Striato-Thalamo-Cortical Neurocircuitry in Response Inhibition and Emotional Responding in Attention Deficit Hyperactivity Disorder with Comorbid Disruptive Behavior Disorder. *Neurosci Bull*. 2018. <https://doi.org/10.1007/s12264-018-0214-x> PMID: 29508250
17. Seidman LJ, Valera EM, Makris N. Structural Brain Imaging of Attention-Deficit/Hyperactivity Disorder. *Biological Psychiatry*. 2005; 57(11):1263–72. <https://doi.org/10.1016/j.biopsych.2004.11.019> PMID: 15949998
18. Depue BE, Burgess GC, Willcutt EG, Ruzic L, Banich MT. Inhibitory control of memory retrieval and motor processing associated with the right lateral prefrontal cortex: Evidence from deficits in individuals with ADHD. *Neuropsychologia*. 2010; 48(13):3909–17. <https://doi.org/10.1016/j.neuropsychologia.2010.09.013> PMID: 20863843
19. Belen R, Aaron DB, Simon L, Alexander R, Danique J, Alvaro P-L. Noninvasive Brain Stimulation in Pediatric Attention-Deficit Hyperactivity Disorder (ADHD): A Review. *Journal of Child Neurology*. 2015; 31(6):784–96. <https://doi.org/10.1177/0883073815615672> PMID: 26661481
20. Langleben DD, Austin G, Krikorian G, Ridlehuber HW, Goris ML, Strauss HW. Interhemispheric asymmetry of regional cerebral blood flow in prepubescent boys with attention deficit hyperactivity disorder. *Nuclear Medicine Communications*. 2001; 22(12):1333–40. 00006231-200112000-00009. PMID: 11711904
21. Christakou A, Murphy CM, Chantiluke K, Cubillo AI, Smith AB, Giampietro V, et al. Disorder-specific functional abnormalities during sustained attention in youth with Attention Deficit Hyperactivity Disorder (ADHD) and with Autism. *Mol Psychiatry*. 2012; 18:236. <https://doi.org/10.1038/mp.2011.185> <https://www.nature.com/articles/mp2011185#supplementary-information>. PMID: 22290121
22. Bush G, Valera EM, Seidman LJ. Functional Neuroimaging of Attention-Deficit/Hyperactivity Disorder: A Review and Suggested Future Directions. *Biological Psychiatry*. 2005; 57(11):1273–84. <https://doi.org/10.1016/j.biopsych.2005.01.034> PMID: 15949999
23. Cai W, Chen T, Szegletes L, Supekar K, Menon V. Aberrant Time-Varying Cross-Network Interactions in Children With Attention-Deficit/Hyperactivity Disorder and the Relation to Attention Deficits. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*. 2018; 3(3):263–73. <https://doi.org/10.1016/j.bpsc.2017.10.005>.
24. Marcos-Vidal L, Martínez-García M, Pretus C, Garcia-Garcia D, Martínez K, Janssen J, et al. Local functional connectivity suggests functional immaturity in children with attention-deficit/hyperactivity disorder. *Hum Brain Mapp*. 2018:n/a-n/a. <https://doi.org/10.1002/hbm.24013> PMID: 29473262
25. Hoogman M, Bralten J, Hibar DP, Mennes M, Zwiers MP, Schweren LSJ, et al. Subcortical brain volume differences in participants with attention deficit hyperactivity disorder in children and adults: a cross-sectional mega-analysis. *The Lancet Psychiatry*. 2017; 4(4):310–9. [https://doi.org/10.1016/S2215-0366\(17\)30049-4](https://doi.org/10.1016/S2215-0366(17)30049-4) PMID: 28219628
26. Sibley MH, Kuriyan AB, Evans SW, Waxmonsky JG, Smith BH. Pharmacological and psychosocial treatments for adolescents with ADHD: An updated systematic review of the literature. *Clinical Psychology Review*. 2014; 34(3):218–32. <https://doi.org/10.1016/j.cpr.2014.02.001> PMID: 24632046
27. Catalá-López F, Hutton B, Núñez-Beltrán A, Mayhew AD, Page MJ, Ridao M, et al. The pharmacological and non-pharmacological treatment of attention deficit hyperactivity disorder in children and adolescents: protocol for a systematic review and network meta-analysis of randomized controlled trials. *Systematic Reviews*. 2015; 4(1):19. <https://doi.org/10.1186/s13643-015-0005-7> PMID: 25875125
28. Cortese S, Ferrin M, Brandeis D, Holtmann M, Aggensteiner P, Daley D, et al. Neurofeedback for Attention-Deficit/Hyperactivity Disorder: Meta-Analysis of Clinical and Neuropsychological Outcomes From Randomized Controlled Trials. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2016; 55(6):444–55. <https://doi.org/10.1016/j.jaac.2016.03.007>.
29. Molina BSG, Hinshaw SP, Swanson JM, Arnold LE, Vitiello B, Jensen PS, et al. The MTA at 8 Years: Prospective Follow-up of Children Treated for Combined-Type ADHD in a Multisite Study. *Journal of*

- the American Academy of Child & Adolescent Psychiatry. 2009; 48(5):484–500. <https://doi.org/10.1097/CHI.0b013e31819c23d0>.
30. Faraone SV, Biederman J, Spencer TJ, Aleardi M. Comparing the Efficacy of Medications for ADHD Using Meta-analysis. *Medscape General Medicine*. 2006; 8(4):4–. PMID: [17415287](https://pubmed.ncbi.nlm.nih.gov/17415287/)
 31. Hsu S, Piedra AN, Froehlich TE, Brinkman WB, Epstein J. 6.18 Caregiver-Reported Side Effects of Methylphenidate and Comorbid Oppositional Defiant Disorder Diagnosis in Children With Attention-Deficit/Hyperactivity Disorder (ADHD). *Journal of the American Academy of Child & Adolescent Psychiatry*. 2017; 56(10, Supplement):S282. <https://doi.org/10.1016/j.jaac.2017.09.363>.
 32. Daley D, van der Oord S, Ferrin M, Danckaerts M, Doepfner M, Cortese S, et al. Behavioral Interventions in Attention-Deficit/Hyperactivity Disorder: A Meta-Analysis of Randomized Controlled Trials Across Multiple Outcome Domains. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2014; 53(8):835–47. <https://doi.org/10.1016/j.jaac.2014.05.013>.
 33. Lefaucheur J-P, Antal A, Ayache SS, Benninger DH, Brunelin J, Cogiamanian F, et al. Evidence-based guidelines on the therapeutic use of transcranial direct current stimulation (tDCS). *Clinical Neurophysiology*. 2017; 128(1):56–92. <https://doi.org/10.1016/j.clinph.2016.10.087> PMID: [27866120](https://pubmed.ncbi.nlm.nih.gov/27866120/)
 34. Vicario CM, Salehinejad MA, Felmingham K, Martino G, Nitsche MA. A systematic review on the therapeutic effectiveness of non-invasive brain stimulation for the treatment of anxiety disorders. *Neuroscience & Biobehavioral Reviews*. 2019; 96:219–31. <https://doi.org/10.1016/j.neubiorev.2018.12.012>.
 35. Harty S, Brem A-K, Cohen Kadosh R. Neurocognitive Effects of tDCS in the Healthy Brain. In: Brunoni A, Nitsche M, Loo C, editors. *Transcranial Direct Current Stimulation in Neuropsychiatric Disorders: Clinical Principles and Management*. Cham: Springer International Publishing; 2016. p. 103–41.
 36. Nitsche MA, Paulus W. Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *The Journal of Physiology*. 2000; 527(3):633–9. <https://doi.org/10.1111/j.1469-7793.2000.t01-1-00633.x> PMID: [10990547](https://pubmed.ncbi.nlm.nih.gov/10990547/)
 37. Kuo M-F, Polanía R, Nitsche M. Physiology of Transcranial Direct and Alternating Current Stimulation. In: Brunoni A, Nitsche M, Loo C, editors. *Transcranial Direct Current Stimulation in Neuropsychiatric Disorders: Clinical Principles and Management*. Cham: Springer International Publishing; 2016. p. 29–46.
 38. Nitsche Boggio, Fregni Pascual-Leone. Treatment of depression with transcranial direct current stimulation (tDCS): a review. *Experimental neurology*. 2009; 219(1):14–9. <https://doi.org/10.1016/j.expneurol.2009.03.038> PMID: [19348793](https://pubmed.ncbi.nlm.nih.gov/19348793/)
 39. Nitsche MA, Cohen LG, Wassermann EM, Priori A, Lang N, Antal A, et al. Transcranial direct current stimulation: State of the art 2008. *Brain Stimulation*. 2008; 1(3):206–23. <https://doi.org/10.1016/j.brs.2008.06.004> PMID: [20633386](https://pubmed.ncbi.nlm.nih.gov/20633386/)
 40. Woods AJ, Martin DM. Clinical Research and Methodological Aspects for tDCS Research. In: Brunoni A, Nitsche M, Loo C, editors. *Transcranial Direct Current Stimulation in Neuropsychiatric Disorders: Clinical Principles and Management*. Cham: Springer International Publishing; 2016. p. 393–404.
 41. Ghanavati E, Salehinejad MA, Nejati V, Nitsche MA. Differential role of prefrontal, temporal and parietal cortices in verbal and figural fluency: Implications for the supramodal contribution of executive functions. *Sci Rep*. 2019; 9(1):3700. <https://doi.org/10.1038/s41598-019-40273-7> PMID: [30842493](https://pubmed.ncbi.nlm.nih.gov/30842493/)
 42. Nejati V, Salehinejad MA, Nitsche MA. Interaction of the Left Dorsolateral Prefrontal Cortex (l-DLPFC) and Right Orbitofrontal Cortex (OFC) in Hot and Cold Executive Functions: Evidence from Transcranial Direct Current Stimulation (tDCS). *Neuroscience*. 2018; 369(Supplement C):109–23. <https://doi.org/10.1016/j.neuroscience.2017.10.042>.
 43. Kuo M-F, Nitsche MA. Effects of Transcranial Electrical Stimulation on Cognition. *Clin EEG Neurosci*. 2012; 43(3):192–9. <https://doi.org/10.1177/1550059412444975> PMID: [22956647](https://pubmed.ncbi.nlm.nih.gov/22956647/).
 44. Ghanavati E, Nejati V, Salehinejad MA. Transcranial Direct Current Stimulation over the Posterior Parietal Cortex (PPC) Enhances Figural Fluency: Implications for Creative Cognition. *Journal of Cognitive Enhancement*. 2018; 2(1):88–96. <https://doi.org/10.1007/s41465-017-0059-7>
 45. Lefaucheur J-P. A comprehensive database of published tDCS clinical trials (2005–2016). *Neurophysiologie Clinique/Clinical Neurophysiology*. 2016; 46(6):319–98. <https://doi.org/10.1016/j.neucli.2016.10.002> PMID: [27865707](https://pubmed.ncbi.nlm.nih.gov/27865707/)
 46. Filmer HL, Dux PE, Mattingley JB. Applications of transcranial direct current stimulation for understanding brain function. *Trends in Neurosciences*. 2014; 37(12):742–53. <https://doi.org/10.1016/j.tins.2014.08.003> PMID: [25189102](https://pubmed.ncbi.nlm.nih.gov/25189102/)
 47. Shah-Basak PP, Hamilton RH, Nitsche MA, Woods AJ. Transcranial Direct Current Stimulation in Cognitive Neuroscience. In: Knotkova H, Nitsche MA, Bikson M, Woods AJ, editors. *Practical Guide to Transcranial Direct Current Stimulation: Principles, Procedures and Applications*. Cham: Springer International Publishing; 2019. p. 597–625.

48. Vicario C, Nitsche M. Transcranial direct current stimulation: a remediation tool for the treatment of childhood congenital dyslexia? *Frontiers in Human Neuroscience*. 2013; 7(139). <https://doi.org/10.3389/fnhum.2013.00139> PMID: 23626530
49. Vicario CM, Nitsche MA. Chapter 9—tDCS in Pediatric Neuropsychiatric Disorders. In: Oberman LM, Enticott PG, editors. *Neurotechnology and Brain Stimulation in Pediatric Psychiatric and Neurodevelopmental Disorders*: Academic Press; 2019. p. 217–35.
50. Ciecchanski P, Kirton A. Chapter 5—Transcranial Direct-Current Stimulation (tDCS): Principles and Emerging Applications in Children. *Pediatric Brain Stimulation*. Oxford: Academic Press; 2016. p. 85–115.
51. Stortelder F, Ploegmakers-Burg M. Adolescence and the Reorganization of Infant Development: A Neuro-Psychoanalytic Model. *The Journal of the American Academy of Psychoanalysis and Dynamic Psychiatry*. 2010; 38(3):503–31. <https://doi.org/10.1521/jaap.2010.38.3.503> PMID: 20849240
52. El-Hagrassy MM, Jones F, Rosa G, Fregni F. CNS Non-invasive Brain Stimulation. In: Gillera JP, Alpert SA, editors. *Adult and Pediatric Neuromodulation*. Cham: Springer International Publishing; 2018. p. 151–84.
53. Sergeant JA, Geurts H, Huijbregts S, Scheres A, Oosterlaan J. The top and the bottom of ADHD: a neuropsychological perspective. *Neuroscience & Biobehavioral Reviews*. 2003; 27(7):583–92. <https://doi.org/10.1016/j.neubiorev.2003.08.004>.
54. Nigg JT, Gustafsson HC, Karalunas SL, Ryabinin P, McWeeney SK, Faraone SV, et al. Working Memory and Vigilance as Multivariate Endophenotypes Related to Common Genetic Risk for Attention-Deficit/Hyperactivity Disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2018; 57(3):175–82. <https://doi.org/10.1016/j.jaac.2017.12.013>.
55. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *International Journal of Surgery*. 2010; 8(5):336–41. <https://doi.org/10.1016/j.ijsu.2010.02.007> PMID: 20171303
56. Sonuga-Barke EJS. Causal Models of Attention-Deficit/Hyperactivity Disorder: From Common Simple Deficits to Multiple Developmental Pathways. *Biological Psychiatry*. 2005; 57(11):1231–8. <https://doi.org/10.1016/j.biopsych.2004.09.008> PMID: 15949993
57. Castellanos FX, Proal E. Large-scale brain systems in ADHD: beyond the prefrontal–striatal model. *Trends in Cognitive Sciences*. 2012; 16(1):17–26. <https://doi.org/10.1016/j.tics.2011.11.007> PMID: 22169776
58. Georgiou G, Essau CA. Go/No-Go Task. In: Goldstein S, Naglieri JA, editors. *Encyclopedia of Child Behavior and Development*. Boston, MA: Springer US; 2011. p. 705–6.
59. Stuss DT, Floden D, Alexander MP, Levine B, Katz D. Stroop performance in focal lesion patients: dissociation of processes and frontal lobe lesion location. *Neuropsychologia*. 2001; 39(8):771–86. [https://doi.org/10.1016/S0028-3932\(01\)00013-6](https://doi.org/10.1016/S0028-3932(01)00013-6). PMID: 11369401
60. Mullane JC, Corkum PV, Klein RM, McLaughlin E. Interference Control in Children with and without ADHD: A Systematic Review of Flanker and Simon Task Performance. *Child Neuropsychology*. 2009; 15(4):321–42. <https://doi.org/10.1080/09297040802348028> PMID: 18850349
61. Conners CK, Staff M, Connelly V, Campbell S, MacLean M, Barnes J. Conners' continuous performance Test II (CPT II v. 5). Multi-Health Syst Inc. 2000; 29:175–96.
62. Brooks BL, Sherman EMS, Strauss E. NEPSY-II: A Developmental Neuropsychological Assessment, Second Edition. *Child Neuropsychology*. 2009; 16(1):80–101. <https://doi.org/10.1080/09297040903146966>
63. Kessels RPC, van Zandvoort MJE, Postma A, Kappelle LJ, de Haan EHF. The Corsi Block-Tapping Task: Standardization and Normative Data. *Appl Neuropsychol*. 2000; 7(4):252–8. https://doi.org/10.1207/S15324826AN0704_8 PMID: 11296689
64. Higgins JPT, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011; 343. <https://doi.org/10.1136/bmj.d5928> PMID: 22008217
65. Larry H, Ingram O. *Statistical Methods for Meta Analysis*. Orlando, FL: Academic Press; 1985.
66. Rosenthal R. The file drawer problem and tolerance for null results. *Psychological Bulletin*. 1979; 86(3):638–41. <https://doi.org/10.1037/0033-2909.86.3.638>
67. Rosenberg MS, Adams DC, Gurevitch J. *MetaWin: statistical software for meta-analysis*. Sunderland, MA: Sinauer Associates Sunderland, MA; 2000.
68. Allenby C, Falcone M, Bernardo L, Wileyto P, Rostain A, Ramsay JR, et al. Transcranial Direct Current Brain Stimulation Decreases Impulsivity in ADHD. *Brain Stimulation*. 2018. <https://doi.org/10.1016/j.brs.2018.04.016>.

69. Bandeira ID, Guimarães RSQ, Jagersbacher JG, Barretto TL, Jesus-Silva JRd, Santos SN, et al. Transcranial Direct Current Stimulation in Children and Adolescents With Attention-Deficit/Hyperactivity Disorder (ADHD): A Pilot Study. *Journal of Child Neurology*. 2016; 31(7):918–24. <https://doi.org/10.1177/0883073816630083> PMID: 26879095.
70. Breitling C, Zaehle T, Dannhauer M, Bonath B, Tegelbeckers J, Flechtner H-H, et al. Improving Interference Control in ADHD Patients with Transcranial Direct Current Stimulation (tDCS). *Front Cell Neurosci*. 2016; 10:72. <https://doi.org/10.3389/fncel.2016.00072> PMC4834583. PMID: 27147964
71. Cosmo C, Baptista AF, de Araújo AN, do Rosário RS, Miranda JGV, Montoya P, et al. A Randomized, Double-Blind, Sham-Controlled Trial of Transcranial Direct Current Stimulation in Attention-Deficit/Hyperactivity Disorder. *PLOS ONE*. 2015; 10(8):e0135371. <https://doi.org/10.1371/journal.pone.0135371> PMID: 26267861
72. Munz MT, Prehn-Kristensen A, Thielking F, Mölle M, Göder R, Baving L. Slow oscillating transcranial direct current stimulation during non-rapid eye movement sleep improves behavioral inhibition in attention-deficit/hyperactivity disorder. *Front Cell Neurosci*. 2015; 9:307. <https://doi.org/10.3389/fncel.2015.00307> PMC4531340. PMID: 26321911
73. Soltaninejad Z, Nejati V, Ekhtiari H. Effect of Anodal and Cathodal Transcranial Direct Current Stimulation on DLPFC on Modulation of Inhibitory Control in ADHD. *Journal of Attention Disorders*. 2015. <https://doi.org/10.1177/1087054715618792> PMID: 26689935
74. Sotnikova A, Soff C, Tagliazucchi E, Becker K, Siniatchkin M. Transcranial Direct Current Stimulation Modulates Neuronal Networks in Attention Deficit Hyperactivity Disorder. *Brain Topogr*. 2017; 30(5):656–72. <https://doi.org/10.1007/s10548-017-0552-4> PMID: 28213645
75. Prehn-Kristensen A, Munz M, Göder R, Wilhelm I, Korr K, Vahl W, et al. Transcranial Oscillatory Direct Current Stimulation During Sleep Improves Declarative Memory Consolidation in Children With Attention-deficit/hyperactivity Disorder to a Level Comparable to Healthy Controls. *Brain Stimulation*. 2014; 7(6):793–9. <https://doi.org/10.1016/j.brs.2014.07.036> PMID: 25153776
76. Soff C, Sotnikova A, Christiansen H, Becker K, Siniatchkin M. Transcranial direct current stimulation improves clinical symptoms in adolescents with attention deficit hyperactivity disorder. *J Neural Transm*. 2017; 124(1):133–44. <https://doi.org/10.1007/s00702-016-1646-y> PMID: 27853926
77. Cosmo C, Ferreira C, Miranda JG, do Rosário R, Baptista A, Montoya P, et al. Spreading effect of tDCS in individuals with attention-deficit/hyperactivity disorder as shown by functional cortical networks: a randomized, double-blind, sham-controlled trial. *Frontiers in Psychiatry*. 2015; 6(111). <https://doi.org/10.3389/fpsyt.2015.00111> PMID: 26300790
78. Brunoni AR, Vanderhasselt M-A. Working memory improvement with non-invasive brain stimulation of the dorsolateral prefrontal cortex: A systematic review and meta-analysis. *Brain and Cognition*. 2014; 86:1–9. <https://doi.org/10.1016/j.bandc.2014.01.008> PMID: 24514153
79. Price AR, McAdams H, Grossman M, Hamilton RH. A Meta-analysis of Transcranial Direct Current Stimulation Studies Examining the Reliability of Effects on Language Measures. *Brain Stimulation*. 2015; 8(6):1093–100. <https://doi.org/10.1016/j.brs.2015.06.013> PMID: 26210573
80. Dedoncker J, Brunoni AR, Baeken C, Vanderhasselt M-A. The effect of the interval-between-sessions on prefrontal transcranial direct current stimulation (tDCS) on cognitive outcomes: a systematic review and meta-analysis. *J Neural Transm*. 2016; 123(10):1159–72. <https://doi.org/10.1007/s00702-016-1558-x> PMID: 27145765
81. Kalu UG, Sexton CE, Loo CK, Ebmeier KP. Transcranial direct current stimulation in the treatment of major depression: a meta-analysis. *Psychological Medicine*. 2012; 42(9):1791–800. Epub 2012/01/12. <https://doi.org/10.1017/S0033291711003059> PMID: 22236735
82. Bastani A, Jaberzadeh S. Does anodal transcranial direct current stimulation enhance excitability of the motor cortex and motor function in healthy individuals and subjects with stroke: A systematic review and meta-analysis. *Clinical Neurophysiology*. 2012; 123(4):644–57. <https://doi.org/10.1016/j.clinph.2011.08.029> PMID: 21978654
83. Shiozawa P, Fregni F, Benseñor IM, Lotufo PA, Berlim MT, Daskalakis JZ, et al. Transcranial direct current stimulation for major depression: an updated systematic review and meta-analysis. *International Journal of Neuropsychopharmacology*. 2014; 17(9):1443–52. <https://doi.org/10.1017/S1461145714000418> PMID: 24713139
84. Hill AT, Fitzgerald PB, Hoy KE. Effects of Anodal Transcranial Direct Current Stimulation on Working Memory: A Systematic Review and Meta-Analysis of Findings From Healthy and Neuropsychiatric Populations. *Brain Stimulation*. 2016; 9(2):197–208. <https://doi.org/10.1016/j.brs.2015.10.006> PMID: 26597929
85. Diamond A. Executive Functions. *Annual Review of Psychology*. 2013; 64:135–68. <https://doi.org/10.1146/annurev-psych-113011-143750> PMC4084861. PMID: 23020641

86. Elliott R. Executive functions and their disorders/Imaging in clinical neuroscience. *British Medical Bulletin*. 2003; 65(1):49–59. <https://doi.org/10.1093/bmb/65.1.49>
87. Hoekzema E, Carmona S, Ramos-Quiroga JA, Fernández VR, Bosch R, Soliva JC, et al. An independent components and functional connectivity analysis of resting state fMRI data points to neural network dysregulation in adult ADHD. *Hum Brain Mapp*. 2014; 35(4):1261–72. <https://doi.org/10.1002/hbm.22250> PMID: 23417778
88. Lei D, Du M, Wu M, Chen T, Huang X, Du X, et al. Functional MRI reveals different response inhibition between adults and children with ADHD. *Neuropsychology*. 2015; 29(6):874–81. <https://doi.org/10.1037/neu0000200> PMID: 25938917
89. McCarthy H, Skokauskas N, Frodl T. Identifying a consistent pattern of neural function in attention deficit hyperactivity disorder: a meta-analysis. *Psychological Medicine*. 2014; 44(4):869–80. Epub 2013/05/13. <https://doi.org/10.1017/S0033291713001037> PMID: 23663382
90. Cepeda NJ, Cepeda ML, Kramer AF. Task Switching and Attention Deficit Hyperactivity Disorder. *Journal of Abnormal Child Psychology*. 2000; 28(3):213–26. <https://doi.org/10.1023/a:1005143419092> PMID: 10885680
91. Aron AR, Poldrack RA. Cortical and Subcortical Contributions to Stop Signal Response Inhibition: Role of the Subthalamic Nucleus. *The Journal of Neuroscience*. 2006; 26(9):2424. <https://doi.org/10.1523/JNEUROSCI.4682-05.2006> PMID: 16510720
92. Miller EK, Cohen JD. An Integrative Theory of Prefrontal Cortex Function. *Annual Review of Neuroscience*. 2001; 24(1):167–202. <https://doi.org/10.1146/annurev.neuro.24.1.167> PMID: 11283309.
93. Mario Vicario C, Martino D. The neurophysiology of magnitude: One example of extraction analogies. *Cognitive Neuroscience*. 2010; 1(2):144–5. <https://doi.org/10.1080/17588921003763969> PMID: 24168285
94. Dedoncker J, Brunoni AR, Baeken C, Vanderhasselt M-A. A Systematic Review and Meta-Analysis of the Effects of Transcranial Direct Current Stimulation (tDCS) Over the Dorsolateral Prefrontal Cortex in Healthy and Neuropsychiatric Samples: Influence of Stimulation Parameters. *Brain Stimulation*. 2016; 9(4):501–17. <https://doi.org/10.1016/j.brs.2016.04.006> PMID: 27160468
95. Datta A. Inter-Individual Variation during Transcranial Direct Current Stimulation and Normalization of Dose Using MRI-Derived Computational Models. *Frontiers in Psychiatry*. 2012; 3(91). <https://doi.org/10.3389/fpsy.2012.00091> PMID: 23097644
96. Shekhawat GS, Sundram F, Bikson M, Truong D, De Ridder D, Stinear CM, et al. Intensity, Duration, and Location of High-Definition Transcranial Direct Current Stimulation for Tinnitus Relief. *Neurorehabil Neural Repair*. 2016; 30(4):349–59. <https://doi.org/10.1177/1545968315595286> PMID: 26180052.
97. Boggio PS, Ferrucci R, Rigonatti SP, Covre P, Nitsche M, Pascual-Leone A, et al. Effects of transcranial direct current stimulation on working memory in patients with Parkinson's disease. *J Neurol Sci*. 2006; 249(1):31–8. <https://doi.org/10.1016/j.jns.2006.05.062> PMID: 16843494
98. Jamil A, Batsikadze G, Kuo H-I, Labruna L, Hasan A, Paulus W, et al. Systematic evaluation of the impact of stimulation intensity on neuroplastic after-effects induced by transcranial direct current stimulation. *The Journal of Physiology*. 2017; 595(4):1273–88. <https://doi.org/10.1113/JP272738> PMID: 27723104
99. Batsikadze G, Moliadze V, Paulus W, Kuo MF, Nitsche MA. Partially non-linear stimulation intensity-dependent effects of direct current stimulation on motor cortex excitability in humans. *The Journal of Physiology*. 2013; 591(7):1987–2000. <https://doi.org/10.1113/jphysiol.2012.249730> PMID: 23339180
100. Brunoni AR, Nitsche MA, Bolognini N, Bikson M, Wagner T, Merabet L, et al. Clinical research with transcranial direct current stimulation (tDCS): Challenges and future directions. *Brain Stimulation*. 2012; 5(3):175–95. <https://doi.org/10.1016/j.brs.2011.03.002> PMID: 22037126
101. Salehinejad MA, Ghanavai E, Rostami R, Nejati V. Cognitive control dysfunction in emotion dysregulation and psychopathology of major depression (MD): Evidence from transcranial brain stimulation of the dorsolateral prefrontal cortex (DLPFC). *Journal of Affective Disorders*. 2017; 210:241–8. <https://doi.org/10.1016/j.jad.2016.12.036> PMID: 28064113
102. Meron D, Hedger N, Garner M, Baldwin DS. Transcranial direct current stimulation (tDCS) in the treatment of depression: Systematic review and meta-analysis of efficacy and tolerability. *Neuroscience & Biobehavioral Reviews*. 2015; 57:46–62. <https://doi.org/10.1016/j.neubiorev.2015.07.012>.
103. Nejati V, Salehinejad MA, Shahidi N, Abedin A. Psychological intervention combined with direct electrical brain stimulation (PIN-CODES) for treating major depression: A pre-test, post-test, follow-up pilot study. *Neurology, Psychiatry and Brain Research*. 2017; 25:15–23. <https://doi.org/10.1016/j.npbr.2017.05.003>.
104. Nitsche MA, Nitsche MS, Klein CC, Tergau F, Rothwell JC, Paulus W. Level of action of cathodal DC polarisation induced inhibition of the human motor cortex. *Clinical Neurophysiology*. 2003; 114(4):600–4. [https://doi.org/10.1016/S1388-2457\(02\)00412-1](https://doi.org/10.1016/S1388-2457(02)00412-1). PMID: 12686268

105. Monte-Silva K, Kuo M-F, Hessenthaler S, Fresnoza S, Liebetanz D, Paulus W, et al. Induction of Late LTP-Like Plasticity in the Human Motor Cortex by Repeated Non-Invasive Brain Stimulation. *Brain Stimulation*. 2013; 6(3):424–32. <https://doi.org/10.1016/j.brs.2012.04.011> PMID: 22695026
106. Polanía R, Nitsche MA, Paulus W. Modulating functional connectivity patterns and topological functional organization of the human brain with transcranial direct current stimulation. *Hum Brain Mapp*. 2011; 32(8):1236–49. <https://doi.org/10.1002/hbm.21104> PMID: 20607750
107. Bikson M, Grossman P, Thomas C, Zannou AL, Jiang J, Adnan T, et al. Safety of Transcranial Direct Current Stimulation: Evidence Based Update 2016. *Brain Stimulation*. 2016; 9(5):641–61. <https://doi.org/10.1016/j.brs.2016.06.004> PMID: 27372845
108. Moliadze V, Andreas S, Lyzhko E, Schmanke T, Gurashvili T, Freitag CM, et al. Ten minutes of 1mA transcranial direct current stimulation was well tolerated by children and adolescents: Self-reports and resting state EEG analysis. *Brain Research Bulletin*. 2015; 119:25–33. <https://doi.org/10.1016/j.brainresbull.2015.09.011> PMID: 26449209
109. Cachoeira CT, Lefa DT, Mittelstadt SD, Mendes LST, Brunoni AR, Pinto JV, et al. Positive effects of transcranial direct current stimulation in adult patients with attention-deficit/hyperactivity disorder A pilot randomized controlled study. *Psychiatry Research*. 2017; 247:28–32. <https://doi.org/10.1016/j.psychres.2016.11.009> PMID: 27863315
110. Wischniewski M, Engelhardt M, Salehinejad MA, Schutter DJLG, Kuo MF, Nitsche MA. NMDA Receptor-Mediated Motor Cortex Plasticity After 20 Hz Transcranial Alternating Current Stimulation. *Cerebral Cortex*. 2018:bhy160–bhy. <https://doi.org/10.1093/cercor/bhy160> PMID: 29992259