



Therapeutic Pearls

Nonsteroid management of residual ocular surface disease on dupilumab (ROSDD), ☆,☆☆



Jodie Raffi, BA^{a,b}, Raagini Suresh, BS^a, Timothy Berger, MD^a, Harvey Fishman, MD, PhD^c, Jenny E. Murase, MD^{a,d,*}

^a Department of Dermatology, University of California, San Francisco, San Francisco, CA, United States

^b University of California, Irvine School of Medicine, Irvine, CA, United States

^c Fishmanvision Medical Practice, Palo Alto, CA, United States

^d Department of Dermatology, Palo Alto Foundation Medical Group, Mountain View, CA, United States

Dear Editors,

Atopic dermatitis is associated with a range of ocular complications. Additionally, an increase in dupilumab-associated conjunctivitis, dry eye, and recently characterized residual ocular surface disease has been demonstrated in multiple studies. Given recent evidence that interleukin-4 and -13 may play a role in goblet cell proliferation and mucous secretion (Garcia-Posadas et al., 2018), blockage of these particular immune pathways by dupilumab may ultimately lead to tear film instability and dry eye.

Individuals who use chronic steroid eyedrops are at risk for the development of adverse effects, including glaucoma, cataracts, decreased wound healing, and orbital fat atrophy, and need to be followed by an ophthalmologist. Significant relief can be achieved with a nonsteroid ophthalmologic regimen after an initial pulse treatment of topical steroids initiated by an ophthalmologist.

Therapeutic pearl

The primary initial goal of treatment is to improve meibomian gland function with eyelid hygiene, warm compresses, oral omega-3 fatty acid supplementation, and azithromycin ophthalmic solution 1% (Azasite), an antibiotic that exhibits anti-inflammatory effects, enhances meibum production, and improves tear stability. After eyelid inflammation is controlled, additional topical ophthalmological treatments to reduce inflammation and enhance the tear film can be introduced. Lifitegrast 5% ophthalmic solution inhibits T-cell-mediated inflammation by blocking the interaction between

Table 1

Non-steroid agents that may be beneficial in patients with dupilumab-related dry eye.

Treatment	Mechanism in dry eye relief
Lifitegrast 5%	Inhibits T-cell-mediated inflammation
Cyclosporine 0.1%	Anti-inflammatory
Bepotastine 1.5%	Blockage of histamine H1 receptors
Azithromycin 1%	Antimicrobial, anti-inflammatory, reduced meibomian gland plugging, increased tear stability

the ligand intercellular adhesion molecule-1 and lymphocyte function-associated antigen-1.

Cyclosporine 0.1% ophthalmic emulsion is another anti-inflammatory agent that can be used safely in conjunction with lifitegrast for treatment of chronic dry eye (Zirwas et al., 2018). Bepotastine 1.5% solution is an antihistamine that relieves the pruritus and redness associated with dry eye by blocking histamine H1 receptors, stabilizing mast cells, and preventing degranulation. Other antihistamines that may be used are azelastine 0.05% solution, epinastine 0.05% solution, alcaftadine 0.25% solution, and olopatadine 0.1% solution (taking into consideration that antihistamine drops may exacerbate dry eye while alleviating ocular pruritus).

We have used a combination of these four agents at twice daily dosing with significant relief in patients with dupilumab-associated dry eye (Table 1).

References

- Garcia-Posadas L, Hodges RR, Diebold Y, Dartt D. Context-dependent regulation of conjunctival goblet cell function by allergic mediators. *Sci Rep* 2018;8:12162.
- Zirwas MJ, Wulff K, Beckman K. Lifitegrast add-on treatment for dupilumab-induced ocular surface disease (DIOSD): a novel case report. *JAAD Case Rep* 2018;5(1):34–6.

* Financial Disclosure(s): The author(s) have no proprietary or commercial interest in any materials discussed in this article. Dr. Murase has participated in advisory boards for Genzyme/Sanofi, Dermira, and UCB, participated in disease statement management talks for Regeneron and UCB, and provided dermatologic consulting services for Ferndale and UpToDate.

** No human subjects were included in this study. No animals were used in this study.

* Corresponding author.

E-mail address: jodieraffi@gmail.com (J.E. Murase).