

# Crystal structure of (perchlorato- $\kappa$ O)(1,4,7,10-tetraazacyclododecane- $\kappa^4$ N)copper(II) perchlorate

Jessica L. Gray, Deidra L. Gerlach and Elizabeth T. Papish\*

Department of Chemistry, University of Alabama, 250 Hackberry Lane, Tuscaloosa, AL 35487-0336, USA.

\*Correspondence e-mail: [etpapish@ua.edu](mailto:etpapish@ua.edu)

Received 30 November 2016

Accepted 6 December 2016



Edited by M. Zeller, Purdue University, USA

**Keywords:** crystal structure; copper(II); cyclen; perchlorate.**CCDC reference:** 1521075**Supporting information:** this article has supporting information at [journals.iucr.org/e](http://journals.iucr.org/e)

The crystal structure of the title salt,  $[\text{Cu}(\text{ClO}_4)(\text{C}_8\text{H}_{20}\text{N}_4)]\text{ClO}_4$ , is reported. The  $\text{Cu}^{\text{II}}$  ion exhibits a square-pyramidal geometry and is coordinated by the four N atoms of the neutral 1,4,7,10-tetraazacyclododecane (cyclen) ligand and an O atom from one perchlorate anion, with the second perchlorate ion hydrogen-bonded to one of the amine N atoms of the cyclen ligand. Additional  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds between the amine H atoms and the coordinating and non-coordinating perchlorate groups create a three-dimensional network structure. Crystals were grown from a concentrated methanol solution at ambient temperature, resulting in no co-crystallization of solvent.


## 1. Chemical context

Azamacrocycle ligands, including 1,4,7,10-tetraazacyclododecane (cyclen), are of significant importance in research due to their ability to form stable metal complexes, allowing for their use in a wide range of applications. Some of these complexes have been studied for their use as chemical sensors, contrast agents in MRI and PET, antimicrobial agents and as biomimetic catalysts (De León-Rodríguez *et al.*, 2010; Yoo *et al.*, 2005). Copper–cyclen complexes have been studied extensively for their ability to perform catalytic DNA cleavage and peptide hydrolysis (Zhang *et al.*, 2016; Li *et al.* 2014; Hormann *et al.*, 2015). Although the synthesis of a similar  $\text{Cu}^{\text{II}}$  complex has been reported previously, no crystal structure of the complex,  $[\text{Cu}(1,4,7,10\text{-tetraazacyclododecane})](\text{ClO}_4)_2$ , has previously been published (Kruppa *et al.*, 2006).



## 2. Structural commentary

In the title complex (Fig. 1), the copper(II) ion coordinated by the four nitrogen atoms of the cyclen ligand and one oxygen




**Figure 1**  
Side (left) and top (right) views, as defined by the cyclen ligand ring, of  $[\text{Cu}(\text{cyclen})](\text{ClO}_4)_2$  represented with ellipsoids at the 50% probability level. Hydrogen bonds are drawn in blue.

atom of a perchlorate ligand. The five-coordinate cupric ion shows a nearly ideal square-pyramidal geometry ( $\tau_5 = 0.049$ ; Addison *et al.*, 1984). The Cu–N bond lengths range from 2.004 (1) to 2.015 (1) Å, which are typical values. The  $\text{Cu}^{\text{II}}$  ion exhibits a tetragonal distortion that leads to a longer apical bond with  $\text{Cu1}–\text{O1} = 2.266$  (1) Å, which is 0.12 Å longer than the average Cu–O distance (Clay *et al.*, 1979; Rohde & Merzweiler, 2010). The average N–Cu–O bond angle is 103.8 (8)°. Three hydrogen bonds are present within the asymmetric unit, with two extending from O2 and O3 of the bound perchlorate anion to N1–H1 and N2–H2, respectively. The third hydrogen bond extends from N2–H2 to O8 of the unbound anion; the numerical details are given in Table 1.

### 3. Supramolecular features

The crystal structure exhibits three unique symmetry elements: an inversion center, a twofold screw axis and a glide plane. The complex cations of two asymmetric units hydrogen-bond across an inversion center, which is clearly visible when viewed along the *a* axis (Fig. 2), creating a dimer. These hydrogen bonds ( $\text{N3}–\text{H3} \cdots \text{O1}$ ,  $\text{N3}–\text{H3} \cdots \text{O4}$ ,  $\text{N4}–\text{H4} \cdots \text{O5}$ ) have an average  $\text{N} \cdots \text{O}$  distance of 3.16 Å (Fig. 3). The complexes assemble in rows parallel to the *b* axis (Fig. 4)



**Figure 2**  
View of the unit cell along the *a* axis. An inversion center (yellow dots) exists between two asymmetric units, creating the dimeric unit defined at the center of the unit cell. Hydrogen bonds are drawn in blue.

**Table 1**  
Hydrogen-bond geometry (Å, °).

$D–H \cdots A$	$D–H$	$H \cdots A$	$D \cdots A$	$D–H \cdots A$
$\text{N1}–\text{H1} \cdots \text{O6}^{\text{i}}$	0.86 (1)	2.50 (2)	3.171 (1)	135 (1)
$\text{N1}–\text{H1} \cdots \text{O2}$	0.86 (1)	2.39 (1)	3.093 (1)	139 (1)
$\text{N2}–\text{H2} \cdots \text{O8}^{\text{ii}}$	0.88 (2)	2.31 (2)	3.050 (1)	142 (1)
$\text{N2}–\text{H2} \cdots \text{O3}$	0.88 (2)	2.44 (2)	3.052 (2)	127 (1)
$\text{N3}–\text{H3} \cdots \text{O1}^{\text{ii}}$	0.86 (2)	2.40 (1)	3.245 (1)	169 (2)
$\text{N3}–\text{H3} \cdots \text{O4}^{\text{ii}}$	0.86 (2)	2.55 (2)	3.132 (1)	126 (1)
$\text{N4}–\text{H4} \cdots \text{O5}$	0.86 (2)	2.36 (1)	3.096 (1)	143 (1)

Symmetry codes: (i)  $-x + 1, -y + 1, -z + 1$ ; (ii)  $-x, -y + 1, -z + 1$ .


due in part to weak electrostatic interactions between the bound perchlorate anion and a neighboring cyclen ligand. A hydrogen bond between the cyclen ligand and a neighboring perchlorate anion ( $\text{N1}–\text{H1} \cdots \text{O3}$ ) allows the building units to assemble parallel to the *a* axis (Fig. 5).

### 4. Database survey

A database survey resulted in several similar Cu–cyclen complexes with five-coordinate copper(II). Four structures chosen for further analysis contained a copper(II) ion coordinated by either five nitrogen atoms or four nitrogen atoms and one oxygen atom (Rohde & Merzweiler, 2010; Sarma *et al.*, 2010; Pérez-Toro *et al.*, 2015; Guo *et al.*, 2008). Where applicable, the complexes have similar Cu–O bond lengths to that of the title complex, with only slight deviations. The title complex and surveyed complexes have similar Cu–N distances with a standard deviation of 0.018 Å.

### 5. Synthesis and crystallization

The title complex was synthesized by a modified method as reported by Kruppa *et al.* (2006). Under a nitrogen atmosphere, 1,4,7,10-tetraazacyclododecane (247 mg, 1.4 mmol) and



**Figure 3**  
A view of hydrogen bonding within a dimer pair. Hydrogen bonds are drawn in blue. Carbon and hydrogen atom labels have been omitted for clarity.

copper(II) perchlorate hexahydrate (527 mg, 1.4 mmol) were separately dissolved in 2.8 mL anhydrous methanol each and combined. The resulting purple solution formed a precipitate. The reaction mixture was heated to reflux for 30 min then filtered. The filtrate was evaporated to dryness to yield a purple amorphous solid. X-ray quality crystals were grown by dissolving the solid in a minimum amount of methanol followed by slow evaporation at ambient temperature. The title complex  $[\text{Cu}(\text{cyclen})](\text{ClO}_4)_2$  was isolated as purple crystals in 84% yield (1.2 mmol, 526 mg). IR [ATR,  $\nu$  ( $\text{cm}^{-1}$ ): 3281, 2939, 1478, 1072, 617. MS (MALDI-TOF, MeOH):  $m/z = 334.2$   $[\text{Cu}(\text{cyclen})]^{2+} + \text{ClO}_4^-$ .

## 6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. H atoms attached to carbon were positioned geometrically and constrained to ride on their parent atoms. The H atoms attached to nitrogen were located in a difference map and restrained to have comparable bond lengths.  $U_{\text{iso}}(\text{H})$  values were set to  $1.2U_{\text{eq}}(\text{C/N})$ .

## Acknowledgements


Special thanks to The University of Alabama Department of Chemistry for funding and facilities. We also thank the Undergraduate Creativity and Research Academy (UCRA) at UA, the Research Grants Committee (RGC) at UA, and

**Table 2**

Experimental details.


Crystal data	
Chemical formula	$[\text{Cu}(\text{ClO}_4)(\text{C}_8\text{H}_{20}\text{N}_4)]\text{ClO}_4$
$M_r$	434.72
Crystal system, space group	Monoclinic, $P2_1/n$
Temperature (K)	173
$a, b, c$ (Å)	8.9387 (2), 15.0607 (4), 11.9235 (3)
$\beta$ (°)	92.949 (1)
$V$ (Å <sup>3</sup> )	1603.05 (7)
$Z$	4
Radiation type	Mo $K\alpha$
$\mu$ (mm <sup>-1</sup> )	1.74
Crystal size (mm)	0.23 × 0.21 × 0.18
Data collection	
Diffractometer	Bruker SMART APEXII CCD
Absorption correction	Multi-scan (SADABS; Sheldrick, 2014)
$T_{\text{min}}, T_{\text{max}}$	0.667, 0.747
No. of measured, independent and observed [ $I > 2\sigma(I)$ ] reflections	43306, 7519, 6655
$R_{\text{int}}$	0.021
$(\sin \theta/\lambda)_{\text{max}}$ (Å <sup>-1</sup> )	0.830
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.025, 0.068, 1.02
No. of reflections	7519
No. of parameters	221
No. of restraints	6
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å <sup>-3</sup> )	0.60, -0.44

Computer programs: APEX2 and SAINT-Plus (Bruker, 2013), SHELXT (Sheldrick, 2008), SHELXL2014 (Sheldrick, 2015) and SHELXLE (Hübschle *et al.*, 2011).



**Figure 4**

Packing of the complex cations, as viewed along the  $c$  axis of the unit cell. The  $a$  axis is drawn in red and the  $b$  axis is drawn in green.



**Figure 5**

Hydrogen bonding between complex cations and anions, as viewed along the  $c$  axis. Hydrogen bonds are drawn in blue. The  $a$  axis is drawn in red and the  $b$  axis is drawn in green.

acknowledge the NSF EPSCoR Track 2 Seed Grant to ETP (PI N. Hammer, grant No. OIA-1539035) for generous financial support.

### References

- Addison, A. W., Rao, N. T., Reedijk, J., van Rijn, J. & Verschoor, G. C. (1984). *J. Chem. Soc. Dalton Trans.* pp. 1349–1356.
- Bruker (2013). *SAINT-Plus* and *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Clay, R., Murray-Rust, P. & Murray-Rust, J. (1979). *Acta Cryst.* **B35**, 1894–1895.
- De León-Rodríguez, L. M., Viswanathan, S. & Sherry, A. D. (2010). *Contrast Media Mol. Imaging*, **5**, 121–125.
- Guo, J.-F., Yeung, W.-F., Gao, S., Lee, G.-H., Peng, S.-M., Lam, M. H.-W. & Lau, T.-C. (2008). *Eur. J. Inorg. Chem.* pp. 158–163.
- Hormann, J., van der Meer, M., Sarkar, B. & Kulak, N. (2015). *Eur. J. Inorg. Chem.* pp. 4722–4730.
- Hübschle, C. B., Sheldrick, G. M. & Dittrich, B. (2011). *J. Appl. Cryst.* **44**, 1281–1284.
- Kruppa, M., Frank, D., Leffler-Schuster, H. & König, B. (2006). *Inorg. Chim. Acta*, **359**, 1159–1168.
- Li, S., Chen, J.-X., Xiang, Q.-X., Zhang, L.-Q., Zhou, C.-H., Xie, J.-Q., Yu, L. & Li, F.-Z. (2014). *Eur. J. Med. Chem.* **84**, 677–686.
- Pérez-Toro, I., Domínguez-Martín, A., Choquesillo-Lazarte, D., Vílchez-Rodríguez, E., González-Pérez, J. M., Castiñeiras, A. & Niclós-Gutiérrez, J. (2015). *J. Inorg. Biochem.* **148**, 84–92.
- Rohde, D. & Merzweiler, K. (2010). *Acta Cryst.* **E66**, m894.
- Sarma, M., Chatterjee, T. & Das, S. K. (2010). *Inorg. Chem. Commun.* **13**, 1114–1117.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2014). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Yoo, S. H., Lee, B. J., Kim, H. & Suh, J. (2005). *J. Am. Chem. Soc.* **127**, 9593–9602.
- Zhang, X., Liu, X., Phillips, D. L. & Zhao, C. (2016). *ACS Catal.* **6**, 248–257.

## supporting information

*Acta Cryst.* (2017). E73, 31-34 [https://doi.org/10.1107/S2056989016019563]

## Crystal structure of (perchlorato- $\kappa$ O)(1,4,7,10-tetraazacyclododecane- $\kappa^4$ N)copper(II) perchlorate

Jessica L. Gray, Deidra L. Gerlach and Elizabeth T. Papish

### Computing details

Data collection: *APEX2* (Bruker, 2013); cell refinement: *SAINTE-Plus* (Bruker, 2013); data reduction: *SAINTE-Plus* (Bruker, 2013); program(s) used to solve structure: *SHELXT* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015) and *SHELXL* (Hübschle *et al.*, 2011).

### (Perchlorato- $\kappa$ O)(1,4,7,10-tetraazacyclododecane- $\kappa^4$ N)copper(II) perchlorate

#### Crystal data

[Cu(ClO<sub>4</sub>)(C<sub>8</sub>H<sub>20</sub>N<sub>4</sub>)]ClO<sub>4</sub>

$M_r = 434.72$

Monoclinic, *P2<sub>1</sub>/n*

$a = 8.9387$  (2) Å

$b = 15.0607$  (4) Å

$c = 11.9235$  (3) Å

$\beta = 92.949$  (1)°

$V = 1603.05$  (7) Å<sup>3</sup>

$Z = 4$

$F(000) = 892$

$D_x = 1.801$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 9899 reflections

$\theta = 2.3$ – $35.9$ °

$\mu = 1.74$  mm<sup>-1</sup>

$T = 173$  K

Block, purple

$0.23 \times 0.21 \times 0.18$  mm

#### Data collection

Bruker SMART APEXII CCD  
diffractometer

Radiation source: fine focus sealed tube

phi and  $\omega$  scans

Absorption correction: multi-scan  
(SADABS; Sheldrick, 2014)

$T_{\min} = 0.667$ ,  $T_{\max} = 0.747$

43306 measured reflections

7519 independent reflections

6655 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.021$

$\theta_{\max} = 36.2$ °,  $\theta_{\min} = 2.2$ °

$h = -14$ → $13$

$k = -24$ → $24$

$l = -19$ → $12$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.025$

$wR(F^2) = 0.068$

$S = 1.02$

7519 reflections

221 parameters

6 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: mixed

H atoms treated by a mixture of independent  
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0357P)^2 + 0.4526P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.003$

$\Delta\rho_{\max} = 0.60$  e Å<sup>-3</sup>

$\Delta\rho_{\min} = -0.44$  e Å<sup>-3</sup>

Extinction correction: SHELXL2014  
(Sheldrick, 2015),

$F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0025 (3)

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.42855 (13)	0.41595 (9)	0.83806 (10)	0.0315 (2)
H1A	0.5240	0.4277	0.8813	0.038*
H1B	0.4043	0.3521	0.8450	0.038*
C2	0.30439 (15)	0.47211 (10)	0.88238 (9)	0.0341 (2)
H2A	0.2850	0.4540	0.9601	0.041*
H2B	0.3344	0.5354	0.8835	0.041*
C3	0.07219 (14)	0.38402 (9)	0.83922 (11)	0.0330 (2)
H3A	0.0198	0.3977	0.9083	0.040*
H3B	0.1353	0.3308	0.8533	0.040*
C4	-0.04050 (12)	0.36717 (8)	0.74254 (13)	0.0353 (3)
H4A	-0.0945	0.3110	0.7554	0.042*
H4B	-0.1148	0.4160	0.7378	0.042*
C5	0.08786 (13)	0.27094 (7)	0.60632 (10)	0.0297 (2)
H5A	0.0006	0.2334	0.5836	0.036*
H5B	0.1415	0.2431	0.6720	0.036*
C6	0.19107 (16)	0.27947 (8)	0.51035 (10)	0.0345 (2)
H6A	0.2356	0.2209	0.4943	0.041*
H6B	0.1335	0.2998	0.4420	0.041*
C7	0.44486 (13)	0.30541 (7)	0.60226 (11)	0.0302 (2)
H7A	0.5046	0.2707	0.5500	0.036*
H7B	0.4126	0.2651	0.6620	0.036*
C8	0.53867 (12)	0.38048 (8)	0.65356 (12)	0.0315 (2)
H8A	0.6201	0.3558	0.7037	0.038*
H8B	0.5847	0.4147	0.5934	0.038*
H1	0.4734 (19)	0.4937 (9)	0.7172 (14)	0.038*
H2	0.1146 (18)	0.5097 (10)	0.8063 (14)	0.038*
H3	-0.0155 (19)	0.3802 (12)	0.5781 (13)	0.038*
H4	0.3389 (19)	0.3719 (11)	0.4820 (12)	0.038*
N1	0.44205 (10)	0.43993 (6)	0.71871 (8)	0.02310 (15)
N2	0.16614 (10)	0.46012 (7)	0.80849 (8)	0.02595 (16)
N3	0.03746 (11)	0.36146 (6)	0.63546 (9)	0.02866 (19)
N4	0.31208 (12)	0.34444 (6)	0.54087 (8)	0.02774 (18)
O1	0.19181 (10)	0.55583 (5)	0.55695 (6)	0.02513 (14)
O2	0.40695 (9)	0.63197 (6)	0.62923 (8)	0.03461 (19)
O3	0.17927 (13)	0.64843 (8)	0.71488 (9)	0.0461 (3)
O4	0.21084 (11)	0.71021 (6)	0.53609 (9)	0.0388 (2)
O5	0.24547 (13)	0.44353 (6)	0.31635 (7)	0.0375 (2)
O6	0.36978 (11)	0.40926 (8)	0.15317 (8)	0.0381 (2)
O7	0.21333 (13)	0.30342 (6)	0.23229 (9)	0.0411 (2)



O8	0.10864 (11)	0.42926 (7)	0.14227 (8)	0.0380 (2)
Cl1	0.24776 (3)	0.63777 (2)	0.61025 (2)	0.02082 (4)
Cl2	0.23354 (3)	0.39588 (2)	0.21099 (2)	0.02216 (5)
Cu1	0.23119 (2)	0.42802 (2)	0.65438 (2)	0.01773 (3)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0265 (5)	0.0390 (6)	0.0278 (5)	−0.0022 (4)	−0.0100 (4)	0.0095 (4)
C2	0.0368 (6)	0.0465 (7)	0.0184 (4)	−0.0037 (5)	−0.0038 (4)	−0.0015 (4)
C3	0.0296 (5)	0.0328 (5)	0.0375 (6)	0.0004 (4)	0.0110 (4)	0.0077 (4)
C4	0.0176 (4)	0.0285 (5)	0.0600 (8)	−0.0005 (4)	0.0049 (5)	0.0049 (5)
C5	0.0299 (5)	0.0191 (4)	0.0386 (5)	−0.0049 (3)	−0.0117 (4)	0.0052 (4)
C6	0.0523 (7)	0.0237 (5)	0.0266 (5)	−0.0087 (5)	−0.0064 (5)	−0.0022 (4)
C7	0.0307 (5)	0.0206 (4)	0.0402 (6)	0.0008 (4)	0.0099 (4)	0.0013 (4)
C8	0.0199 (4)	0.0252 (5)	0.0501 (7)	−0.0006 (3)	0.0081 (4)	0.0014 (4)
N1	0.0190 (3)	0.0217 (3)	0.0282 (4)	−0.0023 (3)	−0.0021 (3)	0.0033 (3)
N2	0.0241 (4)	0.0280 (4)	0.0259 (4)	0.0011 (3)	0.0030 (3)	0.0013 (3)
N3	0.0236 (4)	0.0208 (4)	0.0400 (5)	−0.0022 (3)	−0.0134 (3)	0.0065 (3)
N4	0.0403 (5)	0.0208 (4)	0.0224 (4)	−0.0040 (3)	0.0032 (3)	0.0018 (3)
O1	0.0361 (4)	0.0155 (3)	0.0227 (3)	−0.0025 (3)	−0.0091 (3)	−0.0012 (2)
O2	0.0213 (4)	0.0348 (4)	0.0466 (5)	−0.0019 (3)	−0.0095 (3)	0.0080 (4)
O3	0.0542 (6)	0.0452 (6)	0.0408 (5)	−0.0136 (5)	0.0192 (4)	−0.0216 (4)
O4	0.0420 (5)	0.0181 (3)	0.0538 (5)	−0.0016 (3)	−0.0219 (4)	0.0084 (3)
O5	0.0614 (6)	0.0281 (4)	0.0222 (3)	0.0054 (4)	−0.0054 (4)	−0.0058 (3)
O6	0.0258 (4)	0.0516 (6)	0.0371 (4)	−0.0021 (4)	0.0033 (3)	0.0019 (4)
O7	0.0536 (6)	0.0203 (4)	0.0493 (5)	−0.0032 (4)	0.0031 (5)	−0.0015 (4)
O8	0.0302 (4)	0.0538 (6)	0.0295 (4)	0.0164 (4)	−0.0042 (3)	−0.0005 (4)
Cl1	0.02111 (9)	0.01695 (8)	0.02384 (9)	−0.00122 (7)	−0.00426 (7)	−0.00201 (7)
Cl2	0.02481 (10)	0.02058 (9)	0.02081 (9)	0.00237 (7)	−0.00149 (7)	−0.00166 (7)
Cu1	0.01806 (5)	0.01641 (5)	0.01824 (5)	−0.00136 (3)	−0.00382 (4)	0.00267 (3)

*Geometric parameters (Å, °)*

C1—N1	1.4791 (14)	C7—H7A	0.9900
C1—C2	1.5122 (19)	C7—H7B	0.9900
C1—H1A	0.9900	C8—N1	1.4901 (15)
C1—H1B	0.9900	C8—H8A	0.9900
C2—N2	1.4913 (15)	C8—H8B	0.9900
C2—H2A	0.9900	N1—Cu1	2.0061 (9)
C2—H2B	0.9900	N1—H1	0.858 (14)
C3—N2	1.4781 (16)	N2—Cu1	2.0145 (9)
C3—C4	1.513 (2)	N2—H2	0.876 (14)
C3—H3A	0.9900	N3—Cu1	2.0036 (9)
C3—H3B	0.9900	N3—H3	0.859 (13)
C4—N3	1.4881 (18)	N4—Cu1	2.0099 (10)
C4—H4A	0.9900	N4—H4	0.859 (13)
C4—H4B	0.9900	O1—Cl1	1.4644 (7)

C5—N3	1.4826 (15)	O1—Cu1	2.2664 (7)
C5—C6	1.5118 (19)	O2—C11	1.4320 (8)
C5—H5A	0.9900	O3—C11	1.4267 (10)
C5—H5B	0.9900	O4—C11	1.4321 (9)
C6—N4	1.4898 (15)	O5—C12	1.4459 (9)
C6—H6A	0.9900	O6—C12	1.4441 (10)
C6—H6B	0.9900	O7—C12	1.4285 (9)
C7—N4	1.4835 (16)	O8—C12	1.4409 (9)
C7—C8	1.5174 (17)		
N1—C1—C2	107.30 (9)	C1—N1—C8	115.66 (9)
N1—C1—H1A	110.3	C1—N1—Cu1	102.97 (7)
C2—C1—H1A	110.3	C8—N1—Cu1	107.77 (7)
N1—C1—H1B	110.3	C1—N1—H1	107.1 (11)
C2—C1—H1B	110.3	C8—N1—H1	110.9 (12)
H1A—C1—H1B	108.5	Cu1—N1—H1	112.2 (11)
N2—C2—C1	109.06 (9)	C3—N2—C2	114.31 (10)
N2—C2—H2A	109.9	C3—N2—Cu1	103.55 (7)
C1—C2—H2A	109.9	C2—N2—Cu1	107.38 (7)
N2—C2—H2B	109.9	C3—N2—H2	111.3 (12)
C1—C2—H2B	109.9	C2—N2—H2	109.4 (12)
H2A—C2—H2B	108.3	Cu1—N2—H2	110.7 (11)
N2—C3—C4	107.81 (10)	C5—N3—C4	114.55 (9)
N2—C3—H3A	110.1	C5—N3—Cu1	102.42 (7)
C4—C3—H3A	110.1	C4—N3—Cu1	108.34 (7)
N2—C3—H3B	110.1	C5—N3—H3	106.1 (12)
C4—C3—H3B	110.1	C4—N3—H3	113.6 (12)
H3A—C3—H3B	108.5	Cu1—N3—H3	111.3 (12)
N3—C4—C3	109.92 (9)	C7—N4—C6	114.37 (9)
N3—C4—H4A	109.7	C7—N4—Cu1	102.84 (7)
C3—C4—H4A	109.7	C6—N4—Cu1	107.10 (8)
N3—C4—H4B	109.7	C7—N4—H4	110.1 (12)
C3—C4—H4B	109.7	C6—N4—H4	110.2 (12)
H4A—C4—H4B	108.2	Cu1—N4—H4	112.0 (12)
N3—C5—C6	107.69 (9)	C11—O1—Cu1	116.92 (4)
N3—C5—H5A	110.2	O3—C11—O2	109.65 (7)
C6—C5—H5A	110.2	O3—C11—O4	111.02 (7)
N3—C5—H5B	110.2	O2—C11—O4	109.85 (6)
C6—C5—H5B	110.2	O3—C11—O1	108.78 (6)
H5A—C5—H5B	108.5	O2—C11—O1	109.36 (5)
N4—C6—C5	109.55 (9)	O4—C11—O1	108.14 (5)
N4—C6—H6A	109.8	O7—C12—O8	109.87 (7)
C5—C6—H6A	109.8	O7—C12—O6	109.78 (7)
N4—C6—H6B	109.8	O8—C12—O6	109.12 (6)
C5—C6—H6B	109.8	O7—C12—O5	109.49 (6)
H6A—C6—H6B	108.2	O8—C12—O5	109.95 (6)
N4—C7—C8	108.37 (9)	O6—C12—O5	108.61 (6)
N4—C7—H7A	110.0	N3—Cu1—N1	151.33 (4)



C8—C7—H7A	110.0	N3—Cu1—N4	87.11 (4)
N4—C7—H7B	110.0	N1—Cu1—N4	87.14 (4)
C8—C7—H7B	110.0	N3—Cu1—N2	86.24 (4)
H7A—C7—H7B	108.4	N1—Cu1—N2	86.52 (4)
N1—C8—C7	109.56 (9)	N4—Cu1—N2	153.54 (4)
N1—C8—H8A	109.8	N3—Cu1—O1	104.87 (3)
C7—C8—H8A	109.8	N1—Cu1—O1	103.78 (3)
N1—C8—H8B	109.8	N4—Cu1—O1	103.79 (3)
C7—C8—H8B	109.8	N2—Cu1—O1	102.66 (3)
H8A—C8—H8B	108.2		
N1—C1—C2—N2	54.05 (13)	C6—C5—N3—C4	-168.48 (9)
N2—C3—C4—N3	50.52 (13)	C6—C5—N3—Cu1	-51.42 (9)
N3—C5—C6—N4	53.24 (12)	C3—C4—N3—C5	89.53 (11)
N4—C7—C8—N1	51.48 (13)	C3—C4—N3—Cu1	-24.09 (11)
C2—C1—N1—C8	-169.09 (9)	C8—C7—N4—C6	-165.51 (10)
C2—C1—N1—Cu1	-51.82 (10)	C8—C7—N4—Cu1	-49.77 (10)
C7—C8—N1—C1	89.34 (12)	C5—C6—N4—C7	87.24 (12)
C7—C8—N1—Cu1	-25.21 (11)	C5—C6—N4—Cu1	-26.00 (11)
C4—C3—N2—C2	-166.49 (10)	Cu1—O1—Cl1—O3	-58.89 (8)
C4—C3—N2—Cu1	-50.00 (10)	Cu1—O1—Cl1—O2	60.85 (7)
C1—C2—N2—C3	87.01 (12)	Cu1—O1—Cl1—O4	-179.55 (6)
C1—C2—N2—Cu1	-27.24 (11)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O6 <sup>i</sup>	0.86 (1)	2.50 (2)	3.171 (1)	135 (1)
N1—H1...O2	0.86 (1)	2.39 (1)	3.093 (1)	139 (1)
N2—H2...O8 <sup>ii</sup>	0.88 (2)	2.31 (2)	3.050 (1)	142 (1)
N2—H2...O3	0.88 (2)	2.44 (2)	3.052 (2)	127 (1)
N3—H3...O1 <sup>ii</sup>	0.86 (2)	2.40 (1)	3.245 (1)	169 (2)
N3—H3...O4 <sup>ii</sup>	0.86 (2)	2.55 (2)	3.132 (1)	126 (1)
N4—H4...O5	0.86 (2)	2.36 (1)	3.096 (1)	143 (1)

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $-x, -y+1, -z+1$ .