

Mediterranean Diet Reduces Risk of Incident Stroke in a Population With Varying Cardiovascular Disease Risk Profiles

Katherine E. Paterson, MRes; Phyo K. Myint, MD; Amy Jennings, PhD; Lucy K.M. Bain, PhD; Marleen A.H. Lentjes, PhD; Kay-Tee Khaw, MBBChir, MD; Ailsa A. Welch, PhD

Background and Purpose—Although some evidence has found that the Mediterranean diet (MD) is protective for stroke risk, few studies have investigated whether this relationship differs by sex or cardiovascular disease risk.

Methods—We investigated the relationship between adherence to the MD score, estimated using 7-day dietary diaries and risk of incident stroke in an observational prospective population-based cohort study of 23 232 men and women (54.5% women) aged 40 to 77 years who participated in the European Prospective Investigation into Cancer study in Norfolk, United Kingdom. Risk of incident stroke was calculated using multivariable Cox regression, in the whole population, and also stratified by sex and cardiovascular disease risk profile, using the Framingham risk score.

Results—During 17.0 years of follow-up (395 048 total person-years), 2009 incident strokes occurred. Risk of stroke was significantly reduced with greater adherence to the MD score (quartile 4 versus quartile 1 hazard ratio [HR], 0.83; 95% CI, 0.74–0.94; *P*-trend <0.01) in the whole population and in women (quartile 4 versus quartile 1 HR, 0.78; 95% CI, 0.65, 0.93; *P*-trend <0.01) but not in men (quartile 4 versus quartile 1 HR, 0.94; 95% CI, 0.79–1.12; *P*-trend =0.55). There was reduced risk of stroke in those at high risk of cardiovascular disease and across categories of the MD score (quartile 4 versus quartile 1 HR, 0.87; 95% CI, 0.76–0.99; *P*-trend =0.04). However, this was driven by the associations in women (quartile 4 versus quartile 1 HR, 0.80; 95% CI, 0.65–0.97; *P*-trend =0.02).

Conclusions—Greater adherence to the MD was associated with lower risk of stroke in a UK white population. For the first time in the literature, we also investigated the associations between the MD score in those at both low and high risk of cardiovascular disease. Although the findings in our study were driven by the associations in women, they have implications for the general public and clinicians for prevention of stroke. (*Stroke*. 2018;49:2415–2420. DOI: 10.1161/STROKEAHA.117.020258.)

Key Words: cardiovascular disease ■ diet ■ epidemiology ■ stroke ■ women

The burden of cerebrovascular disease is increasing with stroke being the third leading cause of disability-adjusted life-years, accounting for 6.24 million deaths annually worldwide.^{1,2} Around 90% of stroke risk is preventable and attributable to modifiable risk factors (RFs) including poor diet.³ Conformity to a Mediterranean diet (MD) has been found to be beneficial for stroke prevention, although the results of previous studies have varied because of differences in study design, existing cardiovascular disease (CVD) risk within populations, and different methods of dietary assessment.^{4–6} The beneficial effects of the MD are likely to be because of the overall effect or its constituent nutritional components on lowering blood pressure, lipids, and inflammation and improving metabolic health.⁷ The MD forms part of the

dietary recommendations for guidelines for the primary prevention of stroke.^{8,9} However, only 2 prospective cohort studies have investigated the preventative effect of the MD on stroke in both men and women concurrently, despite recent guidelines suggesting that stroke prevention strategies should differ in men and women.¹⁰ This is because CVD risk profiles are sex specific, for example, women with diabetes mellitus experience a higher risk of stroke than men.^{10,11} Sex-specific research concerning the MD and risk of stroke has also been recommended because of the lack of studies with sufficiently robust outcome data.¹¹

Although understanding the effect of the MD stroke risk is important for population prevention, previous studies have only investigated the effects of the MD either in populations

Received November 27, 2017; final revision received July 9, 2018; accepted August 7, 2018.

From the Department of Population Health and Primary Care (K.E.P., L.K.M.B., A.A.W.) and Department of Nutrition and Preventive Medicine (A.J.), Norwich Medical School, University of East Anglia, United Kingdom; Department of Nutrition and Dietetics, Norfolk & Norwich University NHS Foundation Trust, United Kingdom (K.E.P.); School of Medicine, Medical Sciences and Nutrition, University of Aberdeen, United Kingdom (P.K.M.); and Clinical Gerontology Unit, Department of Public Health and Primary Care, University of Cambridge, United Kingdom (M.A.H.L., K.-T.K.).

The online-only Data Supplement is available with this article at <https://www.ahajournals.org/doi/suppl/10.1161/STROKEAHA.117.020258>.

Correspondence to Ailsa A. Welch, PhD, Department of Population Health and Primary Care, Norwich Medical School, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, United Kingdom. Email a.welch@uea.ac.uk

© 2018 The Authors. *Stroke* is published on behalf of the American Heart Association, Inc., by Wolters Kluwer Health, Inc. This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution, and reproduction in any medium, provided that the original work is properly cited.

Stroke is available at <https://www.ahajournals.org/journal/str>

DOI: 10.1161/STROKEAHA.117.020258

with high risk of CVD or without stratification for CVD risk across whole populations.⁵ We are unaware of previous studies investigating the associations between the MD in those with low and high risk for CVD and the MD.

Key components of a traditional MD include olive oil as the main source of fat, high intakes of fish, fruit, vegetables, nuts, and legumes, and low meat and dairy consumption with moderate alcohol consumption.¹² The CVD preventive effect of MD is conferred by the nutrient and bioactive compound contributions from the foods within this pattern. However, although the traditional MD is well established, the foods that contribute to it vary between Mediterranean and non-Mediterranean countries,¹³ resulting in different nutrient profiles for this dietary pattern. Moreover, food habits differ regionally even within Europe.¹⁴ This may impact on the median cut points of food and nutrient components that are used to form Mediterranean diet scores (MDSs) used in research and, therefore, the comparability between studies.

To our knowledge, although there have been studies that have examined the effect of an association between MD and incident stroke risk,⁴ only one previous study in the United Kingdom used food frequency questionnaire (FFQ) method to estimate food intake.¹⁵ Also, one small prospective cohort study (n=1849) previously used a 7-day diet diary (7DD) method¹⁶ despite this approach being considered more precise than FFQs for estimating food and nutrient intakes.¹⁷ FFQs are known to provide differing estimates of food components in comparison to food diaries, thus potentially distorting population median intake estimates of foods. For example, intakes of vegetables are ≈ 100 g/d higher when derived from FFQs compared with food diaries.¹⁸

The primary aims of our study were to examine whether greater adherence to MD was associated with reduced risk of incident stroke in a prospective cohort study of middle and older aged people. Given the previous research, our aims were first to understand whether the associations with the MD differed between men and women, second to understand these associations according to categories of CVD risk, and finally to understand whether the individual nutritional components of the MD were associated with risk of stroke.

Methods

Data will be available on request from the corresponding author, subject to approval by the European Prospective Investigations into Cancer and Nutrition (EPIC) Norfolk steering committee.

The current prospective cohort study includes 23 232 voluntary participants from a population-based cohort of 25 639 men and women from EPIC-Norfolk, the UK arm of the multicenter European Prospective Investigation into Cancer study (Figure 1 in the [online-only Data Supplement](#)). The EPIC-Norfolk study design and the characteristics of the cohort have been described fully elsewhere.¹⁹ Briefly, all people aged between 40 and 79 years registered with 35 collaborating general practice surgeries in the Norfolk area were invited to the study by mail between 1993 and 1997. Participants who consented to the study completed a Health and Lifestyle Questionnaire and subsequently attended a health examination. The cohort is representative of a UK population, although current smoking levels were marginally lower than that reported in other UK surveys.¹⁹

Baseline Measurements

At baseline, participants were asked to complete 7DDs for all foods and drinks consumed.²⁰ Full details on 7DD methodology in addition

to the collection of health and lifestyle information and anthropometric and biological measurements are available in the [online-only Data Supplement](#). For the current analyses, we defined the MD using the modified MDS, frequently described as a traditional MD.¹³ In calculating this score, one point is given for intakes of each protective item in the score at or above the sex-specific median for the study population, and zero is given for amounts below the sex-specific median. Protective items are fruit and nuts, vegetables, legumes, cereals (including bread and potatoes), fish, and a higher unsaturated fat:saturated fat ratio. For dairy and meat and eggs, intakes at or above the sex-specific median intake received 0 point; for intakes below the sex-specific median, 1 point was given. Women received 1 point for intakes of alcohol between 5 and 25 g/d (4.4–21.9 U/wk) and men who drank between 10 and 50 g/d (8.8–43.7 U/wk) alcohol received 1 point; otherwise, men and women were given 0 points for other amounts of alcohol. Thus, a range of scores between 0 and 9 was possible.

Outcome Ascertainment

Linkage with hospital records and death certificate information determined nonfatal and fatal stroke incidence. Stroke outcome data were available for almost 100% of the participants apart from a few individuals who died abroad (notified by relatives) and for whom we do not have a cause of death. These participants represent 1.4% of deaths within the cohort. *International Classification of Disease* codes ICD-9 430–448/ICD-10, I60–69 were classed as stroke. In the United Kingdom, all individuals have a unique National Health Service ID number, and the National Health Service is used by the whole population. This investigation includes incident strokes during follow-up (FU) until March 31, 2015. This outcome ascertainment method has been validated for stroke in EPIC showing high sensitivity and specificity.²¹

Ethics

All participants gave written informed consent to the study. Norfolk District Health Authority Ethics Committee gave ethical approval.

Statistical Analysis

Statistical analyses were performed using Stata version 11 (StataCorp. 2009. Stata Statistical Software: Release 14. StataCorp LP, College Station, TX).

We compared the baseline characteristics and unadjusted dietary intakes of men and women at baseline using 2 sample *t* tests with equal variance for continuous data or χ^2 tests for categorical data. We also compared the baseline characteristics across sex-specific quartiles of MDS using ANOVA for continuous data or χ^2 tests for categorical data.

Hazard ratios (HRs) and 95% CIs were determined for incident stroke risk in relation to quartiles of the MDS using Cox regression. For these analyses, we report the HRs, stratified by sex, from unadjusted and 3 adjusted models. Model 1: sex, age, body mass index, educational attainment, physical activity, smoking status, material deprivation using the Townsend Index, energy intake, and alcohol intake. Model 2: model 1+baseline serum total cholesterol, baseline myocardial infarction (MI) or diabetes mellitus, and family history of stroke or MI. Model 3: Model 2+systolic blood pressure, antihypertensive use, and aspirin use for >3 months. We also used Cox regression to examine the HRs for incident stroke comparing participants with intakes above and below the median for individual components of the MDS, with covariates as model 3 above.

To investigate the effect of MD adherence in individuals with different existing cardiovascular risk profiles, we performed further stratified analysis. We calculated the 10-year cardiovascular risk of participants using the traditional Framingham risk score²² and stratified participants into 2 groups: low risk (Framingham risk score below the median) and high risk (Framingham risk score above the median). HRs and 95% CIs were determined for incident stroke risk across quartiles of the MDS within these groups using Cox regression. Models

were adjusted for sex, age, body mass index, educational attainment, physical activity, smoking status, material deprivation using Townsend deprivation index, energy intake, alcohol intake, baseline MI, and family history of stroke or MI. We also stratified these analyses by sex.

Results

Of the 23 232 individuals aged 40 to 77 years at baseline (mean: 59.1±9.3 years), 54.5% were women. There were 2009 incident strokes (men n=961 and women n=1048) during FU (mean: 17.0±4.6 years; 395 048 total person-years). At baseline, systolic and diastolic blood pressure was significantly higher in men than in women by 3.6 mmHg (SE: 0.2) and 3.5 mmHg (SE: 0.1; *P*<0.001 for both), respectively. Mean MDS was 4.4 (SD: 1.6) in men and 4.2 (SD: 1.6) in women (*P*<0.01; Table 1). There was a significant trend for lower prevalence of current smoking and higher physical activity levels by increasing quartiles of MDS for both men and women. In women only, higher adherence to the MDS was also associated with lower age (*P*<0.01), body mass index (*P*<0.01), use of antihypertensive medication (*P*<0.01) and deprivation (*P*=0.03), and higher education levels (*P*<0.01; Table II in the [online-only Data Supplement](#)).

In multivariable analysis (Table 2), in the whole study population, there was a significant inverse trend for reduced stroke risk across increasing quartiles of the MDS (quartile 4 versus quartile 1 HR, 0.83; 95% CI, 0.74–0.94; *P*-trend <0.01). After stratification by sex, these associations were only apparent in women (quartile 4 versus quartile 1 HR, 0.78; 95% CI, 0.65–0.93; *P*-trend <0.01). Additionally, moderate MDS (quartile 3 versus quartile 1) was associated with a 20% significant stroke risk reduction in women (HR, 0.80; 95% CI, 0.67–0.95; *P*<0.05) but not in men. Sensitivity analysis with additional adjustment for hormone replacement therapy and menopausal status in women did not markedly change the results (data not shown).

As shown in Table 3, we found higher adherence to the MDS to be associated with reduced stroke risk in those participants at high cardiovascular risk (Framingham risk score above the cohort-specific median). Participants with highest adherence to the MDS had a 13% reduced risk of stroke compared with those in the lowest MDS quartile (quartile 4 versus quartile 1 HR, 0.87; 95% CI, 0.76–0.99; *P*-trend =0.04). No significant associations were observed in the subgroup with low cardiovascular risk (quartile 4 versus quartile 1 HR, 0.77; 95% CI, 0.58–1.02; *P*-trend =0.09). After stratification by sex, it was apparent that the associations found in the high-risk group were driven by the findings in women. There was a reduction of 20% in the women (quartile 4 versus quartile 1 HR, 0.80; 95% CI, 0.65–0.97; *P*-trend =0.02). However, the associations in the low-risk groups were not significant in either sex.

Analysis of the individual components of the MDS highlighted that the reduction in stroke risk seemed to be driven by the vegetable (HR, 0.87; 95% CI, 0.80–0.96) and alcohol (HR, 0.90; 95% CI, 0.82–0.99) components of the score (Table III in the [online-only Data Supplement](#)).

Discussion

Our study found that greater adherence to the MD was associated with lower risk of stroke in a UK white population. We found these significant beneficial associations in the whole

Table 1. Baseline Characteristics in 23 232 Men and Women Aged 40 to 77 Years in the EPIC-Norfolk Cohort

Characteristic	Men	Women	P Value
	n=10 564	n=12 668	
Age, yr	59.5 (9.3)	58.8 (9.3)	<0.01
BMI, kg/m ²	26.4 (3.2)	26.1 (4.3)	<0.01
Reported DM, n (%)	318 (3.0)	179 (1.4)	<0.01
Reported MI, n (%)	543 (5.1)	156 (1.2)	<0.01
Family history stroke, n (%)	2427 (23.0)	3190 (25.2)	<0.01
Family history MI, n (%)	3770 (35.7)	4713 (37.2)	0.02
Family history DM, n (%)	1291 (12.2)	1712 (13.5)	0.01
Systolic blood pressure, mm Hg	137 (17.5)	134 (18.8)	<0.01
Diastolic blood pressure, mm Hg	84.3 (11.0)	80.8 (11.0)	<0.01
Antihypertensive use, n (%)	1889 (17.9)	2335 (18.4)	0.28
Aspirin use, n (%)	1226 (11.6)	846 (6.7)	<0.01
Total cholesterol, mmol/L	6.04 (1.1)	6.30 (1.2)	<0.01
Framingham risk score*	0.24 (0.2)	0.12 (0.1)	<0.01
Smoking, n (%)			<0.01
Current smoker	1323 (12.5)	1521 (12.0)	
Former smoker	5727 (54.2)	4029 (31.8)	
Never	3514 (33.3)	7118 (56.2)	
Physical activity, n (%)			<0.01
Inactive	3195 (30.2)	3760 (29.7)	
Moderately inactive	2588 (24.5)	4079 (32.2)	
Moderately active	2456 (23.3)	2865 (22.6)	
Active	2325 (22.0)	1964 (15.5)	
Education, n (%)			<0.01
No qualifications	3194 (30.2)	5301 (41.9)	
O-level or equivalent	914 (8.7)	1465 (11.6)	
A-level or equivalent	4823 (45.7)	4520 (35.7)	
Degree or equivalent	1633 (15.5)	1382 (10.9)	
TDI, n (%)			0.19
T1 most deprived	3385 (32.0)	804 (33.2)	
T2	3617 (34.2)	827 (33.7)	
T3 least deprived	3562 (33.7)	791 (33.1)	
Incident strokes	961 (9.1)	1048 (8.3)	0.03

Values are represented as mean (SD) for continuous variables and n (%) for categorical variables. *P* value for the difference between males and females calculated using *t* test for continuous variables and χ^2 for categorical variables. DM indicates diabetes mellitus; MI, myocardial infarction; and TDI, Townsend Deprivation Index.

*Framingham risk score calculated for n=22 465 participants with complete data.

population with an HR 0.83 (95% CI, 0.74–0.94) between higher and lower adherence to the MD (quartile 4 versus quartile 1). These associations were also significant in women (HR, 0.78; 95% CI, 0.65–0.93), and although the HR was lower in men (HR, 0.94; 95% CI, 0.79–1.12), the associations

Table 2. Associations Between Baseline MDS and Incident Stroke Risk in 23232 Men and Women, Aged 40 to 77 Years in the EPIC-Norfolk Cohort

	Quartile 1	Quartile 2	Quartile 3	Quartile 4	P-Trend
MDS	0–3	4	5	6–9	
All					
n	7324	5375	5165	5368	...
Stroke events, n (%)	695 (9.5)	468 (8.7)	435 (8.4)	411 (7.7)	...
Unadjusted	1.00 (ref)	0.90 (0.80–1.01)	0.91 (0.80–1.02)	0.85 (0.75–0.96)*	0.01
Model 1	1.00 (ref)	0.91 (0.81–1.02)	0.90 (0.80–1.02)	0.86 (0.76–0.97)*	0.02
Model 2	1.00 (ref)	0.91 (0.81–1.02)	0.90 (0.80–1.01)	0.84 (0.74–0.95)*	<0.01
Model 3	1.00 (ref)	0.91 (0.81–1.02)	0.90 (0.80–1.01)	0.83 (0.74–0.94)*	<0.01
Men					
n	3177	2402	2391	2594	...
Stroke events, n (%)	303 (9.5)	201 (8.4)	219 (9.2)	238 (9.2)	...
Unadjusted	1.00 (ref)	0.84 (0.70–1.01)	0.89 (0.75–1.06)	0.95 (0.80–1.13)	0.61
Model 1	1.00 (ref)	0.86 (0.72–1.03)	0.91 (0.76–1.08)	0.98 (0.82–1.16)	0.81
Model 2	1.00 (ref)	0.86 (0.72–1.03)	0.89 (0.74–1.05)	0.95 (0.80–1.13)	0.57
Model 3	1.00 (ref)	0.86 (0.72–1.03)	0.90 (0.75–1.07)	0.94 (0.79–1.12)	0.55
Women					
n	4184	2983	2717	2784	...
Stroke events, n (%)	406 (9.7)	256 (8.6)	196 (7.2)	190 (6.8)	...
Unadjusted	1.00 (ref)	0.88 (0.75–1.03)	0.79 (0.67–0.94)*	0.78 (0.66–0.93)*	<0.01
Model 1	1.00 (ref)	0.89 (0.76–1.04)	0.81 (0.68–0.96)*	0.80 (0.67–0.95)*	<0.01
Model 2	1.00 (ref)	0.88 (0.75–1.03)	0.80 (0.68–0.95)*	0.79 (0.66–0.94)*	<0.01
Model 3	1.00 (ref)	0.87 (0.74–1.02)	0.80 (0.67–0.95)*	0.78 (0.65–0.93)*	<0.01

Values are represented as unadjusted or adjusted hazard ratios (95% CI). P-trend proportional hazards regression. n (%) represents number and percentage of stroke events within each quartile. MDS indicates Mediterranean diet score; and MI, myocardial infarction.

*Significantly different to quartile 1 ($P < 0.05$). Hazard ratios adjustment: model 1=sex, age, body mass index, educational attainment, physical activity, and smoking status; material deprivation using Townsend index, energy, and alcohol intake. Hazard ratios adjustment model 2=model 1+baseline serum total cholesterol, MI, or diabetes mellitus, and family history of stroke, or MI; Hazard ratios adjustment model 3=model 2+systolic blood pressure, antihypertensive medication use, and aspirin use for >3 mo.

were not significant. We also found that risk of stroke differed by risk of CVD, classified by the Framingham risk score, as well as by the MD in the whole population (quartile 4 versus quartile 1 HR, 0.87; 95% CI, 0.76–0.99; P -trend =0.04) and in women (quartile 4 versus quartile 1 HR, 0.80; 95% CI, 0.65–0.97; P -trend =0.02). To our knowledge, our study is the first to investigate and find differences in risk of stroke, not only according to the MD but also in those with low and high risk for CVD, and in men and women. Although our findings of a protective effect of the MD on risk of stroke were driven by women, they have implications for the general public and clinicians for the prevention of stroke.

We observed few significant associations with individual MD components, indicating that the effects of the MD may be attributable to the synergistic or additive effects of the individual foods or nutrients within the MD¹² and that the overall MD pattern was most important in relation to stroke risk.

Although previous systematic reviews and meta-analyses of RCTs and observational studies have shown significant inverse associations in relation to MD adherence and risk of

incident stroke, with studies finding a relative risk of 0.66 (95% CI, 0.48–0.92) and 0.73 (95% CI, 0.59–0.91), only 2 observational studies investigated men and women separately in the same cohort.^{5,6,23,24} It is important to understand sex differences in exposure to the MD because women have unique stroke RFs,¹¹ which include postmenopausal hormone use, hormone status changes, pregnancy, preeclampsia, gestational diabetes mellitus, and oral contraceptive use. The prevalence of other RFs differs between men and women with hypertension, diabetes mellitus, atrial fibrillation, and migraine with aura¹¹ being more prevalent in women.

Of the 2 other studies investigating sex differences in associations of stroke with the MD, one observational study in Hong Kong Chinese showed a significant inverse association with higher adherence to MD in men but not in women,²³ whereas the other, in Greece, found reductions in incident stroke²⁴ in women, only. It is not clear why we found differences in the associations between the MD and risk of stroke in men and women. However, we postulate that the components of the MD may influence the risk of

Table 3. Associations Between Baseline MDS and Incident Stroke Risk in 22 465 Men and Women, Aged 40 to 77 Years in the EPIC-Norfolk Cohort Stratified by Framingham Risk Score

	Quartile 1	Quartile 2	Quartile 3	Quartile 4	P-Trend
All: low cardiovascular risk					0.09
n	3448	2575	2547	2662	
Stroke events, n (%)	139 (4.0)	103 (4.0)	98 (3.9)	80 (3.0)	
HR (95% CI)	1.00 (ref)	0.95 (0.74–1.23)	0.96 (0.74–1.24)	0.77 (0.58–1.02)	
All: high cardiovascular risk					0.04
n	3876	2800	2618	2706	
Stroke events, n (%)	556 (14.3)	365 (13.0)	337 (12.9)	331 (12.2)	
HR (95% CI)	1.00 (ref)	0.90 (0.79–1.03)	0.89 (0.78–1.02)	0.87 (0.76–0.99)	
Men: low cardiovascular risk					0.92
n	1392	1138	1986	513	
Stroke events, n (%)	55 (4.0)	34 (3.0)	76 (3.8)	20 (3.9)	
HR (95% CI)	1.00 (ref)	0.77 (0.50–1.18)	0.97 (0.68–1.38)	0.91(0.55–1.53)	
Men: high cardiovascular risk					0.27
n	1663	1267	1266	1339	
Stroke events, n (%)	240 (14.4)	172 (13.6)	177 (14.0)	187 (14.0)	
HR (95% CI)	1.00 (ref)	0.90 (0.74–1.09)	0.90 (0.74–1.10)	0.89 (0.74–1.09)	
Women: low cardiovascular risk					0.21
n	1982	1411	1362	1448	
Stroke events, n (%)	63 (3.2)	46 (3.3)	33 (2.4)	39 (2.7)	
HR (95% CI)	1.00 (ref)	0.98 (0.67–1.44)	0.73 (0.48–1.11)	0.84 (0.56–1.25)	
Women: high cardiovascular risk					0.02
n	2287	1559	1352	1267	
Stroke events, n (%)	337 (14.7)	216 (13.9)	170 (12.6)	144 (11.4)	
HR (95% CI)	1.00 (ref)	0.92 (0.78–1.10)	0.87 (0.72–1.05)	0.80 (0.65–0.97)	

Values are represented as adjusted HRs (95% CI). P-trend represents proportional hazards regression. n (%) represents number and percentage of stroke events within each quartile. Model adjustment: sex, age, BMI, educational attainment, physical activity, smoking status, material deprivation using Townsend index, energy and alcohol intakes, baseline MI, and family history of stroke or MI. Low cardiovascular risk represents below median Framingham risk score, and high cardiovascular risk represents at or above median Framingham risk score. Mean±SD, median, and range for Framingham Risk Score within each category were as follows: low cardiovascular risk = 0.07±0.03, 0.07, 0.01–0.13; high cardiovascular risk = 0.28±0.13, 0.24, 0.13–0.97. BMI indicates body mass index; CI, confidence interval; HR, hazard ratio; MDS, Mediterranean diet score; and MI, myocardial infarction.

stroke in women more than in men. We are also aware that the different subtypes of stroke may differ in men between sexes and may lead to heterogeneity of response to the MDS.²⁴ We were unable to test this hypothesis in our cohort because of low incidence. However, this deserves further study in the future.

One previous study in the United Kingdom found a nonsignificant inverse association (HR, 0.96; 95% CI = 0.90–1.02) with greater MD adherence and incident stroke.¹⁵ However, this study did not stratify by sex, the MD was based on FFQs, and average FU was 12.2 years. We build on their analyses by including sex stratified analyses, using 7DDs with a longer FU period.

Study Strengths

The strengths of our study include the prospective design, the use of a comprehensive 7DD, robust outcome

assessment method, with few individuals lost to FU, and, besides smoking status, the population is representative of the United Kingdom.¹⁹ Another of our study’s advantages include longer FU of 17 years than all except one study in women, only. In addition, using 7DD to measure dietary/nutritional intake is more precise than FFQs, which may overestimate the important components of MD.¹⁷ We controlled for and measured potential nondietary confounding factors that affect stroke risk, namely, directly measured BP and cholesterol at baseline. Also, we had sufficient power to detect an effect of adherence to the MD on incident stroke in the population and in women when measured separately.

Limitations

Dietary assessment was based on one assessment at baseline, which does not exclude the possibility that dietary changes or

other treatments may occur subsequently, which would alter incident stroke risk. In addition, we did not control for some stroke RFs, for example, atrial fibrillation. Nevertheless, we controlled for major stroke RFs such as previous MI, BP, and vascular diseases. The MDS uses relative rather than absolute values for MD components. This limits comparison with other studies using the same score, as median intakes may be distinct in different populations. Our population was predominantly white, and, therefore, our results may not apply to other ethnic groups.

Summary

Greater adherence to the MD was associated with lower risk of stroke in a UK white population, which was attributable to the overall MD pattern, not its individual components. For the first time in the literature, we also investigated the potential benefits of the MD in people at low and high risk of CVD, in men and in women. Although the findings in our study were driven by the associations in women, they have implications for the general public and clinicians for prevention of stroke.

Acknowledgments

We thank participants, general practitioners, and EPIC-Norfolk staff.

Sources of Funding

EPIC-Norfolk is supported by the Medical Research Council program grants (G0401527 and G1000143) and Cancer Research UK program grants (C864/A8257 and C864/A14136).

Disclosures

None.

References

1. Feigin VL, Krishnamurthi RV, Parmar P, Norrving B, Mensah GA, Bennett DA, et al; GBD 2013 Writing Group; GBD 2013 Stroke Panel Experts Group. Update on the global burden of ischemic and hemorrhagic stroke in 1990-2013: the GBD 2013 study. *Neuroepidemiology*. 2015;45:161-176. doi: 10.1159/000441085
2. World Health Organisation. Burden of Disease by Cause, Age, Sex, by Country and by Region, 2000-2015. Global health estimates 2015. http://www.who.int/healthinfo/global_burden_disease/estimates/en/. Accessed August 9, 2018.
3. Feigin VL, Roth GA, Naghavi M, Parmar P, Krishnamurthi R, Chugh S, et al; Global Burden of Diseases, Injuries and Risk Factors Study 2013 and Stroke Experts Writing Group. Global burden of stroke and risk factors in 188 countries, during 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet Neurol*. 2016;15:913-924. doi: 10.1016/S1474-4422(16)30073-4
4. Grosso G, Marventano S, Yang J, Micek A, Pajak A, Scalfi L, et al. A comprehensive meta-analysis on evidence of Mediterranean diet and cardiovascular disease: are individual components equal? *Crit Rev Food Sci Nutr*. 2017;57:3218-3232. doi: 10.1080/10408398.2015.1107021
5. Liyanage T, Ninomiya T, Wang A, Neal B, Jun M, Wong MG, et al. Effects of the Mediterranean diet on cardiovascular outcomes—a systematic review and meta-analysis. *PLoS One*. 2016;11:e0159252. doi: 10.1371/journal.pone.0159252
6. Rosato V, Temple NJ, La Vecchia C, Castellan G, Tavani A, Guercio V. Mediterranean diet and cardiovascular disease: a systematic review and meta-analysis of observational studies [published online November 25, 2017]. *Eur J Nutr*. doi: 10.1007/s00394-017-1582-0
7. Tosti V, Bertozzi B, Fontana L. Health benefits of the Mediterranean diet: metabolic and molecular mechanisms. *J Gerontol A Biol Sci Med Sci*. 2018;73:318-326. doi: 10.1093/geron/glx227
8. Meschia JF, Bushnell C, Boden-Albala B, Braun LT, Bravata DM, Chaturvedi S, et al; American Heart Association Stroke Council; Council on Cardiovascular and Stroke Nursing; Council on Clinical Cardiology; Council on Functional Genomics and Translational Biology; Council on Hypertension. Guidelines for the primary prevention of stroke: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2014;45:3754-3832. doi: 10.1161/STR.0000000000000046
9. United States Department of Agriculture (USDA). *Report of the Dietary Guidelines Advisory Committee on the Dietary Guidelines for Americans, 2015*. Washington, DC, USA: United States Department of Agriculture; 2015.
10. Peters SA, Huxley RR, Woodward M. Diabetes as a risk factor for stroke in women compared with men: a systematic review and meta-analysis of 64 cohorts, including 775,385 individuals and 12,539 strokes. *Lancet*. 2014;383:1973-1980. doi: 10.1016/S0140-6736(14)60040-4
11. Bushnell C, McCullough LD, Awad IA, Chireau MV, Fedder WN, Furie KL, et al; American Heart Association Stroke Council; Council on Cardiovascular and Stroke Nursing; Council on Clinical Cardiology; Council on Epidemiology and Prevention; Council for High Blood Pressure Research. Guidelines for the prevention of stroke in women: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2014;45:1545-1588. doi: 10.1161/01.str.0000442009.06663.48
12. Trichopoulos A, Costacou T, Bamia C, Trichopoulos D. Adherence to a Mediterranean diet and survival in a Greek population. *N Engl J Med*. 2003;348:2599-2608. doi: 10.1056/NEJMoa025039
13. Trichopoulos A, Martínez-González MA, Tong TY, Forouhi NG, Khandelwal S, Prabhakaran D, et al. Definitions and potential health benefits of the mediterranean diet: views from experts around the world. *BMC Medicine*. 2014;12:112. doi: 10.1186/1741-7015-12-112
14. Freisling H, Fahey MT, Moskal A, Ocké MC, Ferrari P, Jenab M, et al. Region-specific nutrient intake patterns exhibit a geographical gradient within and between European countries. *J Nutr*. 2010;140:1280-1286. doi: 10.3945/jn.110.121152
15. Tong TY, Wareham NJ, Khaw KT, Imamura F, Forouhi NG. Prospective association of the Mediterranean diet with cardiovascular disease incidence and mortality and its population impact in a non-Mediterranean population: the EPIC-Norfolk study. *BMC Med*. 2016;14:135. doi: 10.1186/s12916-016-0677-4
16. Tognon G, Lissner L, Sæbye D, Walker KZ, Heitmann BL. The Mediterranean diet in relation to mortality and CVD: a Danish cohort study. *Br J Nutr*. 2014;111:151-159. doi: 10.1017/S0007114513001931
17. Bingham SA, Gill C, Welch A, Day K, Cassidy A, Khaw KT, et al. Comparison of dietary assessment methods in nutritional epidemiology: weighed records v. 24 h recalls, food-frequency questionnaires and estimated-diet records. *Br J Nutr*. 1994;72:619-643.
18. Bingham SA, Welch AA, McTaggart A, Mulligan AA, Runswick SA, Luben R, et al. Nutritional methods in the European Prospective Investigation of Cancer in Norfolk. *Public Health Nutr*. 2001;4:847-858.
19. Day N, Oakes S, Luben R, Khaw KT, Bingham S, Welch A, et al. EPIC-Norfolk: study design and characteristics of the cohort. *European Prospective Investigation of Cancer*. 1999;80 Suppl 1:95-103.
20. Welch AA, McTaggart A, Mulligan AA, Luben R, Walker N, Khaw KT, et al. DINER (Data Into Nutrients for Epidemiological Research) - a new data-entry program for nutritional analysis in the EPIC-Norfolk cohort and the 7-day diary method. *Public Health Nutr*. 2001;4:1253-1265.
21. Sinha S, Myint PK, Luben RN, Khaw KT. Accuracy of death certification and hospital record linkage for identification of incident stroke. *BMC Med Res Methodol*. 2008;8:74. doi: 10.1186/1471-2288-8-74
22. D'Agostino RB Sr, Vasan RS, Pencina MJ, Wolf PA, Cobain M, Massaro JM, et al. General cardiovascular risk profile for use in primary care: the Framingham Heart Study. *Circulation*. 2008;117:743-753. doi: 10.1161/CIRCULATIONAHA.107.699579
23. Chan R, Chan D, Woo J. The association of a priori and a posterior dietary patterns with the risk of incident stroke in Chinese older people in Hong Kong. *J Nutr Health Aging*. 2013;17:866-874. doi: 10.1007/s12603-013-0334-y
24. Misirli G, Benetou V, Lagiou P, Bamia C, Trichopoulos D, Trichopoulou A. Relation of the traditional Mediterranean diet to cerebrovascular disease in a Mediterranean population. *Am J Epidemiol*. 2012;176:1185-1192. doi: 10.1093/aje/kws205