



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

distancing interventions, including nationwide school closures and cancellation of public gatherings, was associated with a 30% reduction in the transmission rate.¹⁰ However, it increased after the mandatory school suspension resumed and before summer vacation started.¹⁰ Given the uncertainties surrounding the extent of the COVID-19 pandemic across the country, efforts to sustain online learning and remote working in combination with enhanced infection control measures in health-care settings and nursing homes could aid in mitigating the spread of the virus and gaining control of the epidemic's chains of transmission.

In the USA, past influenza pandemics have spread as a series of waves of varying durations and intensities, which have been associated with seasons and school cycles.^{10,11} Because respiratory viruses, including other coronaviruses, are known to respond to seasonal variation,¹² we might expect that increasing temperatures in the summer could reduce the intrinsic transmissibility of the novel coronavirus to some extent. In the context of a novel virus, warmer weather alone is unlikely to sufficiently mitigate the transmission rate and interrupt community transmission in the absence of social distancing strategies. Perhaps the most reasonable scenario ahead of us is that the transmission rate will decline during the next few months, partly driven by social distancing measures, a scenario that is reminiscent of the influenza 2009 A/H1N1 pandemic.¹⁰ Close monitoring of the transmission potential of the virus on the basis of reliable and publicly available data in near real-time will be key to short-term forecasts and sound public health decisions.

We declare no competing interests.

*Gerardo Chowell, Kenji Mizumoto
gchowell@gsu.edu

Department of Population Health Sciences, School of Public Health, Georgia State University, Atlanta, GA 30033, USA (GC, KM); Graduate School of Advanced Integrated Studies in Human Survivability, Kyoto University Yoshida-Nakaadachi-cho, Sakyo-ku, Kyoto, Japan (KM); and Hakubi Center for Advanced Research, Kyoto University, Yoshidahonmachi, Sakyo-ku, Kyoto, Japan (KM)

- 1 WHO. Coronavirus disease 2019 (COVID-19): situation report—58. World Health Organization, 2020. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200318-sitrep-58-covid-19.pdf?sfvrsn=20876712_2 (accessed March 18, 2020).
- 2 Ghinai I, McPherson TD, Hunter JC, et al. First known person-to-person transmission of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in the USA. *Lancet* 2020; published online March 12. [https://doi.org/10.1016/S0140-6736\(20\)30607-3](https://doi.org/10.1016/S0140-6736(20)30607-3).
- 3 Mizumoto K, Kagaya K, Zarebski A, et al. Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020. *Euro Surveill* 2020; **25**: 2000180.
- 4 CDC. Weekly U.S. influenza surveillance report. Centers for Disease Control and Prevention, 2020. <https://www.cdc.gov/flu/weekly/index.htm> (accessed March 16, 2020).
- 5 Johns Hopkins CSSE. 2019 novel coronavirus COVID-19 (2019-nCoV) data repository by Johns Hopkins CSSE. GitHub, 2020. https://github.com/CSSEGISandData/COVID-19/blob/master/csse_covid_19_data/csse_covid_19_time_series/time_series_19-covid-Confirmed.csv (accessed March 15, 2020).
- 6 Murray CJ, Lopez AD, Chin B, Feehan D, Hill KH. Estimation of potential global pandemic influenza mortality on the basis of vital registry data from the 1918–20 pandemic: a quantitative analysis. *Lancet* 2006; **368**: 2211–18.
- 7 Dahal S, Jenner M, Dinh L, Mizumoto K, Viboud C, Chowell G. Excess mortality patterns during 1918–1921 influenza pandemic in the state of Arizona, USA. *Ann Epidemiol* 2018; **28**: 273–80.
- 8 Olson DR, Simonsen L, Edelson PJ, Morse SS. Epidemiological evidence of an early wave of the 1918 influenza pandemic in New York City. *Proc Natl Acad Sci USA* 2005; **102**: 11059–63.
- 9 Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020; S0140-6736(20)30566-3.
- 10 Chowell G, Echevarría-Zuno S, Viboud C, et al. Characterizing the epidemiology of the 2009 influenza A/H1N1 pandemic in Mexico. *PLoS Med* 2011; **8**: e1000436.
- 11 Chao DL, Halloran ME, Longini IM Jr. School opening dates predict pandemic influenza A(H1N1) outbreaks in the United States. *J Infect Dis* 2010; **202**: 877–80.
- 12 Neher RA, Dyrda R, Druelle V, Hodcroft EB, Albert J. Potential impact of seasonal forcing on a SARS-CoV-2 pandemic. *medRxiv* 2020; published online March 3. DOI:10.1101/2020.02.13.20022806 (preprint).



Preparedness is essential for malaria-endemic regions during the COVID-19 pandemic

Published Online
March 16, 2020
[https://doi.org/10.1016/S0140-6736\(20\)30561-4](https://doi.org/10.1016/S0140-6736(20)30561-4)
For the Chinese translation
see Online for appendix

The coronavirus disease 2019 (COVID-19) pandemic that first emerged in Wuhan in China's Hubei province¹ has quickly spread to the rest of China and many other countries. Within 3 months, more than 125 000 people have been infected and the death toll had reached over 4600 worldwide on March 12, 2020.² In an attempt to contain the virus, the Chinese

Government has made unprecedented efforts and invested enormous resources and these containment efforts have stemmed the spread of the disease.³ As of March 12, 2020, malaria-endemic regions in Africa have reported a few imported COVID-19 cases including in Nigeria, Senegal, and the Democratic Republic of the Congo.² Africa needs to be prepared

to deal with COVID-19, given the infectious potential of the disease and its capacity to undermine malaria control efforts.⁴ In addition to the shared vigilance that countries around the world should maintain, regions need to consider their local malaria epidemic and take additional measures for preparation.

There are relevant lessons from the 2014–16 outbreak of Ebola virus disease in west Africa. The emergence of Ebola in malaria-endemic countries, including Guinea, Liberia, and Sierra Leone, led to a public health emergency and dealt a heavy blow to malaria control efforts. In Guinea alone, an estimated 74 000 fewer malaria cases than expected were seen at health facilities compared with years without Ebola because of decreases in the number of patients with malaria seeking appropriate health care and the volume of malaria treatments being dispensed.⁵ Contributing factors to this situation were the close resemblance of early Ebola symptoms with malaria, leading to difficulties in early diagnosis, and the fear on the part of community members of contracting Ebola in the health-care facilities. As Ebola overwhelmed health-care infrastructure, insufficient resources for malaria control in these regions led to increased mortality and morbidity. In Guinea, the official number of reported deaths from malaria in 2014 was 1067 (WHO estimate 9428) compared with 108 reported in 2013, and there were 2446 deaths from Ebola virus disease in 2014.⁶ More alarmingly, it was estimated that there were about 7000 additional malaria-associated deaths among children younger than 5 years in Guinea, Liberia, and Sierra Leone due to the Ebola outbreak.⁷ There is, therefore, a real and pressing danger for malaria-endemic regions when faced with the threat of a novel infectious disease outbreak.

While our knowledge of COVID-19 is still developing, it is a highly contagious disease that is thought to spread primarily from human to human through direct contact and inhalation of respiratory droplets. Carriers with mild or no symptoms can probably transmit the virus.⁸ In addition to China, Italy, Iran, and South Korea are among the countries with local outbreaks that could be exporting the disease and increasing exposure risks. With Africa's increasing global connectivity, the unfortunate likelihood of a continental outbreak cannot be ruled out.⁹ Much like Ebola, the early symptoms of COVID-19, including



Gallo Images/Getty Images

fever, myalgia, and fatigue, might be confused with malaria and lead to challenges in early clinical diagnosis.¹ These features of COVID-19 and the previous experiences of the Ebola outbreak point to the need for malaria-endemic countries to consider preventive measures against not only the COVID-19 threat but also its likely impact on existing malaria control efforts. The containment efforts and research impetus being taken by China and other affected countries have bought valuable time for the rest of the world, and this time window should be used effectively by vulnerable regions.

WHO is monitoring the fast-evolving situation of the COVID-19 epidemic and needs to advise the countries in the malaria-endemic regions on how to establish and effectively execute public health policies. Preventive measures for COVID-19, including case and contact tracing, quarantine and screening, as well as education to encourage good hand hygiene practices, should be in place. Additional and pre-emptive measures must be taken for malaria control in these countries, anticipating the potential challenge that would be faced by the public health system during an outbreak of COVID-19. In the case of Ebola, it was estimated that malaria cases in Guinea, Liberia, and Sierra Leone could have increased by up to 1 million in 2014 as a result of a cessation of distribution of insecticide-treated bednets (ITNs).¹⁰ Governments and health leaders in malaria-endemic regions must ensure that such stresses to medical infrastructure are minimised in the event of an outbreak of COVID-19. Resource allocation should

be optimised whenever possible to ensure minimal disruption to malaria control should COVID-19 management become necessary. Management of medical supplies and stockpiling of surgical masks and other protective equipment should be done in advance and medical staff should be adequately trained in their use. In cases of emergency, mass drug administration and the distribution of ITNs might be considered for short-term malaria relief in hyperendemic areas. Such measures would also aid efforts in COVID-19 management by reducing the strain on medical resources and minimising confounding factors in diagnosis. Previous successful implementation of such measures occurred during Ebola outbreaks in Sierra Leone in 2014–15 and in the Democratic Republic of the Congo in 2018, in accordance with WHO guidelines.^{11,12} In malaria-endemic regions, malaria diagnostics should be systematically added to fever management, including for suspected cases of COVID-19, and health-care facilities should be well stocked with artemisinin combination therapy drugs. Infection management protocols, such as social distancing, mask-wearing, and prompt seeking of diagnostic testing and necessary treatment, should be communicated in advance. These measures will require collective political will and unity in a coordinated effort by African countries.

Although an outbreak of COVID-19 in malaria-endemic regions might not happen, we must nevertheless advocate caution and recognise that such pre-emptive measures are ultimately worthwhile. Preparedness is the key to navigating any public health crisis, and malaria-endemic countries must be prepared for the challenges that COVID-19 might bring while minimising disruption to malaria control.

We declare no competing interests.

Jigang Wang, Chengchao Xu, Yin Kwan Wong, Yingke He, Ayôla A Adegnik, Peter G Kremsner, Selidji T Agnandji, Amadou A Sall, Zhen Liang, Chen Qiu, Fu Long Liao, Tingliang Jiang, Sanjeev Krishna, *Youyou Tu
yytu@icmm.ac.cn

ShenZhen People's Hospital, ShenZhen, China (JW, CX, ZL, CQ); Artemisinin Research Center and the Institute of Chinese Materia Medica, China Academy of Chinese Medical Sciences, Beijing, China (JW, CX, YKW, FLL, TJ, YT); Central People's Hospital of Zhanjiang, Zhanjiang, Guangdong, China (JW); Department of Anaesthesiology, Singapore General Hospital, Singapore (YH, YKW); Institut für Tropenmedizin, Universitätsklinikum Tübingen, Tübingen, Germany (AAA, PGK, STA, SK); Centre de Recherches Médicales de Lambaréné, Lambaréné, Gabon (AAA, PGK, STA, SK); Institut Pasteur de Dakar, Dakar, Senegal (AAS); Centre for Diagnostics and Antimicrobial Resistance, Institute for Infection Immunity, St George's University of London, London, UK (SK); and St George's University Hospitals NHS Foundation Trust, London, UK (SK)

- 1 Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med* 2020; **382**: 727–33.
- 2 WHO. Coronavirus disease 2019 (COVID-19) situation report—52. March 12, 2020. https://www.who.int/docs/default-source/coronaviruse/20200312-sitrep-52-covid-19.pdf?sfvrsn=e2bfc9c0_2 (accessed March 13, 2020).
- 3 WHO. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19). Geneva: World Health Organization, 2020.
- 4 Nkengasong JN, Mankoula W. Looming threat of COVID-19 infection in Africa: act collectively, and fast. *Lancet* 2020; published online Feb 27. [https://doi.org/10.1016/S0140-6736\(20\)30464-5](https://doi.org/10.1016/S0140-6736(20)30464-5).
- 5 Plucinski MM, Guilavogui T, Sidikiba S, et al. Effect of the Ebola-virus-disease epidemic on malaria case management in Guinea, 2014: a cross-sectional survey of health facilities. *Lancet Infect Dis* 2015; **15**: 1017–23.
- 6 WHO. World malaria report 2015. Geneva: World Health Organization, 2015.
- 7 Parpia AS, Ndeffo-Mbah ML, Wenzel NS, Galvani AP. Effects of response to 2014–2015 Ebola outbreak on deaths from malaria, HIV/AIDS, and tuberculosis, west Africa. *Emerg Infect Dis* 2016; **22**: 433–41.
- 8 Hoehl S, Berger A, Kortenbusch M, et al. Evidence of SARS-CoV-2 infection in returning travelers from Wuhan, China. *N Engl J Med* 2020; published online Feb 18. DOI:10.1056/NEJMc2001899.
- 9 Gilbert M, Pullano G, Pinotti F, et al. Preparedness and vulnerability of African countries against importations of COVID-19: a modelling study. *Lancet* 2020; published online Feb 19. [https://doi.org/10.1016/S0140-6736\(20\)30411-6](https://doi.org/10.1016/S0140-6736(20)30411-6).
- 10 Walker PGT, White MT, Griffin JT, Reynolds A, Ferguson NM, Ghani AC. Malaria morbidity and mortality in Ebola-affected countries caused by decreased health-care capacity, and the potential effect of mitigation strategies: a modelling analysis. *Lancet Infect Dis* 2015; **15**: 825–32.
- 11 Aregawi M, Smith SJ, Sillah-Kanu M, et al. Impact of the Mass Drug Administration for malaria in response to the Ebola outbreak in Sierra Leone. *Malar J* 2016; **15**: 480.
- 12 WHO. Malaria control campaign launched in Democratic Republic of the Congo to save lives and aid Ebola response. Nov 28, 2018. <https://www.who.int/malaria/news/2018/malaria-control-campaign-drc/en/> (accessed March 4, 2020).



Mass gathering events and reducing further global spread of COVID-19: a political and public health dilemma

Published Online
March 19, 2020
[https://doi.org/10.1016/S0140-6736\(20\)30681-4](https://doi.org/10.1016/S0140-6736(20)30681-4)

The coronavirus disease 2019 (COVID-19) pandemic¹ presents countries with major political, scientific, and public health challenges. Pandemic preparedness and reducing risk of global spread of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) are key concerns. Mass gathering (MG) events² pose

considerable public health challenges to health authorities and governments. Historically, sporting, religious, music, and other MGs have been the source of infectious diseases that have spread globally.³ However, the scale of the problem has declined over the years as better public health measures have been implemented