



OPEN ACCESS

CASE REPORT

When a patient with depression is feeling sleepy, be aware of sleep apnoea

Carlijn Wiersema, Willeke Van Zelst, Richard Oude Voshaar

University Center for Psychiatry,
University Medical Center
Groningen, Groningen, The
Netherlands

Correspondence to
Carlijn Wiersema,
c.wiersema@umcg.nl

Accepted 18 June 2018

SUMMARY

A 67-year-old man was referred to an outpatient clinic of geriatric psychiatry because of persistent symptoms of depression and anxiety, accompanied by sleepiness. The latter had been evaluated multiple times in the general practice over several years; each time it was considered to be a symptom of depression. After referral, the patient was diagnosed with severe obstructive sleep apnoea (OSA), comorbid to a depressive and anxiety disorder. Retrospectively, we conclude that affective symptoms accompanying OSA and sleepiness were wrongfully interpreted as depression, but after having led to problems at work, they have triggered psychiatric comorbidity. Treatment of OSA in addition to the psychiatric disorders resulted in a full recovery over time. The delayed diagnosis of OSA has certainly diminished the patient's quality of life and might have precipitated the depressive disorder. Moreover, OSA poses patients at an increased risk of cardiovascular disease, hypertension, stroke and traffic accidents.

BACKGROUND

Obstructive sleep apnoea (OSA) is the most common sleep disorder worldwide, affecting 2%–6% of the adult population, increasing up to 15%–26% in those aged 70–100 years old.¹ OSA is characterised by repetitive complete (apnoea) and partial (hypopnoea) obstruction of the upper airway during sleep, which decreases arterial oxygen saturation.² Daytime functioning may be impaired due to loss of energy, depressed mood, irritability and concentration difficulties, which are core symptoms of affective disorders. Therefore, the diagnosis of OSA may easily be missed in patients with depression. Moreover, sleepiness itself is one of the Diagnostic and Statistical Manual of Mental Disorders, fifth edition (DSM-5) criteria of depressive disorder and also overlaps with the criterion of fatigue or loss of energy.¹ These diagnostic complexities may delay proper diagnosis of OSA in patients with depression. The aim of this case report is to increase awareness of the possibility of OSA in patients with depression.

CASE PRESENTATION

A 67-year-old man was referred to our outpatient clinic of geriatric psychiatry because of persistent affective symptoms for years. The onset of symptoms coincided with an unpleasant work situation that, after a year of sick leave, had led to demotion at 49 years of age. From that moment on, he

suffered from depressed mood, anhedonia, loss of initiative, increased appetite, worrying, catastrophising, feelings of anxiety with excessive sweating, difficulties staying asleep, nightmares, increased need of sleep, loss of energy and tiredness. Symptoms fluctuated over the years but increased during autumn/winter, after life events and during vacation periods. Over the years, he had gradually withdrawn from social activities.

His general practitioner (GP) had prescribed benzodiazepine hypnotics, but without any improvement of symptoms. Six weeks prior to his referral, he was started on citalopram 20 mg one time a day, a first-line serotonergic antidepressant. He had lost 7 kg bodyweight due to nausea. This weight loss increased his physical fitness, which made him feel slightly better.

His medical history did not reveal other mental or physical symptoms.

The patient was monitored, but not treated, for mild hypertension by his GP. He had no other previous medical or psychiatric history. His family history revealed no psychiatric problems or neurodegenerative diseases like M. Parkinson, M. Alzheimer or M. Huntington. The patient did not smoke or drink alcohol. Besides citalopram, the patient used simvastatin 40 mg one time a day.

Heteroanamnestic information of his wife and son revealed a change in character since the onset of his symptoms; he became more emotional and insecure about himself. Since his retirement, he tended to lose his temper over seemingly insignificant events. His wife explained that they slept apart for years because of his snoring. She nevertheless confirmed that his breathing paused regularly during sleep. Although they consulted their GP several times for his conspicuous sleepiness, they accepted the explanation that it was an inevitable part of his depression.

At psychiatric examination, we saw a tired man with bags under his eyes and reduced facial expression. Higher cognitive functions were intact. He had a depressed mood and flat affect, but denied suicidal ideation. Evaluation of his sleep pattern revealed a rapid sleep onset, but strikingly, he never woke up well rested, regularly dozed off during the day and he was often at risk of falling asleep while driving his car.

A full physical and neurological examination was remarkable only for an overweight body mass index (28.9 kg/m²) and hypertension (130/100 mm Hg). Routine blood chemistry did not reveal any abnormalities.



© BMJ Publishing Group Limited 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Wiersema C, Van Zelst W, Oude Voshaar R. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-224873

INVESTIGATIONS

In his next visits to our clinic, the patient repeatedly brought forward his work situation as being a traumatic event. This made him anxious in situations where he did not feel in control, like during his children's holiday. He did not meet DSM-5 criteria of post-traumatic stress disorder, but he met the criteria for a depressive disorder, partly seasonal, with a superimposed generalised anxiety disorder.

The patient scored 38 points on the 30-item Inventory of Depressive Symptomatology (IDS), which is indicative of a severe depression. The scores on the Mini-Mental State Exam (MMSE) and the Cambridge Examination for Mental Disorders of the Elderly/Cambridge Cognition test (CAMDEX/CAMCOG) were not indicative of cognitive problems (MMSE: 30/30, CAMDEX/CAMCOG 90/120). Based on his wife's information, he was additionally referred to a sleep clinic at our hospital. The score on the Epworth Sleepiness Scale was 16/24, which is indicative of excessive daytime sleepiness. Polysomnography showed 356 apnoeas in 6 hours of sleep and an Apnoea-Hypopnoea Index (AHI) of 57. The arterial oxygen saturation varied between 76% and 95%. The sleep architecture was distorted, as no slow wave sleep was present. No significant periodic leg movements or other unusual movements were observed. The patient was diagnosed with severe OSA.

DIFFERENTIAL DIAGNOSIS

Depressive disorder

Depressed mood, anhedonia, increased appetite, constant worrying, loss of initiative, withdrawal of social activities, sleeping problems, loss of energy, increasing symptoms during autumn/wintertime and lesser facial expression.

Anxiety disorder

Feelings of anxiety with excessive sweating, catastrophising, nightmares, withdrawal of social situations and constant worrying.

Post-traumatic stress disorder

As a result of the unpleasant work situation, social withdrawal, nightmares and feelings of anxiety with excessive sweating.

Neurodegenerative disease

Change in character, mood disruptions and temper issues.

Obstructive sleep apnoea

Severe loss of energy/tiredness, falling asleep while driving, dozing off during daytime, wake up not fully rested, snoring, breathing pauses during sleep, irritability, change of character, obesity and hypertension.

TREATMENT

Citalopram, indicated for depressive as well as anxiety disorders, slightly improved his depressed mood and reduced anxiety symptoms over the course of 8 weeks. The patient accepted cognitive-behavioural therapy (CBT), to improve coping with his work situation and to increase his self-esteem. Four weeks after starting CBT, the patient started with continuous positive airway pressure (CPAP) at night to treat his OSA.

OUTCOME AND FOLLOW-UP

Since starting citalopram, the patient experienced a small improvement of his affective symptoms. With CBT, his self-image

became more positive. Over the course of 3 months, prior to the start of CPAP, his affective symptoms further improved, without any effect on his sleep (related) problems and fatigue. After four nights of CPAP treatment, the patient felt more energetic and started rebuilding his life. Full recovery was achieved after 4 months of treatment. His IDS score decreased from 38 to 8 (no depression present).

DISCUSSION

Although (hetero)anamnestic information may point to OSA, assessing the AHI by polysomnography is the gold standard for proper diagnosis. The AHI is the number of apnoeas or hypopnoeas recorded per hour of sleep. An AHI of 5–14 per hour indicates a mild, 15–30 a moderate and ≥ 30 a severe OSA. Hypopnoeas and apnoeas interrupt deep sleep and Rapid Eye Movement (REM) sleep and cause a fragmented sleeping pattern.² Patients wake up feeling unrefreshed and they stay tired during the day. Night-time symptoms include snoring, breathing pauses, a feeling of choking, excessive salivation, excessive sweating, gastro-oesophageal reflux, nocturia, waking up with a dry mouth and/or a headache.^{1 3} During the day, the patient experiences excessive sleepiness, loss of energy, irritability,⁴ withdrawal of social activities, difficulties with concentrating,⁵ cognitive dysfunction,^{6 7} loss of interest in daily activities,⁸ temper issues,⁵ psychomotor changes⁴⁻⁸ and anxiousness and depressed mood.⁹ These symptoms bear a striking resemblance to symptoms of depression.^{3 10 11} Indeed, up to 63% of patients with OSA report depressive symptoms.^{9 11 12} Most sleep clinics therefore regularly evaluate depressive symptoms of their patients with screening questionnaires.³ The similarity in phenotypic expression of OSA and depressive disorders might be enough to misdiagnose sleep apnoea for a depressive disorder and wrongfully start antidepressant treatment.^{13 14} In our case, the affective symptoms accompanying OSA have been wrongfully attributed to a depression by his GP and even temporarily treated with benzodiazepines. When his work problems due to OSA triggered a full-blown depressive disorder, he was referred for specialised mental healthcare.

Of particular interest for this case is that depressive disorders and sleep disorders co-occur more often than can be expected by chance.^{15 16} This association is considered to be bidirectional as depression can evoke sleep disturbances and vice versa. Overall, 90% of patients with depression experience sleep disturbances. Also, sleep apnoea is found more often in patients primarily diagnosed with depressive disorders compared with the general population.^{3 4 17-22} Because sleep disturbances are common in mood disorders, further examination is rarely employed.^{3 17} Several authors have advocated screening for OSA in psychiatric patients, as this will hasten appropriate diagnosis.^{5 17 19 22-26} Treatment of OSA will alleviate hypopnoeas and apnoeas and may improve affective symptoms.^{18 23 27-32} Especially in patients with therapy-resistant depression and/or patients with risk factors for OSA like male sex, obesity and older age, OSA must be considered.^{17 19 23} Our patient noticed some improvement after starting on citalopram, losing 7 kg of bodyweight and starting CBT, but the most significant improvement was following start of CPAP therapy.

Aside from improving psychiatric symptoms, the treatment of OSA offers other health benefits by reducing the risk of cardiovascular disease³³⁻³⁵ and neurological damage.^{36 37} The hypertension our patient presented with might be related to his OSA. Untreated OSA is associated with a twofold to threefold increased risk of traffic accidents because of daytime

sleepiness.^{38 39} Our patient also described almost falling asleep while driving his car. Ascribing symptoms of OSA to a psychiatric disorder may result in inappropriate prescriptions of benzodiazepines, as was the case in our patient. Whereas benzodiazepines are regularly used for treatment of anxiety and sleep(related) problems, benzodiazepines worsen hypopnoeas and apnoeas,²⁴ which can be life threatening for a patient with OSA. Apart from depression, our patient also suffered from an anxiety disorder. For this patient, it is quite possible that the anxiety was exacerbated by the OSA, as anxiety disorders are also associated with OSA.^{30 40} Moreover, daytime tiredness and anxiety are positively correlated in patients with OSA.⁴¹ CPAP treatment also has a positive effect on anxiety,³² as our patient experienced.

With this case report, we underline the importance of diagnosing and treating OSA in patients with a depressive disorder, even when the depression is preceded by psychosocial stressors. As illustrated by our patient, proper diagnostics may improve quality of life and prevent misdiagnosing affective symptoms as a treatment-resistant depression.

Patient's perspective

"The whole thing has actually been a positive experience for me, the discovery of the sleep apnoea has changed my life for the better. In hindsight I already had sleep problems and loss of energy for years, before the start of my depressive symptoms. I did not want to complain so I tried to live with it. I am very happy the doctors discovered my sleep apnoea, although it wasn't what I expected when my general practitioner referred me to the psychiatric clinic! The medication and CBT really helped me, but the CPAP completed it. I am well now. I have my moments of feeling a little bit low, but that's only around the fall/wintertime and is always of short duration. I think that's just part of who I am. It has been never as bad as at that time. I realise how important it is to be healthy; the impact of the sleep apnoea, my depression and anxiety has been immense. But now I have my life back. I hope that physicians will learn from my case and will think of sleep apnoea as a cause of depression at an earlier stage."

Learning points

- ▶ The similarity between daytime symptoms of obstructive sleep apnoea and depressive disorders presents a diagnostic challenge.
- ▶ It is important to be vigilant regarding symptoms of excessive daytime sleepiness and risk factors of obstructive sleep apnoea in patients with depressive disorder.
- ▶ Good history taking of snoring, breathing pauses during sleep, waking up unrefreshed (sometimes with dry mouth and headaches) as well as a thorough physical examination (obesity, hypertension and crowded oropharynx) can point in the direction of obstructive sleep apnoea.
- ▶ Treatment of obstructive sleep apnoea should complement psychiatric or psychological treatment of depression to optimise quality of life, as well as physical health.

Contributors The manuscript was written by CW and supervised by WVZ and ROV.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- 1 Rey de Castro J, Rosales-Mayor E. Depressive symptoms in patients with obstructive sleep apnea/hypopnea syndrome. *Sleep Breath* 2013;17:615–20.
- 2 American Academy of Sleep Medicine/American Academy of Sleep Medicine. *International classification of sleep disorders. Diagnostic and coding manual*. 2nd edn. Westchester, 2005.
- 3 Schröder CM, O'Hara R. Depression and Obstructive Sleep Apnea (OSA). *Ann Gen Psychiatry* 2005;4:13–20.
- 4 Deldin PJ, Phillips LK, Thomas RJ. A preliminary study of sleep-disordered breathing in major depressive disorder. *Sleep Med* 2006;7:131–9.
- 5 Kaplan R. Obstructive sleep apnoea and depression--diagnostic and treatment implications. *Aust N Z J Psychiatry* 1992;26:586–91.
- 6 O'Hara R, Schröder CM, Kraemer HC, O'Hara R, Kryla N, et al. Nocturnal sleep apnea/hypopnea is associated with lower memory performance in APOE epsilon4 carriers. *Neurology* 2005;65:642–4.
- 7 Kim HC, Young T, Matthews CG, et al. Sleep-disordered breathing and neuropsychological deficits. A population-based study. *Am J Respir Crit Care Med* 1997;156:1813–9.
- 8 Chen YH, Keller JK, Kang JH, et al. Obstructive sleep apnea and the subsequent risk of depressive disorder: a population-based follow-up study. *J Clin Sleep Med* 2013;9:417–23.
- 9 Saunamäki T, Jehkonen M. Depression and anxiety in obstructive sleep apnea syndrome: a review. *Acta Neurol Scand* 2007;116:277–88.
- 10 Aloia MS, Arnedt JT, Smith L, et al. Examining the construct of depression in obstructive sleep apnea syndrome. *Sleep Med* 2005;6:115–21.
- 11 Harris M, Glozier N, Ratnavadivel R, et al. Obstructive sleep apnea and depression. *Sleep Med Rev* 2009;13:437–44.
- 12 Ejaz SM, Khawaja IS, Bhatia S, Veteran O, et al. Obstructive sleep apnea and depression: a review. *Innov Clin Neurosci* 2011;8:17–25.
- 13 Smith R, Ronald J, Delaive K, et al. What are obstructive sleep apnea patients being treated for prior to this diagnosis? *Chest* 2002;121:164–72.
- 14 Stores G. Misdiagnosing sleep disorders as primary psychiatric conditions. *Advances in Psychiatric Treatment* 2003;9:69–77.
- 15 Franzen PL, Buysse DJ. Sleep disturbances and depression: risk relationships for subsequent depression and therapeutic implications. *Dialogues Clin Neurosci* 2008;10:473–81.
- 16 Bao YP, Han Y, Ma J, et al. Cooccurrence and bidirectional prediction of sleep disturbances and depression in older adults: Meta-analysis and systematic review. *Neurosci Biobehav Rev* 2017;75:257–73.
- 17 Ong JC, Gress JL, San Pedro-Salcedo MG, et al. Frequency and predictors of obstructive sleep apnea among individuals with major depressive disorder and insomnia. *J Psychosom Res* 2009;67:135–41.
- 18 Nasr S, Wendt B, Kora S. Increased incidence of sleep apnea in psychiatric outpatients. *Ann Clin Psychiatry* 2010;22:29–32.
- 19 Hattori M, Kitajima T, Mekata T, et al. Risk factors for obstructive sleep apnea syndrome screening in mood disorder patients. *Psychiatry Clin Neurosci* 2009;63:385–91.
- 20 Talih FR, Ajaltouni JJ, Tamim HM, et al. Risk of obstructive sleep apnea and excessive daytime sleepiness in hospitalized psychiatric patients. *Neuropsychiatr Dis Treat* 2017;13:1193–200.
- 21 Gupta MA, Simpson FC. Obstructive sleep apnea and psychiatric disorders: a systematic review. *J Clin Sleep Med* 2015;11:165–75.
- 22 Pan ML, Tsao HM, Hsu CC, et al. Bidirectional association between obstructive sleep apnea and depression: A population-based longitudinal study. *Medicine* 2016;95:e4833.
- 23 Habukawa M, Uchimura N, Kakuma T, et al. Effect of CPAP treatment on residual depressive symptoms in patients with major depression and coexisting sleep apnea: Contribution of daytime sleepiness to residual depressive symptoms. *Sleep Med* 2010;11:552–7.
- 24 Cheng P, Casement MD, Chen C, et al. Breathing and depression Sleep-disordered breathing in major depressive disorder. *J Sleep Res* 2013;22:459–62.
- 25 Wheaton AG, Perry GS, Chapman DP, et al. Sleep Disordered Breathing and Depression among U.S. Adults: National Health and Nutrition Examination Survey, 2005–2008. *Sleep* 2012;35:461–7.
- 26 Stubbs B, Vancampfort D, Veronese N, et al. The prevalence and predictors of obstructive sleep apnea in major depressive disorder, bipolar disorder and schizophrenia: A systematic review and meta-analysis. *J Affect Disord* 2016;197:259–67.

Reminder of important clinical lesson

- 27 Diamanti C, Manali E, Ginieri-Coccosis M, *et al.* Depression, physical activity, energy consumption, and quality of life in OSA patients before and after CPAP treatment. *Sleep Breath* 2013;17:1159–68.
- 28 Yamamoto H, Akashiba T, Kosaka N, *et al.* Long-term effects nasal continuous positive airway pressure on daytime sleepiness, mood and traffic accidents in patients with obstructive sleep apnoea. *Respir Med* 2000;94:87–90.
- 29 Joseph S, Zuriqat M, Husari A. Sustained improvement in cognitive and emotional status of apneic patients after prolonged treatment with positive airway pressure. *South Med J* 2009;102:589–94.
- 30 Sánchez AI, Buela-Casal G, Bermúdez MP, *et al.* The effects of continuous positive air pressure treatment on anxiety and depression levels in apnea patients. *Psychiatry Clin Neurosci* 2001;55:641–6.
- 31 Edwards C, Mukherjee S, Simpson L, *et al.* Depressive Symptoms before and after Treatment of obstructive sleep apnea in men and women. *J Clin Sleep Med* 2015;11:1029–38.
- 32 Li YY, Mazarakis T, Shen YC, Yang M-C, *et al.* Anxiety and depression are improved by continuous positive airway pressure treatments in obstructive sleep apnea. *Int J Psychiatry Med* 2016;51:554–62.
- 33 Butt M, Dwivedi G, Khair O, *et al.* Obstructive sleep apnea and cardiovascular disease. *Int J Cardiol* 2010;139:7–16.
- 34 Bradley TD, Floras JS. Obstructive sleep apnoea and its cardiovascular consequences. *Lancet* 2009;373:82–93.
- 35 Wolf J, Lewicka J, Narkiewicz K. Obstructive sleep apnea: an update on mechanisms and cardiovascular consequences. *Nutr Metab Cardiovasc Dis* 2007;17:233–40.
- 36 Harper RM, Kumar R, Ogren JA, a OJ, *et al.* Sleep-disordered breathing: effects on brain structure and function. *Respir Physiol Neurobiol* 2013;188:383–91.
- 37 Cross RL, Kumar R, Macey PM, *et al.* Neural alterations and depressive symptoms in obstructive sleep apnea patients. *Sleep* 2008;31:1103–9.
- 38 Wells RD, Day RC, Carney RM, *et al.* Depression predicts self-reported sleep quality in patients with obstructive sleep apnea. *Psychosom Med* 2004;66:692–7.
- 39 Strohl KP, Brown DB, Collop N, *et al.* An official american thoracic society clinical practice guideline: sleep apnea, sleepiness, and driving risk in noncommercial drivers. an update of a 1994 statement. *Am J Respir Crit Care Med* 2013;187:1259–66.
- 40 Hrubos-Strøm H, Einvik G, Nordhus IH, Randby A, *et al.* Sleep apnoea, anxiety, depression and somatoform pain: a community-based high-risk sample. *Eur Respir J* 2012;40:400–7.
- 41 Bardwell WA, Ancoli-Israel S, Dimsdale JE. Comparison of the effects of depressive symptoms and apnea severity on fatigue in patients with obstructive sleep apnea: a replication study. *J Affect Disord* 2007;97(1-3):181–6.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow