

# Artificial gametes from stem cells

Inmaculada Moreno<sup>1</sup>, Jose Manuel Míguez-Forjan<sup>2</sup>, Carlos Simón<sup>1,2,3</sup>

<sup>1</sup>Department of Research and Development, Igenomix S.L., Paternam; <sup>2</sup>Fundación Instituto Valenciano de Infertilidad (FIVI), Valencia, Spain;

<sup>3</sup>Department of Obstetrics and Gynecology, Stanford University School of Medicine, Stanford, CA, USA

The generation of artificial gametes is a real challenge for the scientific community today. *In vitro* development of human eggs and sperm will pave the way for the understanding of the complex process of human gametogenesis and will provide with human gametes for the study of infertility and the onset of some inherited disorders. However, the great promise of artificial gametes resides in their future application on reproductive treatments for all these people wishing to have genetically related children and for which gamete donation is now their unique option of parenthood. This is the case of infertile patients devoid of suitable gametes, same sex couples, singles and those fertile couples in a high risk of transmitting serious diseases to their progeny. In the search of the best method to obtain artificial gametes, many researchers have successfully obtained human germ cell-like cells from stem cells at different stages of differentiation. In the near future, this field will evolve to new methods providing not only viable but also functional and safe artificial germ cells. These artificial sperm and eggs should be able to recapitulate all the genetic and epigenetic processes needed for the correct gametogenesis, fertilization and embryogenesis leading to the birth of a healthy and fertile newborn.

**Keywords:** Artificial gametes; Cell-based therapy; Gametogenesis; Germ cells; Human germ cell specification; Pluripotent stem cells; Primordial germ cell-like cells

## Relevance of artificial gametes

Artificial gametes are mature germ cells (sperm and eggs) generated *in vitro* by specification and maturation of their natural diploid precursors, the primordial germ cells (PGCs), or by directed differentiation of pluripotent cells to the germ-cell lineage. These cells, after completing meiosis, should be able to undergo fertilization and subsequent embryogenesis, transmitting their genetic and epigenetic information to the next generation as their *in vivo* counterparts do [1].

The generation of human artificial gametes are of outstanding in-

terest in the context of assisted reproductive medicine as they would offer a clinical solution for different patients seeking for *in vitro* fertilization (IVF) treatments, but they would also help to understand the complex mechanism of human gametogenesis that is currently poorly studied due to the technical and ethical limitations associated to the use of those human samples for research.

### 1. Clinical use of artificial gametes

#### 1) Infertile couples

Infertility is a clinical condition that concerns 15% of couples in reproductive age. Among them, 28% are affected by different pathologies causing absence of available gametes [2] including patients with premature ovarian failure [3], post-menopausal women, and male patients suffering from non-obstructive azoospermia or any other pathology leading to the absence of either spermatozoa or elongated spermatids available to be used in assisted reproductive treatments (Guidelines on Male Infertility. European Association of Urology 2009. Available from: [http://www.uroweb.org/fileadmin/tx\\_eau-guidelines/2007/Full/Male\\_Infertility.pdf](http://www.uroweb.org/fileadmin/tx_eau-guidelines/2007/Full/Male_Infertility.pdf)). These pathologies produce infertility in up to 1% of women [4] and 0.63% of men [5] in the

Received: Jun 16, 2015 · Revised: Jun 18, 2015 · Accepted: Jun 18, 2015

Corresponding author: **Carlos Simón**

Fundación IVI, Parc Científic Universitat de València, C\Catedrático Agustín

Escardino nº 9. Edificio 3, 46980- Paterna, Spain

Tel: +34-963903305 ext.18530-18102 Fax: +34-963902522

E-mail: Carlos.Simon@ivi.es, csimonv@stanford.edu

\*This research was supported by Grant PTQ-13-06133 to IM. KY Cha Award in Stem Cell Technology to CS.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

general population. For these patients the use of donated gametes, either ovum or sperm, is mandatory. Also couples bearing genetic disorders with a high risk of transmitting serious diseases to their progeny or patients with repeated IVF failure usually go to donated gametes to improve their chances to have a healthy baby. Another group of patients interested in the therapeutic use of artificial gametes in assisted reproductive treatments rely on patients that have overcome a cancer in their early infancies and are infertile due to the anticancer treatments received while they were unable to cryopreserve their own gametes [6-12].

## 2) Fertile couples

On the other hand, people who are fertile would also benefit from the use of artificial gametes, as same sex couples, single parents and post-menopausal women. For these kind of patients, donor gametes are the sole option to have their own genetic related descendants. Within this group, same sex couples would be able to have their own artificial gametes, either sperm or eggs, independently of their sex, so that both partners of the couple could be genetic parents of the newborn.

## 2. Scientific use of artificial gametes

The generation of artificial gametes will not only provide therapeutic advantages for those people wishing to have genetically related progeny and for whom donor gametes is their only current chance to have babies, but it will also provide new knowledge in the field of germ cell biology. The *in vitro* generation of human gametes will help to understand the biological process of germ cell specification and maturation. Nowadays, developmental studies regarding human germ cell development are performed by using animal models due to the lack of available human samples. However, it has been widely shown that gametogenesis is species-specific, and although the general developmental process is conserved among mammals, there are some differences that hampers the direct translation of the knowledge acquired in mouse into human [13,14]. The main interests on artificial gametes production are focused on the establishment of a reproducible method to generate fully committed haploid sperm and oocytes that will offer large amounts of cells at different stages of germ cell formation and that will be the basis for further experiments. This is of outstanding interest for the understanding of the different molecular and cellular events that take place during normal human gametogenesis and subsequent study of human disease models related to germ cell defects or other diseases producing infertility and lack of functional gametes. If obtaining artificial gametes from disease patients, the mechanisms of some kinds of infertility could be unraveled and thus potential treatments could be developed for those patients devoid of functional gametes. Also, by com-

binning artificial gamete production from stem cells with new gene editing techniques, the correction of diseases could be potentially carried out in those patients suffering from inherited genetic disorders, so that their offspring would not suffer, nor transmit the disease to their progeny. On the other hand, the information obtained from *in vitro* gametogenesis models could be used to avoid the maturation of germ cells, leading to new and effective contraceptives.

As an example of the scientific strength of artificial gamete production from pluripotent cells, Dominguez and collaborators have recently published a study in which induced pluripotent stem cells from Turner syndrome individuals were differentiated *in vitro* into germ cell-like cells (GCLC) and compared to GCLC from control individuals [15]. The results of this work showed that aneuploidies in X chromosome do not impair germ cell formation but the correct dose of X chromosome is critical for the maintenance or functionality of those GCLC until adulthood, uncovering the origin of infertility in Turner syndrome patients.

## Gametes generation: the *in vivo* process

Most of the knowledge accumulated in the development of mammalian germline comes from animal models, mainly with the well-known murine model, while the process is still poorly understood in humans.

Gametes are highly specialized cells responsible for transmitting genetic and epigenetic information through generations [1]. The differentiation between germ and somatic cells occurs very early in development when a group of mesodermal cells escape from their somatic fate during gastrulation, and is characterized by two fundamental facts: reacquisition of the pluripotency and extensive epigenetic remodeling.

Based on the mouse model, during gastrulation the founder population of primordial germ cells (PGCs) are specified by extrinsic signals driven by bone morphogenetic proteins (BMP4 and BMP8b); BMPs induce *Prdm1* and *Prdm14* expression in mouse PGCs probably through the expression of mesodermal factor *T (Brachyury)* [16]. These events have been also observed in human PGC specification with few exceptions, such as overexpression of PRDM14 which is down-regulated in humans [13,14]. *Tcfap2c*, which encodes AP2 $\gamma$ , has also been described in mouse as a critical element that acts downstream of *Prdm1*. AP2 $\gamma$  might also be involved in the repression of somatic genes, including early mesodermal markers, such as *Hoxb1* [17]. After their specification, PGCs are characterized by the expression of several pluripotency genes such as *Oct4* (also known as *Pou5f1*), *Nanog* and *Sox2* [18], as well as other markers such as *Fragilis* and *Stella* [17,19]. Again there are small differences between mice and humans, as the lack of *SOX2* expression and a delay in the ex-

pression of *PRDM14* [14,20].

At the moment of their specification, PGCs are epigenetically indistinguishable from surrounding epiblastic cells, and show inherited epigenetic modifications such as DNA methylation and X chromosome inactivation that represent epigenetic barriers to totipotency acquisition in somatic tissues. PGCs begin the first wave of a profound epigenetic remodeling in order to acquire the epigenetic state necessary to form functional gametes [21]. The first epigenetic changes in PGC reprogramming take place at the histone modification level, leading to a large chromatin remodeling. These changes in histone modifications occur in parallel to an overall reduction in DNA methylation during PGC migration [21,22]. A deeper reset of methylated DNA occurs in a second wave of epigenetic remodeling events [23]. This process involves a global erasure of DNA methylation patterns, including imprinted genes, while PGCs reach a basal epigenetic state [21,24,25].

## Artificial gametes generation: the *in vitro* counterpart

With the development of *in vitro* models we can recapitulate human gametogenesis to understand the mechanisms behind meiotic control and the underlying cross-talk between germ cells and surrounding somatic cells. Two kinds of stem cells can be used as starting material for the generation of artificial gametes: (1) adult stem cells from both male and female gonads, and (2) pluripotent stem cells such as embryonic stem cells (ESCs), obtained by *in vitro* derivation of the embryonic inner cell mass [26], or induced pluripotent stem cells (iPSCs), obtained from somatic cells by overexpression of exogenous defined factors [27-29].

### 1. From adult stem cells

There are numerous adult tissues, if not all, that have the ability to regenerate themselves due to the existence of a specific population of adult stem cells with capacity for self-renewal and specification within their cell lineages [30-32]. The existence of this type of stem cells in the testis, the so-called spermatogonial stem cells (SSCs), has been widely demonstrated [33]. They are located in the basal lamina of the seminiferous tubule and are responsible for the continued production of sperm during the life of males. SSCs start to asymmetrically divide when sexual maturity is reached, giving rise to mitotically active spermatogonia, whilst also self-renewing [34-37], always under the control of a very specific microenvironment directed by Sertoli cells by building the balance between self-renewal and differentiation. It is possible to identify and purify human SSC by using different membrane markers such as  $\alpha 6$ -integrin (CD49f) and  $\beta 1$ -integrin, two laminin receptors [38], CD9, a basal membrane cell at-

tachment protein [39] and other markers [37,40-44]. SSC culture is challenging due to the need of blocking differentiation, maintenance of long-term survival, and promoting of proliferation. Attempts of *in vitro* spermatogenesis demonstrate promising results in humans, where SSCs isolated from human biopsies and co-cultured with Sertoli cells have shown expression of markers for meiotic progression and haploid cell formation [45]. However they have not fared better, due in part to the need for a co-culture system that mimics the microenvironment of testicular niche. Perhaps for this reason neither fertilization nor healthy offspring have been reported to date with gametes generated by this mechanism (Table 1) [13-15, 45-58].

In the female counterpart, it is generally accepted that human ovaries contain a fixed number of non-growing follicles established before birth that decreases with female age and is depleted in menopause [59-68]. Some researchers very early on struggled against the growing dogma about the finite number of follicles and oocytes at birth [69]. In the mouse model, the existence of ovarian stem cells leading to viable offspring was first reported by Zou et al. [70] in 2009. In spite of the fact that there is no evidence of persistence of oogonia in adult human ovaries, a new idea about germinal (oogonial) stem cells in adult ovaries has been developed [71-74]. The existence of stem cells in the human ovary has been surrounded by controversy. Some works in mouse and human have described a population of rare ovarian stem cells able to generate oocytes *in vitro* [55,75], representing an invaluable promise in regenerative medicine to treat infertile women. However, these works have been widely refuted, and some other authors consider that the presence of ovarian stem cells has been overestimated [76-79]. The ovarian stem cells are the most interesting population of cells for potential autologous *de novo* oogenesis and regeneration of non-functional ovaries in infertile women. In the human adult ovaries, putative ovarian stem cells (OSC) can be isolated by Fluorescence-activated cell sorting (FACS) as small rounded VASA positive-cells from the ovarian surface epithelium. These cells can grow and develop into oocyte-like cells *in vivo* and *in vitro* [55,80-82]. Interestingly there appears to be more positive results for the *in vitro* generation of female gametes rather to sperm (Table 1). This may be due to the greater ease to perform co-culture of OSC with the surrounding somatic cells; because follicle-like structure is easier to reproduce *in vitro* [55]. However, this starting material is limited or even absent in infertile patients so they do not seem to be the best option for regenerative medicine purposes on female infertile patients. For this reason pluripotent stem cells (PSCs) seem to be a better option, according to its greater availability in those kind of patients.

### 2. From human pluripotent stem cells (hESCs and hiPSCs)

First evidences of *in vitro* germ line formation from pluripotent stem

**Table 1.** Human germ cell-like cells *in vitro* derivation

Origin	Derivation conditions	<i>In vivo</i> assays	Epigenetic features	Germ cell markers	Outcome	Reference
hESC (XY, XX)	Adherent culture; BMP4, BMP8b, RA; <i>DAZ2, DAZL, BOULE</i> overexpression	Not analyzed	mC low global levels. <i>H19, PEG1, SNRPN</i> , and <i>KCNQ</i> ICR hypomethylation	PRDM1, VASA, DAZL, STELLA, $\gamma$ H2AX, SYCP3	Germ cell-like cells (1N, TEKT1, ACROSIN)	[46]
hiPSC-fetal (XX) Adult (XY)	Adherent culture; BMP4, BMP8b, RA <i>DAZ2, DAZL, BOULE</i> overexpression	Not analyzed	Not analyzed	PRDM1A, VASA, DAZL, STELLAR, FITM1, PELOTA, GCNF, SYCP3, CENP-A	Germ cell-like cells (1N, ACROSIN)	[47]
hESC (XX, XY) hiPSC (XX)	Adherent culture; BMP4, BMP8b, RA; <i>VASA</i> overexpression	Not analyzed	<i>H19</i> ICR demethylation	PRDM1, VASA, DAZL, IFITM1, GCN, GDF, cKIT, PELOTA, GDF9, ZP4, SYCP3, CENP-A	Germ cell-like cells	[48]
hiPSC (XX, XY)	BMP4, BMP8b; <i>VASA</i> overexpression	Transplantation into mouse testis	<i>KCNQ1OT1, PEG1, H19</i> , and <i>H19</i> ICR hypomethylation; Global demethylation; 5mC to 5hmC conversion	PRDM14, VASA, STELLA, NANOS2/3, DAZ2, SYCP3	Induced PGCs (2N)	[49]
hiPSC (XY) Y microdeletions	BMP4, BMP8b	Transplantation into mouse testis	Global DNA demethylation	PRDM1, PRDM14, VASA, DAZ, STELLA, IFITM3, NANOS3, PLZF, UTF1, SALL4	Induced PGCs (2N)	[50]
hiPSC (X0)	BMP4, BMP8b	Transplantation into mouse testis	Not analyzed	Not analyzed	Induced PGCs (2N)	[15]
hESCs (XX, XY) iPSCs (XX, XY)	4i medium; bFGF, BMP2, BMP4, SCF, LIF, EGF, ROCK inhibitor.	Not analyzed	Upregulation <i>TET1</i> and <i>TET2</i> ; 5mC to 5hmC conversion; Nucleus translocation of PRMT5; Downregulation of <i>UHRF1, DNMT3A</i> , and <i>DNMT3B</i>	BLIMP1, TFAP2C, DND1, KIT, SOX17, CD38	Induced hPGCLCs (2N)	[13]
hESC (XX, XY)	Embryoid body formation; BMP4, RA, medium from NMT	Not analyzed	Not analyzed	PGC markers: cKIT, SSEA1; Meiosis marker: SCYP3	Spermatid-like cells: PRMI, 1.97, 1N DNA Follicle-like structures: GDF9 and ZPI	[51]
hESC (XY) hiPSC (XY)	SSC medium (MEM, 0.2% iFBS, bFGF)	Not analyzed	<i>H19</i> and <i>IGF2</i> normal imprinting patterns.	Spermatogonia: UTF1, PLZF, CDH1 Spermatocyte: HIWI/HILI	Male germ cell, (postmeiotic spermatid-like cells)	[52]

(Continued to the next page)

Table 1. Continued

Origin	Derivation conditions	<i>In vivo</i> assays	Epigenetic features	Germ cell markers	Outcome	Reference
hESC (XX, XY)	hESC without bFGF;	Not analyzed	<i>H19</i> methylation levels;	1N cells, Acrosin, TNP1 and PRM1 VASA, SYCP3, $\gamma$ H2AX	Sperm-like phenotype	[53]
hiPSC (XX, XY)	RA, LIF, FRSK, bFGF, CYP26 inhibitor.		<i>CDKN1C</i> , <i>PHLDA2</i> (maternal allele);  <i>MEST</i> , <i>IGF2</i> , <i>NNAT</i> , <i>SNRPN</i> (paternal allele).			
hESC (XX)	EB and monolayer differentiation	Not analyzed	CpG sites of <i>H19/IGF2</i> DMRs show a decrease of methylation;  H3K4 dimethylation increase;  H3K9 dimethylation decrease	SSEA1, VASA, STELLA, OCT4, SCP1, SCP3.	Putative PGCs	[54]
Oogonial stem cells (XX)	Isolation by immunomagnetic sorting from human ovarian cortical tissue.	YES (Oocyte generation)	Not analyzed	VASA, KIT, BLIMP1, STELLA, FRAGILIS, DAZL, NOBOX, ZP1, ZP2, ZP3, GDF9, DMC1, SYCP3.	Oocyte-like cells (1N)	[55]
Human hepatic cell line (HL7702) (X0)	<i>In vitro</i> culture differentiation to PGC-like cells (DMEM + 10% FBS).	Not analyzed	Not analyzed	AP, OCT4, C-KIT, NANOS3, VASA, DAZL, GDF9, SYCP3, ZP3	Follicle-like structure  Oocyte-like cells.  Spontaneous embryo-like structures	[56]
Human endometrial cells (XX)	Somatic cell haploidization using GV enucleated mouse oocytes.	Not analyzed	Not analyzed	Not analyzed	MII oocytes	[57]
hESC, hiPSC (XX, XY)	bFGF, ActA, BMP4  20% KSR, BMP4, LIF, ROCK inhibitor	YES  (PGCLC Survive but not progress)	<i>PEG1</i> , <i>PEG10</i> and <i>NESP55</i> ;  Decreased in methylation levels.	OCT4, NANOG, SOX2, BLIMP1, STELLA, NANOS3, cKIT, T	PGCLCs	[14]
SSC (azoospermic patients)	Co-cultured with Sertoli cells  KSR, FSH, testosterone, GDNF, RA.	Not analyzed	Not analyzed	SCP3, CREST MLH1  Haploid for chromosomes 13, 18, 21, X and Y by FISH.	Meiotic-like cells	[45]
SSCs (cryptorchid patients)	RA, SCF.	YES (Round spermatids generated and ROSI)	Not analyzed	SYCP1, SYCP3, ACR, TNP1, TNP2, PRM1, PRM2, BOULE, CREST, DMC1	Haploid spermatids; early embryos.	[58]

hESC, human embryonic stem cell; BMP, bone morphogenetic protein; RA, retinoic acid; ICR, imprinted control region; hiPSC, human induced pluripotent stem cell; PGC, primordial germ cell; PGCLC, PGC-like cell; LIF, leukemia inhibitory factor; EGF, epidermal growth factor; ROCK, Rho-kinase; NMT, neonatal mouse testes; bFGF, basic fibroblast growth factor; KSR, knockout serum replacement; FSH, follicle-stimulating hormone; GDNF, glial derived neurotrophic factor; SSC, spermatogonial stem cells; SCF, stem cell factor; ROSI, round spermatid injection.

cells in humans were reported by spontaneous differentiation [83,84] of hESCs embryoid bodies [51,54]. Undifferentiated hESCs expressed some early germ cell markers such as c-KIT and DAZL, but not later markers such as VASA (a RNA helicase involved in germ cell maturation in both sexes) or SYCP3 (a structural protein of the synaptonemal complex critical for meiosis) [85], suggesting that hESCs could be a heterogeneous pluripotent population in which some cells had a predisposition towards a germ cell fate. Addition of several molecules have been shown to improve the differentiation efficiency of pluripotent cells to germ cells, this is the case of BMP cytokines [51,52,56], and retinoic acid (RA) that has been demonstrated to induce meiosis [15,50,86]. However, meiotic progression remained as one of the main obstacles for artificial gametes derivation [87]; most of the works published to date report spontaneous differentiation towards putative PGCs or PGC-like cells (PGCLC) with diploid (2N) content of DNA (Table 1), but the completion of meiosis to give haploid (1N) cells remains a challenge when spontaneous differentiation is used.

In fact, complete meiotic progression of *in vitro* derived human germ cells was achieved by inducing the ectopic expression of the DAZ gene family members (*DAZ2*, *DAZL*, and *BOULE*) not only in hESC but also in human iPSCs (hiPSCs) lines subjected to spontaneous differentiation [46,47]. The expression of these highly conserved RNA-binding proteins lead the correct meiotic progression of human germ cells *in vitro* in the absence of a gonadal niche. Also RNA-binding proteins, like VASA, could have a possible role in meiotic entry control [48]. A recent study have provided a more accurate process, in which pluripotent stem cells were first cultured in 4i conditions to induce the naïve pluripotent state, this step make cells prone to response to specification signals [88]. Then, the derivation process was performed in culture media supplemented with BMPs, leukemia inhibitory factor (LIF), Rho kinase (ROCK) inhibitor and Knockout serum replacement, to obtain the PGC precursors and then PGCLCs [13,14]. This system allows for the generation of large numbers of cells for use in research procedures. In summary, human artificial meiotic or sperm-like cells had been generated from both ESCs and iPSCs, assessed by the expression of meiotic and post-meiotic markers and DNA ploidy (Table 1).

On the female side, less studies have been performed with hESC and hiPSC for oocytes generation, partially due to the difficulties derived from the complexity of the own oocyte cell. Nonetheless previous data has shown that oocyte-like cells, expressing oocyte-specific markers, can be developed from ESCs and iPSCs [53,89-93]. These oocyte-like cells were developed from stem cells that expressed at least a degree of pluripotency, and the resulting oocyte-like cells expressed some germinal markers like the structural meiotic protein SYCP3 [90] as well as c-KIT, ZP1, ZP2, ZP3, VASA, and DMC1 [81] but were still far from being real and competent oocytes. Finally, the

most challenged goal in the artificial gamete generation seems to be the production of sperm and eggs from the opposite sex. This would be of great importance to gay couples who wish to have genetically related offspring. However, little has been achieved in this direction and only a couple of papers reporting generation of sperm from female cells exist, albeit this sperm was not fully functional and presented incorrect or unknown epigenetic status [94,95].

## Clinical use of artificial gametes in assisted reproductive treatments

Human artificial gametes are likely to be developed in the following years and they hold a great hope for the treatment of infertile patients devoid of gametes, allowing them to have genetically related progeny. However, clinical applications of *in vitro* derived gametes are unlikely to be used for therapeutic use until their fully functionality and safety will be guaranteed.

### 1. Functionality of artificial sperm and eggs- viability and long-term health of derived offspring

Determining functionality of derived gametes will, therefore, require establishing their capacity for fertilization and early embryogenesis. In order to consider artificial sperm for clinical purposes, they should be tested to be able to produce the main biological processes leading to fertilization as the initiation of oocyte activation, nuclear remodelling and subsequent fusion to the oocyte nucleus, and loss of the own nuclear envelop [96]. On the other side, when artificially derived eggs needs to be assessed for clinical use they should be able to escape from meiotic arrest, undergo pronuclear formation, initiate embryonic gene activation while recapitulating nuclear maturation events (i.e., germinal vesicle breakdown, chromosomal segregation, polar body extrusion) and cytoplasmic reorganization (i.e., adenosine triphosphate (ATP) supply by mitochondria, protein and mRNA storage) [97]. Finally, to fully address the generation of functional artificial gametes they need to be proved to successfully give a healthy embryo upon fecundation, showing correct genetic and epigenetic reprogramming, cleavage divisions and zygotic genome activation [98].

Ethical restrictions make difficult to carry out functional assays in humans with germ cells obtained from stem cells. Moreover, differences in the pluripotent status of mESCs (naïve-like state) and hESCs (primed state) make difficult to compare their potential to fully develop into functional germ cells [99]. The main obstacle to develop fully committed sperm and eggs from stem cells is the lack of knowledge regarding molecules and factors, which are important in the gamete maturation process in the gonad niche and are not present when the immature gametes (OSC or SSC) are exposed to *in vitro* cul-

ture conditions. For all these reasons, the functionality and safety of the obtained GCLC needs to be first tested in animal models.

## 2. Transplantation of artificial PGC into animal models

The most similar to functional assays are xenotransplantation of human germ cell-like cells into seminiferous tubules of immunosuppressed mice, in the male counterpart. These animal models are very helpful to test the ability of human germ cell-like cells to behave like human germ cell progenitors. In this context, the role of RNA-binding proteins has also been tested, demonstrating that hiPSCs improved their ability to colonize the lumen of the seminiferous tubules of sterilized immunodeficient mice upon *VASA* overexpression in combination with pluripotency factors. These results highlight the role of *VASA* in making cells competent to form germ cell development and in controlling the pluripotency-state in a combined *in vitro/in vivo* model [49].

This murine/human heterologous model represents a very interesting analysis method from a clinical point of view, and has already been used to analyze the ability of hiPSCs-GCLC derived from azospermic men with different deletions in the Y chromosome to colonize the spermatogonial niche, demonstrating how the genetic background affects their capability to differentiate into germ cells and colonize the seminiferous niche *in vivo* [49]. Finally, this same approach has been recently used to demonstrate that GCLC can be formed *in vivo* independently of X chromosome dosage [15]. In this case, hiPSCs from Turner syndrome patients were obtained, differentiated to GCLC and xenotransplanted into the seminiferous tubules of sterilized immunodeficient mice, demonstrating that the two X chromosomes are not required for human germ cell formation but for their maintenance until adulthood [15].

Also *in vitro* maturation of human oocytes is one of the most difficult tasks in the IVF programs. It has been proposed that transplantation of stem cells or stem cell-derived "oocytes" into the ovary or other organs would be of great advantage because it would provide a natural niche for maturation of oocytes and bypass several obstacles related to oocyte *in vitro* maturation [100]. In the mouse model it was confirmed that maturation of ESC-derived oocytes ultimately fails *in vitro*, and to overcome this obstacle the transplantation of ESC-derived oocyte-germ cells into an ovarian niche has been proposed to direct their natural and functional maturation. To demonstrate this, ESC-derived germ cells were enclosed into ovarian follicles, and some of them reached the primordial or primary stage of development after the two-step protocol by performing *in vitro* specification from mESCs and subsequent maturation in an *in vivo* model [100].

## 3. Epigenetic remodelling in artificial gametes

In 2006, the birth of murine offspring obtained by fertilization of

murine oocytes with artificial mESCs-derived sperm was achieved [101] proving for the first time that stem cells could be developed *in vitro* to obtain autologous germ cells for reproductive treatments. However, the offspring presented phenotypic abnormalities as well as reduced life-span, mainly because of epigenetic defects accumulated during germline *in vitro* specification and maturation [101]. Later on, Hayashi and collaborators reported the generation of healthy and fertile pups by fertilization of mouse oocytes with artificial sperm obtained by two-step differentiation of mESC through an intermediate epiblastic-like cells [102]. In this work, the authors assessed not only the cellular events and transcriptomics along the generation of gametes, but also their epigenetic profiles, showing that the *in vitro*-derived germ cells closely resemble the physiological process [102], revealing the importance of the acquisition of a new epigenetic program in parallel to transcriptomic changes during differentiation from stem to germ cells. It has been also reported that erroneous epigenetic imprinting can produce diseases in humans [103] emphasizing the relevance of epigenetics in the developmental process from germ cells to adult life.

As discussed before, one of the crucial facts in PGC specification is an extensive epigenetic remodeling. This epigenetic remodeling is mainly characterized by an overall reduction in DNA methylation during migration of PGCs to the gonadal ridge [22]. DNA methylation is a repressive mark that mainly targets cytosines within CpG islands (palindromic CpG dinucleotides in the genome) [24] producing a complete erase of the methylation patterns and thus the acquisition of totipotency in gametes of both sexes [104]. Then, during sex determination, PGCs must establish their sex-specific epigenetic patterns, which include paternal or maternal imprinted marks in order to acquire the epigenetic state necessary to form functional gametes [21]. The imprinting genes involve genomic sequences that exhibit differences in CpG methylation according to the parental origin. These differentially methylated regions (DMRs) can influence the allele-specific expression of one or more genes [21]. After fertilization, a second wave of epigenetic reprogramming is produced in all the genome but in the imprinted regions, which in turn regulates fetal and placental growth, differentiation, development and other important functions after birth [105].

It has been recently suggested that *in vitro* manipulation of cells could lead to epigenetic alterations affecting either stem cells or the obtained artificial gametes [106-109]. Not all the studies performed in the field have analyzed the epigenetic status of generated cells, but when authors assessed epigenetic features of these cells, they found similar patterns than their *in vivo* counterparts, with a sharp decrease in global methylation levels, and methylation patterns in DMRs of imprinting genes according to their fate [46,48]. Nonetheless, the correct epigenetic state of artificial gametes is an important

issue that will determine the successful development of the offspring. Especial attention should be paid to the epigenetic analysis on DNA methylation and histone modification to ensure the complete imprinting of artificial gametes and avoid partially reprogrammed artifacts [53].

#### 4. Safety of artificial gametes

Finally, the long-term safety of the obtained artificial gametes will be a critical issue to assess the quality and applicability of derived artificial gametes into the clinical practice. Observational and biochemical tests can be performed to demonstrate the safety of artificial sperm or eggs, but these assays are insufficient to judge whether the cells would support normal development in humans upon fertilization until the adult life. The most robust and stringent studies reporting human PCGs generated *in vitro* have shown that these cells present normal germ cell genetic and epigenetic profiles and are able to colonize [14] and proliferate when xenotransplanted into mice testes [15,49]. However, some other works have reported incomplete imprinting of the generated germ-like cells [53]. These findings highlight the need for more consistent research that should be focused to the following main directions: (1) tumorigenicity of the cells as *in vitro* culture of stem cells for long periods as well as reprogramming of patient-specific somatic cells to hiPSCs have been shown to constitute risks factors for accumulation of chromosomal aneuploidies [110,111]; (2) immunogenicity of *in vitro* generated gametes, which remains a challenge for their use as potential progenitors to be transplanted into the host as they could produce immunological rejection problems even if they are used in autologous therapy, as it has been reported for some hiPSCs cell lines [112]; (3) the future use of artificial gametes in regenerative medicine programs would require the obtention of therapeutic grade sperm and eggs attending to strict production conditions that guarantee that these cells do not present additional risks acquired during their manipulation.

## Conclusion

Generation of artificial gametes is a great promise for all those couples that, for many different reasons, currently need to go to gamete donation for fulfilling their wish to have children but would prefer to have genetically related offspring due to ethical, social or emotional reasons. Also, the successful generation of artificial gametes will help to understand the complex biological process of gametogenesis in humans, leading to the improvement in the knowledge and potential treatments of many disorders related to infertility and those diseases that are genetically transmitted to the progeny.

Despite the great interest that this field has arisen in the last years, there is still a long way to go until artificial germ cells will be used in

the clinical practice. First, from the technological point of view, the accumulated knowledge learnt from the mouse model needs to be translated to humans considering the important differences between both species. Once we are able to obtain germ cells *in vitro* from stem cells (either adult or pluripotent) the feasibility of using artificial sperm or oocytes for IVF treatments should be carefully analyzed attending to different criteria. In this regard, functionality and safety of artificial gametes needs to be assessed genetically and epigenetically to prove the health of the offspring derived from artificial eggs or sperm.

## Conflict of interest

No potential conflict of interest relevant to this article was reported.

## References

1. Johnson AD, Richardson E, Bachvarova RF, Crother BI. Evolution of the germ line-soma relationship in vertebrate embryos. *Reproduction* 2011;141:291-300.
2. Kashir J, Jones C, Child T, Williams SA, Coward K. Viability assessment for artificial gametes: the need for biomarkers of functional competency. *Biol Reprod* 2012;87:114.
3. Sinha P, Kuruba N. Premature ovarian failure. *J Obstet Gynaecol* 2007;27:16-9.
4. Coulam CB, Adamson SC, Annegers JF. Incidence of premature ovarian failure. *Obstet Gynecol* 1986;67:604-6.
5. Jarow JP, Espeland MA, Lipshultz LI. Evaluation of the azoospermic patient. *J Urol* 1989;142:62-5.
6. Hwang K, Lamb DJ. New advances on the expansion and storage of human spermatogonial stem cells. *Curr Opin Urol* 2010;20:510-4.
7. Levine J, Canada A, Stern CJ. Fertility preservation in adolescents and young adults with cancer. *J Clin Oncol* 2010;28:4831-41.
8. Silber SJ. Sperm retrieval for azoospermia and intracytoplasmic sperm injection success rates: a personal overview. *Hum Fertil (Camb)* 2010;13:247-56.
9. Woodruff TK. The Oncofertility Consortium: addressing fertility in young people with cancer. *Nat Rev Clin Oncol* 2010;7:466-75.
10. Wyns C, Curaba M, Vanabelle B, Van Langendonck A, Donnez J. Options for fertility preservation in prepubertal boys. *Hum Reprod Update* 2010;16:312-28.
11. Jahnukainen K, Ehmcke J, Hou M, Schlatt S. Testicular function and fertility preservation in male cancer patients. *Best Pract Res Clin Endocrinol Metab* 2011;25:287-302.
12. Wallace WH. Oncofertility and preservation of reproductive capacity in children and young adults. *Cancer* 2011;117:2301-10.



13. Irie N, Weinberger L, Tang WW, Kobayashi T, Viukov S, Manor YS, et al. SOX17 is a critical specifier of human primordial germ cell fate. *Cell* 2015;160:253-68.
14. Sugawa F, Arauzo-Bravo MJ, Yoon J, Kim KP, Aramaki S, Wu G, et al. Human primordial germ cell commitment in vitro associates with a unique PRDM14 expression profile. *EMBO J* 2015;34:1009-24.
15. Dominguez AA, Chiang HR, Sukhwani M, Orwig KE, Reijo Pera RA. Human germ cell formation in xenotransplants of induced pluripotent stem cells carrying X chromosome aneuploidies. *Sci Rep* 2014;4:6432.
16. Aramaki S, Hayashi K, Kurimoto K, Ohta H, Yabuta Y, Iwanari H, et al. A mesodermal factor, T, specifies mouse germ cell fate by directly activating germline determinants. *Dev Cell* 2013;27:516-29.
17. Weber S, Eckert D, Nettersheim D, Gillis AJ, Schafer S, Kuckenberger P, et al. Critical function of AP-2 gamma/TCFAP2C in mouse embryonic germ cell maintenance. *Biol Reprod* 2010;82:214-23.
18. Niwa H. How is pluripotency determined and maintained? *Development* 2007;134:635-46.
19. Aflatoonian B, Moore H. Human primordial germ cells and embryonic germ cells, and their use in cell therapy. *Curr Opin Biotechnol* 2005;16:530-5.
20. Perrett RM, Tumpenny L, Eckert JJ, O'Shea M, Sonne SB, Cameron IT, et al. The early human germ cell lineage does not express SOX2 during in vivo development or upon in vitro culture. *Biol Reprod* 2008;78:852-8.
21. Hackett JA, Zylitz JJ, Surani MA. Parallel mechanisms of epigenetic reprogramming in the germline. *Trends Genet* 2012;28:164-74.
22. Godin I, Wylie C, Heasman J. Genital ridges exert long-range effects on mouse primordial germ cell numbers and direction of migration in culture. *Development* 1990;108:357-63.
23. Seisenberger S, Andrews S, Krueger F, Arand J, Walter J, Santos F, et al. The dynamics of genome-wide DNA methylation reprogramming in mouse primordial germ cells. *Mol Cell* 2012;48:849-62.
24. Lee HJ, Hore TA, Reik W. Reprogramming the methylome: erasing memory and creating diversity. *Cell Stem Cell* 2014;14:710-9.
25. Martinez-Arroyo AM, Medrano JV, Remohi J, Simon C. Germ line development: lessons learned from pluripotent stem cells. *Curr Opin Genet Dev* 2014;28:64-70.
26. Aguilar-Gallardo C, Poo M, Gomez E, Galan A, Sanchez E, Marques-Mari A, et al. Derivation, characterization, differentiation, and registration of seven human embryonic stem cell lines (VAL-3, -4, -5, -6M, -7, -8, and -9) on human feeder. *In Vitro Cell Dev Biol Anim* 2010;46:317-26.
27. Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 2006;126:663-76.
28. Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, et al. Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 2007;131:861-72.
29. Yu J, Vodyanik MA, Smuga-Otto K, Antosiewicz-Bourget J, Frane JL, Tian S, et al. Induced pluripotent stem cell lines derived from human somatic cells. *Science* 2007;318:1917-20.
30. Rompolas P, Greco V. Stem cell dynamics in the hair follicle niche. *Semin Cell Dev Biol* 2014;25-26:34-42.
31. Watt FM. Mammalian skin cell biology: at the interface between laboratory and clinic. *Science* 2014;346:937-40.
32. Wabik A, Jones PH. Switching roles: the functional plasticity of adult tissue stem cells. *EMBO J* 2015;34:1164-79.
33. Izadyar F, Wong J, Maki C, Pacchiarotti J, Ramos T, Howerton K, et al. Identification and characterization of repopulating spermatogonial stem cells from the adult human testis. *Hum Reprod* 2011;26:1296-306.
34. Kolasa A, Misiakiewicz K, Marchlewicz M, Wiszniewska B. The generation of spermatogonial stem cells and spermatogonia in mammals. *Reprod Biol* 2012;12:5-23.
35. Waheeb R, Hofmann MC. Human spermatogonial stem cells: a possible origin for spermatocytic seminoma. *Int J Androl* 2011;34:e296-305.
36. Phillips BT, Gassei K, Orwig KE. Spermatogonial stem cell regulation and spermatogenesis. *Philos Trans R Soc Lond B Biol Sci* 2010;365:1663-78.
37. Hermann BP, Sukhwani M, Hansel MC, Orwig KE. Spermatogonial stem cells in higher primates: are there differences from those in rodents? *Reproduction* 2010;139:479-93.
38. Conrad S, Renninger M, Hennenlotter J, Wiesner T, Just L, Bonin M, et al. Generation of pluripotent stem cells from adult human testis. *Nature* 2008;456:344-9.
39. Kanatsu-Shinohara M, Toyokuni S, Shinohara T. CD9 is a surface marker on mouse and rat male germline stem cells. *Biol Reprod* 2004;70:70-5.
40. Kubota H, Avarbock MR, Brinster RL. Spermatogonial stem cells share some, but not all, phenotypic and functional characteristics with other stem cells. *Proc Natl Acad Sci U S A* 2003;100:6487-92.
41. Ryu BY, Orwig KE, Kubota H, Avarbock MR, Brinster RL. Phenotypic and functional characteristics of spermatogonial stem cells in rats. *Dev Biol* 2004;274:158-70.
42. Buaas FW, Kirsh AL, Sharma M, McLean DJ, Morris JL, Griswold MD, et al. Plzf is required in adult male germ cells for stem cell self-renewal. *Nat Genet* 2004;36:647-52.

43. Hofmann MC, Braydich-Stolle L, Dym M. Isolation of male germline stem cells; influence of GDNF. *Dev Biol* 2005;279:114-24.
44. von Schonfeldt V, Wistuba J, Schlatt S. Notch-1, c-kit and GFRalpha-1 are developmentally regulated markers for premeiotic germ cells. *Cytogenet Genome Res* 2004;105:235-9.
45. Riboldi M, Rubio C, Pellicer A, Gil-Salom M, Simon C. In vitro production of haploid cells after coculture of CD49f+ with Sertoli cells from testicular sperm extraction in nonobstructive azoospermic patients. *Fertil Steril* 2012;98:580-90.e4.
46. Kee K, Angeles VT, Flores M, Nguyen HN, Reijo Pera RA. Human DAZL, DAZ and BOULE genes modulate primordial germ-cell and haploid gamete formation. *Nature* 2009;462:222-5.
47. Panula S, Medrano JV, Kee K, Bergstrom R, Nguyen HN, Byers B, et al. Human germ cell differentiation from fetal- and adult-derived induced pluripotent stem cells. *Hum Mol Genet* 2011;20:752-62.
48. Medrano JV, Ramathal C, Nguyen HN, Simon C, Reijo Pera RA. Divergent RNA-binding proteins, DAZL and VASA, induce meiotic progression in human germ cells derived in vitro. *Stem Cells* 2012;30:441-51.
49. Durruthy Durruthy J, Ramathal C, Sukhwani M, Fang F, Cui J, Orwig KE, et al. Fate of induced pluripotent stem cells following transplantation to murine seminiferous tubules. *Hum Mol Genet* 2014;23:3071-84.
50. Ramathal C, Durruthy-Durruthy J, Sukhwani M, Arakaki JE, Turek PJ, Orwig KE, et al. Fate of iPSCs derived from azoospermic and fertile men following xenotransplantation to murine seminiferous tubules. *Cell Rep* 2014;7:1284-97.
51. Aflatoonian B, Ruban L, Jones M, Aflatoonian R, Fazeli A, Moore HD. In vitro post-meiotic germ cell development from human embryonic stem cells. *Hum Reprod* 2009;24:3150-9.
52. Easley CA, Phillips BT, McGuire MM, Barringer JM, Valli H, Hermann BP, et al. Direct differentiation of human pluripotent stem cells into haploid spermatogenic cells. *Cell Rep* 2012;2:440-6.
53. Eguizabal C, Montserrat N, Vassena R, Barragan M, Garreta E, Garcia-Quevedo L, et al. Complete meiosis from human induced pluripotent stem cells. *Stem Cells* 2011;29:1186-95.
54. Tilgner K, Atkinson SP, Golebiewska A, Stojkovic M, Lako M, Armstrong L. Isolation of primordial germ cells from differentiating human embryonic stem cells. *Stem Cells* 2008;26:3075-85.
55. White YA, Woods DC, Takai Y, Ishihara O, Seki H, Tilly JL. Oocyte formation by mitotically active germ cells purified from ovaries of reproductive-age women. *Nat Med* 2012;18:413-21.
56. Ma Z, Liu R, Wang X, Huang M, Gao Q, Lu Y, et al. Spontaneous germline potential of human hepatic cell line in vitro. *Mol Hum Reprod* 2013;19:216-26.
57. Palermo GD, Takeuchi T, Rosenwaks Z. Technical approaches to correction of oocyte aneuploidy. *Hum Reprod* 2002;17:2165-73.
58. Yang S, Ping P, Ma M, Li P, Tian R, Yang H, et al. Generation of haploid spermatids with fertilization and development capacity from human spermatogonial stem cells of cryptorchid patients. *Stem Cell Reports* 2014;3:663-75.
59. Block E. Quantitative morphological investigations of the follicular system in women; variations at different ages. *Acta Anat (Basel)* 1952;14:108-23.
60. Block E. A quantitative morphological investigation of the follicular system in newborn female infants. *Acta Anat (Basel)* 1953;17:201-6.
61. Baker TG. A Quantitative and Cytological Study of Germ Cells in Human Ovaries. *Proc R Soc Lond B Biol Sci* 1963;158:417-33.
62. Gougeon A, Chainy GB. Morphometric studies of small follicles in ovaries of women at different ages. *J Reprod Fertil* 1987;81:433-42.
63. Richardson SJ, Senikas V, Nelson JF. Follicular depletion during the menopausal transition: evidence for accelerated loss and ultimate exhaustion. *J Clin Endocrinol Metab* 1987;65:1231-7.
64. Hansen KR, Knowlton NS, Thyer AC, Charleston JS, Soules MR, Klein NA. A new model of reproductive aging: the decline in ovarian non-growing follicle number from birth to menopause. *Hum Reprod* 2008;23:699-708.
65. Bendsen E, Byskov AG, Andersen CY, Westergaard LG. Number of germ cells and somatic cells in human fetal ovaries during the first weeks after sex differentiation. *Hum Reprod* 2006;21:30-5.
66. Forabosco A, Sforza C. Establishment of ovarian reserve: a quantitative morphometric study of the developing human ovary. *Fertil Steril* 2007;88:675-83.
67. Faddy MJ, Gosden RG, Gougeon A, Richardson SJ, Nelson JF. Accelerated disappearance of ovarian follicles in mid-life: implications for forecasting menopause. *Hum Reprod* 1992;7:1342-6.
68. Faddy MJ, Gosden RG. A model conforming the decline in follicle numbers to the age of menopause in women. *Hum Reprod* 1996;11:1484-6.
69. Zuckerman. The number of oocytes in the mature ovary. *Rec Prog Horm Res.* 1951;6:63-108.
70. Zou K, Yuan Z, Yang Z, Luo H, Sun K, Zhou L, et al. Production of offspring from a germline stem cell line derived from neonatal ovaries. *Nat Cell Biol* 2009;11:631-6.
71. Tilly JL. Apoptosis and ovarian function. *Rev Reprod* 1996;1:162-72.
72. Oatley J, Hunt PA. Of mice and (wo)men: purified oogonial stem cells from mouse and human ovaries. *Biol Reprod* 2012;86:196.
73. Vogel G. Reproductive biology. Potential egg stem cells reignite debate. *Science* 2012;335:1029-30.
74. Ghazal S. Oogonial stem cells: do they exist and may they have

- an impact on future fertility treatment? *Curr Opin Obstet Gynecol* 2013;25:223-8.
75. Johnson J, Canning J, Kaneko T, Pru JK, Tilly JL. Germline stem cells and follicular renewal in the postnatal mammalian ovary. *Nature* 2004;428:145-50.
  76. Byskov AG, Faddy MJ, Lemmen JG, Andersen CY. Eggs forever? *Differentiation* 2005;73:438-46.
  77. Eggan K, Jurga S, Gosden R, Min IM, Wagers AJ. Ovulated oocytes in adult mice derive from non-circulating germ cells. *Nature* 2006;441:1109-14.
  78. Begum S, Papaioannou VE, Gosden RG. The oocyte population is not renewed in transplanted or irradiated adult ovaries. *Hum Reprod* 2008;23:2326-30.
  79. Liu Y, Wu C, Lyu Q, Yang D, Albertini DF, Keefe DL, et al. Germline stem cells and neo-oogenesis in the adult human ovary. *Dev Biol* 2007;306:112-20.
  80. Virant-Klun I, Zech N, Rozman P, Vogler A, Cvjeticanin B, Klemenc P, et al. Putative stem cells with an embryonic character isolated from the ovarian surface epithelium of women with no naturally present follicles and oocytes. *Differentiation* 2008;76:843-56.
  81. Virant-Klun I, Skutella T, Hren M, Gruden K, Cvjeticanin B, Vogler A, et al. Isolation of small SSEA-4-positive putative stem cells from the ovarian surface epithelium of adult human ovaries by two different methods. *Biomed Res Int* 2013;2013:690415.
  82. Szotek PP, Chang HL, Brennand K, Fujino A, Pieretti-Vanmarcke R, Lo Celso C, et al. Normal ovarian surface epithelial label-retaining cells exhibit stem/progenitor cell characteristics. *Proc Natl Acad Sci U S A* 2008;105:12469-73.
  83. Clark AT, Bodnar MS, Fox M, Rodriguez RT, Abeyta MJ, Firpo MT, et al. Spontaneous differentiation of germ cells from human embryonic stem cells in vitro. *Hum Mol Genet* 2004;13:727-39.
  84. Gkountela S, Li Z, Vincent JJ, Zhang KX, Chen A, Pellegrini M, et al. The ontogeny of cKIT+ human primordial germ cells proves to be a resource for human germ line reprogramming, imprint erasure and in vitro differentiation. *Nat Cell Biol* 2013;15:113-22.
  85. Syrjanen JL, Pellegrini L, Davies OR. A molecular model for the role of SYCP3 in meiotic chromosome organisation. *Elife* 2014;3:e02963.
  86. Bowles J, Koopman P. Retinoic acid, meiosis and germ cell fate in mammals. *Development* 2007;134:3401-11.
  87. Sun YC, Cheng SF, Sun R, Zhao Y, Shen W. Reconstitution of gametogenesis in vitro: meiosis is the biggest obstacle. *J Genet Genomics* 2014;41:87-95.
  88. Gafni O, Weinberger L, Mansour AA, Manor YS, Chomsky E, Ben-Yosef D, et al. Derivation of novel human ground state naive pluripotent stem cells. *Nature* 2013;504:282-6.
  89. Hubner K, Fuhrmann G, Christenson LK, Kehler J, Reinbold R, De La Fuente R, et al. Derivation of oocytes from mouse embryonic stem cells. *Science* 2003;300:1251-6.
  90. Novak I, Lightfoot DA, Wang H, Eriksson A, Mahdy E, Hoog C. Mouse embryonic stem cells form follicle-like ovarian structures but do not progress through meiosis. *Stem Cells* 2006;24:1931-6.
  91. West FD, Machacek DW, Boyd NL, Pandiyan K, Robbins KR, Stice SL. Enrichment and differentiation of human germ-like cells mediated by feeder cells and basic fibroblast growth factor signaling. *Stem Cells* 2008;26:2768-76.
  92. West FD, Roche-Rios MI, Abraham S, Rao RR, Natrajan MS, Bacanamwo M, et al. KIT ligand and bone morphogenetic protein signaling enhances human embryonic stem cell to germ-like cell differentiation. *Hum Reprod* 2010;25:168-78.
  93. Richards M, Fong CY, Bongso A. Comparative evaluation of different in vitro systems that stimulate germ cell differentiation in human embryonic stem cells. *Fertil Steril* 2010;93:986-94.
  94. Kerkis A, Fonseca SA, Serafim RC, Lavagnoli TM, Abdelmassih S, Abdelmassih R, et al. In vitro differentiation of male mouse embryonic stem cells into both presumptive sperm cells and oocytes. *Cloning Stem Cells* 2007;9:535-48.
  95. Kono T, Obata Y, Wu Q, Niwa K, Ono Y, Yamamoto Y, et al. Birth of parthenogenetic mice that can develop to adulthood. *Nature* 2004;428:860-4.
  96. Publicover S, Harper CV, Barratt C. [Ca<sup>2+</sup>]<sub>i</sub> signalling in sperm: making the most of what you've got. *Nat Cell Biol* 2007;9:235-42.
  97. Li Q, McKenzie LJ, Matzuk MM. Revisiting oocyte-somatic cell interactions: in search of novel intrafollicular predictors and regulators of oocyte developmental competence. *Mol Hum Reprod* 2008;14:673-8.
  98. Niakan KK, Han J, Pedersen RA, Simon C, Pera RA. Human preimplantation embryo development. *Development* 2012;139:829-41.
  99. MagnUSDottir E, Surani MA. How to make a primordial germ cell. *Development* 2014;141:245-52.
  100. Nicholas CR, Haston KM, Grewall AK, Longacre TA, Reijo Pera RA. Transplantation directs oocyte maturation from embryonic stem cells and provides a therapeutic strategy for female infertility. *Hum Mol Genet* 2009;18:4376-89.
  101. Nayernia K, Nolte J, Michelmann HW, Lee JH, Rathack K, Drusenheimer N, et al. In vitro-differentiated embryonic stem cells give rise to male gametes that can generate offspring mice. *Dev Cell* 2006;11:125-32.
  102. Hayashi K, Ogushi S, Kurimoto K, Shimamoto S, Ohta H, Saitou M. Offspring from oocytes derived from in vitro primordial germ cell-like cells in mice. *Science* 2012;338:971-5.

103. Grossniklaus U, Kelly WG, Ferguson-Smith AC, Pembrey M, Lindquist S. Transgenerational epigenetic inheritance: how important is it? *Nat Rev Genet* 2013;14:228-35.
104. Allegrucci C, Thurston A, Lucas E, Young L. Epigenetics and the germline. *Reproduction* 2005;129:137-49.
105. Messerschmidt DM, Knowles BB, Solter D. DNA methylation dynamics during epigenetic reprogramming in the germline and preimplantation embryos. *Genes Dev* 2014;28:812-28.
106. Geijsen N, Horoschak M, Kim K, Gribnau J, Eggan K, Daley GQ. Derivation of embryonic germ cells and male gametes from embryonic stem cells. *Nature* 2004;427:148-54.
107. Hajkova P, Erhardt S, Lane N, Haaf T, El-Maarri O, Reik W, et al. Epigenetic reprogramming in mouse primordial germ cells. *Mech Dev* 2002;117:15-23.
108. Manipalviratn S, DeCherney A, Segars J. Imprinting disorders and assisted reproductive technology. *Fertil Steril* 2009;91:305-15.
109. Hiura H, Okae H, Miyauchi N, Sato F, Sato A, Van De Pette M, et al. Characterization of DNA methylation errors in patients with imprinting disorders conceived by assisted reproduction technologies. *Hum Reprod* 2012;27:2541-8.
110. Ben-David U, Benvenisty N. The tumorigenicity of human embryonic and induced pluripotent stem cells. *Nat Rev Cancer* 2011;11:268-77.
111. Amps K, Andrews PW, Anyfantis G, Armstrong L, Avery S, Baharvand H, et al. Screening ethnically diverse human embryonic stem cells identifies a chromosome 20 minimal amplicon conferring growth advantage. *Nat Biotechnol* 2011;29:1132-44.
112. Zhao T, Zhang ZN, Rong Z, Xu Y. Immunogenicity of induced pluripotent stem cells. *Nature* 2011;474:212-5.