

SCIENTIFIC REPORTS



OPEN

The effect of oxygen concentration and temperature on nitrogenase activity in the heterocystous cyanobacterium *Fischerella* sp.

Lucas J. Stal^{1,2}

Heterocysts are differentiated cells formed by some filamentous, diazotrophic (dinitrogen-fixing) cyanobacteria. The heterocyst is the site of dinitrogen fixation providing the oxygen-sensitive nitrogenase with a low-oxygen environment. The diffusion of air into the heterocyst is a compromise between the maximum influx of dinitrogen gas while oxygen is kept sufficiently low to allow nitrogenase activity. This investigation tested the hypothesis that the heterocyst is capable of controlling the influx of air. Here, the thermophilic heterocystous cyanobacterium *Fischerella* sp. was analysed for the effects of oxygen concentration and temperature on nitrogenase activity. Dark nitrogenase activity is directly related to aerobic respiration and was therefore used as a measure of the influx of oxygen into the heterocyst. Above 30% O₂, the influx of oxygen was proportional to its external concentration. Below this concentration, the influx of oxygen was higher than expected from the external concentration. A higher or lower temperature also triggered the heterocyst to increase or decrease, respectively, dark nitrogenase activity while the external concentration of oxygen was kept constant. A higher dark nitrogenase activity requires a higher rate of respiration and therefore a higher flux of oxygen. Hence, the heterocyst of *Fischerella* sp. is capable of controlling the influx of air.

Nitrogenase reduces N₂ to two molecules of ammonium - and simultaneously produces at least one molecule of hydrogen (H₂) - at the expense of protons, electrons that are usually derived from reduced ferredoxin, and 2 ATP per electron¹. Another property that is shared by any nitrogenase is the extreme sensitivity of the enzyme for oxygen. Nitrogenase is irreversibly inactivated by exposure to even low concentrations of O₂². Nitrogenase evolved during the pre-oxygenated state of the earth's atmosphere³. With the appearance of oxygenic photosynthesis and the resulting oxygenation of the atmosphere and large parts of the earth's biosphere, aerobic diazotrophic organisms had to evolve mechanisms to protect nitrogenase from inactivation by oxygen.

Cyanobacteria are oxygenic photoautotrophic microorganisms. Many species are capable of fixing dinitrogen but this seems enigmatic in organisms that not only thrive in aerobic environments but also evolve O₂ as a consequence of their photosynthetic mode of life, even though photosynthesis can satisfy the demand of energy and low-potential electrons for N₂ fixation⁴. Diazotrophic cyanobacteria evolved a variety of strategies in order to provide a micro-oxic environment for nitrogenase and allow for the incompatible processes of oxygenic photosynthesis and N₂ fixation. One of the most elegant solutions is the differentiation of a special cell, the heterocyst, for the fixation of N₂ in certain filamentous cyanobacteria⁵. The heterocyst differentiates from a vegetative cell at semi-regular distances along the trichome through a complex chain of events⁶.

The heterocyst develops a special glycolipid layer which serves as a gas diffusion barrier⁷. It is important to note that this glycolipid layer does not distinguish between O₂ and N₂⁸ and does not serve as a specific O₂ barrier. This means that the glycolipid layer should not stop gas diffusion completely because it would then also prevent N₂ from coming in. Hence, the glycolipid layer of the heterocyst, serving as a gas diffusion barrier, must be a compromise between the maximum influx of N₂ and the capacity to scavenge the O₂ by respiration or otherwise. When the capacity of scavenging O₂ is insufficient, oxygen will accumulate in the heterocyst and nitrogenase will

¹NIOZ Royal Institute for Sea Research, Department of Marine Microbiology and Biogeochemistry and Utrecht University, PO Box 59, 1790 AB, Den Burg, The Netherlands. ²Department of Freshwater and Marine Ecology, Institute of Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, The Netherlands. Correspondence and requests for materials should be addressed to L.J.S. (email: lucas.stal@nioz.nl)

be inactivated. Another property of the heterocyst is that it has lost the oxygenic photosystem II but retained photosystem I. The heterocyst is therefore capable of using light as energy source but depends on the neighbouring vegetative cells for reducing equivalents, which are imported as sucrose^{9–11}. The imported reducing equivalents are partly used for respiratory scavenging of O₂ and partly used to reduce ferredoxin, which serves as the electron donor to nitrogenase¹².

It has long been known that N₂ fixation in the heterocyst can acclimate to environmental O₂ concentrations^{13,14}. It is assumed that this requires an alteration of the glycolipid layer of the heterocyst to become a more or less efficient gas diffusion barrier¹⁵. Indeed, depending on the environment from which they originated, heterocystous cyanobacteria exhibit different nitrogenase activity characteristics that are attributed to the gas diffusion property of the glycolipid layer of the heterocyst^{16,17}. Theoretical considerations have indicated that a heterocystous cyanobacterium would maximize its daily N₂ fixation by increasing its dark nitrogenase activity¹⁷. Although this might seem counter-intuitive it can be understood when considering that it would allow N₂ fixation at any time and any place (not only in the light but also in the non-illuminated parts of the ecosystem and at night). Dark nitrogenase activity is driven by respiration. Hence, in order to maximize N₂ fixation the heterocyst should maximize gas diffusion (to increase the influx of O₂) to the extent of its maximum capacity to scavenge the O₂. While the acclimation of the gas diffusion characteristics of the heterocyst glycolipid layer would be a slow process^{13,14}, a more dynamic process would represent an advantage, especially under fluctuating environmental conditions. Walsby¹⁸ hypothesized that the regulation of the rate of gas exchange into the heterocyst could be accomplished through the opening of the pores that connects the heterocyst with the vegetative cell, analogous to stomata in plants. The aim of this work was to test whether the heterocyst is able to acclimate the influx of gas in order to optimize nitrogen fixation.

Results

The effect of O₂ on nitrogenase activity (acetylene reduction) was tested at a fixed temperature of 30 °C. The total chlorophyll-specific nitrogenase activity (N_{tot}) (the sum of the maximum light-dependent (N_m) and light-independent (N_d) acetylene reduction) remained constant at $88 \pm 4 \mu\text{mol mg}^{-1} \text{h}^{-1}$ in the concentration range of 0–30% O₂ (Fig. 1A). However, the relative contributions of the maximum light-dependent and light-independent fractions, respectively N_m (Fig. 1B) and N_d (Fig. 1C), changed markedly. With increasing O₂ concentrations, N_d increased to its maximum at 30% O₂, while N_m decreased accordingly to compensate N_{tot} to its maximum. Therefore, total nitrogenase activity (N_{tot}) in the concentration range of 0–30% O₂ was at its maximum rate at the given set of conditions. Above 30% O₂, nitrogenase activity (N_m as well as N_d) decreased linearly with O₂ concentration (Fig. 1B,C).

The effect of temperature on nitrogenase activity was investigated at constant O₂ concentration of 20%. *Fischerella* is a thermophilic diazotroph allowing the investigation of a wide temperature range of 12–60 °C. The optimum temperature for nitrogenase activity in *Fischerella* was approximately 40 °C (Fig. 2). This was the case for N_m and N_d and therefore also for the total activity (N_{tot}). At 12 and 60 °C there was low but still detectable nitrogenase activity (see Material and Methods for the control). Examining the N_{tot}/N_d revealed that it remained constant in the temperature range of 21–39 °C at a value of $\sim 2.2 \pm 0.1$ (Fig. 3). N_{tot}/N_d is dimensionless and, hence, a convenient parameter to describe nitrogenase activity in the heterocyst. Below 21 °C N_m became almost zero and the N_{tot}/N_d approached 1. A value of N_{tot}/N_d of 2 means that both light-dependent and light-independent (dark) nitrogenase activity contribute equally to the total. The total nitrogenase activity increased in the temperature range 21–39 °C. Above 39 °C N_m rapidly declined while N_d remained fairly constant up to 50 °C. This resulted in N_{tot}/N_d approaching 1, meaning that nitrogenase activity was largely driven by respiration. Above 50 °C nitrogenase activity collapsed and no meaningful analyses of N_m and N_d could be made.

Discussion

Diazotrophic cyanobacteria are exposed to a suite of environmental factors that affect N₂ fixation. For the fixation of dinitrogen, O₂ concentration and temperature are particularly important and these two environmental parameters are also strongly interrelated. Temperature affects the solubility and diffusion rate of oxygen and also influences the rate of physiological (enzymatic) processes. Nitrogenase is irreversibly inactivated by O₂ and therefore the organism has to provide an environment that is sufficiently low in O₂ in order to be able to fix N₂. This investigation aimed at testing the hypothesis that the heterocyst of *Fischerella* is capable of modulating the influx of gas in order to optimize N₂ fixation. It was previously shown¹⁷ that natural populations of diazotrophic heterocystous cyanobacteria can improve the fixation of N₂ best by increasing nitrogenase activity that occurs independently of light (N_d). It should be noted that increasing dark nitrogenase activity is only possible when a higher influx and respiration of O₂ occurs in the heterocyst. However, this must be precisely controlled so that all O₂ that enters the heterocyst is fully and instantaneously respired in order to prevent the inactivation of nitrogenase. Detection of nitrogenase activity is a proof that this condition is fulfilled. Maximizing the influx of O₂ into the heterocyst also maximizes the influx of N₂, guaranteeing the highest possible fixation of N₂.

Interpretation of the oxygen experiment. In the range of 0–30% O₂ nitrogenase activity was more or less constant at $88 \pm 4 \mu\text{mol ethylene (mg chlorophyll } a)^{-1} \text{h}^{-1}$ (Fig. 1). This must have been the maximum achievable nitrogenase activity under the given set of conditions for the following reasons. Nitrogenase activity in the range of 0–30% O₂ was not limited by energy because N_m would not have decreased with increasing concentration of O₂ and increasing N_d . A shift in the energy source from light to respiration occurred. When energy would have been limiting an increase of N_{tot} would have been expected when O₂ concentration increased from 0 to 30%. Nitrogenase activity in this range of O₂ concentrations was also not limited by reducing equivalents. Otherwise N_{tot} was expected to decrease since increasing amounts of electrons are required for scavenging O₂ when the concentration increased from 0 to 30%. Hence, the heterocyst was capable of providing all the reducing equivalents




Figure 1. Nitrogenase activity (chlorophyll-specific rate of acetylene reduction) at different external O_2 concentrations and a fixed temperature of $30^\circ C$. (A) N_{tot} , the total nitrogenase activity at light saturation; equals $N_m + N_d$. (B) N_m , the maximum light-dependent nitrogenase activity at light saturation. (C) N_d , the light-independent (dark) nitrogenase activity.

for nitrogenase and for respiring O_2 in the range of 0–30% O_2 . The O_2 influx into the heterocyst in this range of O_2 concentrations as derived from the increase of N_d followed a dose-response relationship (Fig. 4). Likewise, the decrease of N_m in the range of 0–30% O_2 could also be described by a dose-response equation. This means that the influx of O_2 in the range of 0–30% was not simply a linear function of the external concentration.

The reason for the decrease of nitrogenase activity above 30% O_2 is a limitation of reducing equivalents that are required for scavenging the increased influx of O_2 . The decrease in nitrogenase activity could not be attributed to an inactivation of the enzyme because controls that lowered the O_2 concentration back to 20% restored the original activity (results not shown). The linear decrease of N_d above 30% O_2 (Fig. 4) indicated that the influx (diffusion) of O_2 was a function of the external O_2 concentration. Assuming that the transport of reducing equivalents from the vegetative cells to the heterocysts was not limiting¹⁹, the linear fit of N_d in the range of 30–90% O_2 gives the O_2 -specific utilization of reducing equivalents. The extrapolation of this linear fit to 0% O_2 gives the theoretical availability of reducing equivalents for nitrogen fixation when respiration does not consume reducing equivalents (see argumentation below). The transport of sucrose into the heterocyst must be sufficient to scavenge O_2 , otherwise we would not see nitrogenase activity. It is therefore not a limiting factor.

The N_d data obtained in the range of 30–90% O_2 were linearly fitted using equation (1)

$$y = ax + b \quad (1)$$

where y = chlorophyll a -specific acetylene reduction ($\mu\text{mol mg}^{-1} \text{h}^{-1}$) and $x = O_2$ (%), and the constants are $a = -0.61 \mu\text{mol mg}^{-1} \text{h}^{-1} \%^{-1}$ and $b = 67.1 \mu\text{mol mg}^{-1} \text{h}^{-1}$.




Figure 2. Nitrogenase activity (chlorophyll-specific rate of acetylene reduction) at different temperatures at a fixed O_2 concentration of 20%. (A) N_{tot} , (B) N_m , (C) N_d , as in Fig. 1 legend.

Hence, at 0% O_2 the theoretical supply of electrons (expressed as the rate of chlorophyll-specific acetylene reduction) would equal $67.1 \mu\text{mol mg}^{-1} \text{h}^{-1}$ (because no electrons would be required to reduce O_2), and at 100% O_2 the theoretical residual rate of chlorophyll-specific C_2H_2 reduction would equal $6.1 \mu\text{mol mg}^{-1} \text{h}^{-1}$. Per % O_2 the electron supply (expressed as the rate of chlorophyll-specific acetylene reduction) required was $0.61 \mu\text{mol mg}^{-1} \text{h}^{-1}$ ($(67.1 - 6.1)/100\%$), assuming only gas diffusion through the cell envelope proportional to the external O_2 concentration.

While the relationship of acetylene reduction and O_2 concentration above 30% was linear and, hence, O_2 diffusion was a function of its external concentration, below 30% O_2 N_d followed a dose-response model; equation (2):

$$y = c + d/(1 + 10^{f(g-x)}) \quad (2)$$

where y and x are as in equation (1). The constants c and d are -174.8 and $226 \mu\text{mol mg}^{-1} \text{h}^{-1}$, respectively. The constants f and g are $0.052\%^{-1}$ and -11.6% , respectively.

While the diffusion through the cell envelope will still be a function of the external O_2 concentration, an additional O_2 influx component must be conceived in order to explain the dose response relationship. This additional component is depicted in Fig. 4 and represents the difference between the extrapolated activity calculated by equation (1) and the observed activity (Fig. 1C). It is highest at 0% O_2 , gradually decreasing to zero at 30% O_2 . The data in Fig. 4 followed a first order exponential decay model; equation (3)




Figure 3. The N_{tot}/N_d ratio at different temperatures. This ratio is dimensionless and describes nitrogenase activity in the heterocyst. The ratio indicates the contribution of photosynthesis to nitrogenase activity. A ratio of 2 means that photosynthesis and respiration contribute equally to nitrogenase activity, while a ratio of 1 means that respiration alone satisfies nitrogenase activity.




Figure 4. The difference between the observed N_d (chlorophyll-specific rate of acetylene reduction) and the values in the range 0–30% O_2 , calculated from the linear extrapolation of the linear function of N_d in the range 30–90% O_2 . The deviation from the linear relationship is largest at 0% O_2 , gradually decreasing to zero at 30% O_2 . The observed curve represents the variable gas flux into the heterocyst.

$$y = h + je^{-x/k} \quad (3)$$

where y and x are as in equation (1). The constants h and j are -8.94 and $69.7 \mu\text{mol mg}^{-1} \text{h}^{-1}$, respectively, and k 14.4%.

N_m behaved in the same but opposite manner as N_d . The data above 30% O_2 were linearly fitted; equation (4):

$$y = mx + n \quad (4)$$

where y and x are as in equation (1). The constants m and n are $-0.5 \mu\text{mol mg}^{-1} \text{h}^{-1} \%^{-1}$ and $48.4 \mu\text{mol mg}^{-1} \text{h}^{-1}$, respectively.

The ratio N_{tot}/N_d decreased with increasing O_2 concentration indicating the increasing contribution of N_d to total nitrogenase activity (Fig. 5). The decrease of N_{tot}/N_d was greatest in the range of 0–30% O_2 . N_m compensated for the decreasing N_d to the maximum nitrogenase activity in the range of 0–30% O_2 . From equation (4) it can be calculated that at 97% O_2 N_m would be zero and N_{tot}/N_d would have reached 1 meaning that respiration would provide all of the energy for nitrogenase activity.

Interpretation of the temperature experiment. The constancy of N_{tot}/N_d at ~ 2.2 in the temperature range 21–39 °C means that N_m and N_d increased in the same proportion. The value of ~ 2.2 for N_{tot}/N_d is regularly observed in heterocystous cyanobacteria and may be optimal¹⁶. It may represent an optimal compromise between the influx of gas into the heterocyst and its capacity to scavenge O_2 . It is important to note that N_d can increase




Figure 5. The N_{tot}/N_d ratio at different external O_2 concentrations. This ratio is dimensionless and describes the characteristics of nitrogenase activity in the heterocyst. The ratio indicates the contribution of photosynthesis to nitrogenase activity. Extrapolation shows that at 97% O_2 this ratio equals 1, meaning that light is no longer required to achieve maximum nitrogenase activity. At low O_2 concentration this ratio becomes very large because nitrogenase activity is fully driven by photosynthesis and N_d approaches zero.

only when more O_2 enters the heterocyst, allowing more respiratory energy generation. While the external concentration of O_2 was held constant, the only possibility was that the heterocyst increased the influx of O_2 . The consequent draw on reducing equivalents for respiration resulted in a lower total nitrogenase activity, realized by a markedly lower N_m .

In the temperature range 21–39 °C the Q_{10} could be calculated. The Q_{10} gives the increase of the rate of a process for a 10 °C rise in temperature. For biochemical processes the Q_{10} equals approximately 2, meaning a doubling for each 10 °C rise in temperature (or halving at each 10 °C decrease in temperature). For the total nitrogenase activity (N_{tot}) as well as for N_m a value of 2.3 was calculated and 2.2 for N_d , hence Q_{10} was well within the expected range for a biochemical process. However, if N_d would only depend on the rate of diffusion of O_2 into the heterocyst the Q_{10} is expected to be only marginally above 1 (1.084), that results from the temperature effects on O_2 solubility and diffusion constant¹⁷. The Q_{10} of 2.2 for N_d means that respiration was not limited by O_2 diffusion or availability and it reflects the temperature (kinetic) effect on the enzyme nitrogenase. In order to keep the heterocyst micro-oxic the influx of O_2 must have been precisely controlled.

It is important to note that the heterocyst glycolipid layer forms a gas diffusion barrier that does not discriminate between O_2 and N_2 ⁸. While on the one hand the heterocyst glycolipid layer will limit the rate of O_2 influx, it must also allow sufficient influx of N_2 for N_2 fixation; on the other hand, influx of O_2 is essential for generating energy through respiratory electron transport when light is unavailable and hence extends the period during which the organism is able to fix N_2 . It is also important to understand that by measuring nitrogenase activity one looks specifically at the processes in the heterocyst, rather than in the vegetative cells. For instance, by measuring the light-independent part of nitrogenase activity, this can be translated to respiration. In the dark, respiration is the only known source of energy for nitrogenase in the heterocyst, and, hence, dark nitrogenase activity can be interpreted as the influx of O_2 into the heterocyst. When nitrogenase activity is detected one must conclude that the heterocyst is low in O_2 . The residual O_2 concentration in the heterocyst can be as low as 0.6 μ M, which is 0.2% of air saturation¹⁸ and depends on the $K_m(O_2)$ of the O_2 -scavenging process.

In this investigation, I demonstrate in two ways that *Fischerella* is capable of adjusting the influx of air into the heterocyst. First, the effect of the external O_2 concentration on nitrogenase activity was measured at a constant temperature. Second, the effect of temperature on nitrogenase activity was measured at a constant external O_2 concentration. Temperature affects the rate of metabolic processes, including respiration, but has a small effect on the rate of gas influx by diffusion. The gas influx is the combination of O_2 solubility and diffusion. Whereas diffusion increases with temperature, solubility decreases and the two factors almost compensate each other ($Q_{10} = 1.084$)¹⁷.

I observed that the light independent part of nitrogenase activity increased with temperature in the range from 12–42 °C. The Q_{10} in the range of 21–39 °C was 2.2. N_d can only increase when respiratory energy generation increases and this is only possible when the O_2 influx increases. Since these experiments were done under a constant O_2 concentration in the gas mixture, it was concluded that the heterocyst possesses a mechanism to increase the influx of O_2 . It is unlikely that the heterocyst glycolipid layer possesses the capacity to alter its gas diffusion properties in the short term of the experiment. One possibility that was proposed by Walsby¹⁸ is that the heterocyst terminal pores might perform this function (analogous to stomata in plant leaves) (Fig. 6). Canini *et al.*²⁰ proposed that cyanophycin serves as a plug that closes the neck of the heterocyst connection to the vegetative cell. However, until now hardly anything is known about the physical properties of this insoluble polymer and it is therefore unknown whether this material is gas tight. Nevertheless, these are attractive concepts waiting for experimental proof. Cytoplasmic connections have also been postulated as the site of transport between cells of heterocystous cyanobacteria^{11, 19}. The exchange of solutes (such as sucrose) between the vegetative cell and the




Figure 6. Model of a heterocyst with junctions to a vegetative cell at each side. The model emphasizes the situation for N_d (see Fig. 1 legend). The heterocyst is characterized by an envelope comprising a glycolipid layer, which forms the principal barrier to diffusion of gas (thin arrows), and a polysaccharide layer. The envelope surrounds a cell wall with an outer membrane, peptidoglycan layer, and plasma membrane. Whether or not air enters the heterocysts via the vegetative cells may be regulated by an open/close mechanism (conceptually depicted by orange arrows). O_2 is respired, producing ATP. N_2 and O_2 compete for electrons that are imported as sucrose from the vegetative cells. The pores at the ends of the heterocysts may sometimes contain cyanophycin (polyaspartate-multiarginine polymer). Nase: nitrogenase; CytOx: cytochrome oxidase; [H]: reducing equivalents (NAD(P)H, FdH) derived from the metabolism of sucrose.

heterocyst is through the septal junctions (previously named microplasmodesmata) by molecular diffusion²¹. However, although gas molecules could diffuse together with the solutes, they may additionally also diffuse through the cell wall and cytoplasmic membrane which are impermeable for larger molecules such as sucrose.

The Q_{10} is in the physiological range which means that it is related to the enzyme activity solely and that N_d is not limited (by energy or reducing equivalents) in the temperature range of 21–39°C. Unless nitrogenase was replaced continuously and instantaneously, it was also unlikely that nitrogenase was inactivated by O_2 , because control experiments during which the cells were brought back to the initial conditions reproduced the original nitrogenase activity. This would require that the gas influx is tightly controlled to ensure that all O_2 entering the heterocyst is respired. Another remarkable observation was that N_d was less sensitive to temperature than was the light-dependent part of nitrogenase activity, N_m . This could indicate that the photosystem is more thermosensitive than the respiratory system. Photosystem II is known as the most thermolabile component that determines the upper thermal tolerance of photosynthesis²². It has been shown that the D1 protein degrades faster at high temperature and is not re-synthesized²³. It is not known whether a similar process affects photosystem I in the heterocyst and makes it more thermosensitive than the respiratory system²⁴.

The experiments with different O_2 concentrations at a constant temperature confirmed the presence of a control mechanism that determines the influx of gas into the heterocyst. Only above 30% O_2 the supposed influx was a linear function of the external concentration and determined by the diffusion characteristics of the heterocyst envelope. Below 30% O_2 a dose-response relationship was found. The lower the external concentration of O_2 the higher the relative influx became. The gas diffusion through the heterocyst envelope would be an intrinsic property of the envelope and would be a linear function of the external O_2 concentration at any O_2 level. Subtracting this envelope gas diffusion from the N_d below 30% O_2 , I found an exponential decay of the additional (in excess of the envelope gas diffusion) chlorophyll-specific dark nitrogenase activity, hence O_2 influx (measured as acetylene reduction), from $\sim 60 \mu\text{mol mg}^{-1} \text{h}^{-1}$ at 0% O_2 to $\sim 0 \mu\text{mol mg}^{-1} \text{h}^{-1}$ at 30% O_2 (Fig. 4). Hence, when the terminal pores would serve as a variable gas inlet for the heterocyst they would be ‘open’ at 0% O_2 and ‘closed’ at 30% O_2 (Fig. 6).

The lower N_{tot} with increasing external O_2 concentration above 30% O_2 is explained by an increasing demand of reducing equivalents for respiration that are consequently unavailable for the reduction of N_2 . Extrapolation of the data indicated that only at 97% O_2 N_{tot}/N_d becomes 1, meaning that respiration would theoretically supply all energy for nitrogenase activity. Hence, light remains an important energy source, also perhaps because it generates the electron donor of nitrogenase, reduced ferredoxin, through photosystem I-mediated electron transport¹².

The measurements reported here were carried out by the acetylene reduction assay. The reduction of acetylene to ethylene is an accepted measure of nitrogenase activity. It should be noted that the reduction of acetylene cannot be converted quantitatively directly into fixation of N_2 . Using the electron stoichiometry, the ratio acetylene reduction to N_2 fixation is 4:1. If one supposes that the H_2 produced during N_2 fixation is recycled by a hydrogenase, the ratio would be 3:1. However, this would only make a difference when reducing equivalents limit N_2 fixation, which was not the case. Also, the concentration of 10% acetylene does not saturate nitrogenase and the saturation of nitrogenase activity by acetylene *in vivo* also depends on the light intensity²⁵. At 10% acetylene, light intensity has only a marginal effect on the saturation of nitrogenase by acetylene and therefore the rates of acetylene reduction in the light response curves do not have to be corrected for this effect. I have not taken into account the solubility and diffusion coefficients of acetylene and ethylene or of N_2 . The diffusion coefficients of

acetylene and ethylene may be different from N_2 (and thus from O_2). As proposed here, the influx of a gas into the heterocyst through the pores would be equally controlled for acetylene and ethylene as it would be for air. Hence, it is unlikely that the use of the acetylene reduction assay instead of directly measuring N_2 fixation would have led to different conclusions.

In conclusion, although it is known that heterocystous cyanobacteria can modify their heterocyst glycolipid layer in response to external O_2 concentration, such acclimation cannot be instantaneous and also depends on the properties of this glycolipid layer which should be genetically determined. For instance, it has been observed that heterocystous cyanobacteria that thrive in colder waters possess a heterocyst envelope with a stronger gas diffusion barrier than those that occur in warmer environments where respiration is faster. The results of this investigation support the idea of Walsby¹⁸ that heterocysts can also respond instantaneously to O_2 and temperature as depicted in Fig. 6. It is likely that this is a general property of heterocystous cyanobacteria that contribute to their ecological success.

Methods

Organism and growth conditions. *Fischerella* sp. strain MV11 (Culture Collection Yerseke; CCY0604) was obtained from Prof. dr. W.R. Hess, University of Freiburg, Germany, and has been described in detail²⁶. This is a true-branching, thermophilic, N_2 -fixing, heterocystous cyanobacterium belonging to the Stigonematales (section V) and was originally isolated from a geothermal site at a hot spring in Costa Rica. The organism was routinely grown at 27 °C in BG11_o medium (devoid of any combined nitrogen)²⁷ under a 12:12 hour light-dark cycle and 30 $\mu\text{mol m}^{-2} \text{s}^{-1}$ warm-white fluorescent light.

On-line acetylene reduction assay. Nitrogenase activity was monitored using the acetylene reduction assay²⁸. Here, I used the on-line set-up as described previously^{25,29}. The set-up allows a highly sensitive measurement of ethylene (0.3 ppbv, parts per billion by volume) at a time resolution of 3–5 seconds and therefore provides almost real-time monitoring of nitrogenase activity. The sensitive and rapid monitoring of ethylene was done using a laser photoacoustic trace gas detector (ETD-300, Sensor Sense, Nijmegen, The Netherlands). A low amount (0.4–0.8 μg chlorophyll *a*) of *Fischerella* culture was gravity filtered on a Whatman GF/F glass fibre filter (47 mm). This low amount of culture prevented self-shading and also excessive ethylene production (which would result in an ‘overload’ of the ETD-300). The filter containing *Fischerella* was put in a custom-made measuring cell²⁹. The cell was equipped with a Peltier temperature regulation unit (Supercool DA-075-24-02-00-00, Göteborg, Sweden). The filter was placed on a supporting grid underneath which growth medium was present that ensured that the filter was saturated with medium and that prevented desiccation. The incubation chamber was connected to a continuous flow of gas at 2 L h^{-1} that was led over the filter with *Fischerella*. The gas was a mixture of N_2 , O_2 , CO_2 and C_2H_2 , made up by mass flow controllers (Brooks Instruments, Ede, The Netherlands). Acetylene was kept at 10%. CO_2 was premixed at 0.04% in the N_2 and O_2 . The concentration of O_2 was varied as required and the balance was made up with N_2 . O_2 concentrations ranged from 0–90% (the highest possible concentration since 10% C_2H_2 was always present). The gas was washed through water prior to entering the measuring cell. This served to humidify the gas and removed any acetone from the acetylene (which would interfere with the ethylene detection). Gas mixtures (O_2 and N_2 with 0.04% CO_2) were purchased from Hoek-Loos (The Netherlands) and acetylene was obtained from Messer (The Netherlands). The gas mixture contained a very low background level of ethylene of ~10 ppbv due to contamination of the acetylene. Controls were run without *Fischerella* in the incubation chamber. Ethylene recordings of >100 ppbv were taken as positive for ethylene production. The whole set-up was placed in a temperature-controlled climate room. Illumination was provided by a cold-light 250 W halogen lamp and glass fibre light shower (model 460-F, Heinz Walz GmbH, Effeltrich, Germany). A computer program written in Testpoint (Capital Equipment Corporation, New Hampshire, USA) controlled the gas mixtures, light intensity and temperature.

Light-response curves. At any given temperature and concentration of O_2 the response of nitrogenase activity to light was measured as follows. Ten different photon irradiances were tested: 0, 1.4, 4, 7.1, 10.9, 15.5, 20.5, 26.5, 43.5 and 68.2 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Nitrogenase activity was monitored for 5 minutes at each light level. The measurement started with 15 minutes in the dark, followed by the series of light levels until the highest level tested which saturated nitrogenase activity at all conditions tested (light up), after which the same measurements were made while light levels decreased (light down). This was done to observe any hysteresis effects and to check that the original measurements could be reproduced. The system is robust and measurements were reproducible with high accuracy. A full light response measurement (light up and light down) took 105 minutes after which the computer adjusted the measuring cell to the next condition. The whole experiment ran unattended.

Temperature experiments. The effect of temperature was measured in two series of experiments. In one series of experiments the temperature was decreased from 27 °C (growth temperature) in steps of 3 °C to 12 °C and in the second experiment the temperature was increased from 27 °C to 60 °C, also in steps of 3 °C. Both experiments were carried out immediately one after the other, using cells from the same stock culture. In the first experiment (decreasing temperature) the filter contained 0.4 μg chlorophyll *a*, and in the second experiment (increasing temperature) it contained 0.7 μg chlorophyll *a*. The nitrogenase parameters at 27 °C from both experiments were similar and hence the data of the two experiments overlapped. In test runs it was demonstrated that cells that were lowered to 18 °C and subsequently brought back to 27 °C, reproduced the original nitrogenase activity and the same was true for cells that were increased to 54 °C and brought back to 27 °C. Below 18 °C (incubations at 15 °C and 12 °C) and above 54 °C (incubations at 57 °C and 60 °C) the cultures were to some extent negatively affected and also possessed low nitrogenase activity.

Oxygen experiments. The effect of O₂ was measured in one series of measurements from 0–90% O₂. The filter contained 0.8 µg chlorophyll *a*. The following O₂ concentrations were tested and in this order: 0, 2.5, 5, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, and 90%. The experiment was run at 30 °C. Before starting the experiment the culture was run at 0% O₂, 20.5 µmol m⁻² s⁻¹ light and 30 °C for 30 minutes by which time a constant rate of nitrogenase activity was achieved. The system did not achieve a fully anoxic state. Previously, it was estimated that the setting 0% O₂ may in fact represent a level of ~1% O₂³⁰. This is in part because of leakage through the tubing used and the large volume of water (1 L) needed for washing the gas, as well as due to photosynthetic O₂ production (in the light). Fully anoxic conditions in the dark would not have allowed any activity (since there would not be a possibility to generate energy). At each O₂ concentration the light response of nitrogenase activity was recorded as described above (light up, light down) at the same 10 light levels as mentioned above for the recording of light-response curves. Test runs were performed before carrying out the final experiment reported here.

Fitting the light response curves. The light response curves were fitted with the rectangular hyperbola model³¹; equation (5) using the software package Origin. The fitted parameters derived from this model are N_d , N_m , and α , respectively the light-independent part of the nitrogenase activity (dark activity, which is always present), the maximum light-dependent nitrogenase activity at light saturation, and the slope of the curve. From these fitted parameters, the total maximum nitrogenase activity, N_{tot} , was calculated as the sum of N_d and N_m . N_{tot}/N_d is a convenient parameter that is dimensionless and therefore biomass independent (and, hence, refers only to the cells that possess nitrogenase activity, in this case the heterocysts) and describes the contribution of light to nitrogenase activity. $N_{tot}/N_d \geq 1$; when this ratio equals 1 no light-dependent nitrogenase activity is present. The light saturation coefficient I_k equals N_m/α , and represents the photon irradiance at which 0.5 N_m would have been reached when nitrogenase activity increases linearly with irradiance (K_s in Michaelis-Menten kinetics).

$$N_I = N_m(\alpha I / (N_m + \alpha I)) + N_d \quad (5)$$

where N_I is the actual nitrogenase activity at light level I .

Temperature coefficient. The temperature coefficient of nitrogenase activity (Q_{10}) was fitted from the linear fit of the natural log transformed data of N_d , N_m , and N_{tot} with temperature³² using the software package Origin. Meaningful values (*i.e.* linear fits) of Q_{10} were obtained in the temperature range 18–39 °C. The temperature coefficient of O₂ diffusion in the temperature range 18–39 °C was calculated from O₂ solubility and diffusion coefficients that can be found in the tables provided by the company Unisense and can be downloaded from: <http://www.unisense.com/files/PDF/Diverse/Seawater%20&%20Gases%20table.pdf>.

Chlorophyll. Chlorophyll *a* was extracted from the GF/F filters in 2 ml 96% ethanol overnight at room temperature in the dark, immediately after the experiment was finished. The absorption coefficient of 72.3 ml mg⁻¹ cm⁻¹ was used to calculate the concentration of chlorophyll *a*.

References

- Burgess, B. K. & Lowe, D. J. Mechanism of molybdenum nitrogenase. *Chem. Rev.* **96**, 2983–3011 (1996).
- Gallon, J. R. Reconciling the incompatible: N₂ fixation and O₂. *New Phytol* **122**, 571–609 (1992).
- Raymond, J., Siefert, J. L., Staples, C. R. & Blankenship, R. E. The natural history of nitrogen fixation. *Mol. Biol. Evol.* **21**, 541–554 (2004).
- Bergman, B., Gallon, J. R., Rai, A. N. & Stal, L. J. N₂ fixation by non-heterocystous cyanobacteria. *FEMS Microbiol. Rev.* **19**, 139–185 (1997).
- Adams, D. G. Heterocyst formation in cyanobacteria. *Curr. Opin. Microbiol.* **3**, 618–624 (2000).
- Zhang, C.-C., Laurent, S., Sakr, S., Peng, L. & Bédou, S. Heterocyst differentiation and pattern formation in cyanobacteria: a chorus of signals. *Mol. Microbiol.* **59**, 367–375 (2006).
- Awai, K., Lechno-Yossef, S. & Wolk, C. P. Heterocyst envelope glycolipids. (eds Wada, H. & Murata, N.) *Lipids In Photosynthesis: Essential And Regulatory Functions*. 179–202. (Springer Verlag, New York, 2009).
- Walsby, A. E. The permeability of heterocysts to the gases nitrogen and oxygen. *Proc. Royal Soc. Lond. B* **226**, 345–366 (1985).
- Curatti, L., Flores, E. & Salerno, G. Sucrose is involved in the diazotrophic metabolism of the heterocyst-forming cyanobacterium *Anabaena* sp. *FEBS Lett.* **513**, 175–178 (2002).
- López-Igual, R., Flores, E. & Herrero, A. Inactivation of a heterocyst-specific invertase indicates a principal role of sucrose catabolism in heterocysts of *Anabaena* sp. *J. Bacteriol.* **192**, 5526–5533 (2010).
- Nürnberg, D. J. *et al.* Intercellular diffusion of a fluorescent sucrose analog via the septal junctions in a filamentous cyanobacterium. *MBio* **6**, 02109–14 (2015).
- Razquin, P. *et al.* Differential activities of heterocyst ferredoxin, vegetative cell ferredoxin, and flavodoxin as electron carriers in nitrogen fixation and photosynthesis in *Anabaena* sp. *Photosynth. Res.* **43**, 35–40 (1995).
- Murry, M. A., Horne, A. J. & Benemann, J. R. Physiological studies of oxygen protection mechanisms in the heterocysts of *Anabaena cylindrica*. *Appl. Environ. Microbiol.* **47**, 449–454 (1984).
- Kangatharalingam, N., Prisco, J. C. & Paerl, H. W. Heterocyst envelope thickness, heterocyst frequency and nitrogenase activity in *Anabaena flos-aquae*: influence of exogenous oxygen tension. *J. Gen. Microbiol.* **138**, 2673–2678 (1992).
- Bauersachs, T. *et al.* Distribution of heterocyst glycolipids in cyanobacteria. *Phytochem.* **70**, 2034–2039 (2009).
- Staal, M., Meysman, F. J. R. & Stal, L. J. Temperature excludes N₂-fixing heterocystous cyanobacteria in the tropical oceans. *Nature* **425**, 504–507 (2003).
- Stal, L. J. Is the distribution of nitrogen-fixing cyanobacteria in the oceans related to temperature? *Environ. Microbiol.* **11**, 1632–1645 (2009).
- Walsby, A. E. Cyanobacterial heterocysts: terminal pores proposed as sites of gas exchange. *Trends Microbiol.* **15**, 340–349 (2007).
- Mullineaux, C. W. *et al.* Mechanism of intercellular molecular exchange in heterocyst-forming cyanobacteria. *EMBO J.* **27**, 1299–1308 (2008).
- Canini, A., Galiazzo, F. & Grilli Caiola, M. Superoxide dismutase in the symbiont *Anabaena azollae* Strasb. *Plant Physiol.* **97**, 34–40 (1991).
- Flores, E., Herrero, A., Forchhammer, K. & Maldener, I. Septal junctions in filamentous heterocyst-forming cyanobacteria. *Trends Microbiol.* **24**, 79–82 (2016).

22. Davison, I. R. Environmental effects on algal photosynthesis: temperature. *J. Phycol.* **27**, 2–8 (1991).
23. Singh, M., Satoh, K., Yamamoto, Y., Kanervo, E. & Aro, E.-M. *In vivo* quality control of photosystem II in cyanobacteria *Synechocystis* sp. PCC 6803: D1 protein degradation and repair under the influence of light, heat and darkness. *Ind. J. Biochem. Biophys.* **45**, 237–243 (2008).
24. Balme, A. *et al.* A comparative study of the thermal stability of plastocyanin, cytochrome c_6 and photosystem I in thermophilic and mesophilic cyanobacteria. *Photosynth. Res.* **70**, 281–289 (2001).
25. Zuckermann, H. *et al.* On-line monitoring of nitrogenase activity in cyanobacteria by sensitive laser photoacoustic detection of ethylene. *Appl. Environ. Microbiol.* **63**, 4243–4251 (1997).
26. Finsinger, K. *et al.* Characterization of true-branching cyanobacteria from geothermal sites and hot springs of Costa Rica. *Environ. Microbiol.* **10**, 460–473 (2008).
27. Rippka, R., Deruelles, J., Waterbury, J. B., Herdman, M. & Stanier, R. Y. Generic assignments, strain histories and properties of pure cultures of cyanobacteria. *J. Gen. Microbiol.* **111**, 1–61 (1979).
28. Hardy, R. W. F., Holsten, R. D., Jackson, E. K. & Burns, R. C. The acetylene-ethylene assay for N_2 fixation: laboratory and field evaluation. *Plant Physiol.* **43**, 1185–1207 (1968).
29. Staal, M. te Lintel Hekkert, S., Harren, F. & Stal, L. Nitrogenase activity in cyanobacteria measured by the acetylene reduction assay: a comparison between batch incubation and on-line monitoring. *Environ. Microbiol.* **3**, 343–351 (2001).
30. Staal, M., Rabouille, S. & Stal, L. J. On the role of oxygen for nitrogen fixation in the marine cyanobacterium *Trichodesmium* sp. *Environ. Microbiol.* **9**, 727–736 (2007).
31. Staal, M. te Lintel Hekkert, S., Herman, P. & Stal, L. J. Comparison of models describing light dependence of N_2 fixation in heterocystous cyanobacteria. *Appl. Environ. Microbiol.* **68**, 4679–4683 (2002).
32. Hopkins, D. W. *et al.* Carbon, nitrogen and temperature controls on microbial activity in soils from an Antarctic dry valley. *Soil Biol. Biochem.* **38**, 3130–3140 (2006).

Acknowledgements

I like to thank Jan Peene for writing the program in Testpoint and Lowie Haazen for constructing the temperature controlled measuring cell.

Author Contributions

The author designed and carried out the experiment, interpreted the results and wrote the manuscript.

Additional Information

Competing Interests: The author declares that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017