

**Conclusion.** No local or disseminated HSV infections were encountered in the study cohort. Bacterial skin and soft-tissue infection were uncommon. Most other infections were unrelated to TVEC therapy. Real-world review of the use of an HSV-derived oncolytic viral vector therapy mimics reported infectious complications from clinical trials.

**Disclosures.** All authors: No reported disclosures.

### 2299. A London Hospital's Experience of Confirmed Measles Infections and Re-Infections Between 2009 and 2019

Ann Sturdy, MBBS<sup>1</sup>; Tanzina Haque, MBBS<sup>1</sup>; Paul Griffiths, MBBS<sup>1</sup>; Kevin Brown, MBBS<sup>2</sup>; Dianne Irish, MBBS<sup>1</sup>; Tabitha Mahungu, BM BS<sup>1</sup>; <sup>1</sup>Royal Free London NHS Trust, London, UK; <sup>2</sup>Public Health England, London, UK

**Session:** 247. Clinical Virology/Viral Epidemiology

*Saturday, October 5, 2019: 12:15 PM*

**Background.** 2018 to 2019 has seen a global resurgence in measles cases, with the UK seeing an over 3-fold increase in cases in 2018 compared with 2017. In this context, our center saw a cluster of geographically linked measles cases presenting in the first quarter of 2019. We present this with comparative data of confirmed measles cases from our center over the preceding 10 years. Given the growing recognition of measles "re-infection" in fully vaccinated individuals, we also present our first confirmed cases of reinfection within this cohort.

**Methods.** Retrospective analysis of confirmed measles cases (positive Measles IgM or detectable Measles RNA) between 2009 and 2019. Laboratory and demographic data were obtained from electronic patient records.

**Results.** 18 cases (of which 14 were adults) of measles (all genotype D8 of those tested) were confirmed in the first 4 months of 2019. 12 presented within a 14 day period from a geographically linked part of North London. There were 4 confirmed measles re-infections (detectable measles RNA on a buccal swab with either a high Measles IgG avidity or previous documented measles immunity). From the 10 year data, cases peaked in 2011 and 2016 consistent with national trends. Of the 89 cases identified, 60 (67%) were adults, who were over twice as likely to be admitted as children, had a longer median length of stay in hospital (2 days vs. 1.5 days) and were more likely to develop a hepatitis (1/10 pediatric cases vs. 26/48 adults,  $p \leq 0.01$ ) or other complications. The majority of adults (68%) were unsure of their vaccination status.

**Conclusion.** 4 cases of 18 in the first 4 months of 2019 were confirmed re-infections. Re-infections in fully vaccinated individuals are described in the literature, typically presenting with a milder course; however, these are the first cases we have identified at our center. Overall, adult cases were more likely to be admitted to hospital and to have a complicated course compared with children. Vaccination history in adults was of limited clinical utility due to lack of reliable documentation and the potential for reinfection.

**Disclosures.** All authors: No reported disclosures.

### 2300. Incidence, Complications, and Recurrence of Herpes Zoster in Unvaccinated Adults ≥50 Years of Age

Bradley Ackerson, MD<sup>1</sup>; Katia Bruxvoort, PhD, MPH<sup>2</sup>; Lina S. Sy, MPH<sup>2</sup>; Yi Luo, PhD<sup>2</sup>; Hilary Tanenbaum, PhD, MS<sup>2</sup>; Yun Tian, MS<sup>2</sup>; Chengyi Zheng, PhD, MS<sup>2</sup>; Bianca P. Cheung, MS<sup>2</sup>; Brandon J. Patterson, PharmD, PhD<sup>3</sup>; Desiree Van Oorschot, MSc<sup>4</sup>; Hung Fu Tseng, PhD, MPH<sup>2</sup>; <sup>1</sup>Kaiser Permanente, South Bay Medical Center, Harbor City, California; <sup>2</sup>Kaiser Permanente Southern California, Pasadena, California; <sup>3</sup>GSK, Philadelphia, Pennsylvania; <sup>4</sup>GSK Vaccines, Wavre, Vlaams-Brabant, Belgium

**Session:** 247. Clinical Virology/Viral Epidemiology

*Saturday, October 5, 2019: 12:15 PM*

**Background.** More recent baseline epidemiological data for Herpes Zoster (HZ) in adults ≥ 50 years of age, obtained before the introduction of the adjuvanted Recombinant Zoster Vaccine (RZV), are needed for future evaluations of the impact of RZV on HZ epidemiology.

**Methods.** The study comprised five elements: (1) The incidence of HZ was estimated from immunocompetent adults ≥ 50 years of age not vaccinated with Zoster Vaccine Live who had incident HZ between 2011–2015. HZ was identified by International Classification of Diseases (ICD) codes from electronic health records (EHR) of 4.6 million Kaiser Permanente Southern California members; (2) Postherpetic neuralgia (PHN) was identified by validated survey and medical record review of laboratory-confirmed incident HZ cases recruited during 2012–2015 for HZ-related pain ≥ 90 days after initial HZ diagnosis; (3) HZ Ophthalmicus (HZO) with ocular complications was identified by ICD codes and keyword search in EHR among patients identified with HZO using a validated natural language processing algorithm; (4) The proportion of HZ-related non-PHN and non-HZO cutaneous, neurological or other complications was assessed by double abstraction of EHRs from a sample of 600 incident HZ cases; (5) Recurrent HZ was identified by having an HZ diagnosis with HZ antiviral medication ≥ 6 months after the most recent HZ diagnosis with HZ antiviral medication in a cohort initially diagnosed with HZ between 2007 and 2008 and followed through 2016.

**Results.** We identified 40,893 incident HZ cases with an overall incidence of 9.92 (95% confidence interval [CI]: 9.82–10.01) per 1000 person-years. The proportion of incident HZ cases with PHN and HZO with ocular involvement was 18.37% (95% CI: 14.90–21.84%) and 8.06% (95% CI: 7.80–8.32%), respectively. The proportion of cutaneous, neurological, and other complications was 7.20% (95% CI: 5.44–8.96%), 0.87%

(95% CI: 0.79–0.95%), and 1.24% (95% CI: 1.15–1.33%), respectively. The incidence of recurrent HZ was 10.96/1000 person-years (95% CI: 10.18–11.79).

**Conclusion.** HZ is common among unvaccinated US adults ≥ 50 years of age, with PHN and HZO occurring most frequently among incident HZ cases.

**Funding:** GlaxoSmithKline Biologicals SA, GSK study identifier: HO-17-18378.

**Disclosures.** All authors: No reported disclosures.

### 2301. Increased Risk of Varicella-Associated Hospitalizations Among Adult Immigrants From Temperate and Tropical Countries After the Introduction of a Childhood Varicella Vaccination Program in Quebec, Canada

Jordan Mah, MD<sup>1</sup>; Anthony Lieu, MD<sup>2</sup>; Zoë R. Greenwald, MSc<sup>3</sup>; Arash Akaberi, MSc<sup>3</sup>; Sunny Song, MSc<sup>3</sup>; Laurent Azoulay, PhD<sup>3</sup>; Marc Brisson, PhD<sup>4</sup>; Caroline Quach, MD<sup>5</sup>; Christina Greenaway, MD, MSc<sup>6</sup>; <sup>1</sup>Faculty of Medicine, McGill University, Montreal, QC, Canada; <sup>2</sup>McGill University, Montreal, QC, Canada; <sup>3</sup>Centre for Clinical Epidemiology, Lady Davis Institute for Medical Research, Montreal, QC, Canada; <sup>4</sup>Research Group in Mathematical Modeling and Health Economics of Infectious Disease, Université Laval, Quebec, QC, Canada; <sup>5</sup>Department of Microbiology and Immunology, CHU Sainte-Justine, Montreal, QC, Canada; <sup>6</sup>Jewish General Hospital, McGill University, Montreal, QC, Canada

**Session:** 247. Clinical Virology/Viral Epidemiology

*Saturday, October 5, 2019: 12:15 PM*

**Background.** Varicella zoster virus (VZV) hospitalizations are an uncommon, severe and costly consequence of VZV. Childhood VZV vaccination leads to decreased VZV rates across all age groups through herd immunity but increases the age of VZV acquisition and the potential risk of severe VZV in non-immune adults. A large proportion (~15%) of young adult immigrants from tropical regions are susceptible to VZV due to different transmission dynamics in their countries of origin and lack of vaccination. We aimed to describe the impact of the childhood VZV program introduced in 2006 in Quebec on VZV hospitalizations in immigrants and nonimmigrants.

**Methods.** A population-based cohort of all medically-attended VZV cases in Quebec, Canada (1996–2014) were identified in administrative health databases and linked to immigration data. VZV-attributable hospitalizations included those with primary or secondary ICD-9 or ICD-10 codes for VZV. Overall age-standardized and age-specific rates of hospitalizations were calculated during pre- (1996–98), private (1999–2005) and public vaccination (2006–14) periods and by immigrant status and pregnancy. Relative risk (RR<sub>i,NI</sub>) and 95% CI for immigrants vs. nonimmigrants were estimated.

**Results.** 5873 hospitalizations occurred among 230,052 VZV cases. Hospitalization rates decreased dramatically in the pre to public vaccination period (6.6 to 1.3/100,000 population); however, the proportion of hospitalized varicella cases increased from 1.7% to 3.9% ( $P < 0.01$ ). Immigrants only accounted for 3.6% of hospitalizations ( $N = 213$ ) however, the proportion of all hospitalizations among immigrants increased in the pre- vs. public-vaccination periods in those aged 10–19 years (2.9% to 13.7%) and 20–39 years (8.8% to 22.7%). The RR was higher in these age groups in the public vaccination period [RR<sub>i,NI</sub> 1.96 and RR<sub>i,NI</sub> 1.67] (Table 1). Adults (>20 years) accounted for 52% (CI: 45–59%) and pregnant women 18% (13–25%) of all hospitalizations among immigrants compared with only 14% (13–15%) and 1.6% (1.3–2.0%) in nonimmigrants, respectively.

**Conclusion.** Young adult and pregnant immigrants bore a disproportionate burden of VZV hospitalizations after the introduction of childhood VZV vaccination. Susceptible immigrant adults would benefit from targeted VZV vaccination.

**Table 1: Overall age-standardized and age-specific rates of varicella attributable hospitalizations and proportion of cases by immigrant status in Quebec, Canada (1996-2014)**

Age-group	Vaccination Period	Immigrants			Non-Immigrants			Immigrants vs non-Immigrants Rate Ratio <sub>95%CI</sub>
		Cases N	% cases [row] (95% CI)	Rate per 100,000	Cases	% cases [row] (95% CI)	Rate per 100,000	
0-9 years	1996-1998	16	1.2 (0.6-1.8)	23.0 (14.1 - 37.6)	1322	98.8 (98.2-99.4)	50.7 (48.0 - 53.5)	0.45 (0.28 - 0.74)
	1999-2005	49	1.8 (1.3-2.3)	26.2 (19.8 - 34.7)	2665	98.2 (97.7-98.7)	48.0 (46.2 - 49.8)	0.55 (0.41 - 0.72)
	2006-2014	20	2.9 (1.6-4.1)	6.1 (4.0 - 9.5)	678	97.1 (95.9-98.4)	9.6 (8.9 - 10.3)	0.64 (0.41 - 1.00)
10-19 years	1996-1998	2	2.9 (0.7-10)	1.3 (0.3 - 5.3)	66	97.1 (93.0-100.0)	2.5 (1.9 - 3.1)	0.53 (0.13 - 2.17)
	1999-2005	8	9.6 (3.3-16.0)	2.1 (1.0 - 4.2)	75	90.4 (84.0-96.7)	1.2 (1.0 - 1.5)	1.70 (0.82 - 3.53)
	2006-2014	7	13.7 (4.3-23.2)	1.1 (0.5 - 2.4)	44	86.3 (76.8-95.7)	0.6 (0.4 - 0.8)	1.96 (0.88 - 4.34)
20-39 years	1996-1998	18	8.8 (4.9-12.7)	2.8 (1.8 - 4.5)	187	91.2 (87.3-95.1)	3.3 (2.8 - 3.8)	0.87 (0.54 - 1.42)
	1999-2005	44	13.3 (9.7-17.0)	2.8 (2.1 - 3.7)	286	86.7 (83.0-90.3)	2.3 (2.1 - 2.6)	1.18 (0.86 - 1.62)
	2006-2014	20	22.7 (14.0-31.5)	0.8 (0.5 - 1.2)	68	77.3 (68.5-86.0)	0.5 (0.4 - 0.6)	1.67 (1.02 - 2.76)
40+ years	1996-1998	5	9.3 (1.5-17.0)	0.4 (0.2 - 1.0)	49	90.7 (83.0-98.5)	0.6 (0.5 - 0.8)	0.70 (0.28 - 1.76)
	1999-2005	13	10.5 (5.1-15.9)	0.4 (0.2 - 0.7)	111	89.5 (84.1-94.9)	0.5 (0.4 - 0.6)	0.81 (0.45 - 1.44)
	2006-2014	11	9.2 (4.0-14.3)	0.2 (0.1 - 0.4)	109	90.8 (85.7-96.0)	0.4 (0.3 - 0.4)	0.61 (0.33 - 1.14)
Total (ASR)	1996-1998	41	2.5 (1.7-3.2)	4.2 (2.5 - 5.8)	1624	97.5 (96.8-98.3)	6.7 (6.4 - 7.0)	0.63 (0.42 - 0.93)
	1999-2005	114	3.5 (2.9-4.1)	4.9 (3.8 - 6.0)	3137	96.5 (95.9-97.1)	6.1 (5.9 - 6.4)	0.79 (0.63 - 0.99)
	2006-2014	58	6.1 (4.6-7.6)	1.0 (0.7 - 1.4)	899	93.9 (92.4-95.5)	1.3 (1.3 - 1.4)	0.78 (0.57 - 1.06)

\* 1996-1998 = pre-vaccination, 1999-2005 = private vaccination, 2006-2014 = public vaccination, ASR=age-standardized rate

**Disclosures.** All authors: No reported disclosures.

### 2302. A Meta-Analysis of Risk Factors for Herpes Zoster Infection

Kamalpreet K. Parhar; Bill Huang, BSc (Pharm); Nirma K. Vadlamudi, BA (Bio), BS (Chem), MPH; Fawziah Marra, BSc (Pharm), PharmD; University of British Columbia, Vancouver, BC, Canada

**Session:** 247. Clinical Virology/Viral Epidemiology

*Saturday, October 5, 2019: 12:15 PM*

**Background.** The burden of herpes zoster (HZ) is significant worldwide, with millions affected and the incidence rising. Current literature has identified some risk factors for this disease; however, there is yet to be a comprehensive study that pools