

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-(4-Chloroanilino)-3-(2-hydroxyethyl)-quinazolin-4(3H)-one

Hong-Ling Wang, Xu-Hong Yang and Ming-Hu Wu*

Faculty of Chemistry and Life Science, Xianning University, Xianning 437100, People's Republic of China

Correspondence e-mail: minghuwu@hotmail.com


Received 14 October 2008; accepted 7 November 2008

Key indicators: single-crystal X-ray study; $T = 273$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.037; wR factor = 0.108; data-to-parameter ratio = 13.8.

In the title molecule, $\text{C}_{16}\text{H}_{14}\text{ClN}_3\text{O}_2$, the dihedral angle between the chlorophenyl and pyrimidinone rings is $14.8(1)^\circ$, while the dihedral angle between the fused benzene ring and the pyrimidinone ring is $3.8(1)^\circ$. In the crystal structure, intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, together with intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen-bonding interactions, are present.

Related literature

For the biological activities and applications of 4(3H)-quinazolinone, see: Armarego (1963); Fisnerova *et al.* (1986); Gravier *et al.* (1992). For details of our ongoing heterocyclic synthesis and drug discovery project, see: Yang *et al.* (2008).



Experimental

Crystal data

 $\text{C}_{16}\text{H}_{14}\text{ClN}_3\text{O}_2$ $M_r = 315.75$

Monoclinic, $P2_1/n$
 $a = 9.0707(18)$ Å
 $b = 11.345(2)$ Å
 $c = 14.143(3)$ Å
 $\beta = 96.98(3)^\circ$
 $V = 1444.6(5)$ Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.28$ mm⁻¹
 $T = 273(2)$ K
 $0.20 \times 0.20 \times 0.10$ mm

Data collection

Bruker SMART APEX CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2001)
 $T_{\min} = 0.947$, $T_{\max} = 0.973$

8113 measured reflections
 2824 independent reflections
 2300 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.108$
 $S = 1.05$
 2824 reflections
 205 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.16$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.24$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1A}\cdots\text{O2}$	0.839 (18)	1.993 (19)	2.8017 (19)	161.8 (17)
$\text{O2}-\text{H2A}\cdots\text{O1}^1$	0.86 (2)	1.86 (2)	2.7180 (18)	174 (2)

Symmetry code: (i) $-x - \frac{1}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: SMART (Bruker, 2001); cell refinement: SAINT-Plus (Bruker, 2001); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: PLATON (Spek, 2003); software used to prepare material for publication: PLATON.

We gratefully acknowledge financial support of this work as a project of the Natural Science Foundation of Hubei Province under grant No. 2006ABA334.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: EZ2147).

References

- Armarego, W. L. F. (1963). *Advanced Heterocyclic Chemistry*, Vol. 1, pp. 253–309. New York: Academic Press.
- Bruker (2001). SMART and SAINT-Plus. Bruker AXS Inc., Madison, Wisconsin, USA.
- Fisnerova, L., Grimova, J., Roubal, Z., Maturova, E. & Brunova, B. (1986). *Cesk. Farm.* **3**, 447–450.
- Gravier, D., Dupin, J. P., Casadebaig, F., Hou, G., Boisseau, M. & Bernard, H. (1992). *Pharmazie*, **47**, 91–94.
- Sheldrick, G. M. (2001). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Yang, X. H., Wu, M. H., Sun, S. F., Ding, M. W., Xie, J. L. & Xia, Q. H. (2008). *J. Heterocycl. Chem.* **5**, 1365–1369.

supplementary materials

Acta Cryst. (2008). E64, o2325 [doi:10.1107/S1600536808036623]

2-(4-Chloroanilino)-3-(2-hydroxyethyl)quinazolin-4(3H)-one

H.-L. Wang, X.-H. Yang and M.-H. Wu

Comment

One of the most frequently encountered heterocyclic molecules in medicinal chemistry is 4(3H)-quinazolinone, which has wide application as a result of antibacterial, antifungal, anticonvulsant, and anti-inflammatory activities (Armarego, 1963; Gravier *et al.*, 1992; Fisnerova *et al.*, 1986). In our ongoing heterocyclic synthesis and drug discovery project (Yang *et al.*, 2008) we have focused on the synthesis of quinazolinones and pyrazolo pyrimidinones. Herein, the title compound was synthesized and determined by single-crystal X-ray diffraction.

In the molecule (Fig. 1), the dihedral angle between the chlorophenyl and pyrimidinone rings is 14.8 (1)°, and the dihedral angle between the fused benzene and pyrimidinone rings is 3.8 (1)°.

In the crystal structure, molecules are linked by intramolecular N1–H1A···O2 hydrogen-bonds together with O2–H2A···O1ⁱ intermolecular hydrogen-bonding interactions (symmetry code: $i, -1/2 - x, 1/2 + y, 3/2 - z$) (Fig. 2).

Experimental

To a solution of 2-ethoxycarbonyliminophosphorane (1.27 g, 3 mmol) in 10 ml absolute anhydrous CH₂Cl₂, 4-chlorophenyl-isocyanate (0.46 g, 3 mmol) was added dropwise at room temperature. The reaction mixture was left unstirred for 6 h at 273–278 K, whereafter a solution of 2-hydroxyethylamine (0.18 g, 3 mmol) in 5 ml absolute anhydrous CH₂Cl₂ was added. The reaction mixture was then stirred overnight, the solution cooled and the reaction product recrystallized from CH₃OH to give colorless crystals of the title compound suitable for X-ray analysis in 58% yield.

Refinement

H atoms bonded to C atoms were placed in calculated positions (C–H = 0.93–0.97 Å) and included in the riding model approximation. The positional parameters of H atoms bonded to N and O atoms were refined independently. For all H atoms $U_{\text{iso}}(\text{H}) = 1.2U_{\text{iso}}(\text{C}, \text{N})$ or $1.5U_{\text{iso}}(\text{O})$.

Figures




Fig. 1. View of the molecule with the atom-labeling scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. Crystal packing viewed down the *a* axis. Hydrogen bonds are shown as dashed lines.

2-(4-Chloroanilino)-3-(2-hydroxyethyl)quinazolin-4(3H)-one

Crystal data

$C_{16}H_{14}ClN_3O_2$

$M_r = 315.75$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 9.0707$ (18) Å

$b = 11.345$ (2) Å

$c = 14.143$ (3) Å

$\beta = 96.98$ (3)°

$V = 1444.6$ (5) Å³

$Z = 4$

$F_{000} = 656$

$D_x = 1.452$ Mg m⁻³

Melting point = 432–434 K

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 3170 reflections

$\theta = 2.3$ – 26.2 °

$\mu = 0.28$ mm⁻¹

$T = 273$ (2) K

Block, colourless

$0.20 \times 0.20 \times 0.10$ mm

Data collection

Bruker SMART APEX CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 273$ (2) K

φ and ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 2001)

$T_{\min} = 0.947$, $T_{\max} = 0.973$

8113 measured reflections

2824 independent reflections

2300 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.019$

$\theta_{\text{max}} = 26.0$ °

$\theta_{\text{min}} = 2.3$ °

$h = -11 \rightarrow 9$

$k = -11 \rightarrow 13$

$l = -17 \rightarrow 17$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.037$

$wR(F^2) = 0.108$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0587P)^2 + 0.2177P]$

$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
2824 reflections	$(\Delta/\sigma)_{\max} = 0.001$
205 parameters	$\Delta\rho_{\max} = 0.16 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.24 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.24608 (18)	1.06958 (14)	0.90802 (11)	0.0460 (4)
H1	0.2103	1.0780	0.8439	0.055*
C2	0.34154 (18)	1.15287 (15)	0.95164 (11)	0.0495 (4)
H2	0.3699	1.2173	0.9174	0.059*
C3	0.39474 (17)	1.13990 (14)	1.04660 (12)	0.0460 (4)
C4	0.35073 (19)	1.04650 (15)	1.09827 (12)	0.0524 (4)
H4	0.3857	1.0395	1.1626	0.063*
C5	0.25416 (18)	0.96263 (15)	1.05448 (11)	0.0502 (4)
H5	0.2242	0.8995	1.0895	0.060*
C6	0.20225 (16)	0.97305 (13)	0.95836 (11)	0.0397 (3)
C7	0.03780 (15)	0.79644 (13)	0.93558 (10)	0.0396 (3)
C8	-0.13344 (17)	0.63934 (14)	0.88093 (11)	0.0465 (4)
C9	-0.13969 (17)	0.61551 (14)	0.98083 (11)	0.0460 (4)
C10	-0.04823 (16)	0.68016 (14)	1.04878 (11)	0.0427 (4)
C11	-0.05282 (18)	0.65638 (16)	1.14562 (12)	0.0520 (4)
H11	0.0091	0.6974	1.1915	0.062*
C12	-0.1479 (2)	0.57314 (17)	1.17282 (13)	0.0608 (5)
H12	-0.1512	0.5588	1.2373	0.073*
C13	-0.2398 (2)	0.50952 (18)	1.10526 (15)	0.0672 (5)
H13	-0.3044	0.4533	1.1247	0.081*
C14	-0.2352 (2)	0.52964 (16)	1.01032 (13)	0.0607 (5)
H14	-0.2956	0.4862	0.9652	0.073*
C15	-0.00780 (19)	0.74122 (15)	0.76189 (11)	0.0477 (4)
H15A	-0.0208	0.6648	0.7312	0.057*
H15B	0.0951	0.7643	0.7613	0.057*
C16	-0.1053 (2)	0.82925 (16)	0.70402 (12)	0.0572 (4)

supplementary materials

H16A	-0.1005	0.8163	0.6367	0.069*
H16B	-0.2076	0.8194	0.7161	0.069*
C11	0.51722 (6)	1.24474 (4)	1.10084 (4)	0.06868 (19)
N1	0.10834 (15)	0.89234 (12)	0.90499 (9)	0.0450 (3)
H1A	0.0750 (19)	0.9182 (16)	0.8510 (13)	0.054*
N2	-0.03685 (14)	0.72897 (11)	0.86188 (9)	0.0422 (3)
N3	0.03988 (14)	0.77194 (11)	1.02486 (9)	0.0436 (3)
O1	-0.20317 (14)	0.58621 (11)	0.81420 (9)	0.0639 (4)
O2	-0.05636 (14)	0.94472 (12)	0.72940 (9)	0.0606 (3)
H2A	-0.129 (3)	0.992 (2)	0.7132 (17)	0.091*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0555 (9)	0.0443 (9)	0.0388 (8)	-0.0021 (7)	0.0088 (7)	0.0014 (7)
C2	0.0578 (10)	0.0417 (9)	0.0510 (9)	-0.0077 (7)	0.0148 (7)	0.0012 (7)
C3	0.0454 (8)	0.0386 (8)	0.0536 (9)	-0.0025 (7)	0.0049 (7)	-0.0039 (7)
C4	0.0591 (10)	0.0470 (9)	0.0480 (9)	-0.0069 (8)	-0.0063 (7)	0.0057 (7)
C5	0.0551 (10)	0.0435 (9)	0.0493 (9)	-0.0087 (7)	-0.0040 (7)	0.0097 (7)
C6	0.0385 (8)	0.0355 (8)	0.0450 (8)	0.0015 (6)	0.0043 (6)	-0.0006 (6)
C7	0.0374 (8)	0.0374 (8)	0.0431 (8)	0.0016 (6)	0.0016 (6)	-0.0023 (6)
C8	0.0476 (9)	0.0403 (9)	0.0512 (9)	-0.0021 (7)	0.0043 (7)	-0.0092 (7)
C9	0.0481 (9)	0.0372 (8)	0.0531 (9)	-0.0018 (7)	0.0078 (7)	-0.0045 (7)
C10	0.0421 (8)	0.0391 (8)	0.0465 (8)	0.0006 (6)	0.0035 (6)	0.0020 (7)
C11	0.0526 (9)	0.0539 (10)	0.0482 (9)	-0.0061 (8)	0.0008 (7)	0.0042 (8)
C12	0.0657 (11)	0.0616 (12)	0.0556 (10)	-0.0088 (9)	0.0093 (9)	0.0124 (9)
C13	0.0748 (13)	0.0568 (12)	0.0714 (12)	-0.0225 (10)	0.0149 (10)	0.0049 (10)
C14	0.0681 (11)	0.0491 (10)	0.0654 (11)	-0.0179 (9)	0.0099 (9)	-0.0083 (9)
C15	0.0560 (10)	0.0468 (9)	0.0403 (8)	0.0042 (7)	0.0065 (7)	-0.0050 (7)
C16	0.0626 (11)	0.0598 (11)	0.0466 (9)	0.0048 (9)	-0.0035 (8)	-0.0017 (8)
C11	0.0759 (4)	0.0539 (3)	0.0731 (3)	-0.0218 (2)	-0.0037 (3)	-0.0055 (2)
N1	0.0489 (7)	0.0433 (8)	0.0408 (7)	-0.0053 (6)	-0.0027 (6)	0.0043 (6)
N2	0.0466 (7)	0.0386 (7)	0.0411 (7)	0.0003 (5)	0.0039 (5)	-0.0046 (5)
N3	0.0445 (7)	0.0434 (7)	0.0421 (7)	-0.0048 (5)	0.0018 (5)	0.0005 (6)
O1	0.0715 (8)	0.0628 (8)	0.0562 (7)	-0.0195 (6)	0.0029 (6)	-0.0179 (6)
O2	0.0660 (8)	0.0522 (8)	0.0596 (7)	0.0103 (6)	-0.0079 (6)	0.0050 (6)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.376 (2)	C9—C14	1.400 (2)
C1—C6	1.390 (2)	C10—N3	1.3796 (19)
C1—H1	0.9300	C10—C11	1.402 (2)
C2—C3	1.379 (2)	C11—C12	1.365 (2)
C2—H2	0.9300	C11—H11	0.9300
C3—C4	1.374 (2)	C12—C13	1.391 (3)
C3—C11	1.7408 (16)	C12—H12	0.9300
C4—C5	1.387 (2)	C13—C14	1.368 (3)
C4—H4	0.9300	C13—H13	0.9300
C5—C6	1.389 (2)	C14—H14	0.9300

C5—H5	0.9300	C15—N2	1.476 (2)
C6—N1	1.406 (2)	C15—C16	1.508 (2)
C7—N3	1.2907 (19)	C15—H15A	0.9700
C7—N1	1.3593 (19)	C15—H15B	0.9700
C7—N2	1.400 (2)	C16—O2	1.415 (2)
C8—O1	1.2283 (19)	C16—H16A	0.9700
C8—N2	1.390 (2)	C16—H16B	0.9700
C8—C9	1.446 (2)	N1—H1A	0.839 (18)
C9—C10	1.398 (2)	O2—H2A	0.86 (2)
C2—C1—C6	120.99 (15)	C12—C11—H11	119.9
C2—C1—H1	119.5	C10—C11—H11	119.9
C6—C1—H1	119.5	C11—C12—C13	120.77 (17)
C1—C2—C3	119.35 (15)	C11—C12—H12	119.6
C1—C2—H2	120.3	C13—C12—H12	119.6
C3—C2—H2	120.3	C14—C13—C12	120.04 (17)
C4—C3—C2	120.70 (15)	C14—C13—H13	120.0
C4—C3—C11	120.26 (13)	C12—C13—H13	120.0
C2—C3—C11	119.04 (13)	C13—C14—C9	120.15 (17)
C3—C4—C5	120.01 (15)	C13—C14—H14	119.9
C3—C4—H4	120.0	C9—C14—H14	119.9
C5—C4—H4	120.0	N2—C15—C16	114.90 (14)
C4—C5—C6	119.95 (15)	N2—C15—H15A	108.5
C4—C5—H5	120.0	C16—C15—H15A	108.5
C6—C5—H5	120.0	N2—C15—H15B	108.5
C5—C6—C1	118.97 (14)	C16—C15—H15B	108.5
C5—C6—N1	125.54 (14)	H15A—C15—H15B	107.5
C1—C6—N1	115.48 (14)	O2—C16—C15	109.27 (14)
N3—C7—N1	122.16 (14)	O2—C16—H16A	109.8
N3—C7—N2	123.99 (14)	C15—C16—H16A	109.8
N1—C7—N2	113.85 (13)	O2—C16—H16B	109.8
O1—C8—N2	119.21 (15)	C15—C16—H16B	109.8
O1—C8—C9	125.53 (15)	H16A—C16—H16B	108.3
N2—C8—C9	115.25 (13)	C7—N1—C6	128.91 (13)
C10—C9—C14	119.79 (15)	C7—N1—H1A	116.0 (12)
C10—C9—C8	118.88 (14)	C6—N1—H1A	112.9 (13)
C14—C9—C8	121.33 (15)	C8—N2—C7	121.00 (13)
N3—C10—C9	122.70 (14)	C8—N2—C15	116.40 (13)
N3—C10—C11	118.15 (14)	C7—N2—C15	122.37 (13)
C9—C10—C11	119.01 (14)	C7—N3—C10	117.54 (13)
C12—C11—C10	120.22 (16)	C16—O2—H2A	107.7 (16)
C6—C1—C2—C3	0.2 (2)	C12—C13—C14—C9	-1.0 (3)
C1—C2—C3—C4	-1.5 (3)	C10—C9—C14—C13	0.4 (3)
C1—C2—C3—C11	179.23 (12)	C8—C9—C14—C13	-179.47 (17)
C2—C3—C4—C5	1.3 (3)	N2—C15—C16—O2	75.96 (19)
C11—C3—C4—C5	-179.42 (13)	N3—C7—N1—C6	6.0 (2)
C3—C4—C5—C6	0.1 (3)	N2—C7—N1—C6	-174.53 (14)
C4—C5—C6—C1	-1.4 (2)	C5—C6—N1—C7	7.1 (3)
C4—C5—C6—N1	177.75 (15)	C1—C6—N1—C7	-173.72 (14)

supplementary materials

C2—C1—C6—C5	1.2 (2)	O1—C8—N2—C7	176.07 (14)
C2—C1—C6—N1	-178.01 (14)	C9—C8—N2—C7	-5.2 (2)
O1—C8—C9—C10	177.05 (16)	O1—C8—N2—C15	-9.3 (2)
N2—C8—C9—C10	-1.6 (2)	C9—C8—N2—C15	169.47 (13)
O1—C8—C9—C14	-3.1 (3)	N3—C7—N2—C8	9.8 (2)
N2—C8—C9—C14	178.30 (15)	N1—C7—N2—C8	-169.68 (13)
C14—C9—C10—N3	-174.93 (15)	N3—C7—N2—C15	-164.55 (14)
C8—C9—C10—N3	5.0 (2)	N1—C7—N2—C15	16.0 (2)
C14—C9—C10—C11	0.8 (2)	C16—C15—N2—C8	94.27 (17)
C8—C9—C10—C11	-179.30 (14)	C16—C15—N2—C7	-91.15 (19)
N3—C10—C11—C12	174.43 (16)	N1—C7—N3—C10	173.12 (13)
C9—C10—C11—C12	-1.5 (3)	N2—C7—N3—C10	-6.3 (2)
C10—C11—C12—C13	1.0 (3)	C9—C10—N3—C7	-1.1 (2)
C11—C12—C13—C14	0.3 (3)	C11—C10—N3—C7	-176.85 (14)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A \cdots O2	0.839 (18)	1.993 (19)	2.8017 (19)	161.8 (17)
O2—H2A \cdots O1 ⁱ	0.86 (2)	1.86 (2)	2.7180 (18)	174 (2)

Symmetry codes: (i) $-x-1/2, y+1/2, -z+3/2$.

Fig. 1




Fig. 2

