

The Colonic Single Stripe Sign: A Case of Ischemic Colitis

Malav P. Parikh¹, Jinendra Satiya², Miguel Berger-Saunderson³, Niyati M. Gupta³, Madhusudhan R. Sanaka¹

1. Gastroenterology and Hepatology, Cleveland Clinic Foundation, Cleveland, USA 2. Internal Medicine, University of Miami, John F Kennedy Regional Campus, Atlantis, USA 3. Internal Medicine, Cleveland Clinic Foundation, Cleveland, USA

✉ **Corresponding author:** Jinendra Satiya, jinen19@gmail.com

Disclosures can be found in Additional Information at the end of the article

Abstract

Sudden, transient reduction in the blood flow especially in the “watershed” regions of the colon can lead to intestinal ischemia causing a decrease in the delivery of oxygen and nutrients to the intestinal wall. Patients with ischemic colitis often have elevated white blood cell counts, serum lactate, and serum amylase levels. Colonoscopy can confirm the diagnosis of ischemia and the findings include edematous, friable mucosa, erythema with interspersed pale areas, scattered hemorrhagic erosions or linear ulcerations. A single, linear ulcer running along the anti-mesenteric colonic wall, “single-stripe sign”- favors the diagnosis of ischemic colitis. Management of mild and moderate colonic ischemia includes supportive care with bowel rest, gastric suction for associated ileus, fluid-electrolyte balance, and broad-spectrum antibiotics. Patients with severe colonic ischemia may require abdominal exploration and colectomy.

Categories: Gastroenterology

Keywords: colonic ischemia, colonoscopy, colitis

Introduction

Ischemic colitis is the most common form of vascular injury to the gastrointestinal tract [1]. The most common reason for colonic ischemia is hypoperfusion as compared to thrombotic and embolic events accounting for most vascular insults to the small bowel [2]. Occurrence is typically segmental with left-sided lesions involving the water-shed regions such as the splenic flexure, descending colon, and sigmoid colon. The spectrum of injury ranges from transient colitis, gangrene to fulminant pancolitis. Differential diagnoses for a patient presenting with abdominal pain and bright red bleeding per rectum include inflammatory bowel disease, acute mesenteric ischemia, and infectious colitis. We present the case of a 56-year-old female who presented with bright red bleeding per rectum and was found to have the characteristic single-stripe sign, pathognomonic for ischemic colitis.

Case Presentation

A 62-year-old female with a past medical history of tobacco abuse, hypertension, constipation, and chronic headaches managed by ibuprofen, presented to the emergency room with complaints of diffuse abdominal pain and bright red blood per rectum since two days. On physical examination, the patient was afebrile, the abdomen was soft, non-tender, with normal bowel sounds. The rectal exam was unrevealing except for bright red blood on the gloved finger. Laboratory studies showed hemoglobin of 15 g/dl, white blood cell count of 23,700/ μ l with normal serum lactate, amylase, and lipase levels. Colonoscopy showed discontinuous areas of

Received 04/26/2019

Review began 04/27/2019

Review ended 05/05/2019

Published 05/08/2019

© Copyright 2019

Parikh et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 3.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

How to cite this article

Parikh M P, Satiya J, Berger-Saunderson M, et al. (May 08, 2019) The Colonic Single Stripe Sign: A Case of Ischemic Colitis. Cureus 11(5): e4622. DOI 10.7759/cureus.4622

non-bleeding ulcerated mucosa with no stigmata of recent bleeding in the sigmoid colon and descending colon for a total of 20 cms, consistent with the “single-stripe sign”, characteristic of colonic ischemia (Figure 1A). Further, microscopic evaluation of the biopsied tissue revealed characteristic features of surface epithelial injury, crypt epithelial atrophy, crypt loss, lamina propria hemorrhage, and lamina propria hyalinization consistent with the colonoscopy findings of colonic ischemia (Figure 1B).




FIGURE 1: Colonoscopy demonstrating discontinuous areas of non-bleeding ulcerated mucosa with no stigmata of recent bleeding in the sigmoid colon and descending colon; microscopic evaluation revealed characteristic features of surface epithelial injury, crypt epithelial atrophy, crypt loss, lamina propria hemorrhage, and lamina propria hyalinization

The patient was discharged on a seven-day course of oral ciprofloxacin and metronidazole, managed for constipation with polyethylene glycol, and was advised to quit smoking and to avoid non-steroidal anti-inflammatory drugs.

Discussion

Sudden, transient reduction in the blood flow especially in the “watershed” regions of the colon can lead to intestinal ischemia causing a decrease in the delivery of oxygen and nutrients to the intestinal wall [3]. The etiology of colonic ischemia can be broadly divided into five categories: 1) non-occlusive ischemia due to hypotension; 2) thromboembolic occlusion of arteries; 3) mesenteric vein thrombosis; 4) mechanical obstruction due to cancer, adhesions or fecal impaction; 4) procedures like aortoiliac reconstruction, abdominal aortic aneurysm repair, renal transplant, and 5) drug-induced ischemia. Common clinical manifestations include abdominal cramping and bright red blood per rectum [4-5]. Transmural infarction can result in gangrenous colitis, peritoneal signs, fever, and ileus. Patients with ischemic colitis often have elevated white blood cell counts, serum lactate, and serum amylase levels. Serum lactate can be used as a predictor of bowel ischemia and is elevated in severe cases of ischemic colitis [6]. Its sensitivity increases in the event of a bowel infarction.

Marston et al. first coined the term ischemic colitis in 1966 to characterize the clinical syndrome associated with vascular compromise to the colon [7]. Zuckerman et al. were the first to report the association of the colonic single-stripe sign with a diagnosis of mild ischemic colitis [8]. They reported the presence of the colonic single-stripe sign greater than 5 cms in length, with 89% of the lesions identified in an isolated segment of the left colon. A preceding ischemic event was noted in 62% of these patients. They believed the linear appearance of the

ulcer to be a result of segmental vascular compromise at the mucosal level, occurring due to hypoxemia and small vessel inadequacy. It has been postulated that this edge or demarcation represents involvement of the mesenteric vascular border. Colonoscopy can confirm the diagnosis of ischemic colitis and findings include edematous, friable mucosa, erythema with interspersed pale areas, scattered hemorrhagic erosions or linear ulcerations. A single, linear ulcer running along the anti-mesenteric colonic wall, "single-stripe sign"- favors a diagnosis of mild ischemic colitis.

A plain abdominal radiograph is frequently nonspecific unless there is advanced ischemia, which shows pneumatosis or distension. Computed tomography (CT) of the abdomen with contrast can show edema and thickening of the bowel wall in a segmental pattern, irregular bowel contour, inflamed mesentery with stranding of the fat, or presence of free peritoneal fluid. Management of mild and moderate colonic ischemia includes supportive care with bowel rest, gastric suction for associated ileus, fluid-electrolyte balance and broad-spectrum antibiotics. Patients with severe colonic ischemia require abdominal exploration and colectomy [4-5].

Conclusions

The increase in longevity of the population has brought about a consequent increase in the incidence of colonic ischemia. Patients commonly present with crampy abdominal pain and rectal bleeding. Colonoscopy is the gold standard for diagnosis. Left-sided lesions in the distribution of the inferior mesenteric artery are more common, while right-sided lesions portend a worse prognosis with a higher risk for surgical intervention. In the majority of patients, treatment is supportive with bowel rest and intravenous fluids.

Additional Information

Disclosures

Human subjects: Consent was obtained by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Feuerstadt P, Brandt LJ: Update on colon ischemia: recent insights and advances . *Curr Gastroenterol Rep.* 2015, 17:45. [10.1007/s11894-015-0469-6](https://doi.org/10.1007/s11894-015-0469-6)
2. Oglat A, Quigley EM: Colonic ischemia: usual and unusual presentations and their management. *Curr Opin Gastroenterol.* 2017, 33:34-40. [10.1097/MOG.0000000000000325](https://doi.org/10.1097/MOG.0000000000000325)
3. Brandt LJ, Feuerstadt P, Longstreth G, Boley S: ACG clinical guideline: epidemiology, risk factors, patterns of presentation, diagnosis, and management of colon ischemia (CI). *Am J Gastroenterol.* 2015, 110:18-44. [10.1038/ajg.2014.395](https://doi.org/10.1038/ajg.2014.395)
4. Moszkowicz D, Mariani A, Trésallet C, Menegaux F: Ischemic colitis: the ABCs of diagnosis and surgical management. *J Visc Surg.* 2013, 150:19-28. [10.1016/j.jvisc.2013.01.002](https://doi.org/10.1016/j.jvisc.2013.01.002)
5. Cotter TG, Bledsoe AC, Sweetser S: Colon ischemia: an update for clinicians . *Mayo Clin Proc.* 2016, 91:671-77. [10.1016/j.mayocp.2016.02.006](https://doi.org/10.1016/j.mayocp.2016.02.006)
6. Kintu-Luwaga R, Galukande M, Owori FN: Serum lactate and phosphate as biomarkers of intestinal ischemia in a Ugandan tertiary hospital: a cross-sectional study. *Int J Emerg Med.* 2013, 6:44. [10.1186/1865-1380-6-44](https://doi.org/10.1186/1865-1380-6-44)
7. Marston A, Pheils MT, Thomas ML, Morson BC: Ischaemic colitis. *Gut.* 1966, 7:1-15.

[10.1136/gut.7.1.1](#)

8. Zuckerman GR, Prakash C, Merriman RB, Sawhney MS, DeSchryver-Kecskemeti K, Clouse RE: The colon single-stripe sign and its relationship to ischemic colitis . Am J Gastroenterol. 2003, 98:2018-2022. [10.1016/S0002-9270\(03\)00544-6](#)