

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-Hydroxy-4-(prop-2-ynoxy)benzaldehyde

V. Selvarani,^a M.A. Neelakantan,^{a*} V. Silambarasan^b and D. Velmurugan^b^aChemistry Research Centre, National Engineering College, K.R. Nagar, Kovilpatti 628 503, India, and ^bCAS in Crystallography and Biophysics, University of Madras, Guindy Campus, Chennai-25, India

Correspondence e-mail: drmaneelakantan@gmail.com


Received 16 November 2012; accepted 3 December 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.044; wR factor = 0.132; data-to-parameter ratio = 17.9.

The asymmetric unit of the title compound, $\text{C}_{10}\text{H}_8\text{O}_3$, contains two independent molecules, both of which are almost planar (r.m.s deviations for all non-H atoms of 0.044 and 0.053 Å). The dihedral angles between the benzene ring and the prop-1-yne group are 3.47 (1) and 3.07 (1)° in the two molecules, and the prop-1-yne groups adopt extended conformations. In each molecule, an intramolecular O—H...O hydrogen bond involving the OH and aldehyde substituents forms an $S(6)$ ring. In the crystal, molecules are linked into cyclic centrosymmetric dimers via C—H...O hydrogen bonds, generating $R_2^2(14)$ ring motifs. The crystal structure is further stabilized by aromatic π – π stacking interactions between the benzene rings [centroid–centroid distances = 3.813 (2) and 3.843 (2) Å]

Related literature

For the biological activity of benzaldehyde derivatives, see: Zhao *et al.* (2007); Ley & Bertram (2001); Delogu *et al.* (2010). For a related structure see: Esakkiammal *et al.* (2012). For standard bond lengths, see: Allen *et al.* (1987) and for hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

 $\text{C}_{10}\text{H}_8\text{O}_3$ $M_r = 176.16$

Triclinic, $P\bar{1}$
 $a = 7.0835$ (5) Å
 $b = 10.4059$ (7) Å
 $c = 12.8461$ (8) Å
 $\alpha = 73.910$ (3)°
 $\beta = 89.756$ (4)°
 $\gamma = 73.436$ (4)°

$V = 869.16$ (10) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 293$ K
 $0.20 \times 0.20 \times 0.20$ mm

Data collection

Bruker SMART APEXII area-detector diffractometer
 15699 measured reflections

4347 independent reflections
 2880 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.132$
 $S = 1.05$
 4347 reflections
 243 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.15$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.20$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O2—H2...O3	0.82	1.92	2.6387 (16)	146
O5—H5...O6	0.82	1.93	2.6441 (16)	146
C10—H10...O3 ⁱ	0.86 (2)	2.51 (2)	3.369 (2)	171.4 (2)
C18—H18A...O5 ⁱⁱ	0.97	2.45	3.281 (2)	144
C20—H20...O6 ⁱ	0.91 (2)	2.37 (2)	3.280 (2)	178.8 (2)

Symmetry codes: (i) $x, y, z + 1$; (ii) $-x + 1, -y + 1, -z + 1$.

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 2012); software used to prepare material for publication: SHELXL97 and PLATON (Spek, 2009).

VS and DV thank the TBI X-ray facility, CAS in Crystallography and Biophysics, University of Madras, India, for the data collection and the UGC SAP for the facilities provided to the department.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5285).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
 Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
 Bruker (2008). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Delogu, G., Podda, G., Corda, M., Fadda, M. B., Fais, A. & Era, B. (2010). *Bioorg. Med. Chem. Lett.* **20**, 6138–6140.
 Esakkiammal, M., Selvarani, V., Neelakantan, M. A., Silambarasan, V. & Velmurugan, D. (2012). *Acta Cryst.* **E68**, o2465.
 Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
 Ley, J. P. & Bertram, H. J. (2001). *Bioorg. Med. Chem. Lett.* **9**, 1879–1885.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Zhao, X., Song, D. K., Radbil, A. B. & Radbil, B. A. (2007). *Russ. J. Appl. Chem.* **80**, 1373–1375.

supplementary materials

Acta Cryst. (2013). E69, o64 [doi:10.1107/S1600536812049598]

2-Hydroxy-4-(prop-2-ynoxy)benzaldehyde

V. Selvarani, M.A. Neelakantan, V. Silambarasan and D. Velmurugan

Comment

Schiff bases derived from amines and substituted benzaldehydes exhibit antibacterial, anticancer and antitumour activities (Zhao *et al.*, 2007). Several benzaldoximes, benzaldehyde-*O*-ethyloximes and acetophenone oximes were synthesized and evaluated as tyrosinase inhibitors (Ley & Bertram, 2001). Bis-salicylaldehydes exhibited greater inhibitory activity than salicylaldehyde (Delogu *et al.*, 2010). In view of these potential applications and in continuation of our work on the crystal structures of benzaldehyde derivatives, the structure of the title compound has been carried out and the results are presented here.

X-ray analysis confirms the molecular structure and atom connectivity as illustrated in Fig. 1. The bond distances are normal (Allen *et al.*, 1987) and are comparable with those found in related structures (Esakkiammal *et al.*, 2012). The asymmetric unit contains two crystallographically independent molecules. In both molecules, the dihedral angles between the benzene rings and the prop-1-yne groups are 3.47 (1)° and 3.07 (1)°, respectively. Both molecules are almost planar, with r.m.s deviations of 0.044 and 0.053 Å from the best fit plane through all non-hydrogen atoms in the molecules. The prop-1-yne groups adopt extended conformations as can be seen from the torsion angles C15—O4—C18—C19 = -177.04 (12)° and C5—O1—C8—C9 = 179.39 (13)°. Atoms O2 and O5 deviate by 0.013 (1) and -0.004 (1) Å from the least squares plane of the benzene rings. Intramolecular O2—H2...O3 and O5—H5...O6 hydrogen bonds form S(6) rings in both molecules (Bernstein *et al.*, 1995).

Molecules are linked into cyclic centrosymmetric dimers *via* C—H...O hydrogen bonds with $R_2^2(14)$ ring motifs (Bernstein *et al.*, 1995). The crystal structure is further stabilized by an aromatic π - π stacking interaction between the benzene rings of adjacent molecules, with centroid to centroid distances $Cg1 \cdots Cg2^i = 3.813$ (2) Å; $Cg2 \cdots Cg1^{ii} = 3.843$ (2) Å [(i) 1 - x, 1 - y, 1 - z and (ii) 2 - x, 1 - y, 1 - z]. ($Cg1$ and $Cg2$ are the centroids of the C1—C6 and C11—C16 rings respectively).

Experimental

Equimolar amounts of 3-bromopropyne (10 mmol), 2,4-dihydroxybenzaldehyde (10 mmol) and potassium carbonate (15 mmol) were suspended in acetonitrile (30 ml) and refluxed for 30 h in presence of KI (0.1 g) as a catalyst. The reaction mixture was filtered while hot to remove insoluble impurities, neutralized with dil.HCl (3 M), extracted with chloroform and dried with Na₂SO₄. The extracts were concentrated to obtain a brown solid which was then purified by column chromatography over SiO₂ by eluting with a mixture of 4% ethyl acetate in n-hexane. Evaporation of the purified extract yielded 2-hydroxy-4-(prop-2-ynoxy)benzaldehyde in the form of a pure white solid in 85% yield. Crystals suitable for X-ray analysis were obtained by the slow evaporation method.

Refinement

The acetylenic H atoms H10 and H20 were located in a difference Fourier map and refined freely. Other H atoms were positioned geometrically (C–H = 0.93–0.97 Å; O–H = 0.82 Å) and allowed to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$ for hydroxyl H atoms and $1.2U_{\text{eq}}(\text{C})$ for all other H atoms.

Computing details

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT* (Bruker, 2008); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 2012); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008) and *PLATON* (Spek, 2009).





Figure 1

The molecular structure of the title compound, showing the atomic numbering and displacement ellipsoids drawn at the 30% probability level.

**Figure 2**

The crystal packing of the molecules viewed down *b* axis. H atoms not involved in hydrogen bonding (dashed lines) have been omitted for clarity.

2-Hydroxy-4-(prop-2-ynoxy)benzaldehyde

Crystal data

$C_{10}H_8O_3$
 $M_r = 176.16$
 Triclinic, $P\bar{1}$
 Hall symbol: $-P\ 1$
 $a = 7.0835\ (5)\ \text{\AA}$
 $b = 10.4059\ (7)\ \text{\AA}$
 $c = 12.8461\ (8)\ \text{\AA}$
 $\alpha = 73.910\ (3)^\circ$
 $\beta = 89.756\ (4)^\circ$
 $\gamma = 73.436\ (4)^\circ$
 $V = 869.16\ (10)\ \text{\AA}^3$

$Z = 4$
 $F(000) = 368$
 $D_x = 1.346\ \text{Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$
 Cell parameters from 4347 reflections
 $\theta = 1.7\text{--}28.4^\circ$
 $\mu = 0.10\ \text{mm}^{-1}$
 $T = 293\ \text{K}$
 Block, colourless
 $0.20 \times 0.20 \times 0.20\ \text{mm}$

Data collection

Bruker SMART APEXII area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω and φ scans
 15699 measured reflections
 4347 independent reflections

2880 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$
 $\theta_{\text{max}} = 28.4^\circ$, $\theta_{\text{min}} = 1.7^\circ$
 $h = -9 \rightarrow 9$
 $k = -13 \rightarrow 13$
 $l = -17 \rightarrow 17$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.132$
 $S = 1.05$
 4347 reflections
 243 parameters
 0 restraints

Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0595P)^2 + 0.0997P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$

$$\Delta\rho_{\max} = 0.15 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.20 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O4	0.23016 (15)	0.81458 (9)	0.55793 (7)	0.0528 (3)
C14	0.2775 (2)	0.69318 (13)	0.41998 (10)	0.0473 (3)
H14	0.3173	0.6058	0.4713	0.057*
O5	0.31664 (18)	0.58412 (11)	0.28152 (9)	0.0699 (3)
H5	0.3065	0.6028	0.2151	0.105*
C16	0.1668 (2)	0.94571 (13)	0.37621 (11)	0.0498 (3)
H16	0.1326	1.0264	0.3987	0.060*
C15	0.22699 (19)	0.81361 (13)	0.45238 (10)	0.0423 (3)
C12	0.2087 (2)	0.83519 (14)	0.23261 (10)	0.0473 (3)
O6	0.2396 (2)	0.74928 (15)	0.07950 (9)	0.0831 (4)
C13	0.2683 (2)	0.70387 (14)	0.31045 (11)	0.0467 (3)
C11	0.1590 (2)	0.95453 (14)	0.26846 (11)	0.0511 (3)
H11	0.1196	1.0423	0.2175	0.061*
C18	0.2822 (3)	0.68218 (15)	0.63841 (11)	0.0639 (4)
H18A	0.4161	0.6290	0.6314	0.077*
H18B	0.1940	0.6296	0.6284	0.077*
C19	0.2680 (2)	0.70303 (15)	0.74570 (12)	0.0546 (4)
C17	0.1995 (2)	0.84683 (19)	0.11911 (12)	0.0643 (4)
H17	0.1594	0.9367	0.0712	0.077*
C20	0.2593 (3)	0.71369 (18)	0.83353 (13)	0.0652 (4)
O1	0.27472 (15)	0.21006 (10)	0.97384 (7)	0.0556 (3)
C4	0.2102 (2)	0.31928 (13)	0.77968 (10)	0.0480 (3)
H4	0.1643	0.4086	0.7881	0.058*
C6	0.3435 (2)	0.06902 (13)	0.85818 (11)	0.0475 (3)
H6	0.3856	-0.0081	0.9192	0.057*
C5	0.27430 (19)	0.20366 (13)	0.86926 (10)	0.0426 (3)
C3	0.2154 (2)	0.30031 (13)	0.67725 (10)	0.0458 (3)
C2	0.28516 (19)	0.16624 (13)	0.66383 (10)	0.0442 (3)
C1	0.3487 (2)	0.05198 (14)	0.75674 (11)	0.0480 (3)
H1	0.3956	-0.0377	0.7491	0.058*
O2	0.15146 (18)	0.41492 (10)	0.59107 (7)	0.0675 (3)
H2	0.1609	0.3914	0.5349	0.101*
O3	0.2389 (2)	0.23901 (13)	0.47217 (8)	0.0756 (4)

C7	0.2915 (2)	0.14587 (18)	0.55809 (12)	0.0594 (4)
H7	0.3396	0.0544	0.5542	0.071*
C9	0.2226 (2)	0.32905 (16)	1.10635 (12)	0.0576 (4)
C8	0.2136 (3)	0.34536 (16)	0.99018 (12)	0.0664 (4)
H8A	0.0797	0.3946	0.9582	0.080*
H8B	0.2996	0.3995	0.9556	0.080*
C10	0.2270 (3)	0.3192 (2)	1.19909 (14)	0.0684 (5)
H10	0.230 (3)	0.3082 (19)	1.2682 (16)	0.088 (6)*
H20	0.253 (2)	0.7250 (17)	0.9014 (15)	0.078 (5)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O4	0.0719 (7)	0.0437 (5)	0.0408 (5)	-0.0109 (4)	0.0031 (4)	-0.0153 (4)
C14	0.0586 (8)	0.0379 (6)	0.0406 (7)	-0.0080 (6)	0.0023 (6)	-0.0100 (5)
O5	0.0999 (9)	0.0533 (6)	0.0531 (6)	-0.0056 (6)	0.0018 (6)	-0.0275 (5)
C16	0.0560 (8)	0.0390 (7)	0.0518 (8)	-0.0098 (6)	0.0033 (6)	-0.0134 (6)
C15	0.0445 (7)	0.0427 (7)	0.0397 (6)	-0.0110 (5)	0.0038 (5)	-0.0137 (5)
C12	0.0456 (7)	0.0545 (8)	0.0400 (7)	-0.0149 (6)	0.0048 (6)	-0.0106 (6)
O6	0.1047 (10)	0.1030 (10)	0.0461 (6)	-0.0269 (8)	0.0113 (6)	-0.0326 (7)
C13	0.0496 (8)	0.0453 (7)	0.0448 (7)	-0.0095 (6)	0.0041 (6)	-0.0170 (6)
C11	0.0547 (8)	0.0428 (7)	0.0473 (7)	-0.0116 (6)	0.0008 (6)	-0.0025 (6)
C18	0.0978 (12)	0.0474 (8)	0.0417 (7)	-0.0147 (8)	-0.0004 (7)	-0.0118 (6)
C19	0.0653 (9)	0.0533 (8)	0.0459 (8)	-0.0192 (7)	0.0014 (7)	-0.0134 (6)
C17	0.0681 (10)	0.0793 (11)	0.0416 (8)	-0.0221 (8)	0.0062 (7)	-0.0110 (8)
C20	0.0841 (12)	0.0716 (10)	0.0446 (8)	-0.0277 (9)	0.0068 (8)	-0.0195 (8)
O1	0.0775 (7)	0.0464 (5)	0.0335 (5)	-0.0063 (5)	-0.0011 (4)	-0.0091 (4)
C4	0.0655 (9)	0.0367 (6)	0.0382 (7)	-0.0110 (6)	-0.0009 (6)	-0.0093 (5)
C6	0.0544 (8)	0.0390 (6)	0.0422 (7)	-0.0094 (6)	0.0008 (6)	-0.0051 (5)
C5	0.0475 (7)	0.0426 (7)	0.0342 (6)	-0.0106 (5)	0.0010 (5)	-0.0083 (5)
C3	0.0587 (8)	0.0418 (7)	0.0359 (6)	-0.0192 (6)	-0.0013 (6)	-0.0051 (5)
C2	0.0495 (7)	0.0477 (7)	0.0412 (7)	-0.0213 (6)	0.0059 (6)	-0.0150 (5)
C1	0.0528 (8)	0.0392 (7)	0.0524 (8)	-0.0121 (6)	0.0050 (6)	-0.0154 (6)
O2	0.1155 (10)	0.0468 (6)	0.0338 (5)	-0.0239 (6)	-0.0080 (5)	-0.0015 (4)
O3	0.1107 (10)	0.0851 (8)	0.0383 (6)	-0.0384 (7)	0.0061 (6)	-0.0197 (6)
C7	0.0737 (10)	0.0662 (9)	0.0500 (8)	-0.0290 (8)	0.0105 (7)	-0.0264 (7)
C9	0.0645 (10)	0.0590 (9)	0.0495 (8)	-0.0133 (7)	0.0022 (7)	-0.0212 (7)
C8	0.0962 (12)	0.0514 (8)	0.0438 (8)	-0.0067 (8)	0.0006 (8)	-0.0167 (6)
C10	0.0784 (12)	0.0844 (12)	0.0474 (9)	-0.0227 (9)	0.0045 (8)	-0.0286 (8)

Geometric parameters (\AA , $^\circ$)

O4—C15	1.3591 (14)	O1—C5	1.3634 (14)
O4—C18	1.4237 (16)	O1—C8	1.4245 (16)
C14—C15	1.3798 (17)	C4—C5	1.3797 (17)
C14—C13	1.3806 (18)	C4—C3	1.3825 (18)
C14—H14	0.9300	C4—H4	0.9300
O5—C13	1.3493 (15)	C6—C1	1.3619 (18)
O5—H5	0.8200	C6—C5	1.3935 (17)
C16—C11	1.3619 (18)	C6—H6	0.9300

C16—C15	1.3974 (17)	C3—O2	1.3481 (15)
C16—H16	0.9300	C3—C2	1.3999 (18)
C12—C11	1.3944 (19)	C2—C1	1.3977 (18)
C12—C13	1.4010 (19)	C2—C7	1.4304 (18)
C12—C17	1.4295 (19)	C1—H1	0.9300
O6—C17	1.221 (2)	O2—H2	0.8200
C11—H11	0.9300	O3—C7	1.2251 (19)
C18—C19	1.4522 (19)	C7—H7	0.9300
C18—H18A	0.9700	C9—C10	1.167 (2)
C18—H18B	0.9700	C9—C8	1.454 (2)
C19—C20	1.164 (2)	C8—H8A	0.9700
C17—H17	0.9300	C8—H8B	0.9700
C20—H20	0.911 (18)	C10—H10	0.862 (19)
C15—O4—C18	116.91 (10)	C5—O1—C8	117.38 (10)
C15—C14—C13	119.27 (12)	C5—C4—C3	119.00 (12)
C15—C14—H14	120.4	C5—C4—H4	120.5
C13—C14—H14	120.4	C3—C4—H4	120.5
C13—O5—H5	109.5	C1—C6—C5	119.09 (11)
C11—C16—C15	118.97 (12)	C1—C6—H6	120.5
C11—C16—H16	120.5	C5—C6—H6	120.5
C15—C16—H16	120.5	O1—C5—C4	123.96 (11)
O4—C15—C14	123.91 (11)	O1—C5—C6	114.78 (10)
O4—C15—C16	115.03 (11)	C4—C5—C6	121.26 (12)
C14—C15—C16	121.06 (12)	O2—C3—C4	117.85 (12)
C11—C12—C13	118.41 (12)	O2—C3—C2	121.27 (11)
C11—C12—C17	120.70 (13)	C4—C3—C2	120.88 (12)
C13—C12—C17	120.89 (13)	C1—C2—C3	118.26 (11)
O5—C13—C14	117.79 (12)	C1—C2—C7	120.55 (12)
O5—C13—C12	121.53 (12)	C3—C2—C7	121.19 (12)
C14—C13—C12	120.67 (12)	C6—C1—C2	121.50 (12)
C16—C11—C12	121.62 (12)	C6—C1—H1	119.2
C16—C11—H11	119.2	C2—C1—H1	119.2
C12—C11—H11	119.2	C3—O2—H2	109.5
O4—C18—C19	109.46 (11)	O3—C7—C2	125.35 (14)
O4—C18—H18A	109.8	O3—C7—H7	117.3
C19—C18—H18A	109.8	C2—C7—H7	117.3
O4—C18—H18B	109.8	C10—C9—C8	178.40 (17)
C19—C18—H18B	109.8	O1—C8—C9	108.67 (12)
H18A—C18—H18B	108.2	O1—C8—H8A	110.0
C20—C19—C18	177.13 (16)	C9—C8—H8A	110.0
O6—C17—C12	125.79 (15)	O1—C8—H8B	110.0
O6—C17—H17	117.1	C9—C8—H8B	110.0
C12—C17—H17	117.1	H8A—C8—H8B	108.3
C19—C20—H20	178.1 (11)	C9—C10—H10	177.5 (13)
C18—O4—C15—C14	-2.4 (2)	C8—O1—C5—C4	-3.0 (2)
C18—O4—C15—C16	177.32 (13)	C8—O1—C5—C6	177.38 (12)
C13—C14—C15—O4	179.71 (12)	C3—C4—C5—O1	-179.63 (12)

C13—C14—C15—C16	0.0 (2)	C3—C4—C5—C6	0.0 (2)
C11—C16—C15—O4	-179.91 (12)	C1—C6—C5—O1	180.00 (12)
C11—C16—C15—C14	-0.2 (2)	C1—C6—C5—C4	0.4 (2)
C15—C14—C13—O5	-179.31 (12)	C5—C4—C3—O2	179.81 (13)
C15—C14—C13—C12	0.0 (2)	C5—C4—C3—C2	-0.3 (2)
C11—C12—C13—O5	179.36 (13)	O2—C3—C2—C1	-179.83 (13)
C17—C12—C13—O5	-0.9 (2)	C4—C3—C2—C1	0.3 (2)
C11—C12—C13—C14	0.0 (2)	O2—C3—C2—C7	0.1 (2)
C17—C12—C13—C14	179.81 (13)	C4—C3—C2—C7	-179.79 (13)
C15—C16—C11—C12	0.3 (2)	C5—C6—C1—C2	-0.4 (2)
C13—C12—C11—C16	-0.2 (2)	C3—C2—C1—C6	0.1 (2)
C17—C12—C11—C16	-179.99 (13)	C7—C2—C1—C6	-179.87 (13)
C15—O4—C18—C19	-177.04 (12)	C1—C2—C7—O3	179.84 (15)
O4—C18—C19—C20	-179 (100)	C3—C2—C7—O3	-0.1 (2)
C11—C12—C17—O6	-179.95 (16)	C5—O1—C8—C9	179.39 (13)
C13—C12—C17—O6	0.3 (3)	C10—C9—C8—O1	-162 (7)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2 \cdots O3	0.82	1.92	2.6387 (16)	146
O5—H5 \cdots O6	0.82	1.93	2.6441 (16)	146
C10—H10 \cdots O3 ⁱ	0.86 (2)	2.51 (2)	3.369 (2)	171.4 (2)
C18—H18 ^A \cdots O5 ⁱⁱ	0.97	2.45	3.281 (2)	144
C20—H20 \cdots O6 ⁱ	0.91 (2)	2.37 (2)	3.280 (2)	178.8 (2)

Symmetry codes: (i) $x, y, z+1$; (ii) $-x+1, -y+1, -z+1$.