

# *Entamoeba histolytica* meningoencephalitis diagnosed by trophozoites in cerebrospinal fluid

L. M. L. Goh<sup>1</sup> and J. R. Marrone<sup>1</sup>

1) Pediatric Department, Lyndon Baines Johnson Tropical Medical Center, Pago Pago, AS 96799, American Samoa

## Abstract

*Entamoeba histolytica* meningoencephalitis has not been described in the modern literature, which is distinct from that caused by free-living amoebae. We report the first case of *E. histolytica* meningoencephalitis without liver or brain abscesses. Cerebrospinal fluid revealed 2+ very motile trophozoites. Our patient was successfully treated with intravenous metronidazole.

**Keywords:** Amoebiasis, cerebrospinal fluid, *Entamoeba histolytica*, meningoencephalitis, trophozoites

**Original Submission:** 31 May 2013; **Accepted:** 5 July 2013

**Article published online:** 18 September 2013

*New Microbe New Infect* 2013; **1**: 16–17

**Corresponding author:** L. M. L. Goh, Pediatric Department, Lyndon Baines Johnson Tropical Medical Center, Pago Pago, AS 96799, American Samoa.

**E-mail:** lucy.goh@bj.as

Poster presentation by LMLG at the 61st Annual Meeting of the American Society of Tropical Medicine and Hygiene in Atlanta, GA, 11–15 November 2012.

## Case Report

A 16-month-old Polynesian boy presented to the emergency room with a 4-day history of cough, dyspnoea, wheezing, anorexia, fever and a 1-day history of non-bloody, mucoid diarrhoea. A month earlier, he had undergone surgical debridement of scabies-induced pyoderma of his feet. He also had recurrent otorrhoea.

Pertinent physical findings were mild respiratory distress with retractions (respiratory rate of 50 breaths/min, pulse oximetry of 98% on room air), wheezing, normal abdominal

and neurological examinations and healing sores on anterior right foot. Laboratory abnormalities included marked leucocytosis of  $45.2 \times 10^9/L$ —89% neutrophils, haemoglobin 1.09 mmol/L, haematocrit 0.228, platelets  $630 \times 10^9/L$  and electrolyte disturbances. He was hospitalized for bronchiolitis, acute gastroenteritis and severe dehydration. Ampicillin, crystalloids, bronchodilators and methylprednisolone were given.

On day 2, his abdomen became very distended with absent bowel sounds. Abdominal X-ray revealed air–fluid levels. He was transferred to the intensive care unit for an acute abdomen and anuria. Ceftriaxone and metronidazole were added. Blood culture grew *Streptococcus pyogenes*; a repeat culture was negative. Computed tomography of abdomen without contrast showed ascites without free air.

On day 3, an exploratory laparotomy revealed ascites of 700 mL, copious peritoneal fibrinous exudates throughout, inflamed liver without abscess and 15 cm of inflamed jejunum with petechiae. Peritoneal fluid culture grew *S. pyogenes*. He remained intubated and critically ill.

On day 4, stool returned positive for *Entamoeba histolytica* 2+ cysts—he was switched to oral metronidazole. Stool cultures were negative. On day 6, he received a dose of dexamethasone on self-extubation.

On day 8, he developed tonic–clonic seizures. Computed tomography with/without contrast of the brain and abdomen revealed no intracranial abnormality or liver abscess. Subsequently, he had prolonged seizures. When he was stabilized, a lumbar puncture revealed bloody cerebrospinal fluid (CSF) with leucocytes  $3/\mu L$ , erythrocytes  $3096/\mu L$ , protein 1.73 g/L, glucose 3.6 mmol/L, negative Gram stain and culture; direct wet mount revealed 2+ very motile trophozoites. He received intravenous metronidazole for 2 weeks. No further seizures were noted. On follow up he was doing well without any sequelae.

Invasive amoebiasis causes extra-intestinal disease via haematogenous spread to the liver, lung, brain and rarely other organs [1]. Legrand described 24 cases and Lombardo *et al.* described 17 cases of cerebral amoebiasis associated with liver abscesses [2,3]; neither mentioned finding trophozoites in the CSF. A case of *E. histolytica* liver abscess with encephalitis was diagnosed by PCR of the CSF, which was negative for trophozoites [4]. Our patient would be the first case of *E. histolytica* meningoencephalitis without liver or brain abscesses, with trophozoites in the CSF.

Computed tomography can be negative in the early stages of encephalitis and cell count in the CSF may be falsely low, if measured more than 30 minutes after the lumbar puncture is performed [5]. Several risk factors undoubtedly contributed to the development of fulminant invasive amoebiasis: very young

age, malnutrition, compromised innate immunity and treatment with high-dose corticosteroids [1].

Hence, in endemic areas, clinicians need to be actively looking for cerebral amoebiasis in a child with intestinal amoebiasis and new-onset seizures. It may be under-recognized because of a lack of clinical suspicion and of laboratory capacity.

## Acknowledgements

---

We thank Lana Flippo, Kamlesh Kumar and Amelita Mejia for caring for the patient in the Intensive Care Unit, Tag'ilima l'atala and Saipale Fuimaono for caring for the patient in the emergency room, Seini Biukoto for reading the CSF wet mount, Winnie Ooi of Lahey Clinic Medical Center, Boston, for the archived copy of the 1912 French article and Eric Mouzin of World Health Organization, Geneva, for interpreting it, Dorothy Jarzabek of VA Hospital, Shreveport, for copies of several references and, Barnett Cline of Tulane School of Public Health and Tropical Medicine, New Orleans, for his invaluable input, suggestions and critique.

## Conflicts of Interest

---

None declared.

## References

---

1. Petri WA Jr, Haque R. *Entamoeba* species, including amebiasis. In: Mandell GL, Bennett JE, Dolin R, eds, *Principles and practice of infectious diseases*, 7th edn. Philadelphia, PA: Elsevier Churchill Livingstone, 2009; 3411–3426.
2. Legrand H. Les abcès dysentériques du cerveau (amibiase encéphalique). *Arch Prov Chirurg* 1912; 75: 213–239.
3. Lombardo L, Alonso P, Arroyo LS, Brandt H, Mateos JH. Cerebral Amebiasis: Report of 17 cases. *J Neurosurg* 1964; 21: 704–709.
4. Soleymani-Mohammadi S, Lam MM, Zunt JR, Petri WA. *Entamoeba histolytica* encephalitis diagnosed by PCR of cerebrospinal fluid. *Trans R Soc Trop Med Hyg* 2007; 101: 311–313.
5. Greenlee JE, Carroll KC. Cerebrospinal fluid in central nervous system infections. In: Scheld WM, Whitley RJ, Marra CM, eds, *Infections of the central nervous system*, 3rd edn. Philadelphia, PA: Lippincott Williams & Wilkins, 2004; 6–30.