

**Methods:** A community sample of 318 parents (64.2% mothers, mean age: 39.48 years SD=5.82) filled out an online questionnaire including the Decision-Making Involvement Scale (DMIS) assessing the parent's and child's behaviour in decision-making processes and the Strength and Difficulties Questionnaire (SDQ). Linear regression analyses were conducted for exploring multivariate associations of DMIS Parent and Child subscales with child's psychopathology and prosocial behaviour, controlling for child's gender and age.

**Results:** With SDQ Internalizing problems subscale as dependent, linear regression analysis did not result in a significant model. In a significant model explaining 21.2% of the variance of the dependent variable, SDQ Externalizing problems score were significantly related to child's age and gender, and to both Child and Parent subscales of the DMIS. When choosing SDQ Prosocial behaviour subscale as dependent, child's gender and DMIS Child subscale were significantly associated to the dependent variables, accounting for 12.2% of the variance.

**Conclusions:** Our results suggest that children's involvement in decisions may be related to less externalizing symptoms and higher levels of prosocial skills. However, longitudinal research is needed to uncover the direction of the relationship and underlying mechanisms.

**Disclosure:** No significant relationships.

**Keywords:** decision-making; externalizing and internalizing symptoms; child and youth; participation

## EPV1117

### Religious End-World delusion: psychopathological types

E. Gedeveni<sup>1</sup>, G. Kopeiko<sup>1\*</sup>, O. Borisova<sup>1</sup>, P. Orekhova<sup>2</sup> and V. Kaleda<sup>3</sup>

<sup>1</sup>FSBSI Mental Health Research Center, Researching Group Of Specific Forms Of Mental Disorders, Moscow, Russian Federation; <sup>2</sup>Federal State Budgetary Scientific Institution «Mental Health Research Center», Researching Group Of Specific Forms Of Mental Disorders, Moscow, Russian Federation and <sup>3</sup>Mental Health Research Centre, Department Of Youth Psychiatry, Moscow, Russian Federation

\*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1810

**Introduction:** Diagnostics of End-World delusion with religious content (EWDRC) is relevant due to its insufficient exploration, difficulty in differential diagnostics and social danger of the delusional behavior.

**Objectives:** To develop a typology based on psychopathological and phenomenological features.

**Methods:** Sixty patients with EWDRC were examined. Psychopathological and statistical methods were applied.

**Results:** Study of EWDRC found heterogeneity of clinical appearances. Two different types were identified: apocalyptic and eschatological. **The apocalyptic type** (51 patients, 85%) was characterized by prevalence of End-World ideas in an acute sensual delusion. Due to heterogeneity of delusion's dynamics two subtypes were identified: - Subtype 1 (31 patients, 61%) was characterized with long period of development (changes in the stages) of different delusion's types: delusion of perception, importance, staging, and the antagonistic one. Psychotic symptoms were quickly reduced with antipsychotic therapy. - Subtype 2 (20 patients, 39%) was charac-

terized with rapid development of delusion's stages up to oneiro-catatonic states which were hardly jugulated. **Eschatological type** (9 patients, 15%) was characterized by the systematized interpretive delusion with individual interpretation of apocalyptic signs. These states evolved within mixed forms of schizophrenia.

**Conclusions:** The analysis of EWDRC revealed the apocalyptic type's acute course. Patients with the apocalyptic type have a premonition of upcoming End-World, and feel themselves engaged in it. The eschatological type is based on the systematized interpretive End-World delusion with "confirmations" found in everyday life. The results showed the high risk of the delusional behavior in patients with EWDRC which requires careful approach to the diagnostics and treatment of these conditions.

**Disclosure:** No significant relationships.

**Keywords:** religious delusion; End-World delusion; delusional behavior; schizofrenia

## EPV1118

### Shame and psychopathology. Its role in the genesis and perpetuation of different disorders

F. Azevedo<sup>1\*</sup>, R. André<sup>2</sup>, A. Quintão<sup>3</sup>, D. Jeremias<sup>1</sup> and C. Almeida<sup>4</sup>

<sup>1</sup>Centro Hospitalar de Lisboa Ocidental, Psychiatry, Lisbon, Portugal;

<sup>2</sup>Centro Hospitalar Universitário Lisboa Norte, Psychiatry, Lisboa,

Portugal; <sup>3</sup>Centro Hospitalar de Lisboa Ocidental, Psychiatry

Department, Lisboa, Portugal and <sup>4</sup>NOVA Medical School, Psychiatry And Mental Health, Lisboa, Portugal

\*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1811

**Introduction:** Shame is a profound negative emotion that can sometimes be covered up by guilt and remain undiagnosed. Shame and guilt have been described as self-conscious and moral emotions as they both involve self-evaluation and lay a role in facilitating moral conduct. They derive from the notion of responsibility, but some authors suggest that while guilt focuses only on the act at hand shame focuses on the one executing it. The self is the object.

**Objectives:** To review the literature on shame and its role in different disorders both as a causing agent and as a perpetuating agent

**Methods:** Non-systematic review of the literature with selection of scientific articles published in the past 20 years; by searching Pubmed and Medscape databases using the combination of MeSH descriptors. The following MeSH terms were used: "shame", "psychopathology".

**Results:** Since shame globally decreases self-esteem and is an awareness of personal flaws it can lead to the feelings of helplessness and the development or worsening of mental disorders. As such it is no wonder to find shame being studied in many different forms, more and less structured with important connections being made with social anxiety, eating disorders, dysmorphic disorders, personality disorders and bereavement.

**Conclusions:** Shame's role, independently from guilt can have an impact on both the genesis and perpetuation of mental disorders. Its study can uncover missing links between different types of experiences and the pathological reactions that may subsequently follow.

**Disclosure:** No significant relationships.

**Keywords:** Psychopathology; Disorders; shame