

# Autophagy and bacterial infectious diseases

Jae-Min Yuk<sup>1,2</sup>, Tamotsu Yoshimori<sup>3</sup>  
and Eun-Kyeong Jo<sup>1,2,4</sup>

<sup>1</sup>Department of Microbiology and

<sup>2</sup>Infection Signaling Network Research Center  
Chungnam National University School of Medicine  
Daejeon 301-747, Korea

<sup>3</sup>Department of Genetics  
Graduate School of Medicine

Osaka University  
Osaka 565-0871, Japan

<sup>4</sup>Corresponding author: Tel, 82-42-580-8243;  
Fax, 82-42-580-8436; E-mail, hayoungj@cnu.ac.kr  
<http://dx.doi.org/10.3858/emmm.2012.44.2.032>

Accepted 30 December 2011

Available Online 19 January 2012

Abbreviations: ATG, autophagy-related protein; eis, enhanced intracellular survival; GABARAP,  $\gamma$ -aminobutyric acid receptor-associated protein; GAS, group A *Streptococcus*; Gbp, guanylate-binding protein; LC3, light chain 3; *M. tuberculosis*, *Mycobacterium tuberculosis*; NF, nuclear factor; PAMP, pathogen associated molecular pattern; ROS, reactive oxygen species reactive oxygen species; SCVs, *Salmonella*-containing vacuoles; SLAPs, spacious *Listeria*-containing phagosomes; *S. Typhimurium*, *Salmonella enterica* serovar Typhimurium; T3SSs, type three secretion systems; Th, T helper; TRAF6, tumor necrosis factor receptor-associated factor 6

## Abstract

**Autophagy is a housekeeping process that maintains cellular homeostasis through recycling of nutrients and degradation of damaged or aged cytoplasmic constituents. Over the past several years, accumulating evidence has suggested that autophagy can function as an intracellular innate defense pathway in response to infection with a variety of bacteria and viruses. Autophagy plays a role as a specialized immunologic effector and regulates innate immunity to exert antimicrobial defense mechanisms. Numerous bacterial pathogens have developed the ability to invade host cells or to subvert host autophagy to establish a persistent infection. In this review, we have summarized the recent advances in our understanding of the interaction between antibacterial autophagy**

**(xenophagy) and different bacterial pathogens.**

**Keywords:** autophagy; cytokines; immunity, Innate; infection; reactive oxygen species

## Introduction

During macroautophagy (herein referred to as autophagy), cytoplasmic contents can be sequestered in a unique double-membrane structure, the autophagosome. Through autophagy, cytoplasmic cargo in autophagosomes can be degraded by fusion of autophagosomes and lysosomes, and activation of the lysosomal degradation pathway (Levine and Deretic, 2007). In addition to degradation of damaged organelles or materials, autophagy has received attention as a crucial component of innate defense against a variety of infectious agents. Autophagy of foreign micro-organisms, i.e., xenophagy, has emerged as a powerful method of eliminating intracellular bacteria (Gutierrez *et al.*, 2004; Nakagawa *et al.*, 2004). However, there is a complex interplay between host autophagy mechanisms and pathogens. Numerous microorganisms have evolved strategies to evade or subvert host autophagy to survive and establish a persistent infection (Campoy and Colombo, 2009b; Orvedahl and Levine, 2009). Thus, identification of mechanisms or virulence factors exploiting autophagy may provide a new strategy for therapeutic intervention in infectious diseases.

Host defense against pathogens requires coordination of multiple innate immune signaling pathways (Jo *et al.*, 2007). During infection, host cells recognize the pathogen-associated molecular patterns of a variety of microbes through the expression of various pattern recognition receptors. The recognition of foreign or danger molecules by innate receptors can trigger an intracellular signaling cascade, leading to activation of antimicrobial effector mechanisms to promote clearance of the infection (Jo *et al.*, 2007; Brodsky and Medzhitov, 2009). Autophagy is considered to be one of the effector mechanisms downstream of these receptors, and plays an integral role in both innate and adaptive immunity to various pathogens (Orvedahl and Levine, 2009).

Autophagy also crosstalks with intracellular sig-

naling molecules and effectors (Brodsky and Medzhitov, 2009; Shahnazari and Brumell, 2011). Through these interactions, autophagy carries out not only direct microbial degradation but also other protective mechanisms, such as lysozyme secretion, the ubiquitin-mediated pathway, and antigen presentation (Brodsky and Medzhitov, 2009; Shahnazari and Brumell, 2011). In particular, the autophagic or cargo receptors have received much attention, since they play an important role in transporting ubiquitinated microbial cargos to the autophagy machinery (Pankiv *et al.*, 2007; Ichimura *et al.*, 2008; Kirkin *et al.*, 2009b; Thurston *et al.*, 2009; von Muhlinen *et al.*, 2010).

Recent studies have provided evidence that autophagy acts as a 'tuning module' in the regulation of innate immunity to prevent excessive inflammatory responses and inflammasome signaling (Sumpter and Levine, 2010). Generation of reactive oxygen species (ROS) of cellular and mitochondrial origin is thought to play an important role in autophagy regulation, thus influencing innate defense (Azad *et al.*, 2009; Scherz-Shouval and Elazar, 2011).

The specific role and regulatory mechanisms of autophagy in connection with innate immune pathways during infection will be discussed here. Additionally, the molecular mechanisms by which multiple bacteria and viruses evade and/or resist autophagy will be covered in this review.

### **Xenophagy: autophagic control of intracellular bacterial pathogens**

Autophagy is a mechanism for adjusting cellular quality and quantity through capture and degradation of portions of the cytosol or organelles in response to diverse stress or stimuli, including infection (Deretic and Levine, 2009). Autophagy is a means of bulk degradation of cytoplasmic components within lysosomes. During this process, cytoplasmic constituents are delivered to lysosomes for degradation (Mizushima and Levine, 2008). Although autophagy is fundamentally a 'self-eating' process, it also facilitates degradation of pathogens as a 'xenophagy' process (Huang and Brumell, 2009).

Accumulating evidence indicates that autophagy acts as a defense mechanism against multiple invading microbes (Deretic and Levine, 2009). During infection, host autophagic responses can target several steps of bacterial invasion. For example, autophagy can target bacteria within vacuolar membranous compartments. Intracellular pathogens, such as *Mycobacterium tuberculosis* (*M. tuberculosis*), can actively survive within host cells and exploit

host defense by inhibition of phagosomal maturation (Gutierrez *et al.*, 2004; Yuk *et al.*, 2009b). Autophagy activation by nutrient starvation or rapamycin treatment leads to co-localization of mycobacterial phagosomes with autophagosomes to overcome the inhibition of phagosomal maturation by mycobacteria, thereby resulting in suppression of intracellular bacterial survival (Gutierrez *et al.*, 2004; Yuk *et al.*, 2009b). Stimulation of host autophagy by treatment with vitamin D significantly enhanced the antimicrobial responses against *M. tuberculosis* in human macrophages. This effect was mediated by cathelicidin, a peptide that is induced by vitamin D and promotes co-localization of bacterial phagosomes and autophagosomes (Yuk *et al.*, 2009b). These data suggest that autophagy contributes to elimination of bacteria by overcoming bacterial resistance to lysosomal access.

Another example is group A *Streptococcus* (GAS). When GAS enters epithelial cells *via* endocytosis, they can escape from the endosomes to the cytoplasm by secretion of streptolysin O, a pore-forming toxin (Nakagawa *et al.*, 2004). GAS is then trapped in autophagosome-like compartments, which mature into autolysosomes and so eliminate the bacteria (Nakagawa *et al.*, 2004). Mechanistically, the host small G proteins, Rab5 and Rab7, are associated with autophagosome formation and the progression of endosome maturation in cells infected with GAS (Sakurai *et al.*, 2010). Two endocytic soluble N-ethylmaleimide-sensitive factor attachment protein receptors, VAMP8 from lysosomes and Vti1b from autophagic compartments, are required for the successful fusion of GAS-containing xenophagosomes with lysosomes, which facilitates xenophagy (Furuta *et al.*, 2010).

The third mode of anti-bacterial action of autophagy is receptor-mediated recruitment of bacteria to autophagosomes. Some populations of *Salmonella enterica* serovar Typhimurium (*S. Typhimurium*) are known to invade mammalian cells and replicate in *Salmonella*-containing vacuoles (SCVs). During this process, two kinds of *Salmonella* pathogenicity island (SPI) encoding type III secretion systems (T3SSs) are involved: the SPI-2 encoded T3SS facilitates bacterial growth, whereas the SPI-1 encoded T3SS causes damage to SCV (Birmingham *et al.*, 2006), which activates the autophagic process. The autophagy system targets the intracellular bacteria present within damaged SCVs, and contributes to restriction of bacterial growth during infection (Birmingham *et al.*, 2006). The autophagy receptor p62/SQSTM1, an adaptor protein for degradation of cargo tagged with ubiquitinated protein, is recruited to autophagy-targeted *S. Typhimurium* and required for the

restriction of intracellular bacterial replication through induction of efficient antibacterial autophagy (Zheng *et al.*, 2009). The autophagy receptor NDP52 can also target distinct compartments in cells infected with *Salmonella*, and is required for autophagy activation (Cemna *et al.*, 2011). These data suggest that xenophagy is required for efficient targeting and killing of cytosolic pathogens. The post-translational modification (ubiquitination) of intracellular pathogens and the roles of cargo receptors will be discussed in detail in the latter part of this review.

### Cross-talk of autophagy with innate immune signaling pathways

Apart from its initial description, which is that autophagy is a response to amino acid deprivation, to recycle cytoplasmic constituents and synthesize energy for cell survival under stressful/starved conditions, autophagy is also being recognized as an innate immune process that is activated upon recognition of microbial infection (Brodsky and Medzhitov, 2009; Shahnazari and Brumell, 2011). Bacteria and viruses possess molecules termed 'pathogen associated molecular patterns' (PAMPs) that bind to pattern-recognition receptors such as toll-like receptors (TLRs) on innate immune cells. Ligation of TLRs with PAMPs is known to activate innate immune cells and subsequently facilitate adaptive immune responses. So far, 10 TLRs in humans and 12 in mice have been identified. On the basis of subcellular localization, TLRs can be divided into extracellular and intracellular receptors: TLR1, 2, 4, 5, 6, and 10 are located on the cell surface. In contrast, TLR3, 7, 8, and 9 are expressed in intracellular endosomal/lysosomal compartments and the endoplasmic reticulum (Manavalan *et al.*, 2011). Bacterial PAMPs are recognized by five TLRs in humans: lipopolysaccharide (LPS) is the main ligand of Gram-negative bacteria for TLR4; lipoteichoic acid and diacylated lipopeptides are sensed by TLR2/6; triacylated lipopeptides are sensed by TLR2/1; CpG motifs are sensed by TLR9; and flagellin is sensed by TLR5 (Takeuchi and Akira, 2010). Interestingly, stimulation of TLR2, 4, and 7 with their ligands can stimulate autophagy, which functions to eliminate intracellular mycobacteria (Xu *et al.*, 2007; Shi and Kehrl, 2008; Delgado and Deretic, 2009; Shin *et al.*, 2010b). Thus, TLR responses, which are well-known immune-activating signals, can also directly activate autophagy resulting in killing of invading bacteria.


In TLR signaling, several TIR domain-containing adaptor molecules, such as MyD88, TIRAP, and

TRIF, lie downstream of TLR and deliver activation signals. (Takeda and Akira, 2004). Upon stimulation of TLR4 with LPS, MyD88, TRIF, and tumor necrosis factor receptor (TNFR)-associated factor 6 (TRAF6), associate with ubiquitinated autophagy-related protein (Atg)6/Beclin-1, which results in activation of autophagy (Shi and Kehrl, 2010). The roles of further downstream signaling molecules of the TLR signaling pathway in the regulation of autophagy remain to be determined. Among them, the role of nuclear factor (NF)- $\kappa$ B, an important downstream mediator of TLR signaling (Takeda and Akira, 2004), in autophagy is still being debated. Although the NF- $\kappa$ B signaling pathway inhibits the autophagy pathways in tumor cells (Djavaheri-Mergny *et al.*, 2006) and keratinocytes (Lee *et al.*, 2011), this has not been confirmed in primary innate cells, such as macrophages or dendritic cells. Mitogen-activated protein kinase signaling pathways may play a role in autophagy, since they are important in the activation of innate immune signaling during bacterial infection (Jo *et al.*, 2007). The stress-activated signaling molecule, c-Jun N-terminal protein kinase 1 (JNK1), pathway has been known to activate starvation- or ceramide-induced autophagy through Bcl-2 phosphorylation and subsequent dissociation of Bcl-2 from Beclin-1 (Wei *et al.*, 2008; Pattingre *et al.*, 2009). More comprehensive data are necessary to gain a complete understanding of the roles and regulatory mechanisms of signaling modules in antibacterial autophagic activation.

Stimulation of other pattern recognition receptors, including the cytosolic receptor NOD2, regulates autophagy. NOD2 receptor recognition by muramyl dipeptide induces autophagy in dendritic cells through receptor-interacting serine-threonine kinase-2, ATG5, ATG7 and ATG16L1 (Cooney *et al.*, 2010). Notably, NOD2-induced autophagy activation is required for both direct antimicrobial activities and antigen presentation to T cells via MHC class II antigen to induce adequate CD4<sup>+</sup> T cell responses (Cooney *et al.*, 2010). These studies suggest that specific PAMP recognition through membrane and cytosolic innate receptors can regulate innate and adaptive immune responses through autophagy activation.

### Immune effectors and autophagy regulation: cytokines, IRG proteins (p47 GTPases), and cathelicidins

Numerous cytokines and chemokines, such as interleukin (IL)-12, IL-23, IL-17, interferon (IFN)- $\alpha$ , IFN- $\beta$ , IFN- $\gamma$ , IL-10, IL-27, IL-6, TNF, lymphotoxin, and IL-1 family cytokines, are produced by innate



**Figure 1.** The role of autophagy in control of innate immunity and the survival mechanisms of intracellular bacteria. Various intracellular bacteria are internalized by phagocytes; these survive in the cytosol or phagosome through various strategies. (1) First, some bacteria are able to escape from the phagosome by secreting factors and replicating in the host cytoplasm (left panel). When autophagy is induced, *Listeria monocytogenes* and *Rickettsia conorii* present in the cytoplasm are captured by the autophagosomal membrane and degraded by the autolysosome; however, *Shigella flexneri* has the ability to avoid import into the autophagosome. (2) Second, some bacteria reside in the phagosomal compartment (right panel). Upon autophagy activation, phagosomes containing *Mycobacterium tuberculosis* are caught by the autophagic vacuole and then degraded by fusion with the lysosome. In contrast, *Brucella abortus* and *Legionella pneumophila* can inhibit autophagosome maturation through inhibition of lysosome fusion, allowing the pathogen to survive and replicate.

immune cells during bacterial infection (Cooper *et al.*, 2011; Deretic, 2011). Notably, several cytokines are key regulators of autophagy pathways (Deretic 2009). For example, the Th1 cytokine IFN- $\gamma$  is essential for host defense against intracellular pathogens. IFN- $\gamma$  can overcome the blocking of phagosomal maturation induced by *M. tuberculosis* through autophagy activation (Gutierrez *et al.*, 2004; Harris *et al.*, 2009). In contrast, the T helper (Th) 2 cytokines IL-4 and IL-13 contribute to inhibition of autophagy and autophagy-mediated suppression of mycobacteria in macrophages (Harris *et al.*, 2007).

The mechanisms by which IFN- $\gamma$  activates autophagy are: IFN- $\gamma$  activates several anti-bacterial proteins involved in autophagy, such as IRG (p47 GTPases) and the 65-kilodalton (kD) guanylate-binding protein (Gbp) gene family members (Gbp1, Gbp6, Gbp7, and Gbp10). The IRG proteins (p47

GTPases), transcription of which is induced by IFN- $\gamma$  stimulation, play an essential role in the elimination of intracellular pathogens in mice (Deretic, 2009). Murine IRG proteins are involved in innate defense against *M. tuberculosis* and *Toxoplasma gondii* (Gutierrez *et al.*, 2004; Ling *et al.*, 2006). Although the single human IRG member, IRGM, is not induced by IFN- $\gamma$ , it is essential for induction of autophagy in response to intracellular bacterial infection (Singh *et al.*, 2006). Moreover, recent studies have revealed that the Gbp gene family members (Gbp1, Gbp6, Gbp7, and Gbp10) are involved in oxidase-dependent killing and autophagy pathways in phagocytes to combat listerial or mycobacterial infection. Thus, 65-kD Gbps function to coordinate a potent oxidative and vesicular trafficking program to protect the host from infection (Kim *et al.*, 2011). Reciprocally, autophagy facilitates IFN- $\gamma$ -activated Jak2-STAT1

pathways and inflammatory responses, suggesting a link between autophagy and cellular inflammation (Chang *et al.*, 2010).

Antimicrobial peptides are major components of the bactericidal machinery of mammalian phagocytes, and play an important role as innate effectors during bacterial and viral infections (Cederlund *et al.*, 2011). Of the antimicrobial peptides, cathelicidins exert antimicrobial activities against mycobacteria (Liu *et al.*, 2006; Yang *et al.*, 2009; Sonawane *et al.*, 2011). Interestingly, vitamin D activates anti-mycobacterial autophagy through induction of cathelicidin, which contributes to the promoter activation of the autophagy-related genes Atg5 and Beclin-1, as well as co-localization of bacterial phagosomes and autophagosomes (Yuk *et al.*, 2009b). Future studies are necessary to fully elucidate the roles of the numerous antimicrobial peptides in the context of autophagy activation leading to collaboration with innate immunity.

## Bacterial tactics to modulate host autophagy

Intracellular bacterial pathogens have developed multiple strategies for exploiting or hijacking eukaryotic functions in order to survive and multiply inside host cells (Knodler *et al.*, 2001). Consistently, numerous pathogens antagonize the initiation and maturation of autophagic processes, evade autophagic recognition, or hijack the autophagy pathway to facilitate intracellular survival or replication (Levine *et al.*, 2011).

The various strategies used by bacterial pathogens are summarized in Figure 1. In this section, we will summarize two strategies by which bacterial pathogens can escape from or interact with autophagic pathways: 1) physical escape from degradative endocytic compartments and replication within the cytoplasm (*Shigella*, *Listeria*, *Rickettsia*); and 2) survival and replication within membrane-bound compartments (*Mycobacteria*, *Salmonella*, *Francisella*, *Legionella*, *Brucella*, *Chlamydia*, *Porphyromonas gingivalis*).

### Bacterial escape into cytosol after breaking the host jail (bacterial endocytic compartments): *Shigella*, *Rickettsia*, *M. marinum*, and *Listeria*

Intracellular bacteria such as *Shigella flexneri* and *Listeria monocytogenes* can evade autophagic recognition, survive, and replicate in phagocytic and non-phagocytic cells by disrupting vacuoles and escaping into the cytoplasm (Ogawa *et al.*, 2005), and acquisition of actin-based motilities

(Cossart, 2000), respectively. *S. flexneri* can block autophagic targeting through the secreted T3SS effector *icsB* that competes with the autophagy protein Atg5 for binding to the *S. flexneri* surface protein VirG (Ogawa *et al.*, 2005). Thus, *icsB* deletion mutants are not able to survive in host cells (Ogawa *et al.*, 2005). When *S. flexneri* breaks the vacuolar membrane and escapes from the phagosome, the vacuolar membrane fragments are polyubiquitinated, and targeted to autophagic degradation through p62 accumulation on membrane remnants (Dupont *et al.*, 2009).

*Rickettsia* are Gram-negative obligate intracellular bacteria, and can be transmitted to humans *via* arthropod vectors (Azad and Beard, 1998). After infection, *Rickettsia* can escape from phagosomes into their eukaryotic host cytoplasm, where they replicate and infect adjacent cells (Silverman and Wisseman, 1979; Walker, 2007). Earlier studies reported that rickettsial infection of polymorphonuclear leukocytes results in autophagosome formation (Rikihisa, 1984). The detailed mechanisms by which *Rickettsia* are able to escape from autophagy machinery remain to be determined. Conversely, the host autophagy induced during rickettsial infection is recognized as an important defense mechanism against this invading pathogen (Walker, 2007; Deretic, 2010). One report indicated that *Rickettsia*-induced proinflammatory cytokines led to increased activation of the autophagy that is required for bacterial inhibition (Walker *et al.*, 1997).

*Mycobacterium marinum*, a natural pathogen, and the closest relative of *M. tuberculosis* with a homology of 99.4% (Stamm and Brown, 2004), has been shown to escape from the phagosome and enter the cytosol *via* an ESX-1-dependent mechanism (Collins *et al.*, 2009). In the cytosol, some *M. marinum* promote motility through actin polymerization, while non-motile bacteria are decorated by ubiquitin and sequestered in the host Lamp1-positive vesicular compartments by autophagosome-like double membranes, formation of which is independent of Atg5 (Collins *et al.*, 2009).

*L. monocytogenes* can replicate in LAMP1-positive spacious *Listeria*-containing phagosomes (SLAPs), the formation of which is dependent on both the bacterial virulence toxin listeriolysin O and host autophagy (Birmingham *et al.*, 2008). Interestingly, the listerial virulence factor *actA* triggers host protein recruitment and actin polymerization, which propels bacteria from one cell to another (Cossart, 2000) and prevents autophagic recognition of bacteria within the cytosol (Yoshikawa *et al.*, 2009). *L. monocytogenes* expressing-*actA* mutants that cannot recruit host proteins for actin polymerization

are eventually ubiquitinated, and then recognized by the autophagy machinery through recruitment of p62 (also known as SQSTM1) and LC3 (Yoshikawa *et al.*, 2009). Notably, *L. monocytogenes* expressing listeriolysin O, the pore-forming cytolysin, can escape into the cytosol, where *L. monocytogenes* can be targeted for antibacterial autophagy (Huang and Brumell, 2009). Moreover, *L. monocytogenes* utilizes several virulence mechanisms to escape from host autophagy and grow slowly in specialized niches such as SLAPs, which results in the promotion of bacterial replication and chronic infection (Birmingham *et al.*, 2008; Huang and Brumell, 2009). Recent studies by Dortet *et al.* have shown that *L. monocytogenes* hijacks the host major vault protein through interaction with InlK, a listerial virulence factor, thus resulting in escape from autophagic recognition and intracellular survival (Dortet *et al.*, 2011).

#### **Bacterial survival and replication inside the pathogen-containing compartments: *Mycobacteria*, *Legionella*, *Porphyromonas*, *Coxiella*, and *Brucella***

*M. tuberculosis* is an enormously successful intracellular pathogen that causes tuberculosis. *M. tuberculosis* is one of the best examples of bacterial survival and replication inside host phagocytes through escape from host immune pathways (Jo, 2010). *M. tuberculosis* can achieve persistent infection within the hostile environment of the macrophage through rapid transcriptional responses that counteract host immune processes, such as antigen presentation and pro-inflammatory cytokine secretion (Huynh *et al.*, 2011). Arrest of phagosomal maturation into a phagolysosome by *M. tuberculosis* is a central pathway to overcome host defense and provides these bacteria with a strategy for survival in host cells (Chua *et al.*, 2004; Huynh *et al.*, 2011). Autophagy pathway activation induced by starvation, inhibition of mTOR pathways, vitamin D, and interferon- $\gamma$  contributes to the elimination of intracellular *M. tuberculosis* and phagosomal maturation (Gutierrez *et al.*, 2004; Jo, 2010; Deretic, 2011). A genome-wide siRNA screening identified host factors that regulate intracellular survival of *M. tuberculosis* (Kumar *et al.*, 2010). Importantly, the core host cell factors are found to be predominantly involved in the regulation of the autophagy process (Kumar *et al.*, 2010). Recent studies showed that the *M. tuberculosis eis* gene negatively modulates host cell autophagy through ROS-dependent mechanisms (Shin *et al.*, 2010a).

Early activation of the autophagy response seems to be crucial for restriction of *Legionella pneumophila*, an intracellular bacterial pathogen

responsible for an acute form of pneumonia, Legionnaire's disease (Dubuisson and Swanson, 2006). Like *M. tuberculosis*, *Legionella* can escape from phagolysosome fusion, and perturb and delay the maturation of autophagosomes into autolysosomes (Campoy and Colombo, 2009a; Joshi and Swanson, 2011). Various pathogens such as *Porphyromonas gingivalis*, *Coxiella burnetii* and *Brucella abortus* use an autophagic transit strategy during infection. These bacteria take some advantage of the autophagic pathway, in terms of intracellular survival and replication (Campoy and Colombo, 2009a). Similarly, the facultative intracellular pathogen *Francisella tularensis* uses a strategy to delay progression along the autophagic pathway, re-enter the endocytic pathway after cytoplasmic replication, and even hijack lysosome exocytosis to exit cells (Checroun *et al.*, 2006; Joshi and Swanson, 2011). Future studies will elucidate more detail of the mechanisms by which bacteria interact with host autophagy pathways, which will shed light not only on bacterial pathogenesis at the molecular level but also assist in the development of new therapeutics against refractory chronic infectious diseases.

#### **Cargo receptor and autophagic targeting to bacteria**

Recent reports have suggested that the autophagy pathway involves selective recognition and degradation of the autophagic cargo (Johansen and Lamark, 2011). In selective autophagy, modification of target proteins or intracellular bacteria with ubiquitin is necessary for its autophagic clearance (Kirkin *et al.*, 2009b). Autophagy receptors including p62/SQSTM1 (sequestome 1), NBR1 (neighbour of Brca1 gene) and NDP52 (nuclear dot protein 52 kDa) can simultaneously bind intracellular ubiquitinated cargos and the autophagy modifiers, such as microtubule-associated protein LC3 and  $\gamma$ -aminobutyric acid receptor-associated proteins (GABARAP) (Kirkin *et al.*, 2009b; Kraft *et al.*, 2010). Thus, autophagic receptors can mediate docking of ubiquitin-marked protein aggregates to the autophagosomes to induce selective autophagic degradation of ubiquitinated proteins, organelles, and intracellular bacteria (Kirkin *et al.*, 2009a, 2009b; Thurston *et al.*, 2009).

As reviewed above and elsewhere (Sumpter and Levine, 2010), the cargo receptors p62 and NDP52 can target intracellular bacteria, such as *S. typhimurium*, *S. flexneri*, and *L. monocytogenes*, by addition of a molecular tag (such as poly-ubiquitin) to autophagic machinery (Thurston *et al.*,

2009; Zheng *et al.*, 2009). Recently, it has been shown that optineurin promotes the selective autophagy of ubiquitin-coated cytosolic *Salmonella enterica* through phosphorylation of optineurin by protein kinase, TANK binding kinase 1 (Wild *et al.*, 2011). Additionally, an ubiquitin-independent pathway mediated by a lipid second messenger, diacylglycerol, generated from SCV membranes, is essential for the initiation of autophagy against *S. typhimurium* (Shahnazari *et al.*, 2010). Ogawa *et al.* (2011) revealed that the Atg5 binding partner Tecpr1 (tectonin domain-containing protein) is a cargo receptor for targeting bacterial pathogens to the selective autophagy system in *Shigella*-infected cells. Therefore, several autophagy receptors, which may interconnect with other pathways or mechanisms involved in autophagy, contribute to the xenophagic elimination of cytosolic bacteria in mammalian cells. The mechanisms by which autophagy receptors activate host defenses against bacterial pathogens other than *S. typhimurium* remain to be determined.

### ROS, keeping a balance between antibacterial autophagy and inflammation

Intracellular ROS, produced by incomplete reduction of oxygen, can oxidize and damage various macromolecules such as proteins, lipids, and DNA, and eventually cause cell death (Azad *et al.*, 2009; Scherz-Shouval and Elazar, 2011). Accumulating evidence points to a key role for ROS as signaling messengers, including the initiation of autophagy and various intracellular processes (Azad *et al.*, 2009; Scherz-Shouval and Elazar, 2011). Several stimuli and stresses, including nutrient starvation, infection, and other pathological conditions, result in ROS generation (Azad *et al.*, 2009). Cellular ROS is involved in innate immune signaling through the TLR-dependent ASK activation (Yuk *et al.*, 2009a) and NADPH oxidase 2 pathways (Yang *et al.*, 2009). Recently, it was revealed that ROS of mitochondrial origin are a critical factor in the antibacterial response, mediated by interaction with TRAF6 and ECSIT (evolutionarily conserved signaling intermediate in Toll pathways), which are themselves involved in the mitochondrial respiratory chain assembly (West *et al.*, 2011).

Interestingly, these are generally accepted as activating signals for autophagy in a variety of physiological and pathological settings (Azad *et al.*, 2009; Scherz-Shouval and Elazar, 2011). Notably, TLR signaling from phagosomes can lead to initiation of antibacterial autophagy through generation of NADPH oxidase NOX2-dependent ROS (Huang and Brumell, 2009). Excessive ROS of cellular and

mitochondrial origin is associated with activation of proinflammatory responses, autophagy, and cell death in macrophages infected with enhanced intracellular survival (*eis*)-deficient mycobacteria (Shin *et al.*, 2010a).

In contrast, recent evidence suggests that autophagy is also essential for prevention of excessive inflammation. Depletion of the autophagy proteins LC3B and beclin 1 significantly enhances NLRP3-dependent inflammasome activation and IL-1 $\beta$  and IL-18 levels, and increases susceptibility to LPS-induced septic shock *in vivo* (Nakahira *et al.*, 2011). Intriguingly, in this case, mitochondrial ROS are also thought to be important for mediating inflammatory responses, even in the autophagy-deficient host (Nakahira *et al.*, 2011). This effect was dependent on the release of mitochondrial DNA into the cytosol and increased mitochondrial ROS generation (Nakahira *et al.*, 2011). Thus, it seems that coordinated interplay between autophagy activation and ROS-dependent signaling is necessary for the homeostatic regulation of innate defense and inflammation.

### Concluding remarks

Since the roles of autophagy have been revealed in host immune responses, a number of recent *in vitro*, *in vivo*, and molecular studies have provided evidence of selective autophagic degradation of bacteria and viruses (xenophagy). Moreover, rapid advances have been made in elucidating the mechanisms underlying the cell's defensive response to intracellular bacteria. In this review we have emphasized that numerous bacterial pathogens possess multiple strategies for avoiding or circumventing host defense pathways to ensure their survival and replication within the host. Thus, it is not surprising that autophagy pathways cross talk with innate immunity to contribute to the combat of invading pathogens. Various immune effectors have been identified as cooperating with and regulating the autophagic responses against invading bacterial infection, although additional host factors and detailed mechanisms remain to be delineated. Here, we propose that the host autophagy system has evolved strategies to manipulate certain autophagic adaptors or cargo receptors to promote autophagy targeting to bacteria, proteins, and damaged organelles. Several important questions related to the extended and distinct roles of immune effectors or cargo receptors, i.e., whether they are involved in defense against specific pathogens or general microbial invaders, should be answered in future studies. Current and future revelations regarding

the interplay between host autophagy and its evasion by microbial pathogens will provide opportunities for the development of novel antibacterial agents.

### Acknowledgements

We are indebted to both current and past members of our laboratory for the discussions and investigations that contributed to this article. This research was supported by a National Research Foundation of Korea (NRF) grant funded by the Korea government (MEST) through the Infection Signaling Network Research Center (2011-0006228) at Chungnam National University.

### References

- Azad AF, Beard CB. Rickettsial pathogens and their arthropod vectors. *Emerg Infect Dis* 1998;4:179-86
- Azad MB, Chen Y, Gibson SB. Regulation of autophagy by reactive oxygen species (ROS): implications for cancer progression and treatment. *Antioxid Redox Signal* 2009;11:777-90
- Birmingham CL, Smith AC, Bakowski MA, Yoshimori T, Brumell JH. Autophagy controls *Salmonella* infection in response to damage to the *Salmonella*-containing vacuole. *J Biol Chem* 2006;281:11374-83
- Birmingham CL, Canadien V, Kaniuk NA, Steinberg BE, Higgins DE, Brumell JH. Listeriolysin O allows *Listeria monocytogenes* replication in macrophage vacuoles. *Nature* 2008;451:350-4
- Brodsky IE, Medzhitov R. Targeting of immune signaling networks by bacterial pathogens. *Nat Cell Biol* 2009;11:521-6
- Campoy E, Colombo MI. Autophagy in intracellular bacterial infection. *Biochim Biophys Acta* 2009a;1793:1465-77
- Campoy E, Colombo MI. Autophagy subversion by bacteria. *Curr Top Microbiol Immunol* 2009b;335:227-50
- Cederlund A, Gudmundsson GH, Agerberth B. Antimicrobial peptides important in innate immunity. *FEBS J* 2011;278:3942-51
- Cemma M, Kim PK, Brumell JH. The ubiquitin-binding adaptor proteins p62/SQSTM1 and NDP52 are recruited independently to bacteria-associated microdomains to target *Salmonella* to the autophagy pathway. *Autophagy* 2011;7:341-5
- Chang YP, Tsai CC, Huang WC, Wang CY, Chen CL, Lin YS, Kai JI, Hsieh CY, Cheng YL, Choi PC, Chen SH, Chang SP, Liu HS, Lin CF. Autophagy facilitates IFN- $\gamma$ -induced Jak2-STAT1 activation and cellular inflammation. *J Biol Chem* 2010;285:28715-22
- Checroun C, Wehrly TD, Fischer ER, Hayes SF, Celli J. Autophagy-mediated reentry of *Francisella tularensis* into the endocytic compartment after cytoplasmic replication. *Proc Natl Acad Sci USA* 2006;103:14578-83
- Chua J, Vergne I, Master S, Deretic V. A tale of two lipids: *Mycobacterium tuberculosis* phagosome maturation arrest. *Curr Opin Microbiol* 2004;7:71-7
- Collins CA, De Mazière A, van Dijk S, Carlsson F, Klumperman J, Brown EJ. Atg5-independent sequestration of ubiquitinated *mycobacteria*. *PLoS Pathog* 2009;5:e1000430
- Cooney R, Baker J, Brain O, Danis B, Pichulik T, Allan P, Ferguson DJ, Campbell BJ, Jewell D, Simmons A. NOD2 stimulation induces autophagy in dendritic cells influencing bacterial handling and antigen presentation. *Nat Med* 2010;16:90-7
- Cooper AM, Mayer-Barber KD, Sher A. Role of innate cytokines in mycobacterial infection. *Mucosal Immunol* 2011;4:252-60
- Cossart P. Actin-based motility of pathogens: the Arp2/3 complex is a central player. *Cell Microbiol* 2000;2:195-205
- Delgado MA, Deretic V. Toll-like receptors in control of immunological autophagy. *Cell Death Differ* 2009;16:976-83
- Deretic V. Multiple regulatory and effector roles of autophagy in immunity. *Curr Opin Immunol* 2009;21:53-62
- Deretic V. Autophagy in infection. *Curr Opin Immunol* 2010;22:252-62
- Deretic V. Autophagy in immunity and cell-autonomous defense against intracellular microbes. *Immunol Rev* 2011;240:92-104
- Deretic V, Levine B. Autophagy, immunity, and microbial adaptations. *Cell Host Microbe* 2009;5:527-49
- Djavaheri-Mergny M, Amelotti M, Mathieu J, Besancon F, Bauvy C, Souquère S, Pierron G, Codogno P. NF-kappaB activation represses tumor necrosis factor- $\alpha$ -induced autophagy. *J Biol Chem* 2006;281:30373-82
- Dortet L, Mostowy S, Louaka AS, Gouin E, Nahori MA, Wiemer EA, Dussurget O, Cossart P. Recruitment of the major vault protein by InlK: a *Listeria monocytogenes* strategy to avoid autophagy. *PLoS Pathog* 2011;7:e1002168
- Dubuisson JF, Swanson MS. Mouse infection by *Legionella*, a model to analyze autophagy. *Autophagy* 2006;2:179-82
- Dupont N, Lacas-Gervais S, Bertout J, Paz I, Freche B, Van Nhieu GT, van der Goot FG, Sansonetti PJ, Lafont F. *Shigella* phagocytic vacuolar membrane remnants participate in the cellular response to pathogen invasion and are regulated by autophagy. *Cell Host Microbe* 2009;6:137-49
- Furuta N, Yoshimori T, Amano A. Mediator molecules that fuse autophagosomes and lysosomes. *Autophagy* 2010;6:417-8
- Gutierrez MG, Master SS, Singh SB, Taylor GA, Colombo MI, Deretic V. Autophagy is a defense mechanism inhibiting BCG and *Mycobacterium tuberculosis* survival in infected macrophages. *Cell* 2004;119:753-66
- Harris J, De Haro SA, Master SS, Keane J, Roberts EA, Delgado M, Deretic V. T helper 2 cytokines inhibit autophagic control of intracellular *Mycobacterium tuberculosis*. *Immunity* 2007;27:505-17



- Harris J, Master SS, De Haro SA, Delgado M, Roberts EA, Hope JC, Keane J, Deretic V. Th1-Th2 polarisation and autophagy in the control of intracellular *Mycobacteria* by macrophages. *Vet Immunol Immunopathol* 2009;128:37-43
- Huang J, Brumell JH. Autophagy in immunity against intracellular bacteria. *Curr Top Microbiol Immunol* 2009; 335:189-215
- Huynh KK, Joshi SA, Brown EJ. A delicate dance: host response to *Mycobacteria*. *Curr Opin Immunol* 2011; 23:464-72
- Ichimura Y, Kumanoide T, Sou YS, Mizushima T, Ezaki J, Ueno T, Kominami E, Yamane T, Tanaka K, Komatsu M. Structural basis for sorting mechanism of p62 in selective autophagy. *J Biol Chem* 2008;283:22847-57
- Jo EK. Innate immunity to *Mycobacteria*: vitamin D and autophagy. *Cell Microbiol* 2010;12:1026-35
- Jo EK, Yang CS, Choi CH, Harding CV. Intracellular signalling cascades regulating innate immune responses to *Mycobacteria*: branching out from Toll-like receptors. *Cell Microbiol* 2007;9:1087-98
- Johansen T, Lamark T. Selective autophagy mediated by autophagic adapter proteins. *Autophagy* 2011;7:279-96
- Joshi AD, Swanson MS. Secrets of a successful pathogen: *Legionella* resistance to progression along the autophagic pathway. *Front Microbiol* 2011;2:138
- Kim BH, Shenoy AR, Kumar P, Das R, Tiwari S, MacMicking JD. A family of IFN- $\gamma$ -inducible 65-kD GTPases protects against bacterial infection. *Science* 2011;332:717-21
- Kirkin V, Lamark T, Sou YS, Bjørkøy G, Nunn JL, Bruun JA, Shvets E, McEwan DG, Clausen TH, Wild P, Bilusic I, Theurillat JP, Øvervatn A, Ishii T, Elazar Z, Komatsu M, Cikic I, Johansen T. A role for NBR1 in autophagosomal degradation of ubiquitinated substrates. *Mol Cell* 2009a; 33:505-16
- Kirkin V, McEwan DG, Novak I, Dikic I. A role for ubiquitin in selective autophagy. *Mol Cell* 2009b;34:259-69
- Knodler LA, Celli J, Finlay BB. Pathogenic trickery: deception of host cell processes. *Nat Rev Mol Cell Biol* 2001;2:578-88
- Kraft C, Peter M, Hofmann K. Selective autophagy: ubiquitin-mediated recognition and beyond. *Nat Cell Biol* 2010;12:836-41
- Kumar D, Nath L, Kamal MA, Varshney A, Jain A, Singh S, Rao KV. Genome-wide analysis of the host intracellular network that regulates survival of *Mycobacterium tuberculosis*. *Cell* 2010;140:731-43
- Lee HM, Shin DM, Yuk JM, Shi G, Choi DK, Lee SH, Huang SM, Kim JM, Kim CD, Lee JH, Jo EK. Autophagy negatively regulates keratinocyte inflammatory responses via scaffolding protein p62/SQSTM1. *J Immunol* 2011;186: 1248-58
- Levine B, Deretic V. Unveiling the roles of autophagy in innate and adaptive immunity. *Nat Rev Immunol* 2007;7:767-77
- Levine B, Mizushima N, Virgin HW. Autophagy in immunity and inflammation. *Nature* 2011;469:323-35
- Ling YM, Shaw MH, Ayala C, Coppens I, Taylor GA, Ferguson DJ, Yap GS. Vacuolar and plasma membrane stripping and autophagic elimination of *Toxoplasma gondii* in primed effector macrophages. *J Exp Med* 2006;203:2063-71
- Liu PT, Stenger S, Li H, Wenzel L, Tan BH, Krutzik SR, Ochoa MT, Schaubert J, Wu K, Meinken C, Kamen DL, Wagner M, Bals R, Steinmeyer A, Zügel U, Gallo RL, Eisenberg D, Hewison M, Hollis BW, Adams JS, Bloom BR, Modlin RL. Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science* 2006;311:1770-3
- Manavalan B, Basith S, Choi S. Similar structures but different roles - an updated perspective on TLR structures. *Front Physiol* 2011;2:41
- Mizushima N, Levine B, Cuervo AM, Klionsky DJ. Autophagy fights disease through cellular self-digestion. *Nature* 2008; 451:1069-75
- Nakagawa I, Amano A, Mizushima N, Yamamoto A, Yamaguchi H, Kamimoto T, Nara A, Funao J, Nakata M, Tsuda K, Hamada S, Yoshimori T. Autophagy defends cells against invading group A *Streptococcus*. *Science* 2004; 306:1037-40
- Nakahira K, Haspel JA, Rathinam VA, Lee SJ, Dolinay T, Lam HC, Englert JA, Rabinovitch M, Cernadas M, Kim HP, Fitzgerald KA, Ryter SW, Choi AM. Autophagy proteins regulate innate immune responses by inhibiting the release of mitochondrial DNA mediated by the NALP3 inflammasome. *Nat Immunol* 2011;12:222-30
- Ogawa M, Yoshimori T, Suzuki T, Sagara H, Mizushima N, Sasakawa C. Escape of intracellular *Shigella* from autophagy. *Science* 2005;307:727-31
- Ogawa M, Yoshikawa Y, Kobayashi T, Mimuro H, Fukumatsu M, Kiga K, Piao Z, Ashida H, Yoshida M, Kakuta S, Koyama T, Goto Y, Nagatake T, Nagai S, Kiyono H, Kawalec M, Reichhart JM, Sasakawa C. A Tecpr1-dependent selective autophagy pathway targets bacterial pathogens. *Cell Host Microbe* 2011;9:376-89
- Orvedahl A, Levine B. Eating the enemy within: autophagy in infectious diseases. *Cell Death Differ* 2009;16:57-69
- Pankiv S, Clausen TH, Lamark T, Brech A, Bruun JA, Outzen H, Øvervatn A, Bjørkøy G, Johansen T. p62/SQSTM1 binds directly to Atg8/LC3 to facilitate degradation of ubiquitinated protein aggregates by autophagy. *J Biol Chem* 2007;282: 24131-45
- Pattingre S, Bauvy C, Carpentier S, Levade T, Levine B, Codogno P. Role of JNK1-dependent Bcl-2 phosphorylation in ceramide-induced macroautophagy. *J Biol Chem* 2009; 284:2719-28
- Rikihisa Y. Glycogen autophagosomes in polymorphonuclear leukocytes induced by rickettsiae. *Anat Rec* 1984;208: 319-27
- Sakurai A, Maruyama F, Funao J, Nozawa T, Aikawa C, Okahashi N, Shintani S, Hamada S, Ooshima T, Nakagawa I. Specific behavior of intracellular *Streptococcus pyogenes* that has undergone autophagic degradation is associated with bacterial streptolysin O and host small G proteins Rab5 and Rab7. *J Biol Chem* 2010;285:22666-75

- Scherz-Shouval R, Elazar Z. Regulation of autophagy by ROS: physiology and pathology. *Trends Biochem Sci* 2011; 36:30-8
- Shahnazari S, Brumell JH. Mechanisms and consequences of bacterial targeting by the autophagy pathway. *Curr Opin Microbiol* 2011;14:68-75
- Shahnazari S, Yen WL, Birmingham CL, Shiu J, Namolovan A, Zheng YT, Nakayama K, Klionsky DJ, Brumell JH. A diacylglycerol-dependent signaling pathway contributes to regulation of antibacterial autophagy. *Cell Host Microbe* 2010;8:137-46
- Shi CS, Kehrl JH. MyD88 and Trif target Beclin 1 to trigger autophagy in macrophages. *J Biol Chem* 2008;283: 33175-82
- Shi CS, Kehrl JH. TRAF6 and A20 regulate lysine 63-linked ubiquitination of Beclin-1 to control TLR4-induced autophagy. *Sci Signal* 2010;3:ra42
- Shin DM, Jeon BY, Lee HM, Jin HS, Yuk JM, Song CH, Lee SH, Lee ZW, Cho SN, Kim JM, Friedman RL, Jo EK. *Mycobacterium tuberculosis* eis regulates autophagy, inflammation, and cell death through redox-dependent signaling. *PLoS Pathog* 2010a;6:e1001230
- Shin DM, Yuk JM, Lee HM, Lee SH, Son JW, Harding CV, Kim JM, Modlin RL, Jo EK. Mycobacterial lipoprotein activates autophagy via TLR2/1/CD14 and a functional vitamin D receptor signaling. *Cell Microbiol* 2010b;12:1648-65
- Silverman DJ, Wisseman CL, Jr. *In vitro* studies of rickettsia-host cell interactions: ultrastructural changes induced by *Rickettsia rickettsii* infection of chicken embryo fibroblasts. *Infect Immun* 1979;26:714-27
- Singh SB, Davis AS, Taylor GA, Deretic V. Human IRGM induces autophagy to eliminate intracellular *mycobacteria*. *Science* 2006;313:1438-41
- Sonawane A, Santos JC, Mishra BB, Jena P, Progida C, Sorensen OE, Gallo R, Appelberg R, Griffiths G. Cathelicidin is involved in the intracellular killing of *mycobacteria* in macrophages. *Cell Microbiol* 2011;13:1601-17
- Stamm LM, Brown EJ. *Mycobacterium marinum*: the generalization and specialization of a pathogenic mycobacterium. *Microbes Infect* 2004;6:1418-28
- Sumpter R Jr, Levine B. Autophagy and innate immunity: triggering, targeting and tuning. *Semin Cell Dev Biol* 2010; 21:699-711
- Takeda K, Akira S. TLR signaling pathways. *Semin Immunol* 2004;16:3-9
- Takeuchi O, Akira S. Pattern recognition receptors and inflammation. *Cell* 2010;140:805-20
- Thurston TL, Ryzhakov G, Bloor S, von MN, Randow F. The TBK1 adaptor and autophagy receptor NDP52 restricts the proliferation of ubiquitin-coated bacteria. *Nat Immunol* 2009; 10:1215-21
- von Muhlinen N, Thurston T, Ryzhakov G, Bloor S, Randow F. NDP52, a novel autophagy receptor for ubiquitin-decorated cytosolic bacteria. *Autophagy* 2010;6:288-9
- Walker DH. Rickettsiae and rickettsial infections: the current state of knowledge. *Clin Infect Dis* 2007;45 (Suppl 1):S39-44
- Walker DH, Popov VL, Crocquet-Valdes PA, Welsh CJ, Feng HM. Cytokine-induced, nitric oxide-dependent, intracellular antirickettsial activity of mouse endothelial cells. *Lab Invest* 1997;76:129-38
- Wei Y, Pattingre S, Sinha S, Bassik M, Levine B. JNK1-mediated phosphorylation of Bcl-2 regulates starvation-induced autophagy. *Mol Cell* 2008;30:678-88
- West AP, Brodsky IE, Rahner C, Woo DK, Erdjument-Bromage H, Tempst P, Walsh MC, Choi Y, Shadel GS, Ghosh S. TLR signalling augments macrophage bactericidal activity through mitochondrial ROS. *Nature* 2011;472:476-80
- Wild P, Farhan H, McEwan DG, Wagner S, Rogov VV, Brady NR, Richter B, Korac J, Waidmann O, Choudhary C, Dötsch V, Bumann D, Dikic I. Phosphorylation of the autophagy receptor optineurin restricts *Salmonella* growth. *Science* 2011;333:228-33
- Xu Y, Jagannath C, Liu XD, Sharafkhaneh A, Kolodziejaska KE, Eissa NT. Toll-like receptor 4 is a sensor for autophagy associated with innate immunity. *Immunity* 2007;27:135-44
- Yang CS, Shin DM, Kim KH, Lee ZW, Lee CH, Park SG, Bae YS, Jo EK. NADPH oxidase 2 interaction with TLR2 is required for efficient innate immune responses to *mycobacteria* via cathelicidin expression. *J Immunol* 2009; 182:3696-705
- Yoshikawa Y, Ogawa M, Hain T, Yoshida M, Fukumatsu M, Kim M, Mimuro H, Nakagawa I, Yanagawa T, Ishii T, Kakizuka A, Sztul E, Chakraborty T, Sasakawa C. *Listeria monocytogenes* ActA-mediated escape from autophagic recognition. *Nat Cell Biol* 2009;11:1233-40
- Yuk JM, Shin DM, Lee HM, Yang CS, Jin HS, Kim KK, Lee ZW, Lee SH, Kim JM, Jo EK. Role of apoptosis-regulating signal kinase 1 in innate immune responses by *Mycobacterium bovis* bacillus Calmette-Guerin. *Immunol Cell Biol* 2009a;87:100-7
- Yuk JM, Shin DM, Lee HM, Yang CS, Jin HS, Kim KK, Lee ZW, Lee SH, Kim JM, Jo EK. Vitamin D3 induces autophagy in human monocytes/macrophages via cathelicidin. *Cell Host Microbe* 2009b;6:231-43
- Zheng YT, Shahnazari S, Brech A, Lamark T, Johansen T, Brumell JH. The adaptor protein p62/SQSTM1 targets invading bacteria to the autophagy pathway. *J Immunol* 2009;183:5909-16